

## Original article

# Cost-effectiveness analysis of highly concentrated n-3 polyunsaturated fatty acids in secondary prevention after myocardial infarction

Adawan Permpnitch<sup>a</sup>, Vithaya Kulsomboon<sup>a,b</sup>, Kamol Udol<sup>c</sup>

<sup>a</sup>Graduate Program in Social and Administrative Pharmacy, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok 10330, Thailand

<sup>b</sup>Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok 10330, Thailand

<sup>c</sup>Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok 10700, Thailand

---

**Background:** Acute myocardial infarction (MI) is a leading cause of cardiovascular (CV) mortality and hospitalization. Survivors of acute MI have higher risk of subsequent CV events and death, compared to individuals without MI. Evidences have demonstrated the CV benefits of n-3 polyunsaturated fatty acids (PUFAs) in patients who experienced MI.

**Objectives:** We assessed the cost-effectiveness of highly concentrated n-3 polyunsaturated fatty acids (PUFAs) in addition to standard therapy compared with standard therapy alone in post-MI patients in Thailand.

**Methods:** A Markov model was constructed to assess costs, life years, and quality-adjusted life years (QALYs) with lifetime horizon in post-MI patients, on the basis of provider perspective. Input data were based on information from the Thai Acute Coronary Syndrome (ACS) Registry, a meta-analysis of mortality data and published articles. Outcomes have been presented as incremental cost-effectiveness ratios of life expectancy and quality-adjusted life expectancy. Deterministic and probabilistic sensitivity analyses were performed for key variables in the model.

**Results:** n-3 PUFAs increased life expectancy by 2.34 life-years at an incremental cost-effectiveness ratio (ICER) of 256,199 Thai baht (THB) per life-year gained (LYG), compared to the standard therapy alone in the base case analysis. The quality-adjusted life years (QALY) increased by 2.01 with ICER of 297,193 THB per QALY from n-3 PUFAs supplementation. Both ICER/QALY and ICER/LYG decreased as the age of patients increased. The incremental cost per QALY gained in post-MI patients aged 45 to 85 years old ranged from 216,200 THB to 414,049 THB.

**Conclusion:** Considering the current willingness-to-pay threshold of 160,000 THB/QALY, highly concentrated n-3 PUFAs as secondary prevention of MI appears not to be cost-effective compared to standard treatment alone in Thailand.

**Keywords:** Cost-effectiveness, economic evaluation, myocardial infarction, n-3 PUFA, omega-3, secondary prevention, supplementation, Thailand

---

Despite several new treatments having been developed for the treatment of acute myocardial infarction (MI), the survivors from acute MI still have higher risk to subsequent cardiovascular (CV) events and mortality. Data from the Thai Acute Coronary Syndrome (ACS) Registry showed that approximately 20% of post-MI patients died, 4% had recurrent MI and 10% experienced non-fatal stroke within 1 year after their first MI. In the Asia-Pacific region,

cardiovascular death accounts for nearly one third of all deaths, whereas ischemic heart disease and stroke comprise approximately 80% of all cardiovascular deaths in Thailand [1].

The significant cardiovascular benefits of highly concentrated n-3 polyunsaturated fatty acids (PUFAs) were demonstrated in people who consumed fish or n-3 PUFAs supplementation [2-6], while non-significant results compared to placebo were also shown in some studies [7-10]. Several meta-analyses of n-3 PUFAs studies in patients at high risk of CV diseases demonstrated inconsistency of cardiovascular benefits. The possible factors which contributed to

---

**Correspondence to:** Vithaya Kulsomboon, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok 10330, Thailand. E-mail: vithaya.k@chula.ac.th

this inconsistency, included differences in study populations, baseline consumption of fish, doses and duration of n-3 PUFAs treatment [11-16]. Dose response and time course of n-3 PUFAs intake contribute to the physiologic effects which are most likely to account for CV risk reduction [17]. As such, a meta-analysis of studies using high concentration of n-3 PUFAs for at least 1-year, which would be long enough for n-3 PUFAs to demonstrate efficacy, would be a suitable evidence of the effectiveness of n-3 PUFAs in specific groups of patients.


Due to the development of medications and technologies, people tend to live longer. As a result, CV events increase over time, together with the costs of treatment of these diseases. Despite the proven efficacy, the limited availability of resources is of concern. As the use of n-3 PUFAs is approved as supplemental treatment in secondary prevention after MI in addition to other standard therapies, this study aimed to assess the cost-effectiveness of adding highly

concentrated n-3 PUFAs to the current treatment in post-MI patients in Thailand.

## Methods

### Model structure

A Markov model was developed in Excel version 2013, starting with the event-free MI (post-MI). During each 1-year cycle of the model, some patients entered one of the four health states (cardiovascular mortality, non-fatal MI, non-fatal stroke, and heart failure (HF)), while the others remained in the event-free state (**Figure 1**). The rate at which people moved through the model was regulated by transition probabilities, which described the likelihood of moving between states over each model cycle. The transition between each state and event was determined by probabilities and adjusted factors obtained from randomized control trials and the local registry database. The model was run until the patients were 100 years old or died.



**Figure 1.** Structure of the Markov model

The main assumptions made in this model were as follows; 1) patients in the acute health state (MI, stroke, HF) would not move to other acute health state; 2) the effectiveness of n-3 PUFAs would be maintained over life-time horizon; 3) the duration in each cycle was 1 year.

The discount rate of 3% was applied for both costs and outcomes. The time horizon of the model and the effects in life expectancy were life-long calculation. The provider perspective was chosen for this study; thus only direct health care costs were used in the analysis.

### **Effectiveness data**


We carried out a meta-analysis to estimate the effectiveness of n-3 PUFAs for the base case analysis. A comprehensive literature search of MEDLINE, EMBASE, and the Cochrane Library (up to September 2013) was conducted. All trials that met the following criteria were included: 1) the trial enrolled adult patients (aged at least 18 years) with a history of myocardial infarction; 2) n-3 PUFAs were consumed in dose of at least 1 gram/day; 3) the follow-up period was at least 1 year, which would be long enough for n-3 PUFAs to demonstrate efficacy; 4) the outcomes included CV diseases or events, cardiovascular death, sudden death, all-cause mortality, coronary heart disease, heart failure, stroke, fatal or nonfatal MI. The exclusion criteria included studies with less than 1-year duration, studies with multiple interventions from which the effects of n-3 PUFAs could not be separated, and studies that did not report CV-related endpoints. If two or more articles reported results from the same group of patients at different follow-up periods, only the article with longer follow-up period was retained in order to prevent data duplication.

Titles and abstracts obtained from the electronic and bibliographic searches were rejected on initial screening if it could be determined that the articles did not meet the eligibility criteria. In case the reviewer was uncertain to reject an article based on its title/abstract, the full text of the article was further evaluated. Data extraction was performed by 2 reviewers independently. In case there were discrepancies, the reviewers discussed or consulted the third reviewer. If the same studies were presented in more than 1 article, we included only the larger or first published study.

From 4,042 articles identified from the database search, 3 studies which met the inclusion criteria and did not meet the exclusion criteria were included in the meta-analysis. The heterogeneity of studies was not statistically significant only in the cardiovascular mortality endpoint, while the endpoint of total mortality and sudden death showed significant heterogeneity. Therefore, the cardiovascular mortality endpoint was selected, which was reported in 2 studies, i.e. GISSI-P (Gruppo Italiano per lo Studio della Sopravvivenza nell' Infarto Miocardico-Prevenzione) study [5, 18], and OMEGA (Effect of Omega 3-Fatty Acids on the Reduction of Sudden Cardiac Death after Myocardial Infarction) study [7]. In total 15,174 patients after MI were included. The results of included studies were pooled using fixed effect model, weighted by inverse of variance. The summary of studies included in the meta-analysis has been shown in **Table 1**. The meta-analysis results in **Figure 2** demonstrated the significant reduction in cardiovascular mortality in patients using n-3 PUFAs compared to those using standard treatment only.

**Table 1.** Summary of studies included in the meta-analysis

Study	Dose of n-3 PUFAs	Study duration (month)	Total population (treatment/control)	Number of patients with CV mortality (treatment/control)
GISSI-P	1 gram/day	42	11,324 (5,666/5,658)	680 (310/370)
OMEGA	1 gram/day	12	3,804 (1,919/1,885)	57 (28/29)



**Figure 2.** Meta-analysis results on cardiovascular mortality in post-MI patients using n-3 PUFAs.

Effectiveness data of n-3 PUFAs on MI and stroke were obtained from the GISSI-P study (5, 18). There was only 1 study using high dose of n-3 PUFAs in 6,975 patients with chronic heart failure, 41% of whom had a history of MI; so the result of this study was used to estimate the effectiveness in HF (19). These effectiveness data are shown in **Table 2**.

### Cost data

As the provider perspective is used for this study, cost data in the study included only direct health care costs, which were defined as the costs of goods and services directly provided by the service provider such as medication cost, hospital cost, etc. All costs were

inflated to 2013 values using Consumer Price Index (CPI) of medical care group from the Bureau of Trade and Economic Indices, Ministry of Commerce [20]. The MI costs were taken from a published literature using data collected from patients in the Thai ACS Registry who were admitted to Ramathibodi Hospital [21]. The stroke costs were obtained from a prospective observational study collecting data in 2 regional hospitals [22]. Since the costs of stroke in subsequent years were not collected, it was assumed to be the same as in the first year except for hospitalization cost. The costs of heart failure were collected from 2 sets of the National Health Security Office (NHSO) of Thailand database, i.e.

**Table 2.** Parameters used in the Markov model

	Distribution	RR	SE	References
<b>Treatment effectiveness data</b>				
Effect of n-3 PUFAs on cardiovascular mortality	Gamma	0.73	0.0740	Meta-analysis
Effect of n-3 PUFAs on nonfatal MI	Gamma	0.91	0.1225	(18)
Effect of n-3 PUFAs on nonfatal stroke	Gamma	1.22	0.3138	(18)
Effect of n-3 PUFAs on nonfatal heart failure	Gamma	0.90	0.0459	(19)
<b>Cost data</b>				
Cost of MI in the first year	Gamma	185,143	11,953	(21)
Cost of MI in the subsequent year	Gamma	13,588	679*	(21)
Cost of stroke in the first year	Gamma	30,642	2,165	(22)
Cost of stroke in the subsequent year	Gamma	15,694	1,773	(22)
Cost of heart failure in the first year	Gamma	19,507	975*	NHSO
Cost of heart failure in the subsequent year	Gamma	8,023	401*	NHSO
Cost of n-3 PUFAs per year	Gamma	15,231	762*	DMSIC website
<b>Utility data</b>				
Utility of MI	Beta	0.870	0.0255	(23)
Utility of post-MI	Beta	0.910	0.0255	(23)
Utility of stroke	Beta	0.550	0.0172	(24)
Utility of post-stroke	Beta	0.663	0.0106	(25)
Utility of heart failure	Beta	0.600	0.0102	(26)
Utility of post-heart failure	Beta	0.680	0.0260	(27)
Utility of death	Beta	0.000	0.0000	

\*Assumption (5% of the mean), Assumption (calculated from the formula  $SE = SD/(8840)^{1/2}$ ;  $n = 8840$ , assumed  $SD = 1$ )

hospitalization cost of inpatient from government hospitals and the cost of patients visiting the out-patient departments of hospitals under Bangkok Metropolitan Administration. As Omacor (Abbott Laboratories) is the only highly concentrated formulation of n-3 PUFAs in Thailand, the price in the Drug and Medical Supply Information Center (DMSIC) website was used in this study. All costs are shown in **Table 2**.

### Utility data

The utility weights of acute state of each condition were lower than those of post state. The utility weight data of MI, stroke and HF were obtained from literature review [23-27], as shown in **Table 2**.

### Transition probability

Data from the Thai ACS Registry collected from 1,525 patients who experienced MI were used to estimate the probability of cardiovascular mortality using survival analysis. The probabilities of transition between other health states were derived from published articles.

### Health outcome

The health outcome measures were quality-adjusted life year (QALY) and life-year gained (LYG). QALY was used to measure overall health-related quality of life, which is the preferred approach in economic evaluation of health intervention. LYG was a modified mortality measure in which remaining life expectancy was taken into account. Life years were calculated as the remaining life expectancy at the point of each averted death.

### Uncertainty analyses

Univariate sensitivity analyses were performed to identify the variables which had significant impacts on the results. The key individual variables which

influence the results of cost-effectiveness have been shown in a Tornado diagram. To examine the uncertainty of inputs in the model, the probabilistic sensitivity analysis using a second-order Monte Carlo simulation was conducted. All parameters used were assigned a probability distribution to take into account the uncertainty in all random quantities. The simulation of 1000 times could provide a range of possible values capturing the specified probability distribution of parameters used in the analysis.

## Results


### Cost-effectiveness

The base case scenario was done in patients aged 63 years old which was the mean age of patients who experienced MI in the Thai ACS Registry. The total life years obtained were 17.82 years for the supplementation of n-3 PUFAs and 15.48 years for no supplementation. The QALYs gained were 15.06 and 13.05 years for n-3 PUFAs supplementation and no supplementation, respectively. Thus incremental effectiveness values for supplementation of n-3 PUFAs were 2.34 life-years gained (LYG) and 2.01 QALYs. The lifetime cost from 1,000 simulations was estimated as 2,600,349 THB in the n-3 PUFAs supplementation cohort, and 2,001,572 THB in the no supplementation group, resulting in the incremental cost of 598,777 THB for the addition of n-3 PUFAs. The incremental cost-effectiveness ratio (ICER) was 297,193 THB per QALY and 256,199 THB per LYG (**Table 3**). ICER/QALY and ICER/LYG reduced when the patients had MI at the older age, indicating that the addition of n-3 PUFAs is more cost-effective in elderly patients than in younger patients (**Figure 3**). The probability of cost-effectiveness at different levels of willingness-to-pay in different ages of the patients is shown in **Figure 4**.


**Table 3.** Lifetime costs, quality-adjusted life years, life-year gained, and incremental cost-effectiveness ratio of adding n-3 polyunsaturated fatty acids compared to standard therapy in post-MI patients

Strategy	Cost (THB)	QALY	ICER/QALY	LYG	ICER/LYG
No n-3 PUFAs	2,001,572	13.05		15.48	
n-3 PUFAs	2,600,349	15.06	297,193	17.82	256,199





**Figure 3.** Incremental effectiveness of n-3 PUFAs supplementation at different ages of post-MI patients



**Figure 4.** Cost-effectiveness acceptability curve of n-3 PUFAs supplementation at different ages of post-MI patients

### Uncertainty analyses

#### Deterministic sensitivity analysis


The effect of n-3 PUFAs on cardiovascular mortality provided greatest impact to the result as shown in the Tornado Diagram (**Figure 5**), followed by discount rate, all treatment costs (not including n-3 PUFAs cost), n-3 PUFAs cost and costs of MI. The cost-effectiveness of n-3 PUFAs is least sensitive to the effect of n-3 PUFAs on stroke and the cost of HF.

#### Probabilistic sensitivity analysis


The scatter plot in **Figure 6** shows the result of 1000 Monte Carlo simulations. The majority of

iterations demonstrated that supplementation of n-3 PUFAs is more effective and more costly than standard therapy. Approximately half of the iterations were above the line with the slope of 160,000 THB/QALY, which is the current willingness-to-pay (WTP) threshold for interventions in Thailand [28].


The probability of cost-effectiveness at different willingness-to-pay threshold is shown in **Figure 7**. For the base case scenario, addition of n-3 PUFAs is considered more cost-effective at the willingness-to-pay threshold of at least 297,000 THB, compared to standard therapy alone.



**Figure 5.** Tornado diagram comparing the relative importance of model parameters on the cost-effectiveness result



**Figure 6.** Cost-Effectiveness Plane of n-3 PUFAs supplementation versus standard therapy alone



**Figure 7.** Cost-effectiveness acceptability curve of n-3 PUFAs supplementation versus standard therapy alone

## Discussion

From the base case analysis of a post-MI patient receiving highly concentrated n-3 PUFAs in addition to the standard treatment, the estimated ICER/QALY was 297,193 THB. This cost was based on provider perspective. Provided that the provider will reimburse this cost from the payer, the Thailand National Health Security Office (NHSO), this cost is higher than the current willingness-to-pay threshold of 160,000 THB [28]. The addition of highly concentrated n-3 PUFAs in patients who have survived from acute myocardial infarction would be considered not cost-effective in Thailand based on the recommended threshold.

Both ICER/QALY and ICER/LYG decreased as the age of patients increased. The lower ICER/QALYs in older patients may be due to the higher incidence rate of cardiovascular events in these age groups compared to younger patients. The increase in QALYs from n-3 PUFAs was larger than the increase in cost in patients with older age.

The ICER/QALY from our evaluation is comparable to the ICER/QALY in Ireland [29], and is slightly higher than the ICER/QALY in the U.K. [30]. The ICER/LYG in other countries varied from 2,812 in the U.K. [30] to 9,048 in Belgium [31], corresponding to approximately 150,000–400,000 THB. Our result showed consistency with data from other countries.

The effectiveness of n-3 PUFAs on mortality used in previous analyses was from the GISSI-P study only, while our analysis used the pooled data of GISSI-P and OMEGA studies. The pooled relative risk used in

our analysis is higher than that from the GISSI-P study. The mortality rate derived from the Thai ACS Registry database was a good representative for the probability of cardiovascular mortality of Thai patients who experienced MI. Moreover, all costs used in the study were collected from Thai patients. These data contributed to the robustness of this analysis.

However, this study has some limitations. Firstly, like other model-based evaluations, data from various sources were used with assumptions when data were unavailable or incomplete. The published data from other countries and assumptions might not reflect the real situation in Thailand. Secondly, the effectiveness data were limited and were obtained from only 2 studies which were conducted in European patients. The number of patients in the OMEGA study was only one-third of the GISSI-P study, resulting in the pooled data from the meta-analysis being dominated by the result of the GISSI-P study. Thirdly, though all the cost data were collected in Thailand, they might not well represent the average costs in the country. The costs of MI and stroke were obtained from only 1 and 2 sources, respectively. Data from these secondary and tertiary care hospitals might be systematically different from those in other level hospitals. In addition, the costs of HF were obtained from the NHSO which were collected using a voluntary reporting system. Fourthly, the 1-year period of the model cycle was used to simplify the discounting process, but it might not reflect the actual manifestation of the disease as the acute state of each disease might last for a few months and less than 1 year. Finally,



the incremental cost-effectiveness ratio that was performed based on payer perspective may be more suitable for the results to be used in the Thai national health insurance system. However, the limitation to obtain real cost data reimbursed from the national health system does not allow the evaluation based on payer perspective. To compromise for the possible methodology weakness, cost data which occurred to the provider would be appropriate to be used for calculating cost-effectiveness related to this intervention (n-3 PUFAs supplementation). We assumed that the provider would reimburse such cost from the payer and it could reflect cost based on payer perspective. As such, the result will be useful for the payer's decision making on reimbursement of this intervention.

## Conclusion

As the willingness-to-pay threshold in Thailand is set at 160,000 THB, the calculated ICER/QALY is higher than the threshold value. Therefore, the supplementation of highly concentrated n-3 PUFAs as secondary prevention in post-MI patients is considered not cost-effective. The intervention might be cost-effective if the threshold is higher or the highly concentrated n-3 PUFAs price is lower than the current conditions.

There was no funding support from any organization for this study. The authors have no conflict of interest to declare.

## Reference

1. OECD/WHO. "Mortality from Cardiovascular Disease", in *Health at a Glance: Asia/Pacific 2012*: OECD Publishing. [online] 2012. [cited 2014 Feb 28]; Available from: <http://dx.doi.org/10.1787/9789264183902-8-en>
2. Kromhout D, Bosschieter E, de Lezenne Coutander C. The inverse relation between fish consumption and 20-year mortality from coronary heart disease. *N Engl J Med*. 1985; 312:1205-9.
3. Singh R, Niaz M, Sharma J, Kumar R, Rastogi V, Moshiri M. Randomized, double-blind, placebo-controlled trial of fish oil and mustard oil in patients with suspected acute myocardial infarction: The Indian Experiment of Infarct Survival. *Cardiovasc Drug Ther*. 1997; 11:485-91.
4. Albert C, Hennekens C, O'Donnell C, Ajani U, Carey V, Willet W, et al. Fish consumption and risk of sudden cardiac death. *JAMA*. 1998; 279:23-8.
5. GISSI-Prevenzione Investigators. Dietary supplementation with n-3 polyunsaturated fatty acids and vitamin E after myocardial infarction: results of the GISSI-Prevenzione trial [errata in *Lancet* 2001;357:642 and *Lancet* 2007;369:106]. *Lancet*. 1999; 354:447-55.
6. Burr M, Gilbert J, Holliday R, Elwood P, Fehily A, Rogers S, et al. Effects of changes in fat, fish, and fibre intakes on death and myocardial reinfarction: Diet and Reinfarction Trial (DART). *Lancet*. 1989; 2: 757-61.
7. Rauch B, Schiele R, Schneider S, Diller F, Victor N, Gohlke H, et al. OMEGA, a randomized, placebo-controlled trial to test the effect of highly purified omega-3 fatty acids on top of modern guideline-adjusted therapy after myocardial infarction. *Circulation*. 2010; 122:2152-9.
8. The ORIGIN Trial Investigators. N-3 fatty acids and cardiovascular outcomes in patients with dysglycemia. *N Engl J Med*. 2012; 367:309-18.
9. Galan P, Kesse-Guyot E, Czernichow S, Briancon S, Blacher J, Hercberg S, et al. Effects of B vitamins and omega 3 fatty acids on cardiovascular diseases: a randomised placebo controlled trial. *BMJ*. 2010; 341: c6273.
10. Kromhout D, Giltay E, Geleijnse J, for the Alpha Omega Trial Group. N-3 fatty acids and cardiovascular events after myocardial infarction. *N Engl J Med*. 2010; 363:2015-26.
11. Hooper L, Harrison R, Summerbell C, Moore H, Worthington H, Ness A, et al. Omega 3 fatty acids for prevention and treatment of cardiovascular disease (review). *Cochrane Database Syst Rev*. 2009; CD003177.
12. Rizos E, Ntzani E, Bika E, Kostapanos M, Elisaf M. Association between omega-3 fatty acid supplementation and risk of major cardiovascular disease events: a systematic review and meta-analysis. *JAMA*. 2012; 308:1024-33.
13. Marik P, Varon J. Omega-3 dietary supplements and the risk of cardiovascular events: a systematic review. *Clin Cardiol*. 2009; 32:365-72.
14. Bucher H, Hengstler P, Schindler C, Meier G. N-3 polyunsaturated fatty acids in coronary heart disease: a meta-analysis of randomized controlled trials. *Am J Med*. 2002; 112:298-304.
15. Leon H, Shibata M, Sivakumaran S, Dorgan M, Chatterley T, Tsuyuki R. Effect of fish oil on arrhythmias and mortality: systematic review. *BMJ*. 2009; 338: a2931.
16. Kwak S, Myung S, Lee Y, Seo H, for the Korean

- Meta-analysis Study Group. Efficacy of omega-3 fatty acid supplements (eicosapentaenoic acid and docosahexaenoic acid) in the secondary prevention of cardiovascular disease. *Arch Intern Med.* 2012; 172:686-94.
17. Mozaffarian D, Rimm E. Fish intake, contaminants, and human health evaluating the risks and the benefits. *JAMA.* 2006; 296:1885-99.
  18. Marchioli R, Schweigerb C, Tavazzi L, Valagussad F. Efficacy of n-3 polyunsaturated fatty acids after myocardial infarction: results of GISSI-Prevenzione trial. *Lipids.* 2001; 36(Suppl.):S119-26.
  19. GISSI-HF Investigators. Effect of n-3 polyunsaturated fatty acids in patients with chronic heart failure (the GISSI-HF trial): a randomised, double-blind, placebo-controlled trial. *Lancet.* 2008; 372:1223-30.
  20. Report of Consumer Price Index of Thailand Year 2003-2013 (BASE YEAR 2011): Bureau of Trade and Economic Indices, Ministry of Commerce. [online] 2013. [cited 2013 Nov 27]; Available from: [http://www.indexpr.moc.go.th/price\\_present/TableIndexG\\_region.asp?table\\_name=cpig\\_index\\_country&province\\_code=&type\\_code=g&check\\_f=i&year\\_base=2554&nyear=2546](http://www.indexpr.moc.go.th/price_present/TableIndexG_region.asp?table_name=cpig_index_country&province_code=&type_code=g&check_f=i&year_base=2554&nyear=2546)
  21. Anukoolsawat P, Sritara P, Teerawattananon Y. Costs of lifetime treatment of acute coronary syndrome at Ramathibodi Hospital. *Thai Heart J.* 2006; 19:132-43.
  22. Khiaochaoen O, Pannarunothai S, Riewpaiboon W, Ingsrisawang L, Teerawattananon Y. Economic evaluation of rehabilitation services for inpatients with stroke in Thailand: a prospective cohort study. *Value Health Region Issues.* 2012; 1:29-35.
  23. Tsevat J, Goldman L, Soukup J, Lamas G, Connors K, Chapin C, et al. Stability of time-tradeoff utilities in survivors of myocardial infarction. *Med Decis Making.* 1993; 13:161-5.
  24. Wannasiri Y, Kapol N. The health utility of stroke patients at Ratchaburi Hospital (in Thai). *Region 4-5 Med J.* 2010; 29:95-103.
  25. Liew D, De Abreu Lourenco R, Adena M, Chim L, Aylward P. Cost-effectiveness of 12-month treatment with ticagrelor compared with clopidogrel in the management of acute coronary syndromes. *Clin Ther.* 2013; 35:1110-7.
  26. Calvert M, Freemantle N, Cleland J. The impact of chronic heart failure on health-related quality of life data acquired in the baseline phase of the CARE-HF study. *Eur J Heart Fail.* 2005; 7:243-51.
  27. Kraai I, Vermeulen K, Luttik M, Hoekstra T, Jaarsma T, Hillege H. Preferences of heart failure patients in daily clinical practice: quality of life or longevity. *Eur J Heart Fail.* 2013; 15:1113-21.
  28. Limwattananon S. Sensitivity analysis for uncertainty variables used in the model. In: Chaikletkaow U, Teerawattananon Y, editors. *Guideline of Health Technology Assessment for Thailand.* 2. Bangkok: Chulalongkorn University Book Center; 2013. p. 105-8.
  29. Gerlier L, Vellopoulou K, Lamotte M, Lacey L, Einroos A, Carr E. A health economic evaluation of omega-3 acid ethyl esters 90 in the secondary prevention post-MI in Ireland and Estonia. Abstract presented in ISPOR 15th Annual European Congress. 2012; 15:A373.
  30. Quilici S, Martin M, McGuire A, Zoellner Y. A cost-effective analysis of n-3 PUFA (Omacor) treatment in post-MI patients. *J Clin Prac.* 2006; 60:922-32.
  31. Lamotte M, Annemans L, Kawalec P, Zoellner Y. A multi-country health economic evaluation of highly concentrated n-3 polyunsaturated fatty acids in secondary prevention after myocardial infarction. *Pharmacoeconomics.* 2006; 24:783-95.