Structures of copper(II) 2-chlorobenzoate monohydrate and copper(II) 3,5-dichlorobenzoate trihydrate

Jana Medvecká^a, Jozef Halaška^a, Klaudia Jomova^b, Jan Moncol^a

^aFaculty of Chemical and Food Technology, Slovak Technical University, Radlinského 9, SK-812 37, Bratislava, Slovakia ^bDepartment of Chemistry, Faculty of Natural Sciences, Constantine The Philosopher University, SK-949 74 Nitra, Slovakia jan.moncol@stuba.sk

Abstract: The crystal structures of the title compounds, $[Cu_2(\mu 2-2-Clbz)_4(H_2O)_2]$ (1) and $[Cu(3,5-Cl_2bz)_2(H_2O)_3]$ (2), have been redetermined at 150 K. The compound 1 forms dimeric complex where two Cu^{2+} cations are bridged through four 2-chlorobenzoate ligands. The Cu^{2+} ion in 1 is pentacoordinated by four oxygen atoms of bridging 2-chlorobenzoate ligands and one coordinated water molecule. The compound 2 forms mononuclear complex, where the Cu^{2+} ion is pentacoordinated by two oxygen atoms of monodentate 3,5-dichlorobenzoate ligands and three coordinated water molecule. Both complexes are connected through $O-H\cdots O$ hydrogen bonds into frameworks. The π - π stacking interactions are also observed in crystal structure of 2.

Keywords: copper, X-ray, chlorobenzoate, dichlorobenzoate, hydrogen bonds

Introduction

Different benzoates, especially salicylates and fenamates are known to play an important role in human medicine (Sorenson, 1976; Weder et al. 2002). Our effort in this field has been targeted to halogeno- and dihalogeno- substituted benzoate complexes. The aim of this paper is to report an X-ray structures of the dinuclear complex [Cu(2-Clbz)₂(H₂O)]₂ and mononuclear complex $[Cu(3,5-Cl_2bz)_2(H_2O)_3]$. These complexes were previously prepared and published (Kawata et al., 1992; Wołodkiewicz et al., 1998). Object of our study is mention to conditions of preparation of reported complexes and re-determined of obtained complexes at 150 K because published results don't contain hydrogen bond networks and π - π stacking interactions in crystal structures of both complexes.

	1	2
Chemical formula	$C_{28}H_{20}Cl_4Cu_2O_{10}$	$C_{14}H_{12}Cl_4CuO_7$
$M_{ m r}$	785.32	497.58
Crystal system, space group	Monoclinic, $P2_1/c$	Monoclinic, $P2_1/c$
Temperature (K)	150	150
<i>a</i> , <i>b</i> , <i>c</i> (Å)	7.1940(4), 19.5140(16), 11.5060(4)	16.8280(4), 8.2930(7), 13.4700(14)
β (°)	113.636(3)	103.882(4)
$V(Å^3)$	1479.75(16)	1824.9(2)
Ζ	2	4
Radiation type	Mo K_{α}	Mo K_{α}
μ (mm ⁻¹)	1.856	1.817
Crystal size (mm)	$0.23 \times 0.20 \times 0.20$	$0.35 \times 0.31 \times 0.15$
Diffractometer	Bruker Kappa APEXII CCD	Bruker Kappa APEXII CCD
Absorption correction	Multi-scan SADABS (Bruker, 2008)	Multi-scan SADABS (Bruker, 2008)
T_{\min}, T_{\max}	0.675, 0.836	0.569, 0.772
No. of measured, independent and observed $[I > 2\sigma(I)]$ reflections	13697, 2976, 2410	14578, 3697, 2991
$R_{\rm int}$	0.0404	0.0379
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.0366, 0.0785, 1.091	0.0305, 0.0649, 1.068
No. of reflections	2976	3697
No. of parameters	199	235
$\Delta \lambda_{\rm max}$, $\Delta \lambda_{\rm min}$ (e Å ⁻³)	0.463, -0.594	0.360, -0.438
CCDC	869080	869081

Tab. 1. Experimental details.

Experimental

Synthesis

A water solution of copper(II) acetate monohydrate (0.200 g, 1 mmol in 40 ml of water) was stirred with nicotinamide (0.061 g, 0.5 mmol) for 1 or N,Ndiethylnicotinamide (0.084 ml, 0.5 mmol) for 2 and then the 2-chlorobenzoic acid (0.313 g, 2 mmol) for 1 or 3,5-dichlorobenzoic acid (0.382 g, 2 mmol) for 2 was added. These mixtures were stirred for 1 day resulting in a light green (1) or turquoise-blue (2) suspensions. The main products of 1 or 2 were filtered off and dried at ambient temperature and a green (1) or turquoise (2) mother liquid were left crystallization. Green block crystals of 1 and turquoise plate crystals of 2 were separated from the mother liquor after several weeks. Powders were obtained from systems without added neutral ligand too.

X-ray Crystallography

Crystal data and conditions of data collection and refinement are reported in Table 1. Data collection and cell refinement were carried out using diffractometer Bruker Kappa APEXII CCD (Bruker, 2008) at 150 K with graphite monochromated Mo K_{α} radiation. The semi-empirical absorption corrections were applied the using multi-scan method. The structures were solved by direct methods using SIR-97 (Altomare et al., 1999) or SHELXS-97 (Sheldrick, 2008) and refined by the full-matrix leastsquares procedure with SHELXL-97 (Sheldrick, 2008). Geometrical analyses were performed with SHELXL-97. The structures were drawn with XP in SHELXTL (Sheldrick, 2008).

Results and discussion

The principle structure features of $[Cu_2(2 Clbz)_4(H_2O)_2](1)$ are illustrated in Fig. 1 and selected bond distances and angles are given in Table 2. The compound 1 crystallizes in the monoclinic system (space group $P2_1/c$). The crystal structure of 1 at 295 K has already been reported (Kawata et al., 1992) and now it is re-determined at 150 K. The complex has been prepared by a different method (see experimental section) as reported previously (Kawata et al., 1992). The crystal structure of 1 consists of centrosymmetric tetracarboxylato-bridged dimmers $[Cu(2-Clbz)_2(H_2O)]_2$, where the Cu···Cu separation is 2.5934(6) Å. This molecular structure is typical of dimeric copper(II) carboxylate complexes, where the Cu···Cu distances are in the range 2.56-2.88 Å (Melnik et al., 1998; Sundberg et al., 1996). The τ value (Addison et al., 1984) of 0.0° for the copper atom is typical for square pyramidal geometry. The basal plane consists of four oxygen atoms O1, O2ⁱ, O3 and O4ⁱ [Symmetry code: (i) -x + 2, -y, -z + 2] at distances of 1.952(2)–1.985(2) Å. The apical site is occupied by the oxygen atom (O1W) from coordinated water molecules. The apical Cu-O1W bond length of 2.134(2) Å, and the Cu—O_{carb} bond lengths are in good agreement with those found in the other dimeric copper(II) carboxylate complexes (Melnik et al., 1998; Sundberg et al., 1996). The copper atom is displaced from the squares plane defined by the basal atoms toward the apical oxygen atom by 0.190(1) Å. The sum of the bond distances $(\Sigma Cu - L)$ 11.29 Å in **1** (the half value of the Cu · · · Cu distance is also included in the sum) agrees will with the values reported for known dimeric structures with the CuO₄O chromophores (Melnik et al., 1998). The values of angles O1—C1—O2 [124.5(3) Å] and O3—C8—O4 [124.4(3) Å] are in the range 11.6-132.0° found for tetracarboxylato dimers with CuO₄O chromophores (Sundberg et al., 1996).





The principal structural feature and atom numbering scheme of complex **2** are depicted in Fig. 2 and selected bond distances and angles are listed in Table 2. The compound **2** crystallizes in the monoclinic system with space group $P2_1/c$ and its structure at 295 K has already been reported (Wołodkiewicz et al., 1998). This report describes re-determination at 150 K complex from crystal prepared by a different method (see experimental section) as reported previously (Wołodkiewicz et al., 1998). The independent part of structure of **2** consists of $[Cu(3,5-Cl_2bz)_2(H_2O)_3]$ molecule elongated

Table 2. Selected geometric parameters (Å, ^o).

1			
Cu—O3	1.952(2)	Cu—O2 ⁱ	1.985(2)
Cu—O4 ⁱ	1.958(2)	Cu—O1W	2.134(2)
Cu—O1	1.960(2)	Cu—Cu ⁱ	2.5934(6)
O3—Cu—O4 ⁱ	168.78(8)	O3—Cu—O1W	94.15(8)
O3—Cu—O1	90.33(10)	O4 ⁱ —Cu—O1W	96.97(8)
O4 ⁱ —Cu—O1	89.92(10)	O1—Cu—O1W	96.11(8)
O3—Cu—O2 ⁱ	90.67(10)	O2 ⁱ —Cu—O1W	94.96(8)
O4 ⁱ —Cu—O2 ⁱ	86.94(10)	O1—Cu—O2 ⁱ	168.78(9)
2			
Cu1—O13	1.911(2)	Cu1—O1W	1.968(2)
Cu1—O12	1.944(2)	Cu1—O3W	2.291(2)
Cu1—O2W	1.966(2)		
O13—Cu1—O12	177.29(7)	O2W—Cu1—O1W	168.29(8)
O13—Cu1—O2W	91.93(7)	O13—Cu1—O3W	83.08(7)
O12—Cu1—O2W	90.22(7)	O12—Cu1—O3W	95.27(7)
O13—Cu1—O1W	89.27(7)	O2W—Cu1—O3W	89.90(7)
O12—Cu1—O1W	88.96(7)	O1W—Cu1—O3W	101.80(7)

Symmetry code: (i) -x + 2, -y, -z + 2.

tetragonal-pyramidal coordination polyhedron around Cu atom. The coordination sphere of copper(II) atom in *trans*-basal plane is formed by two 3,5-dichlorobenzoate anions bonded *via* O12 and O13 carboxylate oxygen atoms [Cu—O12 distance is of 1.944(2) Å and Cu–O13 is of 1.911(2) Å] and two water molecules bonded *via* O1W and O2W atoms [Cu—O1W distance is of 1.968(2) Å and Cu—O2W is of 1.966(2) Å]. Third water molecule is bonded in an apical position of square-pyramidal polyhedron *via* O3W donor atom with slightly longer bond distance Cu—O3W of 2.291(2) Å. The tetragonality parameter value (τ) (Addison et al., 1984) is about 0.15 indicating nearly ideal square-pyramidal coordination sphere. The basal planes of the neighboring complex molecules are coplanar and the distance between the basal planes is 3.518(1) Å. Moreover, the water molecules in basal position O1W form strong hydrogen bonds (Fig. 2 and Table 3) with pairs of carboxylate oxygen atoms O22 and O23 [O1W—H1W···O22ⁱⁱ and O1W—H2W···O23ⁱⁱ (Symmetry codes: (ii) -*x*+1, -*y*+1, -*z*+2) with O···O distances of 2.784(2) and 2.703(2) Å, respectively] which causes that the Cu···Cuⁱⁱ distance of neighboring copper atoms is 4.731(1) Å. These pairs of supramolecular synthons (Bernstein et al., 1995) $R_2^2(10)$ and $R_2^2(12)$ form supramolecular dimmers from complex molecules (**2**) which are to those found in monomeric complex [Cu(3-NO₂bz)₂(nia)(H₂O)₂] (nia=nicotinamide, 3-NO₂bz=3-nitrobenzoate) with



Fig. 2. Perspective view of 2, with the atom numbering scheme. Thermal ellipsoids are drawn at the 30% probability level.[Symmetry code: (ii) -x + 1, -y + 1, -z + 2].

D—H···A	<i>D</i> —H (Å)	$\mathrm{H} \cdot \cdot \cdot A$ (Å)	$D \cdot \cdot \cdot A$ (Å)	D—H · · · A (°)
1				
O1W—H1W···O4iii	0.84	2.41	3.028(3)	131
O1W—H1W···Cl2iii	0.84	2.54	3.333(2)	158
O1W—H2W···O2iii	0.84	2.04	2.827(3)	156
2				
O1W—H1W···O22ii	0.84	1.96	2.784(2)	167
O1W—H2W···O23ii	0.84	1.86	2.703(2)	178
O2W—H3W···O23iv	0.84	1.90	2.725(2)	168
O2W—H4W···O12v	0.84	1.89	2.724(2)	175
O3W—H6W···O22iv	0.84	2.18	2.999(2)	164

Table 3. Hydrogen-bond geometry (Å, ^o).

Symmetry codes: (ii) -x+1, -y+1, -z+2; (iii) x-1, y, z; (iv) -x+1, y+1/2, -z+3/2; (v) -x+1, y-1/2, -z+3/2.

tetragonal-pyramidal coordination of copper(II) atoms (Vaskova et al., 2010). The molecules of [Cu(3- $NO_{9}bz_{9}(nia)(H_{9}O_{9})$ are connected through similar O-H···O hydrogen bonds into supramolecular dimmers, which are shown weak anti-ferromagnetic interactions (Vaskova et al., 2010). The O-H···O hydrogen bonds in supramolecular dimmers of 2 are enriched by π - π stacking interactions (Janiak, 2000) between two 3,5-dichlorobenzoate ligands. The distances between two benzene ring are in the range 3.3–3.8 Å (Fig. 3) and $c_g \cdot \cdot \cdot c_g$ distance (Janiak, 2000) is 4.17 Å. The similar supramolecular synthons $R_2^2(10)$ and $R_2^2(12)$ (Bernstein et al., 1995) have been also observed in crystal structure of one dinuclear complex (Valigura et al., 2006) and several polymeric complexes (Abourahma et al., 2002; Burrows et al., 2008; Cao et al., 2008; Korabik et al., 2011; Maharramov et al., 2012; Vaskova et al., 2011; Xu et al., 2007; Yamada et al., 1998).

The crystal packing of **1** is shown in Fig. 3. The complex molecules of **1** are connected through O—H···O hydrogen bonds between coordinated water molecule (O1W) and carboxylate oxygen atoms of neighboring complex molecules $[O1W - H1W \cdots O4^{iii}]$ and $O1W - H2W \cdots O2^{iii}$ (Symmetry code: (iii) x - 1, y, z) with $O \cdots O$ distances of 3.028(3) and 2.827(3) Å, re-

spectively] (Table 3) into 1-D supramolecular chain in the zaxis (Fig. 3). The O1W—H1W \cdots O4ⁱⁱⁱ hydrogen bond is also supplemented by O1W—H1W \cdots Cl2ⁱⁱⁱ [O \cdots Cl distances is 3.333(2) Å] hydrogen bond (Table 3) and both ones form multicentered hydrogen bonds system (Desiraju & Steiner, 1999; Jeffrey, 1997; Gilli & Gilli, 2009).

The supramolecular dimmers of complex 2 are linked through next O-H···O hydrogen bonds between coordinated water molecules and carboxylate oxygen atoms [O2W—H3W···O23^{iv}; O2W—H4W···O12^{v,} and O3W—H6W···O22^{iv} (Symmetry codes: (iv) -x+1, y+1/2, -z+3/2; (v) -x+1, y-1/2, -z+3/2) with O···O distances of 2.725(2); 2.724(2) and 2.999(2) Å, respectively] (Table 3) forming 2-D framework in bc plane (Fig. 4). The π - π stacking interactions (Janiak, 2000) between two 3,5-dichlorobenzoate ligands are also observed between two supramolecular dimmers (Fig. 5). The distances between two benzene rings (between two supramolecular dimmers) are in the range 3.18–3.50 Å (Fig. 3) and $c_g \cdots c_g$ distance (Janiak, 2000) is 3.61 Å.

Both studied complexes have been prepared similar method, but this method is different that published methods of synthesis of complex **1** (Kawata et al.,



Fig. 3. The two 1-D chains connecting through O—H···O and O—H···Cl hydrogen bond in crystal structure of **1**. [Symmetry code: (iii) x - 1, y, z].



Fig. 4. The 2-D framework connecting through O—H····O hydrogen bond in crystal structure of 2. The benzene rings and chlorine atoms are omitted for clarity.
[Symmetry codes: (ii) -*x*+1, -*y*+1, -*z*+2; (iv) -*x*+1, *y*+1/2, -*z*+3/2; (v) -*x*+1, *y*-1/2, -*z*+3/2].



Fig. 5. The π - π stacking interactions in crystal structure of 2.

1992) or **2** (Wołodkiewicz et al., 1998). The structures of both complexes have been determinated at room temperature, but hydrogen atoms have not been calculated in corrected positions, but Low-temperature re-determination at 150 K allows determinate more better results and the hydrogen atoms are calculated in correct positions. Novelty in present paper is description of hydrogen bond networks and π - π stacking interactions in crystal structures of both complexes. The studies of hydrogen bonds and intermolecular interaction have been absent in original room temperature crystal structures (Kawata et al., 1992; Wołodkiewicz et al., 1998).

At the end we would like to pointed out, presence of neutral ligand in reaction mixture lead to formation of complexes without ligand in coordination sphere. They were prepared by different methods of syntheses in the past. Re-determined of obtained complexes at 150 K we specify basic features and parameters and we specify shape of coordination polyeder of known complexes. Description of hydrogen bond networks and π - π stacking interactions in crystal structures of both complexes have been presented too.

Acknowledgement

This work was supported by the Scientific Grant Agency of the Ministry of Education of the Slovak Republic (VEGA Project Nos. 1/0856/11 and 1/0562/10) and by the Research and Development Agency of the Slovak Republic (contract No. APVV-0202-10).

References

- Abourahma H, Moulton B, Kravtsov V, Zaworotko MJ (2002) J. Am. Chem. Soc. 124: 9990.
- Addison AW, Rao TN, Reedijk J, Rija J, Verchoors GC (1984) J. Chem. Soc. Dalton Trans. pp. 1349.
- Altomare A, Burla MC, Camalli M, Cascarano GL, Giacovazzo C, Guagliardi A, Moliterni AGG, Polidori G, Spagna R (1999) J. Appl. Cryst. 32: 115.
- Bernstein J, Davis RE, Shimoni L, Chang NL (1995) Angew. Chem., Int. Ed. Engl. 34: 1555.
- Bruker (2008) APEX2, SAINT-Plus and SADABS. Bruker AXC Inc., Madison, Wisconsin, USA.
- Burrows AD, Frost CG, Mahon MF, Winsper M, Richardson C, Attfield JP, Rodgers JA (2008) Dalton Trans. pp.6788.
- Cao R, Lu J, Batten SR (2008) CrystEngComm 10: 784.
- Desiraju GR, Steiner T (1999) The Weak Hydrogen Bonds and in Structural Chemistry and Biology, Oxford University Press, Oxford, UK.

Janiak C (2000) J. Chem. Soc., Dalton Trans. pp. 3885.

- Jeffrey GA (1997) An Introduction to Hydrogen Bonding, Oxford University Press, Oxford, UK.
- Kawata T, Ohba S, Tokii T, Muto Y, Kato M (1992) Acta Cryst. C48: 1590.
- Korabik M, Repicka Z, Martiska L, Moncol J, Svorec J, Padelkova Z, Lis T, Mazur M, Valigura D (2011) Z. Anorg. Allg. Chem. 637: 224.
- Gilli G, Gilli P (2009) The Nature of the Hydrogen Bond, Oxford University Press, Oxford, UK.
- Maharramov AM, Mardanova VI, Chyraqov FM, Gurbanov AV, Ng SW (2012) Acta Cryst. E68: m128.
- Melnik M, Kabesova M, Koman M, Macaskova L, Garaj J, Holloway CE, Valent A (1998) J. Coord. Chem. 45: 147.
- Sheldrick GM (2008) Acta Cryst. A64: 112.
- Sorenson [R] (1976) J. Med. Chem. 19: 135.
- Sundberg MR, Uggla R, Melnik M (1996) Polyhedron 15: 1157.
- Valigura D, Moncol J, Korabik M, Pucekova Z, Lis T, Mrozinski J, Melnik M (2006) Eur. J. Inorg. Chem. pp. 3813.
- Vaskova Z, Moncol J, Korabik M, Valigura D, Svorec J, Lis T, Valko M, Melnik M (2010) Polyhedron 29: 154.
- Vaskova Z, Moncol J, Korabik M, Medvecka J, Svorec J, Padelkova Z, Valko M, Valigura D (2011) Polyhedron 30: 86.
- Weder JE, Dillon CT, Hambley TW, Kennedy BJ, Lay PA, Biffin JR, Regtop HL, Davies N M (2002) Coord. Chem. Rev. 232: 95.
- Wołodkiewicz W, Brzyska W, Głowiak T (1998) Monatsh. Chem. 129: 111.
- Xu JY, Chen BL, Ng SW (2007) Acta Cryst. E63: m2200.
- Yamada Y, Ueyama N, Okamura T, Mori W, Nakamura A (1998) Inorg. Chim. Acta 275-276: 43.