

Proteomics of heavy metal toxicity in plants

Petra Cvjetko¹, Mira Zovko², and Biljana Balen^{1,2}

*Department of Molecular Biology, Faculty of Science, University of Zagreb¹, Croatian Environment Agency²,
Zagreb, Croatia*

Received in August 2013

CrossChecked in August 2013

Accepted in November 2013

Plants endure a variety of abiotic and biotic stresses, all of which cause major limitations to production. Among abiotic stressors, heavy metal contamination represents a global environmental problem endangering humans, animals, and plants. Exposure to heavy metals has been documented to induce changes in the expression of plant proteins. Proteins are macromolecules directly responsible for most biological processes in a living cell, while protein function is directly influenced by posttranslational modifications, which cannot be identified through genome studies. Therefore, it is necessary to conduct proteomic studies, which enable the elucidation of the presence and role of proteins under specific environmental conditions. This review attempts to present current knowledge on proteomic techniques developed with an aim to detect the response of plant to heavy metal stress. Significant contributions to a better understanding of the complex mechanisms of plant acclimation to metal stress are also discussed.

KEY WORDS: *heavy metal stress; mass spectrometry; plant proteome; protein biomarkers; two-dimensional electrophoresis*

The ever-increasing human population is in growing demand for agricultural production. Whereas global warming represents a major impediment to the increase of arable surfaces (1-3), the wide variety of human activities inherent to this demand results in an immense burden to the environment from pollutants such as metals and metalloids. For example, smelting operations, fossil fuel combustion (4), and contamination with arsenic (As)-based pesticides (5) are major sources of arsenate in environment, alongside geologically derived activities. Plants are not resistant to arsenate because it is an analogue of phosphate, competing for the same uptake carriers in the root (6). Nitrates and phosphates are also one of the major pollutants leaching to groundwater. The excessive use of nitrogen-based fertilisers and industrial effluents, as well as biological waste and

landfills that have not gone through nitrification, lead to nitrogen pollution (7, 8). In addition, increased levels of phosphate from phosphate fertiliser industries may cause soil contamination, which is potentially hazardous to plants and groundwater (9).

The last century has seen a tremendous rise in heavy metal emission and an enormous release of these elements into soil, water, and air through sources such as industrial activity, wastewater treatment, contaminated sewage sludge, phosphate fertilizers, and various agrochemicals (10, 11). All metals of a specific gravity higher than $>5 \text{ g cm}^{-3}$ (12) and an atomic number over 20 (13) belong to the group of heavy metals. Some, such as zinc (Zn), iron (Fe), manganese (Mn), molybdenum (Mo), and copper (Cu), are micronutrients essential for physiological processes in plants, metalloproteins and ion-dependent

enzymes such as ascorbate oxidase, cytochrome catalase, peroxidase, superoxide dismutase, nitrate reductase, etc. (14). These metals may become toxic if their concentrations exceed the levels necessary for plant metabolism (13). On the other hand, metals such as cadmium (Cd), mercury (Hg), and lead (Pb) are not required for plant metabolism and can cause harmful effects even at very low concentrations. Furthermore, certain non-physiological ions are able to replace physiological ions and still perform physiological functions for a certain time without harming a plant. For instance, strontium (Sr^{2+}) can replace calcium ions during heat stress in moss (15).

Soil and water contaminated with toxic heavy metals pose a major environmental and human health issue. However, the metabolism of certain plants can be used to reduce the content or toxic effects of metals in the environment (16). The phytoremediation of metals refers to the use of metal-accumulating plants and their associated microbiota for environmental clean up. It is widely accepted as an environmentally friendly, non-invasive and cost-effective restoration technology attractive for the remediation of metal-contaminated sites (17).

As sessile organisms, plants are inevitably exposed to environmental concentrations of heavy metals, either geologically or through human activities (18, 19). Heavy metals enter a plant from soil and water through the root system via passive or active transport (20). After the uptake, heavy metal ions can be translocated from the root via the xylem to the aerial parts of a plant such as stems and leaves (21). Plants exposed to heavy metal stress are capable of accumulating high levels of heavy metals following an accumulation gradient in organs: roots > stems > leaves > seeds (22). The toxic effect of certain heavy metals depends on many factors, primarily on the plant species, its stage of development, and the concentration and chemical properties of the metal.

All plants possess a basal tolerance to heavy metals, but elevated concentrations can cause harmful effects and disturbances such as transpiration and photosynthesis inhibition, disturbance of carbohydrate metabolism, nutrition stress, and oxidative stress, all of which collectively yield a general effect on plant development and growth (23). The uptake of non-essential and potentially high toxic heavy metal ions such as Cd^{2+} , Hg^{2+} , and Pb^{2+} , as well as metalloids As (phytoavailable ion forms are arsenate, AsO_4^{3-} and arsenite, AsO_3^{3-}) and selenite (Se) (ion forms selenate, SeO_4^{2-} and selenite, SeO_3^{2-}) (21) is of major concern.

Typical symptoms of heavy metals stress are often similar to those of essential nutrient deficiency (13); weaker growth, lower biomass, necrotic spots in leaves, and damage mainly at the root level (24-26). Accumulation of higher concentrations can cause reductions in crop yields (27), but can also result in a potential threat to human health (28, 29) because heavy metals thus become part of the food chain.

In order to alleviate the stress and re-establish cellular homeostasis and antioxidant capacity, plants have developed highly effective mechanisms to regulate the uptake, accumulation, distribution, and detoxification of heavy metal ions (30). Vital components in these mechanisms are metal transporters responsible for metal uptake and vacuolar transport, chelators for heavy metal detoxification and tolerance, and chaperones which play a significant role in the delivery and trafficking of metal ions (31, 32), all of which are protein molecules. That is why the identification of protein markers could serve as a good starting point for revealing new aspects of heavy metal stress in plants (24).

Proteomes and proteomics

Ever since the completion of genome sequencing projects on many model plant species, such as *Arabidopsis thaliana* (33), moss *Physcomitrella patens* (34), rice (35), and other important crops (36), plant biologists have faced the constant challenge of successfully designating the role and function of each gene as well as their products. They do so in order to complete and improve our understanding of how biological macromolecules interact to shape an organism as a fully functional entity. Due to the extensive number of experimental data from different high-throughput analyses, new terminology has recently been coined. In an attempt to describe a complete dataset of particular biomolecules, two interrelating suffixes were introduced: “-ome” to refer to a particular population of molecules in the cell (e.g., genome, proteome, transcriptome, metabolome, etc.) and “-omics” to highlight a research interest (such as transcriptomics, proteomics, metabolomics, and metabolomics). To obtain deeper insight into the glossary and taxonomy of current “-omes” and “-omics”, readers are encouraged to visit the web page of the genomic glossary (37).

The application of modern “-omics” techniques has been useful in expanding our knowledge of molecular pathways involved in the tolerance toward different environmental stresses. Therefore, if we are

to obtain deeper insight into the plant stress response to biotic or abiotic stresses (38), traditional biochemical and physiological approaches are simply no longer sufficient.

Plant response to abiotic stress, heavy metals included, often occurs as a consequence of changes in protein expression. This response is usually not only accompanied by an alteration in the gene expression pattern, but also by inevitable qualitative and quantitative changes in proteins. Genomic technologies monitor changes at the transcriptional level (i.e., mRNA level), which need not correlate with changes at the translational level (i.e., protein level). Put more simply, a positive correlation would suggest that mRNA transcripts are immediately translated to proteins. According to Anderson and Seilhamer (39), the correlation coefficient in quantity between mRNA and protein abundance is relatively low, approx. 0.5. In reality, the mRNA/protein ratio depends on both the mRNA translation rate and protein stability (*de novo* synthesis vs. degradation). Moreover, mRNA availability can be affected by post-transcriptional processes at the stage of RNA maturation, splicing, transportation or translation initiation as well as degradation (40). Therefore, the mRNA levels in a particular biological sample cannot always be used to predict changes in protein expression (30, 41). In addition, protein expression is also regulated at the posttranslational level via glycosylation, phosphorylation, phosphorylation, sumoylation or ubiquitination (42, 43). Proteins, when perceived as the functional translation portion of a genome, play very important roles throughout the plant organism in terms of numerous, structural and functional roles, as well as via effector molecules, in alleviating the stress response induced by internal or environmental stimuli (44). Protein function can also be perceived depending on the level (biochemical, cellular or higher level of biological organization such as organ or whole organism, as well as developmental or physiological) at which a particular protein is studied.

Based on the expanding knowledge on protein function and structure, new technology for their high-throughput analysis has emerged in the form of proteomics. Since proteins are crucial for various cellular processes, proteomics is a promising tool to assess changes in proteins following a variety of stress conditions. Originally, the concept of the proteome was established by Marc Wilkins in the mid-1990s (45) as a combination of the words *protein* and *genome*. Widely accepted inside the research

community, a proteome is regarded as a complete set of proteins expressed by a genome at a given point in time. More specifically, it represents the entire set of proteins expressed in a specific type of cell or tissue under a specific condition at a specific point of time (46). Proteomics comprises many different aspects of protein roles in the living cell, including the identification and quantification of proteins, patterns of protein expression, posttranslational modification, and protein-protein interactions.

The methodology of proteomics

Since proteomics is focused on the functional translated portion of the genome, its use is expanding rapidly in the field of heavy metal stress biology. However, proteomic studies of metal stress have been far less numerous and comprehensive in comparison with investigations regarding other environmental stresses in plants (30). Although the processes of heavy metal uptake, accumulation, distribution, oxidative stress induction, and detoxification have been investigated in a wide range of studies on plants (47-54), the mechanisms involved are still only partially understood. The proteomic approach can therefore help to elucidate new aspects of plant metal stress.

The differential-expression approach in proteomics is used for describing sets of proteomes differing both in protein quality and quantity and is aimed at protein identification and relative quantitative determination. This methodology is most frequently applied in studies dealing with heavy metal stress in plants. It is based on the comparison of the composition of different proteomes, which originate from non-stressed (control) plants and corresponding plants exposed to heavy metals (24, 25, 55-67). An alternative approach is to compare proteomes from different genotypes with a distinct tolerance for heavy metal stress [e.g., differential responses of proteins in two contrasting Cd-accumulating soybean cultivars (68, 69), comparative analysis of proteomic changes in contrasting flax cultivars upon Cd exposure (70), or a comparative proteomic analysis of roots under Al stress in Al-tolerant and Al-sensitive soybean genotypes (71)].

Protein extraction

At the present, many studies (24-26, 49, 60, 62) investigating heavy metal toxicity-induced differentially expressed proteins or heavy metal stress-responsive proteins in plants apply a classical gel-

based proteomic approach, which includes One- (1-DE) or Two-Dimensional electrophoresis (2-DE) for a high quality protein separation followed by protein identification via mass spectrometry (MS) analysis and database search (Figure 1).

The extraction of proteins and the preparation of samples is one of the most challenging steps in any proteomics study, particularly in those that comprise 2-DE as their first step. Effective protein extraction, solubilisation, and purification from plant tissues are essential for a good and reproducible 2-DE approach, as the amount and quality of extracted proteins ultimately determine the protein spot number, resolution and intensity. Important characteristics of a good protein isolation method are the reproducible capture and solubilisation of a full set of proteins from a certain sample with minimal post-extraction artefact formation, proteolytic degradation and contamination with non-protein molecules (72, 73). In plant proteomics, there are several factors that can affect the method chosen for protein extraction, such as plant species, tissue type, organ, cell, organelle, and the chemical and physical properties of the observed proteins. Generally, in the preparation of protein samples, plants are regarded as “difficult” or recalcitrant tissues due to their low protein content and abundance of cell walls and vacuoles, which account for most of the cell mass (74). Another issue in protein extraction from plant tissues is the presence of phenolic compounds, proteolytic and oxidative enzymes, as well as high levels of polysaccharides, lipids, and other secondary metabolites. These compounds generate horizontal and vertical streaking and smearing and reduce protein resolution on 2-DE gels (30, 75). Therefore, in plant protein extraction and solubilisation, the buffer composition with regard to combinations and concentrations of detergents and reducing and chaotropic agents has a considerable effect on the quality of the extracted proteins (76). The most frequently used protocols for the extraction of plant proteins involve a precipitation step, which should serve to separate proteins from interfering compounds originating from a plant’s secondary metabolism. Chemicals commonly used for this purpose are acetone, trichloroacetic acid (TCA)/acetone, and Tris-buffered phenol. A comparison of acetone, TCA/acetone, and phenol extraction methods applied on three plant species, regarded as recalcitrant for 2-DE protein extraction and downstream processing, has shown that phenol extraction was superior to the other two methods, providing more

protein bands or spots on the gels and less proteolysis (77). Moreover, a comprehensive proteomic study performed on nine organs from soybean plants in various developmental stages revealed that, whereas the use of an alkaline phosphatase buffer followed by TCA/acetone precipitation caused horizontal streaking in 2-DE, the extraction with alkaline phenol and methanol/ammonium acetate produced high-quality proteome maps with well-separated spots, high spot intensities, and high numbers of separate protein spots in 2-DE gels (78). Sarma et al. (79) also reported a similar preference toward the use of alkaline phenol and methanol/ammonium acetate rather than TCA/acetone for the extraction of soybean tissues. Some other proteomic studies dealing with plant response to heavy metal stress have confirmed that phenol-based protocols are the most effective protein extraction methods for obtaining a high-quality proteome map (24, 62, 68, 80-84).

The abovementioned extraction procedures are generally used in the separation of soluble proteins by classical proteomic approaches. In the case of organelle proteomics, particularly those that include membrane proteins, a modified extraction procedure should be applied to dissolve hydrophobic proteins and additional purification steps need to be included (85, 86). Furthermore, when studying protein-protein interactions, it is necessary to extract protein complexes by using buffers with mild non-denaturing and non-ionic detergents, such as digitonin and dodecyl maltoside, which stabilise protein supercomplexes and allow the dissociation and separation of each protein complex, and therefore, keep proteins in their native states (87).

One of the fundamental challenges in proteome analysis is the fact that samples are often dominated by a relatively small number of high-abundance proteins whose presence can obscure that of the less abundant ones, thus limiting the capacity and resolution of the separation technique(s) employed. Among plant tissues, this problem is very apparent in leaves and other photosynthetic tissues, where the majority of proteins is highly hindered by a single protein, ribulose biphosphate carboxylase (RuBisCo) (88). Therefore, a group of methodological strategies for the separation of protein molecules for proteomics, called separomics, is applied for the identification of low-abundance proteins and peptides in plants (89). Different techniques have been developed to remove or reduce a substantial portion of RuBisCo from leaf protein extracts, such as the partial removal of




Figure 1 Outline of the gel-based proteomic approach for studying heavy metal stress in plants

RuBisCo by the increasing dithiothreitol concentration in rice leaves extracts (90), or by applying a fast and simple fractionation technique using 10 mmol L⁻¹ Ca²⁺ and 10 mmol L⁻¹ phytate to precipitate RuBisCo from soluble protein extracts of soybean leaf (91). Moreover, additional methods have been employed to reduce high-abundance proteins in extracts in order to improve the low-abundance protein detection, such as addition of isopropanol to remove storage proteins from soybean seeds extracts (92), division of the proteome into subproteomes in membranes of banana leaves (93), and the use of affinity and immunoaffinity columns for protein purification (94, 95). One particular methodology seems very promising – the use of combinatorial hexapeptide ligand libraries (CPLL; such as ProteoMiner), which have been applied for the detection of low abundance proteins in plant tissues and demonstrated a remarkable decrease in high-abundance proteins as well as an enrichment of less abundant proteins (96, 97). The concomitant enrichment of less abundant proteins may facilitate a deeper insight into the plant proteome, which could lead to novel findings of the changes in plant proteins induced by heavy metal stress.

One- and two-dimensional polyacrylamide gel electrophoresis (1- and 2-DE)

An average protein extract, even from a purified fraction, is a mixture of a large number of individual proteins as well as several other non-protein components. The main goal of the analysis of such a mixture is to acquire as much qualitative and quantitative information on the investigated proteome as possible. One of the most frequently applied approaches is to provide protein separation by means of gel electrophoresis, which is based on the migration of molecules in an electric field in a gel medium. One- and Two-Dimensional polyacrylamide gel Electrophoresis are common separation methods employed in the analysis of proteins. Both electrophoretic techniques are frequently used to profile cell or tissue proteins or determine the purity of native or recombinant proteins (98, 99). One-Dimensional Electrophoresis has been successfully applied for the analysis of protein profile of tobacco seedlings and adult plants experimentally exposed to Cd and Zn (100) as well as in the study of physiological and proteomic changes in germinating rice seeds induced by Cu (58). However, 1-DE has a major disadvantage in resolving protein mixtures according to only one physicochemical attribute, mostly

molecular weight (MW) (101). Therefore, it is very likely that one protein band obtained on a gel comprises numerous proteins of different characteristics and functions, simply because they all possess a very similar MW. On the other hand, 2-DE represents a much more powerful technique for protein separation since it resolves proteins by two properties (102). In classical gel-based proteomics, proteins are resolved in the first dimension by isoelectric focusing (IEF), which separates proteins according to their charge, i.e. isoelectric point (pI), the pH at which an amino acid does not migrate in an electric field (103). This is the most critical step of the 2-DE process. There are two alternative methods to create a pH gradient - carrier ampholytes and immobilized pH gradient (IPG) gels. When IPGs were introduced for the first dimension, they brought about better resolution and reproducibility in protein separation (104). IPG gels have a dried homogeneous polyacrylamide gel matrix covalently bound to a plastic carrier and are available in several overlapping pH ranges (narrow-, medium-, and wide-range) and different strip lengths. Prior to use, IPG gels are rehydrated with an adequate buffer. Protein samples can be included directly in the rehydration solution for efficient loading of dilute samples or for loading larger amounts of protein. Alternatively, samples can be loaded after rehydration directly on the IPG gels. The second dimension is a classical Sodium Dodecyl Sulphate Polyacrylamide Gel Electrophoresis (SDS-PAGE), which separates proteins according to their MW. Each spot on the resulting 2-D gel corresponds to a single protein in the sample. Thousands of different proteins can be separated and information such as the pI, apparent MW and the amount of each protein can be obtained (103, 104).

Methodological advances in 2-DE have led to the development of 2-D Differential In-Gel Electrophoresis (2-D DIGE) (105), a variant of 2-DE that overcomes some of the drawbacks of the original technique, mainly the variation between gels that often leads to multiple repeat runs. It is a high throughput technique that enables several samples to be simultaneously separated and visualized in one gel by labelling proteins with various fluorescent reagents before 2-DE. Different pools of proteins are labelled with different fluorescent dyes (up to three, Cy2, Cy3, or Cy5). The samples are then mixed and resolved on the same 2-D gel. The gel is scanned with the excitation wavelength of each dye one after the other, which allows imaging of each sample separately, or the

images of all samples can be overlapped to allow better comparison of differentially expressed proteins (106). Cy fluorescence is linear over a wide range of concentrations and sensitive down to 125 pg (105). Accordingly, this method successfully overcomes limitations in traditional 2-DE, considerably reduces gel-to-gel variation, and improves the power of quantitative analyses. Two-Dimensional DIGE also has the advantage of running an internal standard as part of each separation, which allows the measurement of the abundance of a protein in each sample relative to the internal standard. This technique can reveal the presence of several hundred proteins in a highly reproducible manner and remains unchallenged as the most efficient method for analysing complex protein mixtures. So far 2-D DIGE has been used to successfully identify heavy metal-induced differentially regulated proteins in Cd-exposed poplar plants (25), for proteomic analysis of soybean roots under aluminium (Al) stress (71) and to affirm Al-induced proteome changes in tomato cotyledons (67) (Table 1). The results obtained by these studies (25, 67, 71) indicate that 2-D DIGE in combination with MS is a powerful tool for the molecular characterization of plant heavy metal stress and identification of stress-specific protein markers.

For a high resolution separation of enzymatically active protein complexes from tissues and cells, Blue Native Polyacrylamide Gel Electrophoresis (BN-PAGE) is the method of choice (107, 108). It is a special form of native electrophoresis, which allows separation of proteins between 10 and 10000 kDa (109), and is particularly suitable for separation of membrane protein complexes after their solubilisation with mild neutral detergents. The separation principle relies on binding of dye Coomassie blue G-250, which provides negative charges to the surface of the protein. During migration to the anode, protein complexes are separated according to MW as blue bands. BN-PAGE allows efficient protein resolution under native conditions and therefore proteins keep their activity. BN-PAGE gel strips or individual protein complex bands can be used for different downstream analyses. BN-PAGE in conjunction with denaturing SDS-PAGE can give insights into the protein complex composition and protein-protein interactions as it provides a better separation of high MW protein complexes associated with the proteomes of organelles such as chloroplasts and mitochondria (110).

The principle of the conjunction of these two electrophoretic methods is that protein complexes are

separated according to their sizes using BN-PAGE; a BN-PAGE gel strip is denatured and placed on the top of a second dimension SDS-PAGE gel; in the second dimension the protein complexes are separated into their individual subunits. Two-dimensional BN-SDS-PAGE (2-D BN-SDS-PAGE) coupled with MS has been used so far to analyse the heavy metal stress responses of multiprotein complexes in the leaf apoplast proteome (59, 111) and thylakoid membrane (112) (Table 1), and also permits direct quantitative assessment of differential changes in a given proteome.

Gel image analysis

Various types of imaging systems and associated software are commercially available for analysing gels stained with any of the commonly used gel stains. ImageQuant TL (GE Healthcare), an automatic and easy-to-use general image analysis software represents a very frequently applied technique, although there are several other programs that can be found on the market like EZQuant-Gel (Microsoft™ Windows™ application for analysis of 1-D gels), Quantity One (Bio-Rad), and BioDocAnalyze (Biometra) that are of equally high quality. Apart from these commercially available programs there is also a number of freeware software for 1-D gel electrophoresis image analysis like GelAnalyzer (113) and GelScape (114), which are free for both educational and commercial usage, although specifically designed for educational institutes and researchers. All 1-D gel analysis software has similar characteristics; they can define lanes and bands, quantify bands, create standard curves and determine MW. In all of these programs, images can be adjusted for contrast, processed in various ways, annotated and exported to other files for publication or document control.

The 2-D analysis software assigns and quantifies protein spots in 2-DE gels. Moreover, these programs include statistical software designed for quantitative comparisons of large numbers of gels and are suitable for analysis of spot patterns derived from differentially expressed proteins. Some can be supported by databases to help protein identification. All 2-D analysis software can be used for image adjustments, annotation and export in a variety of file forms. The most prominent among them are ImageMaster 2D Platinum (GE Healthcare), PDQuest (Bio-Rad), REDFIN (Ludesi), Dymension (Syngene), "Progenesis" (Nonlinear Dynamics), and "Proteovue" (Eprogen). Differential labelling with fluorescent dyes Cy2, Cy3 and Cy5 (DIGE technology, GE Healthcare)

Table 1 Summary of proteomic studies of plant response to heavy metal stress published between 2003 and 2013

Metal	Plant species and tissue	Proteomic methodology	Major findings	Ref.
Cd	<i>Populus tremula</i> L. leaf, root	IPG, 2-D DIGE, MALDI-TOF-TOF MS	Deleterious effect on proteins involved in the primary carbon metabolism and oxidative stress response	25
	<i>Populus tremula</i> L. leaf	IPG, 2-D DIGE, MALDI-TOF-TOF MS	Enhanced accumulation of stress-related proteins	26
	<i>Brassica juncea</i> L. (Acc: PI 173874) root	IPG, 2-D DIGE, iTRAQ, nano LC-MS/MS	Overexpression of sulfite reductase and O-acetylserine sulphydrylase, involved in reduction of sulfate to cysteine	80
	<i>Hordeum vulgare</i> L. var. Baraka leaf mesophyll tonoplast	iTRAQ labelling, MALDI-TOF-TOF MS	MRP-like ABC transporter and two novel CAX transporters (CAX1a and CAX5) assure Cd ²⁺ transport into vacuole	130
	<i>Spinacia oleracea</i> L. thylakoid membrane	IPG, 2D BN-SDS-PAGE, nano LC-MS/MS	Strong reduction of chlorophyll concentration; significant reduction of antenna proteins of PSI, while PSII antennae were affected to a minor extent	112
	<i>Oryza sativa</i> L. cv. Dongjin leaf, root	IPG, 2-DE, MALDI-TOF MS	Activation and up-regulation of ROS scavengers	62
	<i>Lycopersicon esculentum</i> Mill cv. Tres Cantos root	IPG, 2-DE, MALDI-TOF MS, LIFT TOF-TOF	Up-regulation of Hsp70 and higher abundance of glycolytic and TCA cycle enzymes	83
	<i>Linum usitatissimum</i> L. cv. Jitka, cv. Tábor cell suspension culture	IPG, 2-DE, MALDI-TOF MS	Up-regulation of proteins associated with Cd-chelating pathways	70
	<i>Arabidopsis thaliana</i> ecotype Columbia seedlings (aerial parts)	IPG, 2-DE, LC-ESI-MS/MS	Up-regulation of proteins associated with Cd-chelating pathways	66
	<i>Glycine max</i> L. cv. Enrei, Harosoy root microsome	IPG, 2-DE, nano LC-MS/MS	Up-regulation of proteins associated with Cd-chelating pathways	68
Cu	<i>Glycine max</i> L. cv. Enrei, Harosoy, Fukuyutaka, CDH-80 leaf, root	IPG, 2-DE, nano LC-MS/MS, MALDI-TOF MS	Activation and up-regulation of ROS scavengers accumulation of molecular chaperons and heat-shock proteins	61
	<i>Glycine max</i> L. cv. Enrei leaf	IPG, 2-DE, nano LC-MS/MS, MALDI-TOF MS	Abundance of Hsp70 and Peroxiredoxin	69
	<i>Cannabis sativa</i> var. Felina 34 root	IPG, 2-DE, LC-MS/MS	Down-regulation of glycolytic enzyme ENO, the metallo enzyme that catalyses conversion of 2-phosphoglycerate to phosphoenolpyruvate	24
	<i>Oryza sativa</i> L. cv. Hwayeong germinating seeds	1-DE, IEF tube gel, 2-DE, MALDI-TOF MS	Accumulation of antioxidant and stress-related proteins	58
	<i>Oryza sativa</i> L. Wuyunjing germinating embryos	IPG, 2-DE, MALDI-TOF MS	Increased abundance of PR-10a and putative PR proteins as well as of GSTs; down-regulation of Trx and Trx-POD	116
	<i>Elsholtzia splendens</i> L. leaf, root	IPG, 2-DE, MALDI-TOF MS, LTQ-ESI-MS/MS	Up-regulation of proteins with potential ion-binding functions and proteins engaged in vacuolar metal sequestrations	115
	<i>Ectocarpus siliculosus</i> strains Es32, Es524 algal cells	IPG, 2-DE, MALDI-TOF MS	Up-regulation of photosynthesis, glycolysis, and pentose phosphate metabolism	82

Table 1 Continued

Metal	Plant species and tissue	Proteomic methodology	Major findings	Ref.
As	<i>Oryza sativa</i> L. cv. Dongjin root	IPG, 2-DE, MALDI-TOF MS	Enhanced expression of antioxidative enzymes	56
	<i>Agrostis tenuis</i> L. leaf	IPG, 2-DE, MALDI-TOF MS	Degradation of several protein involved in photosynthesis, RuBisCo in particular	117
	<i>Oryza sativa</i> L. cv. Dongjin leaf	IPG, 2-DE, MALDI-TOF MS, ESI-MS/MS	Over-expression of energy and metabolism related proteins	55
	<i>Anabaena</i> sp. PCC7120 algal cells	PG, 2-DE, MALDI-TOF MS, LC-MS	Over-expression of energy and metabolism related proteins	63
Al	<i>Glycine max</i> (L.) Merr cvs. BaXi 10, BenDi 2 root	IPG, 2-DE, MALDI-TOF MS	CS and GSH play essential role in Al adaptation; activation of molecular chaperons	118
	<i>Oryza sativa</i> L. cv. Xiangnuo 1 (XN1) root	IPG, 2-DE, MALDI-TOF MS, MALDI-TOF/TOF MS	CS and GSH play essential role in Al adaptation; enhanced accumulation of several enzymes involved in ROS detoxification	119
	<i>Solanum esculentum</i> Mill cv. Money Maker root	IPG, 2-DE, MALDI-TOF/TOF MS	Induced enzymes with antioxidant activities and detoxification	67
	<i>Glycine max</i> (L.) genotypes PI 416937 and Young root	IPG, 2-D DIGE, MALDI-TOF MS, MALDI-TOF/TOF MS	Tolerant genotype - accumulation of enzymes which catalyse synthesis of citrate, involved in Al ³⁺ detoxification; sensitive genotype – induction of proteins related to general stress response	71
Mn	<i>Vigna unguiculata</i> [L.] Walp. Cvs TVu 91, TVu 1987 leaf apoplast	IPG, 2-DE, 2D BN-SDS-PAGE, nano LC-MS/MS	Role of PODs in the expression of Mn ²⁺ toxicity mediating H ₂ O ₂ production/ consumption and the oxidation of phenols	59
	<i>Vigna unguiculata</i> [L.] Walp. Cvs TVu 91, TVu 1987 leaf symplast	IPG, 2-DE, nano LC-MS/MS, ESI-MS/MS	Down-regulation of chloroplastic proteins involved in photosynthesis and carbon assimilation	60
	<i>Oryza sativa</i> L. var. Guara, <i>Hordeum vulgare</i> L. var. Baroness leaf apoplast	IPG, 2-DE, 2D BN-SDS-PAGE, ESI-MS/MS	Mn toxicity in barley involves apoplastic lesions mediated by PODs; lower Mn tolerance of young rice leaves in comparison to old ones	111
	<i>Pseudokirchneriella subcapitata</i> strain Hindák algal cells	IPG, 2-DE, LC-ESI-MS/MS	Enhanced abundance of RuBisCo activase and modulation of proteins involved in amino acids metabolism	81
Cr	<i>Miscanthus sinensis</i> cv. Kosung root	IPG, 2-DE, MALDI-TOF MS, MALDI-TOF/TOF MS	Accumulation of Cr-responsive proteins linked to heavy metal tolerance and senescence pathways	84
	<i>Zea mays</i> L. cv. Zheng 58 leaf	IPG, 2-DE, MALDI-TOF/TOF MS	Enhanced expression of proteins involved in ROS detoxification, defense responses, photosynthesis, and chloroplast organization	120
B	<i>Hordeum vulgare</i> cvs. GP, Cp, Sh, Cp x Sh DH leaf, root	iTRAQ peptide tagging, MS/MS	Enhanced abundance of enzymes involved in production of proteins with a strong chelating activity, phytosiderophores, which participate in iron up-take in B-tolerant barley	129
	<i>Lupinus albus</i> cv. Rio Maior root	IPG, 2-DE, LC-MS/MS	Under B deficiency proteins involved in energy, cell division, and protein metabolic processes were down-regulated	121

greatly improves comparisons between samples for which the DeCyder 2D Differential Analysis Software was developed.

Mass spectrometry (MS) techniques

In any proteome analysis, a number of proteins is separated. This approach typically comprises 2-DE coupled with MS techniques such as Matrix-Assisted Laser Desorption Ionization Time-of-Flight (MALDI-TOF) MS or nanoscale liquid chromatography (nano-LC) electrospray ionization (ESI)-MS/MS analysis (38). The first step is usually 2-DE, as previously described. The protein spots obtained are excised from the gel and submerged to digestion with an appropriate protease, most frequently trypsin, to obtain a mixture of peptides. In the following step, a peptide map called “peptide mass fingerprint” is constructed by MS in order to identify the protein. Alternatively, the partial amino acid sequence is determined by “*de novo* sequencing” and occasionally by a gas-phase protein sequencer. Then the database constructed by the genome analysis is retrieved and the gene corresponding to the protein of interest can be identified. The entry of the protein sequence database shows which proteins have been assigned particular functions. If the sequence of the analysed protein is identical or similar to other proteins from the database with a known function, the function of the target protein can be identified. This approach has been successfully applied in proteomic studies of different plant species exposed to Cd (25, 26, 61, 62, 66, 68-70, 80, 83, 112), Cu (24, 82, 115, 116), As (55, 56, 63, 117), Al (67, 71, 118, 119), Mn (59, 60, 111), Cr (81, 84, 120), and B (121) (Table 1).

Although classical proteomic approaches can identify several hundred heavy metal-responsive proteins from plant samples, they are still not apt for resolving and identifying certain components of protein complexes composed of multiple subunits. A number of second-generation gel-free MS-based proteomic technologies have been developed that can be implemented not only for protein identification, but also for determination of protein quantity. Techniques such as stable isotope labelling of amino acids in cell culture (SILAC), isotope tagging for relative and absolute protein quantitation (iTRAQ), and intensity-based absolute quantitation (iBAQ) can be applied for identifying almost all types of proteins, including highly acidic, basic, and hydrophobic proteins, as well as protein complexes (122-124). SILAC is a simple and straightforward approach for the *in vivo*

incorporation of a label into proteins for MS-based quantitative proteomics and it relies on metabolic incorporation of a given ‘light’ or ‘heavy’ form of the amino acid with substituted stable isotopic nuclei (e.g., deuterium, ^{13}C , ^{15}N) into proteins. Differential labelling of plant cells by two different versions of the labelled amino acids enables similar quantitation accuracy, precision, and reproducibility as conventional SILAC in animal cells (125). The iTRAQ method is based on isobaric tags, i.e. tags that have the same mass and are primarily designed for chemically labelling the N-terminus of peptides generated from protein digests that have been isolated from cells exposed to, for example, two different environmental conditions. The labelled samples are combined, fractionated by nano LC and analysed by tandem mass spectrometry. Database searches for peptide fragmentation data result in the identification of labelled peptides and corresponding proteins (126). iBAQ is a label-free proteome quantification method which is applicable for an accurate calculation of protein concentrations at the proteome level (127). iBAQ has proven to be suitable for proteomics-based identification of low-abundance signalling and regulatory protein complexes in native plant tissues (124). Currently, SILAC, iTRAQ, and iBAQ are considered to be among the most promising techniques for large-scale quantitative proteome analyses in plant biology (128, 124). However, despite numerous advantages of quantitative shotgun proteomics, there are only a few reports that employed these techniques for analyses of metal toxicity-induced proteins in plants (80, 129, 130). iTRAQ was used by Patterson et al. (129) for quantitative determination of proteins in two barley cultivars with different sensitivity to increased boron concentrations and the results obtained suggested that iTRAQ had a considerable potential for identifying and quantifying the expression of proteins in conjunction with bulked segregant analysis. Schneider et al. (130) applied the same technique to elucidate the role of vacuolar transporters in Cd-detoxification processes in the proteome of barley leaf tonoplast. Comparative proteomic approaches, including both gel-based (2D-DIGE) and gel-free (iTRAQ) techniques were applied by Alvarez et al. (80) to investigate the response of *Brassica juncea* roots to Cd treatment. They found that membrane and low abundant proteins were primarily identified by the iTRAQ method, while 2D-DIGE analysis identified many differentially expressed posttranslationally modified proteins, suggesting that

each method has its own advantages and the best approach would be to apply them both in a complementary manner. However, Finka and Golubinoﬀ (131) managed to demonstrate a robust protocol for the normalisation of iBAQ data for the purpose of analysing protein stoichiometries in complexes.

Current trends in proteomics of heavy metal toxicity in plants

In recent years, due to the advancement in techniques involving the identification and separation of proteins, substantial effort has been made in the clarification of molecular mechanisms underlying heavy metals toxicity in plants, but the importance of protein profiling has still not waned. The number of proteomic studies that have been carried out under different stress conditions in plants is relatively low, but for heavy metals-induced stress, proteomics studies are considerably even less represented (19, 38).

The greatest number of proteomic studies dealing with adverse effects of heavy metals on a proteome of different plant species has focused on the impact of Cd exposure. In a proteomic study on Cd-stressed poplar plants (25), it was found that this heavy metal had a deleterious effect on the expression of proteins involved in the primary carbon metabolism and from the oxidative stress response of stem tissue. Moreover, leaf and root proteome of poplar plants exposed to Cd²⁺ exhibited both an enhanced accumulation of stress-related proteins, like heat shock proteins, chaperones, foldases, proteases and pathogenesis-related (PR) proteins, while for the root tissue significant decrease in most proteins from the primary metabolism was characteristic (26). Several papers reported an activation and up-regulation of ROS scavengers, thus indicating a prompt antioxidative response to oxidative stress damage induced by Cd (61, 62, 69). Cd²⁺ also induced an enhanced accumulation of proteins such as molecular chaperons (61) and heat-shock proteins (69), which are protective proteins that contribute to protein stabilisation, adequate folding, assembly, and translocation under both optimal and stressed growth conditions (132). In many studies, up-regulation of proteins associated with Cd-chelating pathways was reported (25, 49, 66, 68, 70). In order to cope with an excess of heavy metals, plants have developed effective mechanism to synthesize chelators of low molecular weight in

order to minimize the binding of toxic metal ions to functionally important proteins (133).

Exposure to Cu mostly resulted in the up-regulation of photosynthesis, glycolysis, and pentose phosphate metabolism (82) as well as in enhanced accumulation of antioxidant and stress-related proteins (58). Moreover, Li et al. (115) examined the effects of increased Cu concentrations on *Elsholtzia splendens* and found that expression of several proteins with potential ion-binding functions as well as proteins engaged in vacuolar metal sequestrations were up-regulated.

Increased concentrations of arsenite or arsenate induced the over-expression of energy and metabolism related proteins (55, 63) and enhanced the expression of antioxidative enzymes (56), which indicated a higher energy demand and induced oxidative stress during As exposure. In addition, elevated As content led to a partial disruption of photosynthetic processes and degradation of several proteins involved in photosynthesis, RuBisCO in particular (117).

In studies of Al³⁺ toxicity it was found that cysteine synthase (CS) and glutathione (GSH) play an essential role in the Al adaptation of soybean (118) and rice (119). Moreover, it was also reported that Al-exposure induced activation of molecular chaperons in soybean (118), while an enhanced accumulation of several enzymes involved in ROS detoxification was found in rice roots (119). Proteomic analysis on tomato root tissue has shown that Al treatment induced enzymes associated with antioxidant activities and detoxification (67). Comparative analysis of root proteome of aluminium-tolerant and aluminium-sensitive soybean genotypes revealed an enhanced accumulation of enzymes that catalyse the synthesis of citrate, a key organic acid involved in Al³⁺ detoxification in a tolerant genotype, while proteins related to general stress response were induced in sensitive soybean (71).

Effects of increased manganese concentrations were investigated in legume crop cowpea (*Vigna unguiculata*) leaf apoplast (59) and symplast (60). Fecht-Christoffers et al. (59) revealed the particular role of peroxidases (PODs) in the expression of Mn²⁺ toxicity mediating H₂O₂ production/consumption and the oxidation of phenols in the leaf apoplast. The release of PODs in the apoplast was accompanied by the secretion of wound-induced proteins and PR enzymes, e.g. glucanase, chitinase, and thaumatin-like proteins, which appeared to be a late response to excess Mn. Furthermore, Führs et al. (60) reported the down-regulation of chloroplastic proteins involved in

photosynthesis and carbon assimilation in symplast of cowpea exposed to Mn-stress. A comparative proteomic study of Mn-response in rice and barley suggested that Mn toxicity in barley involved apoplastic lesions mediated by PODs, while the lower Mn tolerance of young rice leaves in comparison to old ones could have been related to Mn excess-induced displacement of Mg and Fe from essential metabolic functions (111).

Under chromate treatment of *Pseudokirchneriella subcapitata*, an enhanced abundance of RuBisCO activase and modulation of proteins involved in amino acids metabolism was reported (81). Moreover, Sharmin et al (84) found novel accumulation of Cr-responsive proteins linked to heavy metal tolerance and senescence pathways in roots of *Miscanthus sinensis*. In the study of induced proteomic changes in maize leaf upon Cr treatment the proteins identified were mainly involved in ROS detoxification and defence responses as well as in photosynthesis and chloroplast organization, which indicates that plants modify their metabolism by an altered expression of genes to adapt to Cr stress (120).

In studies dealing with the influence of boron (B) on plant proteome it was found that proteins involved in energy, cell division, and protein metabolic processes were down-regulated under B deficiency (121). In a comparative proteomic study of B-sensitive and B-tolerant barley exposed to increased B concentrations, Patterson et al (129) reported an enhanced abundance of enzymes phyto siderophores involved in production of proteins with a strong chelatin activity, which participate in iron uptake in B-tolerant barley.

Obstacles and future perspectives

What still remains challenging in this field is the implementation of proteomic studies in non-model plant species, i.e. plants for which the whole genome sequence is not fully available. For instance, plant scientists are interested in many different species as study objects, because some biological, as well as physiological processes of a particular research interest might not naturally be inherent to traditional model plants (134, 135). Nevertheless, high proteome coverage in terms of complete proteome profile lacks information even for model plants (136). Further, the identification of proteins in a particular biological sample does not automatically reveal the protein function. Moreover, proteomics studies have come up with a high number of unknown or hypothetical

proteins, up to 50 % of all protein sequences in databases (137). This seems to be even more complex in non-model species for which functional annotation of genes and proteins are at very beginnings (135). However, substantial progress has been made in this field and the results obtained from studies carried out in the last decade have greatly improved our understanding of plant response to heavy metal stress. In the future, proteomic studies of heavy metal stress in plants could pave the way for identifying novel protein candidates that could be used for improving stress tolerance. There is great potential in combining mass spectrometry with Nanostring technology used to quantify particular mRNA, as well as to improve current knowledge of translational and transcriptomics in order to get more information of how plant proteins respond to heavy metal stress. Moreover, new findings could be implemented in the engineering of transgenic plants that could be used for phytomining or phytoremediation of heavy metals from contaminated soils (17, 138, 139).

REFERENCES

- Gornall J, Betts R, Burke E, Clark R, Camp J, Willet K, Wiltshire A. Implications of climate change for agricultural productivity in the early twenty-first century. *Phil. Trans R Soc B* 2010;365:2973-89. doi: 10.1098/rstb.2010.0158
- Alexandrov V, Eitzinger J, Cajic V, Oberforster M. Potential impact of climate change on selected agricultural crops in north-eastern Austria. *Global Change Biol* 2002;8:372-89. doi: 10.1046/j.1354-1013.2002.00484.x
- Mittler R, Finka A, Goloubinoff P. How do plants feel the heat? *Trends Biochem Sci* 2012;37:118-25. doi: 10.1016/j.tibs.2011.11.007.
- Nriagu JO, Pacyna JM. Quantitative assessment of worldwide contamination of air, water and soils by trace metals. *Nature* 1988;333:134-9. doi: 10.1038/333134a0
- Woolson EA, Axley JH, Kearney PC. The chemistry and phytotoxicity of arsenic in soils: I. Contaminated field soils. *Soil Sci Soc Am J* 1971;35:938-43.
- Meharg AA, Rahman M. Arsenic contamination of Bangladesh paddy field soils: implications for rice contribution to arsenic consumption. *Environ Sci Technol* 2003;37:229-34. doi: 10.1021/es0259842
- Camargo JA, Ward JV. Short-term toxicity of sodium nitrate (NaNO₃) to non-target freshwater invertebrates. *Chemosphere* 1992;24:23-8. doi: 10.1016/0045-6535(92)90563-7
- Tilman D. Global environmental impacts of agricultural expansion: The need for sustainable and efficient practices. *Proc Natl Acad Sci USA* 1999;96:5995-6000. doi: 10.1073/pnas.96.11.5995
- Kassir LN, Lartiges B, Ouaini N. Effects of fertilizer industry emissions on local soil contamination: a case study of a phosphate plant on the east Mediterranean coast. *Environ*

- Technol 2012;33:873-85. doi: 10.1080/09593330.2011.601765
10. Nicholson FA, Smith SR, Alloway BJ, Carlton-Smith C, Chambers BJ. An inventory of heavy metals inputs to agricultural soils in England and Wales. *Sci Total Environ* 2003;311:205-19. doi: 10.1016/S0048-9697(03)00139-6
 11. Satarug S, Baker JR, Urbenjapol S, Haswell-Elkims M, Reilly PEB, Williams DJ. A global perspective on cadmium pollution and toxicity in non-occupationally exposed population. *Toxicol Lett* 2003;137:65-83. PMID: 12505433
 12. Di Toopi LS, Gabrielli R. Response to cadmium in higher plants. *Environ Exp Bot* 1999;41:105-30.
 13. Siedlecka A. Some aspects of interactions between heavy metals and plant mineral nutrients. *Acta Soc Bot Pol* 1995;64:265-72.
 14. Hänsch R, Mendel RR. Physiological functions of mineral micronutrients (Cu, Zn, Mn, Fe, Ni, Mo, B, Cl). *Curr Opin Plant Biol* 2009;12:259-66. doi: 10.1016/j.pbi.2009.05.006
 15. Saidi Y, Finka A, Goloubinoff P. Heat perception and signalling. Response to cadmium in plants: a tortuous path to thermotolerance. *New Phytol* 2011;190:556-65.
 16. Salt DE, Smith RD, Raskin I. Phytoremediation. *Annu Rev Plant Physiol Plant Mol Biol* 1998;49:643-68. PMID: 15012249
 17. Rugh CL, Senecoff JF, Meagher RB, Merkle SA. Development of transgenic yellow poplar for mercury phytoremediation. *Nature Biotechnol* 1998;16:925-8. doi: 10.1038/nbt1098-925
 18. Jung MC, Thorton I. Heavy metal contamination of soils and plants in the vicinity of a lead-zinc mine, Korea. *Appl Geochem* 1996;11:53-9. doi: 10.1016/0883-2927(95)00075-5
 19. Liao XY, Chen TB, Xie H, Liu YR. Soil As contamination and its risk assessment in areas near the industrial districts of Chenzhou City, Southern China. *Environ Int* 2005;31:791-8. PMID: 15979720
 20. Schützendübel A, Polle A. Plant responses to abiotic stresses: heavy metal – induced oxidative stress and protection by mycorrhization. *J Exp Bot* 2002;53:1351-65. doi: 10.1093/jxb/53.372.1351
 21. Clemens S. Toxic metal accumulation, responses to exposure and mechanisms of tolerance in plants. *Biochimie* 2006;88:1707-19. doi: 10.1016/j.biochi.2006.07.003
 22. Rogers S, Girolami M, Kolch W, Waters KM, Liu T, Thrall B, Wiley HS. Investigating the correspondence between transcriptomic and proteomic expression profiles using coupled cluster models. *Bioinformatics* 2008;24:2894-900. doi: 10.1093/bioinformatics/btn553
 23. Krämer U, Clemens S. Molecular biology of metal homeostasis and detoxification. In: Tamás M, Martinoia E, editors. *Topics in current genetics*. New York (NY): Springer Verlag; 2005. p. 216-71.
 24. Bona E, Marsano F, Cavaletto M, Berta G. Proteomic characterization of copper stress response in *Cannabis sativa* roots. *Proteomics* 2007;7:1121-30. doi: 10.1002/pmic.200600712
 25. Kieffer P, Dommes J, Hoffmann L, Hausman JF, Renaut J. Quantitative changes in protein expression of cadmium-exposed poplar plants. *Proteomics* 2008;8:2514-30. doi: 10.1002/pmic.200701110
 26. Kieffer P, Planchon S, Oufir M, Ziebel J, Dommes J, Hoffmann L. Combining proteomics and metabolite analyses to unravel cadmium stress-response in poplar leaves. *J Proteome Res* 2009;8:400-17. doi: 10.1021/pr800561r
 27. Giordano PM, Mortvedt J, Mays A. Effect of municipal wastes on crop yields and uptake of heavy metals. *J Environ Qual* 1975;4:394-9. doi: 10.2134/jeq1975.00472425000400030024x
 28. Islam E, Yang X, He Z, Mahmood Q. Assessing potential dietary toxicity of heavy metals in selected vegetables and food crops. *J Zhejiang Univ Sci B* 2007;8:1-13. doi: 10.1631/jzus.2007.B0001
 29. Fu J, Zhou Q, Liu J, Liu W, Wang T, Zhang Q, Jiang G. High levels of heavy metals in rice (*Oryza sativa* L.) from a typical E-waste recycling area in southeast China and its potential risk to human health. *Chemosphere* 2008;71:1269-75. doi: 10.1016/j.chemosphere.2007.11.065
 30. Hossain Z, Komatsu S. Contribution of proteomic studies towards understanding plant heavy metal stress response. *Front Plant Sci* 2013;3:310. doi: 10.3389/fpls.2012.00310
 31. Clemens S. Molecular mechanisms of plant metal tolerance and homeostasis. *Planta* 2001;212:475-86. PMID: 11525504
 32. Sharma SK, Goloubinoff P, Christen P. Heavy metal ions are potent inhibitors of protein folding. *Biochem Biophys Res Commun* 2008;372:341-5. doi: 10.1016/j.bbrc.2008.05.052
 33. The Arabidopsis Genome Initiative. Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* 2000;408:796-815. doi: 10.1038/35048692
 34. Rensing SA, Lang D, Zimmer AD, Terry A, Salamov A, Shapiro H, Nishiyama T, Perroud PF, Lindquist EA, Kamisugi Y, Tanahashi T, Sakakibara K, Fujita T, Oishi K, Shin-I T, Kuroki Y, Toyoda A, Suzuki Y, Hashimoto S, Yamaguchi K, Sugano S, Kohara Y, Fujiyama A, Anterola A, Aoki S, Ashton N, Barbazuk WB, Barker E, Bennetzen JL, Blankenship R, Cho SH, Dutcher SK, Estelle M, Fawcett JA, Gundlach H, Hanada K, Heyl A, Hicks KA, Hughes J, Lohr M, Mayer K, Melkozernov A, Murata T, Nelson DR, Pils B, Prigge M, Reiss B, Renner T, Rombauts S, Rushton PJ, Sanderfoot A, Schween G, Shiu SH, Stueber K, Theodoulou FL, Tu H, Van de Peer Y, Verrier PJ, Waters E, Wood A, Yang L, Cove D, Cuming AC, Hasebe M, Lucas S, Mishler BD, Reski R, Grigoriev IV, Quatrano RS, Boore JL. The *Physcomitrella* genome reveals evolutionary insights into the conquest of land by plants. *Science* 2008;319:64-9. doi: 10.1126/science.1150646
 35. International Rice Genome Sequencing Project. The map-based sequence of the rice genome. *Nature* 2005;436:793-800. doi: 10.1038/nature03895
 36. Phytozome [displayed 16 January 2014]. Available at <http://www.phytozome.com>
 37. Cambridge Healhtech Institute. -Omes and -omics glossary & taxonomy [displayed 16 January 2014]. Available at <http://www.genomicglossaries.com/content/omes.asp>
 38. Ahsan N, Renaut J, Komatsu S. Recent developments in the application of proteomics to the analysis of plant responses to heavy metals. *Proteomics* 2009;9:2602-21. doi: 10.1002/pmic.200800935
 39. Anderson L, Seilhamer J. A comparison of selected mRNA and protein abundances in human liver. *Electrophoresis* 1997;18:533-7. PMID: 9150937
 40. Mazzucotelli E, Mastrangelo AM, Crosatti C, Guerra D, Stanca AM, Cattivelli L. Abiotic stress response in plants: When post-transcriptional and post-translational regulations control transcription. *Plant Sci* 2008;174:420-31. doi: 10.1016/j.plantsci.2008.02.00

41. Hakeem KR, Chandna R, Ahmad P, Iqbal M, Ozturk M. Relevance of proteomic investigations in plant abiotic stress physiology. *OMICS* 2012;16:621-35. doi: 10.1089/omi.2012.0041
42. Cobon GS, Verrills N., Papakostopoulos P, Eastwood H, Linnane AW. The proteomics of ageing. *Biogerontology* 2002;3:133-6. doi: 10.1023/A:1015240304287
43. Hirano H, Islam N, Kawasaki H. Technical aspects of functional proteomics in plants. *Phytochemistry* 2004;65:1487-9. PMID: 15276446
44. Timperio AM, Egidi MG, Zolla L. Proteomics applied on plant abiotic stresses: Role of heat shock proteins (HSP). *J Proteomics* 2008;71:391-411. doi: 10.1016/j.jprot.2008.07.005
45. Swinbanks D. Government backs proteome proposal. *Nature* 1995;378:653. doi: 10.1038/378653a0
46. de Hoog CL, Mann M. Proteomics. *Annu Rev Genomics Hum Genet* 2004;5:267-93. doi: 10.1146/annurev.genom.4.070802.110305
47. Cuypers A, Vangronsveld J, Clijsters H. The redox status of plant cells (AsA and GSH) is sensitive to zinc imposed oxidative stress in roots and primary leaves of *Phaseolus vulgaris*. *Plant Physiol Biochem* 2001;39:657-64. doi: 10.1016/S0981-9428(01)01276-1
48. Aravind P, Prasad MNV. Zinc alleviates cadmium-induced oxidative stress in *Ceratophyllum demersum* L.: a free floating freshwater macrophyte. *Plant Physiol Biochem* 2003;41:391-7. doi: 10.1016/S0981-9428(03)00035-4
49. Horvat T, Vidaković-Cifrek Ž, Oreščanin V, Tkalec M, Pevallek- Kozlina B. Toxicity assessment of heavy metal mixtures by *Lemna minor* L. *Sci Total Environ* 2007;384:229-38. doi: 10.1016/j.scitotenv.2007.06.007
50. Gratão PL, Monteiro CC, Antunes AM, Peres LEP, Azevedo RA. Acquired tolerance of tomato (*Lycopersicon esculentum* cv. Micro-Tom) plants to cadmium induced stress. *Ann Appl Biol* 2008;153:321-33. doi: 10.1111/j.1744-7348.2008.00299.x
51. Tkalec M, Prebeg T, Roje V, Pevallek-Kozlina, Ljubešić N. Cadmium induced responses in duckweed *Lemna minor* L. *Acta Physiol Plant* 2008;30:881-90. doi: 10.1007/s11738-008-0194-y
52. Hassan MJ, Zhang G, Wu F, Wie K, Chen Z. Zinc alleviates growth inhibition and oxidative stress caused by cadmium in rice. *J Plant Nutr Soil Sci* 2005;168:255-61. doi: 10.1002/jpln.200420403
53. Cvjetko P, Tolić S, Šikić S, Balen B, Tkalec M, Vidaković-Cifrek Ž, Pavlica M. Effect of copper on the toxicity and genotoxicity of cadmium in duckweed (*Lemna minor* L.) *Arh Hig Rada Toksikol* 2010;61:287-96. doi: 10.2478/10004-1254-61-2010-2059
54. Balen B, Tkalec M, Šikić S, Tolić S, Cvjetko P, Pavlica M, Vidaković-Cifrek Ž. Biochemical responses of *Lemna minor* experimentally exposed to cadmium and zinc. *Ecotoxicology* 2011;20:815-26. doi: 10.1007/s10646-011-0633-1
55. Ahsan N, Lee DG, Kim KH, Alam I, Lee SH, Lee KW, Lee H, Lee BH. Analysis of arsenic stress-induced differentially expressed proteins in rice leaves by two-dimensional gel electrophoresis coupled with mass spectrometry. *Chemosphere* 2010;78:224-31. doi: 10.1016/j.chemosphere.2009.11.004
56. Ahsan N, Lee DG, Alam I, Kim PJ, Lee JJ, Ahn YO, Kwak SS, Lee I-J, Bahk JD, Kang KY, Renaut J, Komatsu S, Lee BH. Comparative proteomic study of arsenic-induced differentially expressed proteins in rice roots reveals glutathione plays a central role during As stress. *Proteomics* 2008;8:3561-76. doi: 10.1002/pmic.200701189
57. Ahsan N, Lee SH, Lee DG, Lee H, Lee SW, Bahk JD, Lee BH. Physiological and protein profiles alternation of germinating rice seedlings exposed to acute cadmium toxicity. *C R Biol* 2007;330:735-46. PMID: 17905393
58. Ahsan N, Lee DG, Lee SH, Kang KY, Lee JJ, Kim PJ, Yoon HS, Kim JS, Lee BH. Excess copper induced physiological and proteomic changes in germinating rice seeds. *Chemosphere* 2007;67:1182-93. doi: 10.1016/j.chemosphere.2006.10.075
59. Fecht-Christoffers MM, Braun HP, Lemaitre-Guillier C, Van Dorsselaer A, Horst WJ. Effect of manganese toxicity on the proteome of the leaf apoplast in cowpea. *Plant Physiol* 2003;133:1935-46. doi: 10.1104/pp.103.029215
60. Führs H, Hartwig M, Molina LE, Heintz D, Van Dorsselaer A, Braun HP, Horst WJ. Early manganese-toxicity response in *Vigna unguiculata* L. – a proteomic and transcriptomic study. *Proteomics* 2008;8:149-59. doi: 10.1002/pmic.200700478
61. Hossain Z, Hajika M, Komatsu S. Comparative proteome analysis of high and low cadmium accumulating soybeans under cadmium stress. *Amino Acids* 2012;43:2393-416. doi: 10.1007/s00726-012-1319-6
62. Lee K, Bae DW, Kim SH, Han HJ, Liu X, Park HC, Lim CO, Lee SY, Chung WS. Comparative proteomic analysis of the short-term responses of rice roots and leaves to cadmium. *J Plant Physiol* 2010;167:161-8. doi: 10.1016/j.jplph.2009.09.006
63. Pandey S, Rai R, Rai LC. Proteomics combines morphological, physiological and biochemical attributes to unravel the survival strategy of *Anabaena* sp. PCC7120 under arsenic stress. *J Proteomics* 2012;75:921-37. doi: 10.1016/j.jprot.2011.10.011
64. Requejo R, Tena M. Maize response to acute arsenic toxicity as revealed by proteome analysis of plant shoots. *Proteomics* 2006;6(Suppl 1):S156-62. PMID: 16534746
65. Sarry JE, Kuhn L, Ducruix C, Lafaye A, Junot C, Hugouvieux V, Jourdain A, Bastien O, Fievet JB, Vailhen D, Amekraz B, Moulin C, Ezan E, Garin J, Bourguignon J. The early responses of *Arabidopsis thaliana* cells to cadmium exposure explored by protein and metabolite profiling analyses. *Proteomics* 2006;6:2180-98. doi: 10.1002/pmic.200500543
66. Semane B, Dupae J, Cuypers A, Noben JP, Tuomainen M, Tervahauta A, Kärenlampi S, Van Belleghem F, Smeets K, Vangronsveld J. Leaf proteome responses of *Arabidopsis thaliana* exposed to mild cadmium stress. *J Plant Physiol* 2010;167:247-54. doi: 10.1016/j.jplph.2009.09.015
67. Zhou S, Sauvé R, Thannhauser TW. Proteome changes induced by aluminium stress in tomato roots. *J Exp Bot* 2009;60:1849-57. doi: 10.1093/jxb/erp065
68. Ahsan N, Nakamura T, Komatsu S. Differential responses of microsomal proteins and metabolites in two contrasting cadmium (Cd) accumulating soybean cultivars under Cd stress. *Amino Acids* 2012;42:317-27. doi: 10.1007/s00726-010-0809-7
69. Hossain Z, Makino T, Komatsu S. Proteomic study of β -aminobutyric acid-mediated cadmium stress alleviation in soybean. *J Proteomics* 2012;75:4151-64. doi: 10.1016/j.jprot.2012.05.037

70. Hradilová J, Rehulka P, Rehulková H, Vrbová M, Griga M, Brzobohatý B. Comparative analysis of proteomic changes in contrasting flax cultivars upon cadmium exposure. *Electrophoresis* 2010;31:421-31. doi: 10.1002/elps.200900477
71. Duressa D, Soliman K, Taylor R, Senwo Z. Proteomic analysis of soybean roots under aluminum stress international. *Int J Plant Genomics* 2011;2011:2825-31. doi: 10.1155/2011/282531
72. Cho K, Torres NL, Subramanyam S, Deepak SA, Sardesai N, Han O, Williams CE, Ishii H, Iwahashi H, Rakwal R. Protein extraction/solubilization protocol for monocot and dicot plant gel-based proteomics. *J Plant Biol* 2006;49:413-20. doi: 10.1007/BF03031120
73. Rose JKC, Bashir S, Giovannoni JJ, Jahn MM, Saravanan RS. Tackling the plant proteome: practical approaches, hurdles and experimental tools. *Plant J* 2004;39:715-33. doi: 10.1111/j.1365-3113X.2004.02182.x
74. Jellouli N, Salem AB, Ghorbel A, Jouira HB. Evaluation of protein extraction methods for *Vitis vinifera* leaf and root. *J Integr Plant Biol* 2010;52:933-40. doi: 10.1111/j.1744-7909.2010.00973.x
75. Isaacson T, Damasceno CM, Saravanan RS, He Y, Catalá C, Saladié M, Rose JK. Sample extraction techniques for enhanced proteomic analysis of plant tissues. *Nat Protoc* 2006;1:769-74. PMID: 17406306
76. Nouri MZ, Komatsu S. Comparative analysis of soybean plasma membrane proteins under osmotic stress using gel-based and LC MS/MS-based proteomics approaches. *Proteomics* 2010;10:1930-45. doi: 10.1002/pmic.200900632
77. Pavoković D, Križnik B, Krsnik-Rasol M. Evaluation of protein extraction methods for proteomic analysis of non-model recalcitrant plant tissues. *Croat Chem Acta* 2012;85:177-83. doi: 10.5562/cca1804
78. Komatsu S, Ahsan N. Soybean proteomics and its application to functional analysis. *J Proteomics* 2009;72:325-36. doi: 10.1016/j.jprot.2008.10.001
79. Sarma AD, Oehrle NW, Emerich DW. Plant protein isolation and stabilization for enhanced resolution of two-dimensional polyacrylamide gel electrophoresis. *Anal Biochem* 2008;379:192-5. doi: 10.1016/j.ab.2008.04.047
80. Alvarez S, Berla BM, Sheffield J, Cahoon RE, Jez JM, Hicks LM. Comprehensive analysis of the *Brassica juncea* root proteome in response to cadmium exposure by complementary proteomic approaches. *Proteomics* 2009;9:2419-31. doi: 10.1002/pmic.200800478
81. Vannini C, Marsoni M, Domingo G, Antognoni F, Biondi S, Bracale M. Proteomic analysis of chromate-induced modifications in *Pseudokirchneriella subcapitata*. *Chemosphere* 2009;76:1372-9. doi: 10.1016/j.chemosphere.2009.06.022
82. Ritter A, Ubertini M, Romac S, Gaillard F, Delage L, Mann A, Cock JM, Tonon T, Correa JA, Potin P. Copper stress proteomics highlights local adaptation of two strains of the model brown alga *Ectocarpus siliculosus*. *Proteomics* 2010;10:2074-88. doi: 10.1002/pmic.200900004
83. Rodríguez-Celma J, Rellán-Alvarez R, Abadía A, Abadía J, López-Millán AF. Changes induced by two levels of cadmium toxicity in the 2-DE protein profile of tomato roots. *J Proteomics* 2010;73:1694-706. doi: 10.1016/j.jprot.2010.05.001
84. Sharmin SA, Alam I, Kim KH, Kim YG, Kim PJ, Bahk JD, Lee BH. Chromium-induced physiological and proteomic alterations in roots of *Miscanthus sinensis*. *Plant Sci* 2012;187:113-26. doi: 10.1016/j.plantsci.2012.02.002
85. Komatsu S, Wada T, Abaléa Y, Nouri MZ, Nanjo Y, Nakayama N, Shimamura S, Yamamoto R, Nakamura T, Furukawa K. Analysis of plasma membrane proteome in soybean and application to flooding stress response. *J Proteome Res* 2009;8:4487-99. doi: 10.1021/pr9002883
86. Agrawal GK, Bourguignon J, Rolland N, Ephritikhine G, Ferro M, Jaquinod M, Alexiou KG, Chardot T, Chakraborty N, Jolivet P, Doonan JH, Rakwal R. Plant organelle proteomics: Collaborating for optimal cell function. *Mass Spectrom Rev* 2011;30:1772-853. doi: 10.1002/mas.20301
87. Eubel H, Braun HP and A Millar H. Blue-native PAGE in plants: a tool in analysis of protein-protein interactions. *Plant Methods* 2005;1:11. PMID: 16287510
88. Xi J, Wang X, Li S, Zhou X, Yue L, Fan J, Hao D. Polyethylene glycol fractionation improved detection of low-abundant proteins by two dimensional electrophoresis analysis of plant proteome. *Phytochemistry* 2006;67:2341-8. doi: 10.1016/j.phytochem.2006.08.005
89. Baracat-Pereira MC, de Oliveira Barbosa M, Magalhães Júnior MJ, Carrijo LC, Games PD, Almeida HO, Sena Netto JF, Rodrigues Pereira M, de Barros EG. Separomics applied to the proteomics and peptidomics of low-abundance proteins: Choice of methods and challenges – A review. *Gen Mol Biol* 2012;35:283-91. doi: 10.1590/S1415-47572012000200009
90. Cho J-H, Hwang H, Cho M-H, Kwon Y-K, Jeon J-S, Bhoo SH, Hahn T-R. The effect of DTT in protein preparations for proteomic analysis: removal of a highly abundant plant enzyme, ribulose biphosphate carboxylase/oxygenase. *J Plant Biol* 2008;51:297-301. doi: 10.1007/BF03036130
91. Krishnan HB, Natarajan SS. A rapid method for depletion of Rubisco from soybean (*Glycine max*) leaf for proteomic analysis of lower abundance proteins. *Phytochemistry* 2009;70:1958-64. doi: 10.1016/j.phytochem.2009.08.020
92. Natarajan SS, Krishnan HB, Lakshman S, Garrett WM. An efficient extraction method to enhance analysis of low abundant proteins from soybean seed. *Anal Biochem* 2009;394:259-68. doi: 10.1016/j.ab.2009.07.048
93. Vertommen A, Møller AL, Cordewenerd JHG, Swennen R, Panis B, Finnie C, America AHP, Carpentiera SC. A workflow for peptide-based proteomics in a poorly sequenced plant: A case study on the plasma membrane proteome of banana. *J Proteomics* 2011;74:1218-29. doi: 10.1016/j.jprot.2011.02.008
94. Azarkan M, Huet J, Baeyens-Volant D, Looze Y, Vandenbussche G. Affinity chromatography: A useful tool in proteomics studies. *J Chromatogr B Analyt Technol Biomed Life Sci* 2007;849:81-90. PMID: 17113368
95. Fang X, Zhang W. Affinity separation and enrichment methods in proteomic analysis. *J Proteomics* 2008;71:284-303. doi: 10.1016/j.jprot.2008.06.011
96. Fröhlich A, Lindermayr C. Deep insights into the plant proteome by pretreatment with combinatorial hexapeptide ligand libraries. *J Proteomics* 2011;74:1182-9. doi: 10.1016/j.jprot.2011.02.019
97. Fröhlich A, Gaupels F, Sarioglu H, Holzmeister C, Spannagl M, Durner J, Lindermayr C. Looking deep inside: Detection of low-abundance proteins in leaf extracts of Arabidopsis

- and phloem exudates of pumpkin. *Plant Physiol* 2012;159:902-14. doi: 10.1104/pp.112.198077
98. Gallagher SR. One-dimensional SDS gel electrophoresis of proteins. *Curr Protoc Mol Biol* 2006;75:10.2.1–10.2A.37. doi: 10.1002/0471142727.mb1002as75
 99. Klose J. From 2-D electrophoresis to proteomics. *Electrophoresis* 2009;30:S142-9. doi: 10.1002/elps.200900118
 100. Peharec Štefanić P, Šikić S, Cvjetko P, Balen B. Cadmium and zinc induced similar changes in protein and glycoprotein patterns in tobacco (*Nicotiana tabacum* L.) seedlings and plants. *Arh Hig Rada Toksikol* 2012;63:321-35. doi: 10.2478/10004-1254-63-2012-2173
 101. Gallagher SR. One-dimensional SDS gel electrophoresis of proteins. *Curr Protoc Cell Biol* 2007;37:6.1.1–6.1.38. doi: 10.1002/0471143030.cb0601s37
 102. O'Farrell PH. High resolution two-dimensional electrophoresis. *J Biol Chem* 1975;250:4007-21. PMID: 236308
 103. Friedman D, Hoving S, Westmeier R. Isoelectric focusing and two-dimensional gel electrophoresis. *Methods Enzymol* 2009;463:515-40. doi: 10.1016/S0076-6879(09)63030-5
 104. Bjellqvist B, Ek K, Righetti GP, Gianazza E, Görg A, Westermeier R, Postel W. Isoelectric focusing in immobilized pH gradients: principle, methodology and some applications. *J Biochem Biophys Methods* 1982;6:317-39. PMID: 7142660
 105. Ünlü M, Morgan ME, Minden JS. Difference gel electrophoresis: a single gel method for detecting changes in protein extracts. *Electrophoresis* 1997;18:2071-7. doi: 10.1002/elps.1150181133
 106. Tonge R, Shaw J, Middleton B, Rowlinson R, Rayner S, Young J, Posgnan F, Hawkins E, Currie I, Davison M. Validation and development of fluorescence two-dimensional differential gel electrophoresis proteomics technology. *Proteomics* 2001;1:377-96. PMID: 11680884
 107. Schägger H, von Jagow G. Blue native electrophoresis for isolation of membrane protein complexes in enzymatically active form. *Anal Biochem* 1991;199:223-31. doi: 10.1016/0003-2697(91)90094-a
 108. Schägger H, Cramer W A, von Jagow G. Analysis of molecular masses and oligomeric states of protein complexes by blue native electrophoresis and isolation of membrane protein complexes by two-dimensional native electrophoresis. *Anal Biochem* 1994;217:220-30. doi: 10.1006/abio.1994.1112
 109. Reisinger V, Eichacker LA. Analysis of membrane protein complexes by Blue Native PAGE. *Proteomics* 2006;6(Suppl 2):6-15. PMID: 17031799
 110. Nijtmans LG, Henderson NS, Holt IJ. Blue native electrophoresis to study mitochondrial and other protein complexes. *Methods* 2002;26:327-34. PMID: 12054923
 111. Führs H, Behrens C, Gallien S, Heintz D, Van Dorsselaer A, Braun HP, Horst WJ. Physiological and proteomic characterization of manganese sensitivity and tolerance in rice (*Oryza sativa*) in comparison with barley (*Hordeum vulgare*). *Ann Bot* 2010;105:1129-40. doi: 10.1093/aob/mcq046
 112. Fagioni M, D'Amici GM, Timperio AM, Zolla L. Proteomic analysis of multiprotein complexes in the thylakoid membrane upon cadmium treatment. *J Proteome Res* 2009;8:310-26. doi: 10.1021/pr800507x
 113. GelAnalyzer [displayed 20 January 2014]. Available at <http://www.gelanalyzer.com>
 114. GelScape [displayed 20 January 2014]. Available at <http://www.gelscape.ualberta.ca>
 115. Li F, Shi J, Shen C, Chen G, Hu S, Chen Y. Proteomic characterization of copper stress response in *Elsholtzia splendens* roots and leaves. *Plant Mol Biol* 2009;71:251-63. doi: 10.1007/s11103-009-9521-y
 116. Zhang H, Lian C, Shen Z. Proteomic identification of small, copper-responsive proteins in germinating embryos of *Oryza sativa*. *Ann Bot* 2009;103:923-30. doi: 10.1093/aob/mcp012
 117. Duquesnoy I, Goupil P, Nadaud I, Branlard G, Piquet-Pissaloux A, Ledoigt G. Identification of *Agrostis tenuis* leaf proteins in response to As(V) and As(III) induced stress using a proteomics approach. *Plant Sci* 2009;176:206-13. doi: 10.1016/j.plantsci.2008.10.008
 118. Zhen Y, Qi JL, Wang SS, Su J, Xu GH, Zhang MS, Miao L, Peng XX, Tian D, Yang YH. Comparative proteome analysis of differentially expressed proteins induced by Al toxicity in soybean. *Physiol Plant* 2007;131:542-54. doi: 10.1111/j.1399-3054.2007.00979.x
 119. Yang Q, Wang Y, Zhang J, Shi W, Qian C, Peng X. Identification of aluminium-responsive proteins in rice roots by a proteomic approach: cysteine synthase as a key player in Al response. *Proteomics* 2007;7:737-49. PMID: 17295357
 120. Wang R, Gao F, Guo BQ, Huang JC, Wang L, Zhou YJ. Short-term chromium-stress-induced alterations in the maize leaf proteome. *Int J Mol Sci* 2013;14:11125-44. doi: 10.3390/ijms140611125
 121. Alves M, Moes S, Jenö P, Pinheiro C, Passarinho J, Ricardo CP. The analysis of *Lupinus albus* root proteome revealed cytoskeleton altered features due to long-term boron deficiency. *J Proteomics* 2011;74:1351-63. doi: 10.1016/j.jprot.2011.03.002
 122. Ong SE, Blagoev B, Kratchmarova I, Kristensen DB, Steen H, Pandey A, Mann M. Stable isotope labelling by amino acids in cell culture, SILAC, as a simple and accurate approach to expression proteomics. *Mol Cell Proteomics* 2002;1:376-86. doi: 10.1074/mcp.M200025-MCP200
 123. Ross PL, Huang YN, Marchese JN, Williamson B, Parker K, Hattan S, Khainovski N, Pillai S, Dey S, Daniels S, Purkayastha S, Juhasz P, Martin S, Bartlett-Jones M, He F, Jacobson A, Pappin DJ. Multiplexed protein quantitation in *Saccharomyces cerevisiae* using amine-reactive isobaric tagging reagents. *Mol Cell Proteomics* 2004;3:1154-69. PMID: 15385600
 124. Smaczniak C, Li N, Boeren S, America T, van Dongen W, Goerdal SS, de Vries S, Angenent GC, Kaufmann K. Proteomics-based identification of low-abundance signalling and regulatory protein complexes in native plant tissues. *Nature Protocols* 2012;7:2144-58. doi: 10.1038/nprot.2012.129
 125. Schütz W, Hausmann N, Krug K, Hampp R, Macek B. Extending SILAC to proteomics of plant cell lines. *Plant Cell* 2011;23:1701-5. doi: 10.1105/tpc.110.082016
 126. Lucker J, Laszczak M, Smith D, Lund ST. Generation of a predicted protein database from EST data and application to iTRAQ analyses in grape (*Vitis vinifera* cv Cabernet Sauvignon) berries at ripening initiation. *BMC Genomics* 2009;10:50. doi: 10.1186/1471-2164-10-50
 127. Arike L, Valgepea K, Peil L, Nahku R, Adamberg K, Vilu R. Comparison and applications of label-free absolute proteome

- quantification methods on *Escherichia coli*. J Proteomics 2012;75:5437-48. doi: 10.1016/j.jprot.2012.06.020
128. Thelen JJ, Peck SC. Quantitative proteomics in plants: choices in abundance. Plant Cell 2007;19:3339-46. doi: 10.1105/tpc.107.053991
129. Patterson J, Ford K, Cassin A, Natera S, Bacic A. Increased abundance of proteins involved in phytosiderophore production in boron-tolerant barley. Plant Physiol 2007;144:1612-31. doi: 10.1104/pp.107.096388
130. Schneider T, Schellenberg M, Meyer S, Keller F, Gehrig P, Riedel K, Lee Y, Eberl L, Martinoia E. Quantitative detection of changes in the leaf-mesophyll tonoplast proteome in dependency of a cadmium exposure of barley (*Hordeum vulgare* L.) plants. Proteomics 2009;9:2668-77. doi: 10.1002/pmic.200800806
131. Finka A, Goloubinoff P. Proteomic data from human cell cultures refine mechanisms of chaperone-mediated protein homeostasis. Cell Stress Chaperones 2013;18:591-605. doi: 10.1007/s12192-013-0413-3
132. Wang W, Vinocur B, Shoseyov O, Altman A. Role of plant heat-shock proteins and molecular chaperones in the abiotic stress response. Trends Plant Sci 2004;9:244-52. doi: 10.1007/s12192-013-0413-3
133. Verbruggen N, Hermans C, Schat H. Mechanisms to cope with arsenic or cadmium excess in plants. Curr Opin Plant Biol 2009;12:364-72. doi: 10.1016/j.pbi.2009.05.001
134. Remmerie N, De Vijlder T, Laukens K, Dang TH., Lemiére F, Mertens I. Next generation functional proteomics in non-model plants: a survey on techniques and applications for the analysis of protein complexes and post-translational modifications. Phytochemistry 2011;72:1192-218. doi: 10.1016/j.phytochem.2011.01.003
135. Champagne A, Boutry M. Proteomics of nonmodel plant species. Proteomics 2013;13:663-73. doi: 10.1002/pmic.201200312
136. Vanderschuren H, Lentz E, Zainuddin I, Gruijssem W. Proteomics of model and crop plant species: Status, current limitations and strategic advances for crop improvement. J Prot 2013;93:5-19 doi: 10.1016/j.jprot.2013.05.036
137. Galperin MY, Koonin E V. From complete genome sequence to 'complete' understanding? Trends Biotechnol 2010;28:398-406. doi: 10.1016/j.tibtech.2010.05.006
138. Eapen S, D'Souza SF. Prospects of genetic engineering of plants for phytoremediation of toxic metals. Biotechnol Adv 2005;23:97-114. PMID: 15694122
139. Aken BV. Transgenic plants for phytoremediation: helping nature to clean up environmental pollution. Trends Biotechnol 2008;26:225-7. doi: 10.1016/j.tibtech.2008.02.001

Sažetak**Proteomika u istraživanjima toksičnosti teških metala u biljaka**

Biljke su neprestano izložene različitim čimbenicima abiotičkog i biotičkog stresa koji nepovoljno utječu na njihovu produktivnost. Teški metali kao česti zagađivači okoliša vrlo nepovoljno djeluju na sva živa bića, uključujući biljke, životinje i ljude. Poznato je da teški metali mogu mijenjati ekspresiju biljnih proteina. Proteine ubrajamo u biološki vrlo važne makromolekule čija je aktivnost u stanici izravno ovisna o posttranslacijskim modifikacijama, koje nije moguće pratiti na razini genoma. Stoga je nužno provoditi proteomska istraživanja kako bi se razotkrila prisutnost i uloga proteina u različitim vrstama okolišnog stresa. U ovom radu sažete su različite tehnike i metode istraživanja učinaka teških metala na biljni proteom, uključujući i sažet osvrt na složene mehanizme odgovora biljke na stres izazvan teškim metalima.

KLJUČNE RIJEČI: *dvodimenzionalna elektroforeza; masena spektrometrija; proteinski biomarkeri; proteom; teški metali*

CORRESPONDING AUTHOR:

Biljana Balen
Associate Professor
Department of Molecular Biology
Faculty of Science, University of Zagreb
Horvatovac 102a, Zagreb, Croatia
E-mail: bbalen@zg.biol.pmf.hr