



Altering distances, some generalizations of Meir - Keeler theorems and applications

Alina-Mihaela Patriciu and Valeriu Popa

Abstract

The purpose of this paper is to prove a general theorem of Meir - Keeler type using the notion of altering distance for occasionally weakly compatible mappings satisfying an implicit relation.

As application, a problem of Meir - Keeler type satisfying a condition of integral type becomes a special case of a problem of Meir - Keeler type with an altering distance.

1 Introduction

Let f and g be self mappings of a metric space (X, d) . We say that $x \in X$ is a coincidence point of f and g if $fx = gx$.

We denote by $C(f, g)$ the set of all coincidence points of f and g .

A point w is a point of coincidence of f and g if there exists an $x \in X$ such that $w = fx = gx$.

Jungck [10] defined f and g to be compatible if $\lim d(fgx_n, gfx_n) = 0$ whenever $\{x_n\}$ is a sequence in X such that $\lim fx_n = \lim gx_n = t$ for some $t \in X$.

In 1994, Pant introduced the notion of pointwise R - weakly commuting mappings. It is proved in [25] that pointwise R - weakly commuting is equivalent to commutativity in coincidence points.

Key Words: Fixed point, Meir - Keeler type, altering distance, integral type, occasionally weakly compatible.

2010 Mathematics Subject Classification: Primary 54H25; Secondary 47H10.

Received: January 2013

Accepted: February 2013

Definition 1.1 ([11]). *Two self mappings of a metric space (X, d) are said to be weakly compatible if $fgu = gfu$ for each $u \in C(f, g)$.*

Al - Thagafi and Naseer Shahzad [3] introduced the notion of occasionally weakly compatible mappings.

Definition 1.2. *Two self mappings f and g of a metric space (X, d) are said to be occasionally weakly compatible (owc) mappings if there exists a point $x \in X$ which is a coincidence point of f and g at which f and g commute.*

Remark 1.1. *Two weakly compatible mappings having coincidence points are owc. The converse is not true, as shown in the Example of [3].*

Some fixed point theorems for occasionally weakly compatible mappings are proved in [2], [12], [34] and in other papers.

Lemma 1.1 ([12]). *Let X be a nonempty set and let f and g be owc self maps of X . If f and g have a unique point of coincidence $w = fx = gx$, then w is the unique common fixed point of f and g .*

2 Preliminaries

In 1969, Meir and Keeler [19], established a fixed point theorem for self mappings of a metric space (X, d) satisfying the following condition:

for each $\varepsilon > 0$, there exists a $\delta > 0$ such that

$$\varepsilon < d(x, y) < \varepsilon + \delta \text{ implies } d(fx, fy) < \varepsilon. \quad (2.1)$$

There exists a vast literature which generalizes the result of Meir-Keeler. In [18], Mati and Pal proved a fixed point theorem for a self mapping of a metric space (X, d) satisfying the following condition which is a generalization of (2.1):

for $\varepsilon > 0$, there exists $\delta > 0$ such that

$$\varepsilon < \max\{d(x, y), d(x, fx), d(y, fy)\} < \varepsilon + \delta \text{ implies } d(fx, fy) < \varepsilon. \quad (2.2)$$

In [29] and [35], Park-Rhoades, respectively Rao-Rao extend this result for two self mappings f and g of a metric space (X, d) satisfying the following condition:

$$\varepsilon < \max \left\{ d(fx, fy), d(fx, gx), d(fy, gy), \frac{1}{2} [d(fx, gy) + d(fy, gx)] \right\} < \varepsilon + \delta \quad (2.3)$$

implies $d(gx, gy) < \varepsilon$.

In 1986, Jungck [10] and Pant [21] extend these results for four mappings. It is known by Jungck [10], Pant [22], [24], [25] and other papers that, in the case of theorems for four mappings A, B, S and $T : (X, d) \rightarrow (X, d)$, a condition of Meir-Keeler type does not assure the existence of a fixed point. The following theorem is stated in [9].

Theorem 2.1. *Let (A, S) and (B, T) be compatible pairs of self mappings of a complete metric space (X, d) such that*

- (1) $AX \subset TX$ and $BX \subset SX$,
- (2) given an $\varepsilon > 0$ there exists a $\delta > 0$ such that for all $x, y \in X$,

$$\varepsilon \leq M(x, y) < \varepsilon + \delta \text{ implies } d(Ax, By) < \varepsilon,$$

where

$$M(x, y) = \max \left\{ d(Sx, Ty), d(Sx, Ax), d(Ty, By), \frac{1}{2} [d(Sx, By) + d(Ty, Ax)] \right\}$$

and

(3)

$$d(Ax, By) < k(d(Sx, Ty) + d(Sx, Ax) + d(Ty, By) + d(Sx, By) + d(Ty, Ax))$$

for all $x, y \in X$, where $k \in [0, \frac{1}{3})$.

If one of mappings A, B, S and T is continuous then A, B, S and T have a unique common fixed point.

Some similar theorems are proved in [8], [27], [28] and in other papers. Recently, Theorem 2.1 was improved and extended for weakly compatible pairs in [4].

Theorem 2.2. *Let (A, S) and (B, T) be weakly compatible pairs of self mappings of a complete metric space (X, d) such that the following conditions hold:*

- 1) $AX \subset TX$ and $BX \subset SX$,
- 2) one of AX, BX, SX and TX is closed,

3) for each $\varepsilon > 0$, there exists a $\delta > 0$ such that

$$\varepsilon < M(x, y) < \varepsilon + \delta \text{ implies } d(Ax, By) \leq \varepsilon,$$

4) $x, y \in X$, $M(x, y) > 0$ implies $d(Ax, By) < M(x, y)$,

5)

$$\begin{aligned} d(Ax, By) &< k [d(Sx, Ty) + d(Sx, Ax) + d(Ty, By) \\ &\quad + d(Sx, By) + d(Ty, Ax)] \end{aligned}$$

for all $x, y \in X$, where $k \in [0, \frac{1}{3})$.

Then A , B , S and T have a unique common fixed point.

Other generalizations of Theorem 2.1 are proved in [5].

Theorem 2.3. Let (A, S) and (B, T) be weakly compatible pairs of self mappings of a complete metric space (X, d) such that

1) $AX \subset TX$ and $BX \subset SX$,

2) one of AX , BX , SX and TX is closed,

3) given an $\varepsilon > 0$, there exists a $\delta > 0$ such that for all x, y in X

$$\varepsilon \leq M(x, y) < \varepsilon + \delta \text{ implies } d(Ax, By) < \varepsilon,$$

4)

$$\begin{aligned} d(Ax, By) &\leq k \max \{ d(Sx, Ty), d(Sx, Ax), d(Ty, By), \\ &\quad d(Sx, By), d(Ty, Ax) \} \end{aligned}$$

for all x, y in X , where $k \in [0, 1)$.

Then A , B , S and T have a unique common fixed point.

Remark 2.1. Because if (X, d) is complete and AX , BX , SX , TX are closed, then AX , BX , SX , TX are complete subspaces of X , in Theorems 2.2, 2.3 the conditions that X is complete and AX , BX , SX , TX are closed should be replaced by the statement that one of AX , BX , SX , TX is a complete subspace of X .

In [6], Branciari established the following result.

Theorem 2.4. *Let (X, d) be a complete metric space, $c \in (0, 1)$ and $f : X \rightarrow X$ such that*

$$\int_0^{d(fx, fy)} h(t)dt \leq c \int_0^{d(x, y)} h(t)dt, \quad (2.4)$$

whenever $h : [0, \infty) \rightarrow [0, \infty)$ is a Lebesgue measurable mapping which is summable (i.e. with a finite integral) on each compact subset of $[0, \infty)$ such that for $\varepsilon > 0$, $\int_0^\varepsilon h(t)dt > 0$. Then, f has a unique fixed point z such that, for each $x \in X$, $\lim f x_n = z$.

Theorem 2.4 is extended to compatible, weakly compatible, occasional weakly compatible in [1], [15], [16], [20], [34] and in other papers.

Quite recently, Gairola and Rawat [7] proved a fixed point theorem for two pairs of maps satisfying a new contractive condition of integral type, using the concept of occasionally weakly compatible mappings, which generalize Theorem 2.1.

Theorem 2.5. *Let A, B, S and T be self mappings of a metric space (X, d) satisfying the following conditions:*

- 1) $AX \subset TX$ and $BX \subset SX$,
- 2) *given $\varepsilon > 0$ there exists $\delta > 0$ such that for all x, y in X ,*

$$\int_0^{M(x, y)} h(t)dt < \varepsilon + \delta \text{ implies } \int_0^{d(Ax, By)} h(t)dt \leq \varepsilon,$$

where $h(t)$ is as in Theorem 2.4 and $\int_0^{M(x, y)} h(t)dt > 0$ implies

$$\int_0^{d(Ax, By)} h(t)dt < \int_0^{M(x, y)} h(t)dt,$$

$$3) \int_0^{d(Ax, By)} h(t)dt < k \left[\int_0^{d(Sx, Ty)} h(t)dt + \int_0^{d(Sx, Ax)} h(t)dt + \int_0^{d(Ty, By)} h(t)dt + \int_0^{d(Sx, By)} h(t)dt + \int_0^{d(Ty, Ax)} h(t)dt \right],$$

for all x, y in X and $k \in [0, \frac{1}{3})$.

If one of AX, BX, SX, TX is a complete subspace of X , then:

- a) A and S have a coincidence point,
- b) B and T have a coincidence point.

Moreover, if the pairs (A, S) and (B, T) are occasionally weakly compatible mappings, then A, B, S and T have a unique common fixed point.

Definition 2.1. An altering distance is a mapping $\psi : [0, \infty) \rightarrow [0, \infty)$ which satisfies

- (ψ_1) : $\psi(t)$ is increasing and continuous,
- (ψ_2) : $\psi(t) = 0$ if and only if $t = 0$.

Fixed point problem involving an altering distance have been studied in [14], [17], [32], [36], [37] and in other papers.

Lemma 2.1. The function $\psi(x) = \int_0^x h(t)dt$, where $h(t)$ is as in Theorem 2.4, is an altering distance.

Proof. By definitions of $\psi(t)$ and $h(t)$ it follows that $\psi(x)$ is increasing and $\psi(x) = 0$ if and only if $x = 0$. By Lemma 2.5 [20], $\psi(x)$ is continuous. \square

In [30] and [31] the study of fixed points for mappings satisfying implicit relations was initiated. A general fixed point theorem of Meir-Keeler type for noncontinuous weakly compatible mappings satisfying an implicit relation, which generalize (2.1) and others is proved in [33].

The purpose of this paper is to prove a general theorem of Meir-Keeler type using the notion of altering distance for occasionally weakly compatible mappings satisfying an implicit relation.

As an application, a problem of Meir-Keeler type satisfying a condition of integral type becomes a special case of Meir-Keeler type with an altering distance.

3 Implicit relation

Let \mathcal{F}_{MK} be the set of all real continuous mappings $\phi(t_1, \dots, t_6) : \mathbb{R}_+^6 \rightarrow \mathbb{R}$, increasing in t_1 satisfying the following conditions:

- (ϕ_1) : $\phi(t, 0, 0, t, t, 0) \leq 0$ implies $t = 0$,
- (ϕ_2) : $\phi(t, 0, t, 0, 0, t) \leq 0$ implies $t = 0$,
- (ϕ_3) : $\phi(t, t, 0, 0, t, t) > 0, \forall t > 0$.

Example 3.1. $\phi(t_1, \dots, t_6) = t_1 - k(t_2 + t_3 + t_4 + t_5 + t_6)$, where $k \in [0, \frac{1}{3})$.

Example 3.2. $\phi(t_1, \dots, t_6) = t_1 - at_2 - b(t_3 + t_4) - c(t_5 + t_6)$, where $a, b, c \geq 0, b + c < 1$ and $a + 2c < 1$.

Example 3.3. $\phi(t_1, \dots, t_6) = t_1 - b(t_3 + t_4) - c \min\{t_5, t_6\}$, where $b, c \geq 0$, $b < 1$ and $a + c < 1$.

Example 3.4. $\phi(t_1, \dots, t_6) = t_1 - h \max\{t_2, t_3, t_4, t_5, t_6\}$, where $h \in (0, 1)$.

Example 3.5. $\phi(t_1, \dots, t_6) = t_1 - h \max\{t_2, t_3, t_4, \frac{1}{2}(t_5 + t_6)\}$, where $h \in (0, 1)$.

Example 3.6. $\phi(t_1, \dots, t_6) = t_1^2 - at_2^2 - t_3t_4 - bt_5^2 - ct_6^2$, where $a, b, c \geq 0$ and $a + b + c < 1$.

Example 3.7. $\phi(t_1, \dots, t_6) = t_1^3 - k(t_2^3 + t_3^3 + t_4^3 + t_5^3 + t_6^3)$, where $k \in [0, \frac{1}{3})$.

Example 3.8. $\phi(t_1, \dots, t_6) = t_1^3 - \frac{t_3^2 \cdot t_4^2 + t_5^2 \cdot t_6^2}{1 + t_2 + t_3 + t_4}$.

Example 3.9. $\phi(t_1, \dots, t_6) = t_1 - \max\{ct_2, ct_3, ct_4, at_5 + bt_6\}$, where $a, b, c \geq 0$, $c < 1$ and $a + b < 1$.

Example 3.10. $\phi(t_1, \dots, t_6) = t_1 - \alpha \max\{t_2, t_3, t_4\} - (1 - \alpha)(at_5 + bt_6)$, where $\alpha \in [0, 1)$, $a, b \geq 0$ and $a + b < 1$.

Example 3.11. $\phi(t_1, \dots, t_6) = t_1 - \max\left\{t_2, \frac{1}{2}(t_3 + t_4), \frac{1}{2}[(t_5 + t_6)k]\right\}$, where $k \in [0, 1)$.

Example 3.12. $\phi(t_1, \dots, t_6) = t_1 - \max\left\{k_1t_2, \frac{k_2}{2}(t_3 + t_4), \frac{t_5 + t_6}{2}\right\}$, where $k_1 \in [0, 1)$, $k_2 \in [1, 2)$.

Example 3.13. $\phi(t_1, \dots, t_6) = t_1 - \max\left\{k_1(t_2 + t_3 + t_4), \frac{k_2}{2}(t_5 + t_6)\right\}$, where $k_1 \in [0, 1)$, $k_2 \in [0, 2)$.

Let $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a function satisfying the following conditions:

- (φ_1) : φ is continuous,
- (φ_2) : φ is nondecreasing on \mathbb{R}_+ ,
- (φ_3) : $0 < \varphi(t) < t$ for $t > 0$.

Example 3.14. $\phi(t_1, \dots, t_6) = t_1 - \varphi \max\left\{t_2, t_3, t_4, t_5, \frac{t_6}{2}\right\}$.

Example 3.15. $\phi(t_1, \dots, t_6) = t_1 - \varphi \max\left\{t_2, \frac{t_3 + t_4}{2}, \frac{t_5 + t_6}{2}\right\}$.

Example 3.16. $\phi(t_1, \dots, t_6) = t_1 - \varphi \max\left\{t_2, t_3, t_4, \frac{k}{2}(t_5 + t_6)\right\}$, where $k \in [0, 2)$.

Example 3.17. $\phi(t_1, \dots, t_6) = t_1 - \varphi \max\left\{t_2, t_3, t_4, \frac{t_5 + t_6}{2}\right\}$.

4 Main results

Theorem 4.1. *Let A, B, S, T be self mappings of a metric space (X, d) satisfying the inequality:*

$$\begin{aligned} &\phi(\psi(d(Ax, By)), \psi(d(Sx, Ty)), \psi(d(Ax, Sx)), \\ &\psi(d(Ty, By)), \psi(d(Sx, By)), \psi(d(Ty, Ax))) \leq 0 \end{aligned} \quad (4.1)$$

for all x, y in X , where ϕ satisfies property (ϕ_3) and ψ is an altering distance. If there exist $u, v \in X$ such that $Su = Au$ and $Tv = Bv$, then A and S have a unique point of coincidence and B and T have a unique point of coincidence.

Proof. First we prove that $Su = Tv$. If $Su \neq Tv$, using (4.1), we have, successively:

$$\begin{aligned} &\phi(\psi(d(Au, Bv)), \psi(d(Su, Tv)), \psi(d(Au, Su)), \\ &\psi(d(Tv, Bv)), \psi(d(Su, Bv)), \psi(d(Tv, Au))) \leq 0, \end{aligned}$$

$$\phi(\psi(d(Su, Tv)), \psi(d(Su, Tv)), 0, 0, \psi(d(Su, Tv)), \psi(d(Su, Tv))) \leq 0,$$

a contradiction of (ϕ_3) if $d(Su, Tv) > 0$. Hence $\psi(d(Su, Tv)) = 0$, which implies that $Su = Tv$.

Assume that there exists a $p \in X$ such that $Ap = Sp$. Then by (4.1) we have successively:

$$\begin{aligned} &\phi(\psi(d(Ap, Bv)), \psi(d(Sp, Tv)), \psi(d(Sp, Ap)), \\ &\psi(d(Tv, Bv)), \psi(d(Sp, Bv)), \psi(d(Tv, Ap))) \leq 0, \end{aligned}$$

$$\phi(\psi(d(Sp, Tv)), \psi(d(Sp, Tv)), 0, 0, \psi(d(Sp, Tv)), \psi(d(Sp, Tv))) \leq 0,$$

a contradiction of (ϕ_3) if $d(Sp, Tv) > 0$. Therefore $Sp = Tv$ and $z = Au = Su$ is the unique point of coincidence of A and S . Similarly, $w = Tv = Bv$ is the unique point of coincidence of B and T . \square

Lemma 4.1. *Let A, B, S, T be self mappings of a metric space (X, d) such that $AX \subset TX$ and $BX \subset SX$ and ψ is an altering distance. For each $\varepsilon > 0$ there exists a $\delta > 0$ such that*

$$\varepsilon < \psi(M(x, y)) < \varepsilon + \delta \text{ implies } \psi(d(Ax, By)) \leq \varepsilon, \quad (4.2)$$

$$\psi(M(x, y)) > 0 \text{ implies } \psi(d(Ax, By)) < \psi(M(x, y)). \quad (4.3)$$

For $x_0 \in X$ and $\{y_n\}$ defined by $y_{2n-1} = Tx_{2n-1} = Ax_{2n-2}$ and $y_{2n} = Sx_{2n} = Bx_{2n-1}$ for $n \in \mathbb{N}^$ and $d_n = d(y_n, y_{n+1})$, then $\lim d_n = 0$.*

Proof. First we prove that if, for some $k \in \mathbb{N}^*$, $d_{k+1} > 0$, then

$$\psi(d_{k+1}) < \psi(d_k). \quad (4.4)$$

a) Assume that $d_{2k} > 0$ for some $k \in \mathbb{N}^*$. Then $M(x_{2k}, x_{2k-1}) > 0$ otherwise $Ax_{2k} = Bx_{2k-1}$, i.e. $y_{2k} = y_{2k+1}$ so $d_{2k} = 0$, a contradiction. Hence $d_{2k} = d(Ax_{2k}, Bx_{2k-1}) < M(x_{2k}, x_{2k-1})$ which implies by (4.3) that $0 < \psi(d_{2k}) < \psi(M(x_{2k}, x_{2k-1})) \leq \psi(\max\{d_{2k-1}, d_{2k}\}) = \psi(d_{k-1})$.

b) If $d_{2k+1} > 0$ for some $k \in \mathbb{N}^*$, using a similar argument as in a), one can verify that $\psi(d_{2k+1}) < \psi(d_{2k})$.

c) Combining the results of a) and b) we may conclude that

$$\psi(d_{k-1}) < \psi(d_{2k}) \text{ for } k \in \mathbb{N}^*. \quad (4.5)$$

Moreover, if for some $k \in \mathbb{N}^*$, $\psi(d_k) = 0$, then $d_k = 0$ which implies $d_{k+1} = 0$ because, if $d_{k+1} > 0$, then $\psi(d_{k+1}) > 0$ which implies by a) and b) that $\psi(d_{k+1}) < \psi(d_k) = 0$, a contradiction. Hence, for $n \geq k$ we have $y_n = y_k$ and hence $\lim d(y_n, y_{n+1}) = 0$.

We prove that $\lim d(y_n, y_{n+1}) = 0$ for $\psi(d_k) > 0$.

By (4.5) it follows that $\psi(d_n)$ is strictly decreasing, hence convergent to some $\ell \in \mathbb{R}_+$. Suppose that $\ell > 0$. Then by (4.2) for $\varepsilon = \ell$, there exists a $\delta > 0$ such that $\ell < \psi(d_n) < \ell + \delta$ for $n \geq k$. In particular $\ell < \psi(M(x_{2k}, x_{2k+1})) < \ell + \delta$, since $M(x_{2k}, x_{2k+1}) = \max\{d_{2k}, d_{2k+1}\} \leq \ell$. Hence $\ell < \psi(d_{2k}) \leq \ell$, a contradiction, and $\ell = 0$. Let $a_n = \psi(d(y_n, y_{n+1}))$, $n \geq 0$. Then by the continuity of ψ we obtain

$$0 = \lim a_n = \lim \psi(d(y_n, y_{n+1})) = \psi(\lim d(y_n, y_{n+1})).$$

Hence $\lim d(y_n, y_{n+1}) = 0$. □

Theorem 4.2. Let A, B, S and T be self mappings of a metric space (X, d) satisfying the following conditions:

a) $AX \subset TX$ and $BX \subset SX$,

b) given an $\varepsilon > 0$ there exists a $\delta > 0$ such that for all $x, y \in X$

$$\varepsilon < \psi(M(x, y)) < \varepsilon + \delta \text{ implies } \psi(d(Ax, By)) \leq \varepsilon,$$

c) $\psi(M(x, y)) > 0$ implies $\psi(d(Ax, By)) < \psi(M(x, y))$,

d) the inequality (4.1) holds for all x, y in X , where $\phi \in \mathcal{F}_{MK}$ and ψ is an altering distance.

If one of AX, BX, SX, TX is a complete subspace of X , then:

e) A and S have a coincidence point,

f) B and T have a coincidence point.

Moreover, if the pairs (S, A) and (T, B) are occasionally weakly compatible mappings, then A, B, S and T have a unique common fixed point.

Proof. First we prove that $\{y_n\}$ is a Cauchy sequence. Since by Lemma 4.1, $\lim d(y_n, y_{n+1}) = 0$ it is sufficient to show that $\{y_{2n}\}$ is a Cauchy sequence. Suppose that $\{y_{2n}\}$ is not a Cauchy sequence. Then there exists an $\varepsilon > 0$ such that for even integer $2k$, there exists even integers $2m(k)$ and $2n(k)$ such that $d(y_{2m(k)}, y_{2n(k)}) > \varepsilon$ with $2m(k) > 2n(k) \geq 2k$. For even integer $2k$, let $2m(k)$ be the least even integer exceeding $2n(k)$ such that $d(y_{2n(k)}, y_{2m(k)}) < \varepsilon$.

As in Theorem 2.2 [12] we deduce that

$$\begin{aligned}\lim d(y_{2n(k)}, y_{2m(k)}) &= \varepsilon, \\ \lim d(y_{2n(k)}, y_{2m(k)-1}) &= \varepsilon, \\ \lim d(y_{2n(k)+1}, y_{2m(k)-1}) &= \varepsilon.\end{aligned}$$

On the other hand we have successively

$$\begin{aligned}d(y_{2n(k)}, y_{2m(k)}) &\leq d_{2n(k)} + d(Ax_{2n(k)}, Bx_{2m(k)-1}), \\ d(y_{2n(k)}, y_{2m(k)}) - d_{2n(k)} &\leq d(y_{2n(k)+1}, y_{2m(k)}).\end{aligned}$$

Hence

$$\psi(d(y_{2n(k)}, y_{2m(k)}) - d_{2n(k)}) \leq \psi(d(y_{2n(k)+1}, y_{2m(k)})).$$

Setting in (4.1) $x = x_{2n(k)}$ and $y = x_{2m(k)-1}$ we obtain

$$\begin{aligned}\phi(\psi(d(y_{2n(k)}, y_{2m(k)})), \psi(d(y_{2n(k)}, y_{2m(k)-1})), \psi(d(y_{2n(k)}, y_{2n(k)+1})), \\ \psi(d(y_{2m(k)-1}, y_{2m(k)})), \psi(d(y_{2n(k)}, y_{2m(k)})), \psi(d(y_{2m(k)-1}, y_{2n(k)}))) \leq 0,\end{aligned}$$

$$\begin{aligned}\phi(\psi(d(y_{2n(k)}, y_{2m(k)}) - d_{2n(k)}), \psi(d(y_{2n(k)}, y_{2m(k)-1})), \psi(d(y_{2n(k)}, y_{2n(k)+1})), \\ \psi(d(y_{2m(k)-1}, y_{2m(k)})), \psi(d(y_{2n(k)}, y_{2m(k)})), \psi(d(y_{2m(k)-1}, y_{2n(k)}))) \leq 0.\end{aligned}$$

Letting n tend to infinity we obtain

$$\phi(\psi(\varepsilon), \psi(\varepsilon), 0, 0, \psi(\varepsilon), \psi(\varepsilon)) \leq 0,$$

a contradiction of (ϕ_3) . Hence, $\{y_{2n}\}$ is a Cauchy sequence. It follows that $\{y_n\}$ is a Cauchy sequence.

Assume that least one of AX or TX is a complete subspace of X .

Since $y_{2n+1} \in AX \subset TX$ and $\{y_{2n+1}\}$ is a Cauchy sequence, there exists a $u \in TX$ such that $\lim y_{2n+1} = u$. The sequence $\{y_n\}$ converges to u since it

is Cauchy and has the subsequence $\{y_{2n+1}\}$ convergent to u . Let $v \in X$ such that $u = Tv$. Setting $x = x_{2n}$ and $y = v$ in (4.1) we get

$$\begin{aligned} & \phi(\psi(d(Ax_{2n}, Bv)), \psi(d(Sx_{2n}, Tv)), \psi(d(Sx_{2n}, Ax_{2n})), \\ & \psi(d(Tv, Bv)), \psi(d(Sx_{2n}, Bv)), \psi(d(Tv, Ax_{2n}))) \leq 0, \\ & \phi(\psi(d(y_{2n+1}, Bv)), \psi(d(y_{2n}, Tv)), \psi(d(y_{2n}, y_{2n+1})), \\ & \psi(d(Tv, Bv)), \psi(d(y_{2n}, Bv)), \psi(d(u, y_{2n+1}))) \leq 0. \end{aligned}$$

Letting n tend to infinity in the above inequality we obtain

$$\phi(\psi(d(u, Bv)), 0, 0, \psi(d(u, Bv)), \psi(d(u, Bv)), 0) \leq 0.$$

From (ϕ_1) , $\psi(d(u, Bv)) = 0$ which implies that $d(u, Bv) = 0$ i.e. $u = Bv$. Hence $u = Tv = Bv$ and v is a coincidence point of T and B . Since $u = Bv \in BX \subset SX$, there exists a $w \in X$ such that $u = Sw$. Using a similar argument as above we obtain $u = Aw$. Hence, $u = Tv = Bv = Sw = Aw$.

Indeed, setting $x = w$ and $y = x_{2n+1}$ in (4.1) we obtain

$$\begin{aligned} & \phi(\psi(d(Aw, y_{2n+2})), \psi(d(u, y_{2n+2})), \psi(d(u, Aw)), \\ & \psi(d(y_{2n+1}, y_{2n+2})), \psi(d(u, y_{2n+2})), \psi(d(y_{2n+1}, Aw))) \leq 0, \end{aligned}$$

and letting n tend to infinity we obtain

$$\phi(\psi(d(Aw, u)), 0, \psi(d(u, Aw)), 0, 0, \psi(d(Aw, u))) \leq 0.$$

By (ϕ_1) it follows that $\psi(d(Aw, u)) = 0$, hence $u = Aw$.

By Theorem 4.1 u is the unique point of coincidence of A , S and B , T .

If the pairs (A, S) and (B, T) are occasionally weakly compatible, then by Lemma 1.1 u is the unique common fixed point of A , B , S and T . \square

Taking $A = B$ and $S = T$ in Theorem 4.2 we obtain

Theorem 4.3. *Let A and S and T be self mappings of a metric space (X, d) satisfying the following conditions:*

- 1) $AX \subset SX$,
- 2) $\varepsilon < \psi(M_1(x, y)) < \varepsilon + \delta$ implies $\psi(d(Ax, By)) \leq \varepsilon$, where $M_1(x, y) = \max\{d(Sx, Sy), d(Sx, Ax), d(Sy, Ty), d(Sx, Ay), d(Sy, Ax)\}$,
- 3) $\psi(M_1(x, y)) > 0$ implies $\psi(d(Ax, By)) < \psi(M_1(x, y))$.
- 4) $\phi(\psi(d(Ax, Ay)), \psi(d(Sx, Sy)), \psi(d(Sx, Ax)), \psi(d(Sy, Ay)), \psi(d(Sx, Ay)), \psi(d(Sy, Ax))) \leq 0$ for all x, y in X , where $\phi \in \mathcal{F}_{MK}$ and ψ is an altering distance.

If one of SX and AX is a complete subspace of X , then:

5) A and S have a coincidence point.

Moreover, if the pair (A, S) is occasionally weakly compatible, then A and S have a unique common fixed point.

For $\psi(t) = t$ by Theorem 4.2 we obtain

Theorem 4.4. *Let A, B, S and T be self mappings of a metric space (X, d) satisfying the following conditions:*

- 1) $AX \subset SX$,
- 2) given an $\varepsilon > 0$, there exists a $\delta > 0$ such that for all x, y in X , $\varepsilon < M(x, y) < \varepsilon + \delta$ implies $d(Ax, By) \leq \varepsilon$,
- 3) $M(x, y) > 0$ implies $d(Ax, By) < M(x, y)$,
- 4) $\phi(d(Ax, By), d(Sx, Ty), d(Sx, Ax), d(Ty, By), d(Sx, By), d(Ty, Ax)) \leq 0$ for all x, y in X and $\phi \in \mathcal{F}_{MK}$.

If one of AX, BX, SX, TX is a complete subspace of X , then:

- 5) A and S have a coincidence point,
- 6) B and T have a coincidence point.

Moreover, if the pairs (A, S) and (B, T) are occasionally weakly compatible, then A, B, S and T have a unique common fixed point.

Remark 4.1. 1) By Example 3.1 and Theorem 4.4 we obtain a generalization of Theorem 2.2 and Theorem 2.1.

2) By Example 3.4 and Theorem 4.4 we obtain a generalization of Theorem 2.1 [5].

3) By Example 3.12 and Theorem 4.4 we obtain a generalization of the result from [27].

4) By Example 3.11 and Theorem 4.4 we obtain a generalization of the result from [28] for $k \in [0, 1)$.

5) By Example 3.16 and Theorem 4.4 we obtain a generalization of Theorem 2.1 [26].

6) Theorem 4.4 is a generalization of Theorem 5 [33] for weakly compatible mappings satisfying an implicit relation.

7) By Examples 3.2, 3.3, 3.5 - 3.10, 3.13 - 3.15 we obtain new fixed point theorems of Meir - Keeler type.

5 Applications

Theorem 5.1. *Let A, B, S and T be self mappings of a metric space (X, d) satisfying conditions (a), (b), (c) of Theorem 4.2. Assume that there exists a $\phi \in \mathcal{F}_{MK}$ such that:*

$$\begin{aligned} & \phi\left(\int_0^{d(Ax, By)} h(t)dt, \int_0^{d(Sx, Ty)} h(t)dt, \int_0^{d(Sx, Ax)} h(t)dt, \right. \\ & \left. \int_0^{d(Ty, By)} h(t)dt, \int_0^{d(Sx, By)} h(t)dt, \int_0^{d(Ty, Ax)} h(t)dt\right) \leq 0 \end{aligned} \quad (5.1)$$

for all x, y in X where $h(t)$ is as in Theorem 2.4.

If one of AX, BX, SX, TX is a complete subspace of X then:

1) A and S have a coincidence point,

2) B and T have a coincidence point.

Moreover, if the pairs (A, S) and (B, T) are occasionally weakly compatible, then A, B, S and T have a unique common fixed point.

Proof. Let

$$\begin{aligned} \psi(d(Ax, By)) &= \int_0^{d(Ax, By)} h(t)dt, \psi(d(Sx, Ty)) = \int_0^{d(Sx, Ty)} h(t)dt, \\ \psi(d(Sx, Ax)) &= \int_0^{d(Sx, Ax)} h(t)dt, \psi(d(Ty, By)) = \int_0^{d(Ty, By)} h(t)dt, \\ \psi(d(Sx, By)) &= \int_0^{d(Sx, By)} h(t)dt, \psi(d(Ty, Ax)) = \int_0^{d(Ty, Ax)} h(t)dt, \end{aligned}$$

where $h(t)$ is as in Theorem 2.4.

By Lemma 2.1 $\psi(x) = \int_0^x h(t)dt$ is an altering distance.

By (5.1) we obtain

$$\begin{aligned} & \phi(\psi(d(Ax, By)), \psi(d(Sx, Ty)), \psi(d(Sx, Ax)), \\ & \psi(d(Ty, By)), \psi(d(Sx, By)), \psi(d(Ty, Ax))) \leq 0 \end{aligned}$$

for all x, y in X , which is inequality (4.1).

Hence, the conditions of Theorem 4.2 are satisfied and the conclusion of Theorem 5.1 it follows from Theorem 4.2. \square

Remark 5.1. *If $h(t) = 1$ we obtain Theorem 4.3.*

Remark 5.2. *By Theorem 5.1 and Example 3.1 we obtain Theorem 2.5.*

Examples 3.2 - 3.17 are new results.

References

- [1] A. Aliouche, *A common fixed point theorem for weakly compatible mappings in symmetric spaces satisfying contractive conditions of integral type*, J. Math. Anal. Appl. **322**(2000), 796 - 802.
- [2] A. Aliouche and V. Popa, *Common fixed point for occasionally weakly compatible mappings via implicit relations*, Filomat **22**(2)(2008), 99 - 107.
- [3] M. A. Al - Thagafi and Naseer Shahzad, *Generalized I - nonexpansive maps and invariant approximations*, Acta Math. Sinica **24**(5)(2008), 867 - 876.
- [4] H. Bouhajdera and A. Djoudi, *On common fixed point theorems of Meir - Keeler type*, An. Șt. Univ. Ovidius Constanța **16**(2)(2008), 39 - 46.
- [5] H. Bouhajdera and B. Fisher, *Common fixed point theorems of Meir - Keeler type for weakly compatible mappings*, Mathematica Moravica **11**(2007), 9 -16.
- [6] A. Branciari, *A fixed point theorem for mappings satisfying a general contractive condition of integral type*, Intern. J. Math. Math. Sci. **29**, 9(2002), 531 - 536.
- [7] U. Gairola and A. R. Rawat, *A fixed point theorem for two pairs of maps satisfying a new contractive condition of integral type* (submitted).
- [8] K. Jha, *Common fixed point for weakly compatible maps in metric spaces*, Kathmandu Univ. J. Sci. Eng. Techn. **1**, 4(2007), 1 - 6.
- [9] K. Jha, R. Pant and S. L. Singh, *Common fixed point for compatible mappings in metric spaces*, Radovi Mat. **12**(2003), 107 - 114.
- [10] G. Jungck, *Compatible mappings and common fixed points*, Intern. J. Math. Math. Sci. **9**(1986), 771 - 779.
- [11] G. Jungck, *Common fixed points for noncontinuous nonself maps on a nonnumeric space*, Far. East J. Math. Sci. **4**(2)(1996), 199 - 215.
- [12] G. Jungck and B. E. Rhoades, *Fixed points for occasionally weakly compatible mappings*, Fixed Point Theory **7**(2)(2007), 287 - 297.
- [13] S. M. Kang, Y. J. Cho and G. Jungck, *Common fixed point of compatible mappings*, Intern. J. Math. Math. Sci. **13**(1)(1990), 61 - 66.
- [14] M. S. Khan, M. Swaleh and S. Sessa, *Fixed point theorems by altering distances between points*, Bull. Austral. Math. Soc. **30**(1984), 1 - 9.

- [15] J. K. Kohli and S. Washistha, *Common fixed point theorems for compatible and weak compatible mappings satisfying general contractive conditions*, Stud. Cerc. St. Ser. Mat. Univ. Bacău **16**(2006), 33 - 42.
- [16] S. Kumar, R. Chugh and R. Kumar, *Fixed point theorem for compatible mappings satisfying a contractive condition of integral type*, Soochow J. Math. **33**(2007), 181 - 185.
- [17] B. Marzuki and A. M. Mbarki, *Multivalued fixed point theorems by altering distances between points*, Southwest J. Pure Appl. Math. **1**(2002), 126 - 134.
- [18] M. Mati and T. K. Pal, *Generalizations of two fixed point theorems*, Bull. Calcutta Math. Soc. Math. Soc. **70**(1978), 57 - 61.
- [19] A. Meir and E. Keeler, *A theorem of contraction mappings*, J. Math. Anal. Appl. **28**(1969), 326 - 329.
- [20] M. Mocanu and V. Popa, *Some fixed point theorems for mappings satisfying implicit relations in symmetric spaces*, Libertas Math. **28**(2008), 1 - 13.
- [21] R. P. Pant, *Common fixed point of two pairs of commuting mappings*, Indian J. Pure Appl. Math. **17**(2) (1986), 187 - 192.
- [22] R. P. Pant, *Common fixed points of weakly commuting mappings*, Math. Student **62**, 1 - 4 (1993), 97 - 102.
- [23] R. P. Pant, *Common fixed points for noncommuting mappings*, J. Math. Anal. Appl. **188**(1994), 436 - 440.
- [24] R. P. Pant, *Common fixed point theorems for contractive maps*, J. Math. Anal. Appl. **226**(1998), 251 - 286.
- [25] R. P. Pant, *Common fixed points for four mappings*, Calcutta Math. Soc. **9**(1998), 281 - 287.
- [26] R. P. Pant, *A new common fixed point principle*, Soochow J. Math. **27**, 3(2001), 287 - 292.
- [27] R. P. Pant and K. Jha, *A generalization of Meir - Keeler type common fixed point for four mappings*, J. Natural and Physical Sciences **16**(1 - 2)(2002), 77 - 84.
- [28] R. P. Pant and K. Jha, *A generalization of Meir - Keeler type fixed point theorems for four mappings*, Ultra Science **15**(1)(2003), 97 - 102.

- [29] S. Park and B. E. Rhoades, *Meir – Keeler contractive conditions*, Math. Japonica **26**(1)(1981), 13 - 20.
- [30] V. Popa, *Fixed point theorems for implicit contractive mappings*, Stud. Cerc. St. Ser. Math. Univ. Bacău **7**(1997), 129 - 133.
- [31] V. Popa, *Some fixed point theorem for four compatible mappings satisfying an implicit relation*, Demonstratio Math. **32**(1999), 157 - 163.
- [32] V. Popa, *A fixed point theorem for four compatible mappings in compact metric spaces*, U. P. B. Bull. Ser. A **63**, 4(2001), 43 - 46.
- [33] V. Popa, *A general common fixed point theorem of Meir – Keeler type for noncontinuous weak compatible mappings*, Filomat **18**(2004), 33 - 40.
- [34] V. Popa and M. Mocanu, *A new viewpoint in the study of fixed points for mappings satisfying a contractive condition of integral type*, Bull. Inst. Politeh. Iași, Sect. Mat. Mec. Teor. Fiz. **53**(57), 5(2007), 269 - 272.
- [35] J. H. N. Rao and K. P. R. Rao, *Generalizations of fixed point theorem of Meir and Keeler type*, Indian J. Pure Appl. Math. **16**(1)(1985), 1249 - 1262.
- [36] K. P. Sastri and G. V. R. Babu, *Fixed point theorems by altering distances*, Bull. Calcutta Math. Soc. **90**(1998), 175 - 182.
- [37] K. P. Sastri and G. V. R. Babu, *Some fixed point theorems by altering distance between the points*, Indian J. Pure Appl. Math. **30**(1999), 641 - 647.

Alina-Mihaela PATRICIU,
Department of Mathematics, Informatics and Education Sciences,
“Vasile Alecsandri” University of Bacău,
157 Calea Mărășești, Bacău, 600115, Romania.
Email: alina.patriciu@ub.ro

Valeriu POPA,
Department of Mathematics, Informatics and Education Sciences,
“Vasile Alecsandri” University of Bacău,
157 Calea Mărășești, Bacău, 600115, Romania.
Email: vpopa@ub.ro