



# RELATION BETWEEN GROUPS WITH BASIS PROPERTY AND GROUPS WITH EXCHANGE PROPERTY

A.Al Khalaf, M. Alkadhi

## Abstract

A group  $G$  is called a group with basis property if there exists a basis (minimal generating set) for every subgroup  $H$  of  $G$  and every two bases are equivalent. A group  $G$  is called a group with exchange property, if  $x \notin \langle X \rangle \wedge x \in \langle X \cup \{y\} \rangle$ , then  $y \in \langle X \cup \{x\} \rangle$ , for all  $x, y \in G$  and for every subset  $X \subseteq G$ .

In this research, we proved the following: Every polycyclic group satisfies the basis property. Every element in a group with the exchange property has a prime order. Every  $p$ -group satisfies the exchange property if and only if it is an elementary abelian  $p$ -group. Finally, we found necessary and sufficient condition for every group to satisfy the exchange property, based on a group with the basis property.

## 1 Introduction

A generating set  $X$  is said to be minimal if it has no proper subset which forms a generating set. The subset  $X$  of a group  $G$  is called independent, if for all  $x \in X$ ,  $x \notin \langle X \setminus \{x\} \rangle$ . Independent set  $X$  is called a basis subgroup  $\langle X \rangle$ . In 1978 Jones [5] introduced an initial study of semigroups with the basis property. Jones [5] states that if  $G$  is an inverse semigroup and  $U \leq V \leq G$  then a  $U$ -basis for  $V$  is a subset  $X$  of  $V$  which is minimal such that  $\langle U \cup X \rangle = G$ .

---

Key Words: Groups, Basis property, Exchange property.  
2010 Mathematics Subject Classification: Primary 20D25; Secondary 20D10.  
Received: 31.07.2014  
Accepted: 20.02.2015

So a minimal generating set for  $V$  is a  $\emptyset$ -basis. A basis property of universal algebra  $A$  means that every two minimal (with respect to inclusion) generating set (basis) of an arbitrary subalgebra of  $A$  have the same cardinality [1].

## 2 Basis property

**Definition 2.1** A group  $G$  is called a group with basis property if there exists a basis minimal (irreducible) generating sets (with respect to inclusion) for every subgroup  $H$  of  $G$  and every two bases are equivalent (i.e. they have the same cardinality) [1].

Notice that finitely generated vector spaces have the property that all minimal generating sets have the same cardinality. Jones [5] introduce another concept which is state for inverse semigroup.

**Definition 2.2** An inverse semigroup  $S$  has the strong basis property if for any inverse subsemigroup  $V$  of  $S$  and inverse subsemigroup  $U$  of  $V$  any two  $U$ -bases for  $V$  have the same cardinality.

Let  $(\mathbb{Z}, +)$  be an additive abelian group, then we can write  $\mathbb{Z} = \langle 1 \rangle = \langle 2, 3 \rangle$  even though  $2 \notin \langle 3 \rangle$  and  $3 \notin \langle 2 \rangle$ . Thus  $\mathbb{Z}$  does not have the basis property. Hence free groups do not have the basis property. The first results on the basis property of groups was in [6]. The author proved that a group with basis property is periodic, all elements of such a group have prime power order, and solvable. Therefore by [1] every finite  $p$ -group has a basis property, and the homomorphic image of every finite group with basis property is again a group with basis property, but in case of infinite group we have the following:

**Remark 2.3** Let  $G = \sum_{i=1}^{\infty} \mathbb{Z}_{p^i}$  be a direct sum of a cyclic  $p$ -group  $P$ , then one of homomorphic image is a quasicyclic group  $K = \mathbb{Z}_{p^\infty}$ , which is not a group with basis property, but the group  $G$  is a group with basis property.

**Lemma 2.4** Let  $G$  be a group in which every element has prime power order, let  $x \in G$  such that  $|x| = p^c$  and  $y \in G$  such that  $|y| = q^b$ ,  $p \neq q$  are primes. Then  $xy \neq yx$ .

*Proof.* Suppose that  $xy = yx$ , then  $xy$  is an element of order  $p^c q^b$ , hence  $xy$  has a composite order in  $G$ . This is contradiction with basis property[1], so  $xy \neq yx$ . □

**Proposition 2.5** Let  $G$  be a finite nilpotent group. Then  $G$  is a group with basis property if and only if  $G$  is a primary group.

*Proof.* Suppose that  $G$  is a finite nilpotent group with basis property. From [11] every finite nilpotent group is decomposable in a direct product of Sylow

subgroups. Then

$$G = G_1 \times G_2 \times \cdots \times G_m,$$

such that  $G_i$  is a  $p_i$ -group for some primes  $p_i$ ,  $p_i \neq p_j$  if  $i \neq j$ . If  $m > 1$ , then in  $G$  there exists two commute elements with a prime power order. Hence we have a contradiction with lemma(2-4). Thus  $G$  is a primary group.

Conversely, if  $G$  is a primary group, then  $G$  is a group with basis property [5].  $\square$

A classification of group with the basis property was announced by Dickson and Jones in [5], but as far as we can see this has yet to be published. However a classification of finite groups with the basis property was given by Al Khalaf [1] exploiting Higman's result, this classification requires a technical condition on the  $p$ -group and he proved the following theorem:

**Theorem 2.6** [1]. Let a finite group  $G$  be a semidirect product of a  $p$ -group  $P = \text{Fit}(G)$  (Fitting subgroup) of  $G$  by a cyclic  $q$ -group  $\langle y \rangle$ , of order  $q^b$ , where  $p \neq q$  ( $p$  and  $q$  are primes),  $b \in \mathbb{N}$ . Then the group  $G$  has basis property if and only if for any element  $y \in \langle y \rangle$ ,  $y \neq e$  and for any invariant subgroup  $H$  of  $P$  the automorphism  $\varphi_u$  must define an isotopic representation on every quotient Frattini subgroup of  $H$ .

In [4], the author used some common results from both group and module theory using Maschke, Clifford and Krull-Schmidt, to classify the group with basis property.

Finally Jones [7] studied basis property from the point of view exchange properties.

**Theorem 2.7** [3] Let  $G$  be a semidirect product of abelian  $p$ -group  $P$  by a cyclic  $q$ -group  $\langle y \rangle$ , of order  $q^b$ , where  $p \neq q$  ( $p$  and  $q$  are primes),  $b \in \mathbb{N}$ , which is defined automorphism  $\varphi$  of order  $q^b$  and  $P$  has an exponent  $p^c$ ,  $c \in \mathbb{N}$ . Then the group  $G$  has basis property if and only if there exists a polynomial  $g(x) \in \mathbb{Z}[x]$  such that satisfy the following conditions:

1. The polynomial  $f(x) = \bar{\theta}(g(x))$  is irreducible over the field  $GF(p)$ ,  
 $f(x) \mid x^{q^b} - 1$  and  $f(x) \nmid x^{q^{b-1}} - 1$ .
2.  $g^m(\varphi) = 0$ .

In this research we study special group with the basis property. The concept of exchange property and continued results as shown in [7] and [8].

**Theorem 2.8** Let  $G$  be a finite polycyclic group such that  $G$  has a presentation [9]:

$$G = \langle x, y : x^{p^c} = y^{q^b} = 1, y^{-1}xy = x^r \rangle, \quad (2-1)$$

such that  $p \neq q$  ( $p$  and  $q$  are primes)  $b, c, r \in \mathbb{Z}^+, (p, r-1) = 1$  and

$$r^{q^b} \equiv 1 \pmod{p^c}, r \not\equiv 1 \pmod{p}, 0 \leq r \leq p^c. \quad (2-2)$$

Then  $G$  is a group with the basis property if and only if it satisfies the following conditions:

$$p \equiv 1 \pmod{q^b}, \quad (2-3)$$

$$r^{q^{b-1}} \not\equiv 1 \pmod{p}. \quad (2-4)$$

*Proof.* Suppose that  $G$  is a group with the basis property. From (2-1) we have that  $G$  is a semidirect product of cyclic  $p$ -group  $\langle x \rangle$ ,  $|\langle x \rangle| = p^c$  by a cyclic  $q$ -group  $\langle y \rangle$ ,  $|\langle y \rangle| = q^b$ , where  $p \neq q$  ( $p$  and  $q$  are primes)  $b, c \in \mathbb{Z}^+$ . Then from [1]  $G$  is a Frobenius group with kernel  $\langle x \rangle$  and complement  $\langle y \rangle$ . Thus by [3] we see that  $p \equiv 1 \pmod{q^b}$ . Thus (2-3) holds.

Assume that

$$r^{q^{b-1}} \equiv 1 \pmod{p}. \quad (2-5)$$

Then  $r^{q^{b-1}} = 1 + mp$  for some  $m \in \mathbb{Z}^+$ . Considering the non trivial elements  $x^{p^{c-1}}, y^{q^{b-1}}$  and using (2-1) and (2-5) then we have:

$$\begin{aligned} y^{-q^{b-1}} x^{p^{c-1}} y^{q^{b-1}} &= \left( y^{-q^{b-1}} x y^{q^{b-1}} \right)^{p^{c-1}} = \left( y^{-q^{b-1}-1} (y^{-1} x y) y^{q^{b-1}-1} \right)^{p^{c-1}} \\ &= \left( y^{-q^{b-1}-1} x^r y^{q^{b-1}-1} \right)^{p^{c-1}} = \dots = \left( x^{r^{q^{b-1}}} \right)^{p^{c-1}} = x^{p^{c-1}(1+mp)} = \\ &= x^{p^{c-1}} (x^{p^c})^m = x^{p^{c-1}}. \end{aligned}$$

Hence the  $p$ -element  $x^{p^{c-1}}$  commutes with the  $q$ -element in  $G$ , so we have a contradiction with lemma (2-4). Thus (2-4) holds.

Conversely, let  $G$  be a polycyclic group satisfying conditions (2-3), and (2-4). Then from [9] we see that  $G$  is an extension of cyclic  $p$ -group  $\langle x \rangle$  of order  $p^c$  by cyclic  $q$ -group  $\langle y \rangle$  of order  $q^b$ ,  $p \neq q$  ( $p$  and  $q$  are primes)  $b, c \in \mathbb{Z}^+$ . Thus  $(|\langle x \rangle|, |\langle y \rangle|) = 1$  and  $|G| = |\langle x \rangle| |\langle y \rangle|$ , then  $\langle x \rangle \cap \langle y \rangle = \{1\}$  and  $G = \langle x \rangle \langle y \rangle$ ,

so  $G = \langle x \rangle \rtimes \langle y \rangle$ . Since  $\langle x \rangle \trianglelefteq G$  and  $\langle x \rangle$  is an abelian  $p$ -group, then by using theorem (2-7) we prove that  $G$  is a group with the basis property.

Now consider the polynomial  $g(x) = x - r$  over the ring  $\mathbb{Z}$ . Denote that  $f(x) = \bar{\theta}(g(x))$ . Then the polynomial  $f(x)$  is an irreducible over the field  $GF(p)$  and has  $\bar{r}$  zeros. Thus by (2-2), and (2-4) we have  $\bar{r}^{q^b} = 1$ ,  $\bar{r}^{q^{b-1}} \neq 1$ , hence by Bezout theorem the polynomial  $f(x)$  divides  $x^{q^b} - 1$  and not divides  $x^{q^{b-1}} - 1$ , i.e. the condition 1) in theorem(2-7) holds for  $g(x)$ . Now consider the automorphism  $\varphi$ , which defines a semidirect product  $\langle x \rangle \rtimes \langle y \rangle$  and induced by  $y$  element, i.e.

$$\varphi : a \rightarrow y^{-1}ay, \quad \forall a \in \langle x \rangle.$$

From (2-1) we get

$$\varphi(a) = a^r, \quad \forall a \in \langle x \rangle.$$

Using additive form in  $\langle x \rangle$ , then we have  $g(\varphi) = 0$ . Thus the condition 2) of theorem(2-7) for  $g(x)$  holds too. Hence  $G$  is a group with the basis property.  $\square$

### 3 Exchange property

The fundamental property of generating operator  $\varphi$  of subspace of the vector space  $V$  over the field  $F$  that this operator satisfies exchange property.

**Definition 3.1** Let  $V$  be a vector space, then  $\forall x, y \in V$  and for every subset  $X \subseteq V$  if  $x \notin \varphi(X)$  and if  $x \in \varphi(X \cup \{y\})$ , then  $y \in \varphi(X \cup \{x\})$ .

**Theorem 3.2** Let  $G$  be a group with the exchange property, i.e.  $\forall x, y \in G$  and for every subset  $X \subseteq G$ ,

$$\text{if } x \notin \langle X \rangle \wedge x \in \langle X \cup \{y\} \rangle, \text{ then } y \in \langle X \cup \{x\} \rangle. \quad (3-1)$$

Then the order of every element  $a \in G$ ,  $a \neq 1$  is a prime.

*Proof.* First, we prove that every cyclic subgroup of  $G$  is simple, i.e. every cyclic subgroup does not contain non trivial normal subgroup.

Suppose that  $\{1\} \leq \langle x \rangle \leq \langle y \rangle$  for  $x, y \in G$ . Then  $x \notin \{1\}$  and  $x \in \langle \{1\} \cup \{y\} \rangle$  such that substituting  $X = \{1\}$  in (3-1) we find  $y \in \langle \{1\} \cup \{x\} \rangle = \langle x \rangle$  and we get a contradiction with our assumption. Thus  $O(x) \in \{p, q\}$ ,  $\forall x \in G \setminus \{1\}$ .

**Theorem 3.3** Let  $G$  be a  $p$ -group such that  $p$  is a prime. Then  $G$  is a group with the exchange property if and only if  $G$  is elementary abelian  $p$ -group.

*Proof.* Suppose that  $G$  is a  $p$ -group with the exchange property. Then by theorem (3-2)

$$x^p = 1, \forall x \in G, \quad (3-2)$$

hence  $G^p = \{1\}$  and by [10]  $\Phi(G) = G^p G'$ . Since  $G$  is a  $p$ -group, then

$$\Phi(G) = G', \quad \Phi^2(G) = G'', \dots$$

If  $G' = \{1\}$ , then  $G$  is an elementary abelian group.

Suppose that  $G' \neq \{1\}$ . Then there exist elements  $a, b, c \in G$  such that

$$[a, b] = a^{-1}b^{-1}ab = c \neq 1. \quad (3-3)$$

Now assume that  $c \in \langle a \rangle$ , then  $a \in \langle c \rangle$ . Let consider the subgroup, which is generated by two elements  $a, b$ , i.e.  $\langle a, b \rangle$ . If  $\langle a, b \rangle$  is a cyclic group, then it is commutative and we have a contradiction with (3-3), then  $a \notin \langle b \rangle$  and  $b \notin \langle a \rangle$ . Hence the set  $\{a, b\}$  forms a basis of group  $\langle a, b \rangle$ . Since  $\langle a \rangle = \langle c \rangle$ , so  $\langle a, b \rangle = \langle c, b \rangle$  and by the basis property of  $G$  [6]. Thus we have that the set  $\{c, b\}$  forms a basis of  $G$  and this is a contradiction with properties of the Frattini subgroup, i.e.  $c \in \Phi(G)$ .

Hence  $c \notin \langle a \rangle$  and  $c \in \langle a, b \rangle$ , and by the exchange property we have  $b \in \langle a, c \rangle$ . But then  $\langle a, b \rangle = \langle a, c \rangle$ . So by the basis property for  $G$  and since  $a \notin \langle b \rangle$ ,  $b \notin \langle a \rangle$  we conclude that the set  $\{a, c\}$  forms a basis for  $G$ . Hence this is a contradiction with properties of the the Frattini subgroup  $\Phi(G)$ , i.e.  $c \in \Phi(G)$ . Thus  $[a, b] = 1$  and the group  $G$  is an elementary abelian  $p$ -group.

Conversely, suppose that a group  $G$  is an elementary abelian  $p$ -group, then we consider  $G$  as an additive group of a vector space over the field  $GF(p)$ .

Hence the exchange property is satisfied for a group  $G$ .

#### 4 Intersection between the basis property and the exchange property

**Example 4.1** Let  $S$  be the semilattice  $\{a, b, 0\}$ , where  $a, b$  are incomparable and  $ab = 0$ . Then  $S$  has unique basis, so  $S$  has basis property. But  $0 \in \langle \langle a \rangle \cup \{b\} \rangle$  and  $0 \notin \langle \langle a \rangle \rangle$ ,  $b \notin \langle \langle a \rangle \cup \{0\} \rangle$ . Hence  $S$  does not satisfy the exchange property.

**Example 4.2** Let  $G = \langle a \rangle$  be a cyclic group such that  $|G| = p^2$ ,  $p$  is a prime. Then  $G$  is a group with the basis property, because it is a  $p$ -group, but it does not satisfy the exchange property.

**Theorem 4.3** Let  $G$  be a finite group. Then  $G$  is a group with the exchange property if and only if one of the following conditions hold:

1.  $G$  is an elementary abelian  $p$ -group,  $p$  is a prime.
2.  $G$  is a semidirect product of an elementary abelian  $p$ -group  $P$  by a cyclic  $q$ -group  $\langle y \rangle$ , of order  $q$ , where  $p \neq q$  ( $p$  and  $q$  are primes). Therefore  $G$  must satisfy the following relations:

$$\begin{aligned} p &\equiv 1 \pmod{q}, \quad y^{-1}ay = a^r, \quad r \in \mathbb{Z}^+, \\ r &\not\equiv 1 \pmod{p}, \quad r^q \equiv 1 \pmod{p}. \end{aligned}$$

*Proof.* Suppose that  $G$  is a group with the exchange property. Then we consider two cases:

Firstly, if  $G$  is a primary group ( $p$ -group),  $p$  is a prime, then by theorem(3-3)  $G$  is an elementary abelian  $p$ -group for a prime  $p$ .

Secondly, if  $G$  is not primary group, then from the basis property in theorem(2-6), we see that  $G$  is a semidirect product (i.e.  $G = P \rtimes \langle y \rangle$ ) of  $p$ -group  $P$  by a cyclic  $q$ -group  $\langle y \rangle$ , where  $p \neq q$  ( $p$  and  $q$  are primes). Since  $P$  is a group with the exchange property, then by theorem(3-3)  $P$  is an elementary abelian  $p$ -group. Therefore by theorem(3-1) the group  $\langle y \rangle$  has order  $q$ ,  $q$  is a prime.

Suppose that  $|P| = p^s$ ,  $s \in \mathbb{Z}^+$ . Since the element  $y$  is regular operator on  $P$ , i.e. the operator  $\varphi$  inducing by element  $y$  is a regular, then

$$p^s \equiv 1 \pmod{q}.$$

Assume that  $a \in P$ ,  $a \neq 1$ . Consider the element  $b = y^{-1}ay$ , since the operator  $\varphi$  induced by element  $y$  is regular, then  $b \neq a$ . Assume that  $b \in \langle a \rangle$ , hence  $b = a^r$ ,  $r \not\equiv 1 \pmod{p}$ . From  $y^q = 1$  we have  $a^{r^q} = 1$ , i.e.  $r^q \equiv 1 \pmod{p}$ .

Now let  $b \notin \langle a \rangle$ , so by the exchange property if  $b \in \langle y, a \rangle$ , then  $y \in \langle a, b \rangle \leq P$ . We get a contradiction with  $y \notin P$ . Thus the automorphism  $\varphi_y : P \rightarrow P$  is regular and act on a group  $\langle a \rangle$  of order  $p$ , hence  $p \equiv 1 \pmod{q}$  and  $p > q$ . Since  $G$  is a group with the basis property, then by theorem(2-6) the representation  $y \rightarrow \varphi_y$  is an isotopic with dimension 1, i.e. the matrix  $A$  of linear operator  $\varphi_y$  in some basis of vector space  $P$  which contains  $s$  elements has the following form:

$$A = \begin{pmatrix} \bar{r} & 0 & \dots & 0 \\ 0 & \bar{r} & \dots & 0 \\ 0 & 0 & & \bar{r} \end{pmatrix},$$

such that  $\bar{r}$  is an image of the element  $r$  under the conical homomorphism  $\bar{\theta}: \mathbb{Z} \rightarrow \mathbb{Z}/p\mathbb{Z}$ , then

$$r \not\equiv 1 \pmod{p}, \text{ and}$$

$$r^q \equiv 1 \pmod{p}.$$

Conversely, If  $G$  is an elementary abelian  $p$ -group for a prime  $p$ , then  $G$  is a group with basis property. Using theorem(3-3), then it remains to prove that if  $P$  is an elementary abelian, and  $\langle y \rangle$  has order  $q$ , where  $p \neq q$  ( $p$  and  $q$  are primes), and if the following conditions hold

$$y^{-1}xy = x^r, \quad \forall x \in P,$$

$$p \equiv 1 \pmod{q},$$

$$r \not\equiv 1 \pmod{p},$$

$$r^q \equiv 1 \pmod{p}. \quad (4-1)$$

Then  $G$  is a group with the exchange property. Suppose that the set  $X \subseteq G$  and  $a, b \in G$  such that  $a \notin \langle X \rangle$  and  $a \in \langle X \cup \{b\} \rangle$ . Now we prove that  $b \in \langle X \cup \{a\} \rangle$ .

Let  $G_1 = \langle X \cup \{b\} \rangle$  and we study the following cases:

If  $\langle X \cup \{b\} \rangle \leq P$ , then  $G_1 \leq P$ ,  $G_1$  satisfies the exchange property, because it is an elementary abelian  $p$ -group (by theorem(3-3)), hence  $b \in \langle X \cup \{a\} \rangle$ .

If  $\langle X \cup \{b\} \rangle \not\leq P$ , then suppose that the set  $X \cup \{b\}$  contains element of order  $q$ . Now if  $X$  contains elements of order  $q$ , and since  $G$  is a semidirect product of  $p$ -group by cyclic  $\langle y \rangle$ . Then we can prove that the set  $X$  contains only one element of order  $q$ , (because if there exist two elements as  $y^{s_1}a_1$ ,  $y^{s_2}a_2$  in  $X$  of order  $q$ , then for some  $w \in \mathbb{Z}$  there exists  $c \in P$  such that

$$y^{s_2}a_2 = (y^{s_1}a_1)^w c.$$

Then  $\langle y^{s_1}a_1, c \rangle = \langle y^{s_1}a_1, y^{s_2}a_2 \rangle$ , hence we consider element  $y^{s_2}a_2$  as  $c \in P$ . Now suppose that the set  $X = \{x_1, x_2, \dots, x_n\}$  such that  $x_2, \dots, x_n \in P$ ,  $x_1 \notin P$ . Then the Fitting subgroup  $F(\langle X \rangle)$  of group  $\langle X \rangle$  is generated by the set  $\{x_2, \dots, x_n\}$  and the image of this set under the automorphism  $\varphi_{x_1}^m$ ,  $m \in \mathbb{Z}$ .

Since the group  $P$  is an abelian group, then the Fitting subgroup  $F(\langle X \rangle)$  is generated by the set  $\{x_2, \dots, x_n\}$  and the image of this set under the automorphism  $\varphi_y^m$  and by (4-1) this is the power of the same elements  $x_2, \dots, x_n$ . In another words, the group  $F(\langle X \rangle)$  is generated by  $x_2, \dots, x_n$  if these elements are exists. So by our assumption  $a \in \langle X \cup \{b\} \rangle$ . Then there exists a word  $u(x_1, x_2, \dots, x_n)$  such that  $a = u(x_1, x_2, \dots, x_n, b)$  and by (4-1) we have

$$a = v(x_1, x_2, \dots, x_n)b^w, \quad (4-2)$$



such that  $v(x_1, x_2, \dots, x_n)$  is a word. If  $b^w = e$ , then by (4-2) we have

$$a = v(x_1, x_2, \dots, x_n) \in \langle X \rangle .$$

Thus we get a contradiction with our assumption for  $a$ , so we assume that  $b^w \neq e$ . Since a group  $P$  is an elementary abelian  $p$ -group, then  $\langle b^w \rangle = \langle b \rangle$ , so by (4-2) we have

$$b \in \langle b^w \rangle = \langle v(x_1, x_2, \dots, x_n)^{-1} a \rangle \subseteq \langle X \cup \{a\} \rangle .$$

Finally, let  $X \subseteq P$ . Since  $X \cup \{b\} \not\subseteq P$ , then  $b$  is element of order  $q$ . Suppose that  $G_1 = \langle X \cup \{b\} \rangle$  is a semidirect product of a group  $\langle X \rangle$  by  $\langle b \rangle$ . Then from  $a \in \langle X \cup \{b\} \rangle$  we have the following for  $w \in \mathbb{Z}$  and  $c \in \langle X \rangle$

$$a = b^w c. \quad (4-3)$$

If an element  $a$  is a  $q$ -element, then  $b^w \neq e$  and since  $\langle b \rangle = \langle b^w \rangle$  we get

$$b \in \langle b^w \rangle = \langle c^{-1} a \rangle \subseteq \langle X \cup \{a\} \rangle .$$

If  $a$  is  $p$ -element, then by (4-3) we have  $b^w = e$  and  $a = c \in \langle X \rangle$  which is a contradiction with  $a \notin \langle X \rangle$ . Thus we study all cases. Hence  $G$  is a group with change property.

## 5 Acknowledgements

The authors gratefully acknowledge the partial financial support from the Deanship of Academic Research (Project No.331201 ) at Al Imam Mohammad Ibn Saud Islamic University (IMSIU), Riyadh, Saudi Arabia.

## References

- [1] A. Al Khalaf, Finite group with basis property, Dok. Acad. Nauk BSSR, n. 11, 1989 (Russian)
- [2] M. Alkadhi, A. Al Khalaf and M. Quick, The Nilpotency Class of Fitting Subgroups of Groups with Basis Property. International Journal of Algebra, Vol. 6, 2012, no. 14, 697 - 704.
- [3] A. Aljouiee, Basis Property Conditions on Some Groups, International Journal of Mathematics and Computer Science, 3, No3, 1-11, (2008), Lebanon.

- [4] Jonathan McDougall-Bagnall, Martyn Quick, Groups with the basis property, *Journal of Algebra* 346 (2011) 332-339.
- [5] P. R. Jones, A basis theorem for free inverse Semigroup, *J. Algebra*, v. 49, 1977. p.172-190.
- [6] P. R. Jones, Basis properties for inverse Semigroups, *J. Algebra*, v. 50, 1978. p.135-152.
- [7] P. R. Jones, Basis properties, exchange properties and embeddings in idempotent-free semigroups, *Semigroups and their applications*, 69-82, Reidel, Dordrecht, 1987.
- [8] P. R. Jones, Exchange properties and basis properties for closure operators, *Colloq. Math.* 57(1989), no. 1, 29-33.
- [9] M. Hall, *The Theory of Groups*, Macmillan, 1959.
- [10] B. Huppert, *Endliche Gruppen*, Springer-Verlag, 1967.
- [11] D. Robinson, *A Course in the Theory of Groups*, Second Edition. Springer-Verlag New York, Inc. (1996)

Al Khalaf Ahmad,  
Department of Mathematics, College of Sciences  
Al-Imam Mohammad Ibn Saud Islamic University  
P.O. Box 90950, Riyadh, Saudi Arabia.  
Email: akhalaf59@gmail.com

Alkadhi Mohammed  
Department of Mathematics, College of Sciences  
Al-Imam Mohammad Ibn Saud Islamic University  
P.O. Box 90950, Riyadh, Saudi Arabia.  
Email: malki2@yahoo.com