



Hypergroups associated with HX -groups

Piergiulio Corsini

Abstract

We have already studied a correspondence between HX -groups and hypergroups. Now we extend this correspondence to semigroups of subsets of a group G (and to begin, we considered $\mathbf{Z}_2 \times \mathbf{Z}_3$ as group G). Moreover these subsets can have non empty intersection, they can be contained, one in another one. One calculates the fuzzy grade and we find always $\partial H = 1$.

1 Introduction

In the classic theory of HX -groups, one supposes that the structure from which one starts, is a group G , which is a subgroup of the semigroup $\mathcal{P}^*(K)$ of nonempty subsets of a group K , see for instance [3], [20], [21], [22], [23], [33], [34] and often one supposes also that

$$(1) \quad \forall (A, B) \in \mathcal{P}^*(K) \times \mathcal{P}^*(K), \quad A \neq B \Rightarrow A \cap B = \emptyset.$$

In this paper one has not done always these hypotheses.

In a precedent paper [8] one considered the HX -groups associated with $\mathbf{Z}/n\mathbf{Z}$ for $n \in \{8, 9, 12, 15, 16\}$.

In this paper, we calculate those associated with $\mathbf{Z}_2 \times \mathbf{Z}_3$, without supposing always satisfied the hypothesis (1). In this new situation, we show that the fuzzy grade is always equal to one.

Let us see what means a "fuzzy grade".

Key Words: HX -group, Fuzzy grade.

2010 Mathematics Subject Classification: Primary 20N20; Secondary 20N25.

Received: March, 2016.

Revised: March, 2016.

Accepted: May, 2016.

Let (H_0, \circ) be a hypergroupoid.

Set $\forall(x, y) \in H^2$, $\forall u \in x \circ y$,

$$m_{x,y}(u) = 1/|x \circ y|, \quad A_1(u) = \sum_{(x,y) \in H^2} m_{x,y}(u)$$

$$Q_1(u) = \{(a, b) \mid u \in a \circ b\}, \quad q_1(U) = |Q_1(U)| \quad \text{and} \quad \mu_1(u) = A_1(u)/q_1(u).$$

Set now $\forall(x, y) \in H^2$,

$$x \circ_1 y = \{z \mid \min\{\mu_1(x), \mu_1(y)\} \leq \mu_1(z) \leq \max\{\mu_1(x), \mu_1(y)\}\}$$

and set denote by H_1 the hypergroupoid (H_0, \circ_1) (which is a join space, see [4]). One goes on in the same way and from H_1 one obtains A_2, q_2, μ_2 defined similarly as A_1, q_1, μ_1 , but starting from H_1 . One obtains another join space (H_2, \circ_2) and so on. The *fuzzy grade* of H_0 , denoted by ∂H_0 is the least k , such that H_k is isomorphic to H_{k+1} .

2 HX -groups on finite groups

If (G, \cdot) is a group and $\mathcal{P}^*(G)$ is the set of all non empty subsets of G then an HX -group is a nonempty subset H of $\mathcal{P}^*(G)$, which is a group, with respect to the hyperoperation (see [3], [20], [22], [21], [23], [33], [34]):

$$\forall(A, B) \in \mathcal{P}^*(G) \times \mathcal{P}^*(G), \quad A \cdot B = \{xy \mid x \in A, y \in B\}.$$

Let us consider $A_0 = \mathbf{Z}_2 \times \mathbf{Z}_3 = \{(0,0), (0,1), (0,2), (1,0), (1,1), (1,2)\}$.
Set

$$B_0 = \{(0,0), (1,0)\}, \quad B_1 = \{(0,0), (0,1), (0,2)\}, \quad B_2 = \{(0,0), (0,2)\},$$

$$B_3 = \{(0,1), (1,1)\}, \quad B_4 = \{(0,2), (1,2)\}, \quad B_5 = \{(1,0), (1,1), (1,2)\}.$$

We find

$B_{1,5}$	B_1	B_5	$B_{0,3,4}$	B_0	B_3	B_4
B_1	B_1	B_5	B_0	B_0	B_3	B_4
B_5	B_5	B_1	B_3	B_4	B_0	B_3

From $B_{1,5}$ we obtain:

$B_{1,5}$	(0, 0)	(0, 1)	(0, 2)	(1, 0)	(1, 1)	(1, 2)
(0, 0)	B_1	B_1	B_1	B_5	B_5	B_5
(0, 1)		B_1	B_1	B_5	B_5	B_5
(0, 2)			B_1	B_5	B_5	B_5
(1, 0)				B_1	B_1	B_1
(1, 1)					B_1	B_1
(1, 2)						B_1

$$B_6 = \{(0, 0), (1, 0), (0, 2), (1, 2)\}, \quad B_7 = \{(0, 1), (0, 2), (1, 1), (1, 2)\},$$

$$B_8 = \{(0, 0), (0, 1), (1, 0), (1, 1)\}.$$

$B_{1,5}$ can be written also

$H_{1,5}^0$	(0, 0)	(0, 1)	(0, 2)	(1, 0)	(1, 1)	(1, 2)
(0, 0)	(0, 0)	(0, 0)	(0, 0))	(1, 0)	(1, 0)	(1, 0)
	(0, 1)	(0, 1)	(0, 1)	(1, 1)	(1, 1)	((1, 1)
	(0, 2)	(0, 2)	(0, 2)	(1, 2)	(1, 2)	(1, 2)
(0, 1)		(0, 0)	(0, 0))	(1, 0)	(1, 0)	(1, 0)
		(0, 1)	(0, 1)	(1, 1)	(1, 1)	((1, 1)
		(0, 2)	(0, 2)	(1, 2)	(1, 2)	(1, 2)
(0, 2)			(0, 0)	(1, 0)	(1, 0)	(1, 0)
			(0, 1)	(1, 1)	(1, 1)	((1, 1)
			(0, 2)	(1, 2)	(1, 2)	(1, 2)
(1, 0)				(0, 0)	(0, 0)	(0, 0)
				(0, 1)	(0, 1)	((0, 1)
				(0, 2)	(1, 2)	(1, 2)
(1, 1)					(0, 0)	(0, 0)
					(0, 1)	((0, 1)
					(0, 2)	(1, 2)
(1, 2)						(0, 0)
						((0, 1)
						(0, 2)

We deduce $A_1(0, 0) = A_1(0, 1) = A_1(0, 2) = 29/3$,

$$q_1(0, 0) = q_1(0, 1) = q_1(0, 2) = 18,$$

$$\mu_1(0, 0) = \mu_1(0, 1) = \mu_1(0, 2) = 1/3.$$

$$A_1(1, 0) = A_1(1, 1) = A_1(1, 2) = 18/3,$$

$$q_1(1, 0) = q_1(1, 1) = q_1(1, 2) = 18,$$

$$\mu_1(1, 0) = \mu_1(1, 1) = \mu_1(1, 2) = 1/3.$$

It follows that $\partial H_{1,5}^0 = 1$.

$B_{2,1,5}$	B_2	B_1	B_5
B_2	B_1	B_1	B_5
B_1	B_1	B_1	B_5
B_5	B_5	B_5	B_1

One obtains $A_1(0,0) = A_1(0,1) = A_1(0,2) = 25/3 + 9/3 = 34/3$,

$$A_1(1,0) = A_1(1,1) = A_1(1,2) = 30/3.$$

$$q_1(0,0) = q_1(0,1) = q_1(0,2) = 34, \quad q_1(1,0) = q_1(1,1) = q_1(1,2) = 30.$$

$H_{2,1,5}$	(0,0)	(0,2)	(0,0)	(0,1)	(0,2)	(1,0)	(1,1)	(1,2)
(0,0)	(0,0)	(0,0)	(0,0))	(0,0)	(0,0)	(1,0)	(1,0)	(1,0)
	(0,2)	(0,2)	(0,2)	(0,1)	(0,1)	((1,1)	(1,1)	(1,1)
	(0,1)	(0,1)	(0,1)	(0,2)	(0,2)	(1,2)	(1,2)	(1,2)
(0,2)	(0,0)	(0,0)	(0,0))	(0,0)	(0,0)	(1,0)	(1,0)	(1,0)
	(0,2)	(0,2)	(0,2)	(0,1)	(0,1)	((1,1)	(1,1)	(1,1)
	(0,1)	(0,1)	(0,1)	(0,2)	(0,2)	(1,2)	(1,2)	(1,2)
(0,0)	(0,0)	(0,0)	(0,0))	(0,0)	(0,0)	(1,0)	(1,0)	(1,0)
	(0,1)	(0,1)	(0,1)	(0,1)	(0,1)	((1,1)	(1,1)	(1,1)
	(0,2)	(0,2)	(0,2)	(0,2)	(0,2)	(1,2)	(1,2)	(1,2)
(0,1)	(0,0)	(0,0)	(0,0))	(0,0)	(0,0)	(1,0)	(1,0)	(1,0)
	(0,1)	(0,1)	(0,1)	(0,1)	(0,1)	((1,1)	(1,1)	(1,1)
	(0,2)	(0,2)	(0,2)	(0,2)	(0,2)	(1,2)	(1,2)	(1,2)
(0,2)	(0,0)	(0,0)	(0,0))	(0,0)	(0,0)	(1,0)	(1,0)	(1,0)
	(0,1)	(0,1)	(0,1)	(0,1)	(0,1)	((1,1)	(1,1)	(1,1)
	(0,2)	(0,2)	(0,2)	(0,2)	(0,2)	(1,2)	(1,2)	(1,2)
(1,0)	(1,0)	(1,0)	(1,0))	(1,0)	(1,0)	(0,0)	(0,0)	(0,0)
	(1,1)	(1,1)	(1,1)	(1,1)	(1,1)	((0,1)	(0,1)	(0,1)
	(1,2)	(1,2)	(1,2)	(1,2)	(1,2)	(0,2)	(0,2)	(0,2)
(1,1)	(1,0)	(1,0)	(1,0))	(1,0)	(1,0)	(0,0)	(0,0)	(0,0)
	(1,1)	(1,1)	(1,1)	(1,1)	(1,1)	((0,1)	(0,1)	(0,1)
	(1,2)	(1,2)	(1,2)	(1,2)	(1,2)	(0,2)	(0,2)	(0,2)
(1,2)	(1,0)	(1,0)	(1,0))	(1,0)	(1,0)	(0,0)	(0,0)	(0,0)
	(1,1)	(1,1)	(1,1)	(1,1)	(1,1)	((0,1)	(0,1)	(0,1)
	(1,2)	(1,2)	(1,2)	(1,2)	(1,2)	(0,2)	(0,2)	(0,2)

By consequence, $\mu_1(i,j) = 1/3$, $\forall (i,j)$ so $H_1 = T$ (the total hypergroup), whence $\partial(H_{2,1,5}) = 1$.

From $B_{0,3,4}$ we obtain

$H_{0,3,4}^1$	(0, 0)	(1, 0)	(0, 1)	(0, 2)	(1, 1)	(1, 2)
(0, 0)	B_0	B_0	B_3	B_4	B_3	B_4
(1, 0)		B_0	B_3	B_4	B_3	B_4
(0, 1)			B_4	B_0	B_4	B_0
(0, 2)				B_3	B_0	B_3
(1, 1)					B_4	B_0
(1, 2)						B_3

$H_{0,3,4}^1$ can be written also in the following way:

$H_{0,3,4}^0$	(0, 0)	(1, 0)	(0, 1)	(0, 2)	(1, 1)	(1, 2)
(0, 0)	(0, 0)	(0, 0)	(0, 1)	(1, 2)	(0, 1)	(1, 2)
	(1, 0)	(1, 0)	(1, 1)	(0, 2)	(1, 1)	((0, 2))
(1, 0)		(0, 0)	(0, 1)	(1, 2)	(0, 1)	(1, 2)
		(1, 0)	(1, 1)	(0, 2)	(1, 1)	((0, 2))
(0, 1)			(1, 2)	(0, 0)	(1, 2)	(0, 0)
			(0, 2)	(1, 0)	(0, 2)	(1, 0)
(0, 2)				(0, 1)	(0, 0)	(0, 1)
				(1, 1)	(1, 0)	(1, 1)
(1, 1)					(0, 2)	(0, 0)
					(1, 2)	(1, 0)
(1, 2)						(0, 1)
						(1, 1)

$$A_1(0, 0) = 2/2 + 8/2 = A_1(1, 0) = 5,$$

$$A_1(0, 1) = 2/2 + 10/2 = A_1(1, 1) = 6,$$

$$A_1(0, 2) = 2/2 + 10/2 = A_1(1, 2) = 6.$$

We obtain $q_1(0, 0) = q_1(1, 0) = 10$, $q_1(0, 1) = q_1(1, 1) = q_1(0, 2) = q_1(1, 2) = 12$, whence

$$\mu_1(0, 0) = 1/2, \mu_1(1, 0) = 1/2$$

$\mu_1(0, 1) = \mu_1(1, 1) = \mu_1(0, 2) = \mu_1(1, 2) = 1/2$. Therefore, H^1 is total, whence $\partial(H_{0,3,4}^0) = 1$.

One considers now the semigroup constructed on the set

$$\{B_i \mid 0 \leq i \leq 8\} \cup \{A_0\}.$$

$$B_1 = \{(0, 0), (0, 1), (0, 2)\},$$

$$B_2 = \{(0, 0), (0, 2)\},$$

$$B_5 = \{(1, 0), (1, 1), (1, 2)\},$$

$$B_0 = \{(0, 0), (1, 0)\},$$

$$B_3 = \{(0, 1), (1, 1)\},$$

$$\begin{aligned}B_4 &= \{(0, 2), (1, 2)\}, \\B_6 &= \{(0, 0), (1, 0), (0, 2), (1, 2)\}, \\B_7 &= \{(0, 1), (0, 2), (1, 1), (1, 2)\}, \\B_8 &= \{(0, 0), (0, 1), (1, 0), (1, 1)\}.\end{aligned}$$

We have

$$\begin{aligned}B_1B_6 &= \{(0, 0), (0, 1), (0, 2)\} \cdot \{(0, 0), (1, 0), (0, 2), (1, 2)\} = \\&\quad \{(0, 0), (1, 0), (0, 2), (1, 1) \mid (0, 1), (1, 1), (0, 0), (1, 0)\} = \\&\quad \{(0, 2), (1, 2), (0, 1), (1, 1)\} = A_0 \\B_1B_7 &= \{(0, 0), (0, 1), (0, 2)\} \cdot \{(0, 1), (0, 2), (1, 1), (1, 2)\} = \\&\quad \{(0, 1), (0, 2), (1, 1), (1, 2) \mid (0, 2), (0, 0), (1, 2), (1, 0)\} = A_0 \\B_1B_8 &= \{(0, 0), (0, 1), (0, 2)\} \cdot \{(0, 0), (0, 1), (1, 0), (1, 1)\} = \\&\quad \{(0, 0), (0, 1), (1, 0), (1, 1) \mid (0, 1), (0, 2), (1, 1), (1, 2)\} = A_0 \\B_2B_5 &= \{(0, 0), (0, 2)\} \cdot \{(1, 0), (1, 1)\} = \\&\quad \{(1, 0), (1, 1) \mid (1, 2), (1, 0)\} = B_5 \\B_2B_0 &= \{(0, 0), (0, 2)\} \cdot \{(0, 0), (1, 0)\} = \\&\quad \{(0, 0), (1, 0) \mid (0, 2), (1, 2)\} = B_6 \\B_2B_3 &= \{(0, 0), (0, 2)\} \cdot \{(0, 1), (1, 1)\} = \\&\quad \{(0, 1), (1, 1) \mid (0, 0), (1, 0)\} = B_8 \\B_2B_4 &= \{(0, 0), (0, 2)\} \cdot \{(0, 2), (1, 2)\} = \\&\quad \{(0, 2), (1, 2) \mid (0, 1), (1, 1)\} = B_7 \\B_2B_6 &= \{(0, 0), (0, 2)\} \cdot \{(0, 0), (1, 0), (0, 2), (1, 2)\} = \\&\quad \{(0, 0), (1, 0), (0, 2), (1, 2) \mid (0, 2), (1, 2), (0, 1), (1, 1)\} = \\&\quad \{(0, 0), (0, 1), (0, 2), (1, 0), (1, 1), (1, 2)\} = A_0 \\B_2B_7 &= \{(0, 0), (0, 2)\} \cdot \{(0, 1), (0, 2), (1, 1), (1, 2)\} = \\&\quad \{(0, 1), (0, 2), (1, 1), (1, 2) \mid (0, 0), (0, 1), (1, 0), (1, 1)\} = A_0 \\B_2B_8 &= \{(0, 0), (0, 2)\} \cdot \{(0, 0), (0, 1), (1, 0), (1, 1)\} = \\&\quad \{(0, 0), (0, 1), (1, 0), (1, 1) \mid (0, 2), (0, 0), (1, 2), (1, 0)\} = A_0 \\B_5B_0 &= \{(1, 0), (1, 1), (1, 2)\} \cdot \{(0, 0), (1, 0)\} = \\&\quad \{(1, 0), (1, 1), (1, 2) \mid (0, 0), (0, 1), (0, 2)\} = A_0 \\B_5B_3 &= \{(1, 0), (1, 1), (1, 2)\} \cdot \{(0, 1), (1, 1)\} = \\&\quad \{(1, 1), (0, 1) \mid (1, 2), (0, 2) \mid (1, 0), (0, 0)\} = A_0 \\B_5B_4 &= \{(1, 0), (1, 1), (1, 2)\} \cdot \{(0, 2), (1, 2)\} \\&\quad \{(1, 2), (0, 2) \mid (1, 0), (0, 0), (1, 1), (0, 1)\} = A_0 \\&\text{Since } B_4 \subseteq B_6, \text{ it follows that } B_5B_6 = A_0. \\&\text{Since } B_3 \subseteq B_7, \text{ it follows that } B_5B_7 = A_0. \\&\text{Since } B_0 \subseteq B_8, \text{ it follows that } B_5B_8 = A_0. \\B_0B_0 &= \{(0, 0), (1, 0)\} \cdot \{(0, 0), (1, 0)\} \\&\quad \{(0, 0), (1, 0) \mid (1, 0), (0, 0)\} = B_0 \\B_0B_3 &= \{(0, 0), (1, 0)\} \cdot \{(0, 1), (1, 1)\} \\&\quad \{(0, 1), (1, 1) \mid (1, 1), (0, 1)\} = \{(0, 1), (1, 1)\} = B_3 \\B_0B_4 &= \{(0, 0), (1, 0)\} \cdot \{(0, 2), (1, 2)\}\end{aligned}$$

$$\begin{aligned}
& \{(0, 2), (1, 2) \mid (1, 2), (0, 2)\} = \{(0, 2), (1, 2)\} = B_4 \\
& B_0B_5 = \{(0, 0), (1, 0)\} \cdot \{(1, 0), (1, 1), (1, 2)\} \\
& \{(1, 0), (1, 1), (1, 2) \mid (0, 0), (0, 1), (0, 2)\} = A_0 \\
& B_0B_6 = \{(0, 0), (1, 0)\} \cdot \{(0, 0), (1, 0), (0, 2), (1, 2)\} \\
& \{(0, 0), (1, 0), (0, 2), (1, 2) \mid (1, 0), (0, 0), (1, 2), (0, 2)\} = B_6 \\
& B_0B_7 = \{(0, 0), (1, 0)\} \cdot \{(0, 1), (0, 2), (1, 1), (1, 2)\} = \\
& \{(0, 1), (0, 2), (1, 1), (1, 2) \mid (1, 1), (1, 2), (0, 1), (0, 2)\} = B_7 \\
& B_0B_8 = \{(0, 0), (1, 0)\} \cdot \{(0, 0), (0, 1), (1, 0), (1, 1)\} = \\
& \{(0, 0), (0, 1), (1, 0), (1, 1) \mid (1, 0), (1, 1), (0, 0), (0, 1)\} = B_8 \\
& B_3B_6 = \{(0, 1), (1, 1)\} \cdot \{(0, 0), (1, 0), (0, 2), (1, 2)\} = \\
& \{(0, 1), (1, 1), (1, 1), (0, 1) \mid (0, 0), (1, 0), (1, 0), (0, 0)\} = B_8 \\
& B_3B_7 = \{(0, 1), (1, 1)\} \cdot \{(0, 1), (0, 2), (1, 1), (1, 2)\} = \\
& \{(0, 2), (0, 0), (1, 2), (1, 0) \mid (1, 2), (1, 0), (0, 2), (0, 0)\} = B_6 \\
& B_6B_6 = \{(0, 0), (1, 0), (0, 2), (1, 2)\} \cdot \{(0, 0), (1, 0), (0, 2), (1, 2)\} = \\
& \{(0, 0), (1, 0), (0, 2), (1, 2) \mid (1, 0), (0, 0), (1, 2), (0, 2)\} = \\
& \{(0, 2), (1, 2), (0, 1), (1, 1) \mid (1, 2), (0, 2), (1, 1), (0, 1)\} = A_0 \\
& B_4B_8 = \{(0, 2), (1, 2)\} \cdot \{(0, 0), (0, 1), (1, 0), (1, 1)\} = \\
& \{(0, 2), (0, 0), (1, 2), (1, 0) \mid (1, 2), (1, 0), (0, 2), (0, 0)\} = B_6 \\
& B_6B_7 = \{(0, 0), (1, 0), (0, 2), (1, 2)\} \cdot \{(0, 1), (0, 2), (1, 1), (1, 2)\} = \\
& \{(0, 0), (0, 1), (1, 0), (1, 1) \mid (1, 0), (1, 1), (0, 0), (0, 1)\} = A_0 \\
& B_3B_8 = \{(0, 1), (1, 1)\} \cdot \{(0, 0), (0, 1), (1, 0), (1, 1)\} = \\
& \{(0, 1), (0, 2), (1, 1), (1, 2) \mid (1, 1), (1, 2), (0, 1), (0, 2)\} = B_7 \\
& B_4B_6 = \{(0, 2), (1, 2)\} \cdot \{(0, 0), (1, 0), (0, 2), (1, 2)\} = \\
& \{(0, 2), (1, 2), (0, 1), (1, 1) \mid (1, 2), (0, 2), (1, 1), (0, 1)\} = B_7 \\
& B_4B_7 = \{(0, 2), (1, 2)\} \cdot \{(0, 1), (0, 2), (1, 1), (1, 2)\} = \\
& \{(0, 0), (0, 1), (1, 0), (1, 1) \mid (1, 0), (1, 1), (0, 0), (0, 1)\} = B_8 \\
& B_6B_8 = \{(0, 0), (1, 0), (0, 2), (1, 2)\} \cdot \{(0, 0), (0, 1), (1, 0), (1, 1)\} = \\
& \{(0, 0), (0, 1), (1, 0), (1, 1) \mid (1, 0), (1, 1), (0, 0), (0, 1) \mid \\
& (0, 2), (0, 0), (1, 2), (1, 0) \mid (1, 2), (1, 0), (0, 2), (0, 0)\} = A_0 \\
& B_7B_7 = \{(0, 1), (0, 2), (1, 1), (1, 2)\} \cdot \{(0, 1), (0, 2), (1, 1), (1, 2)\} = \\
& \{(0, 2), (0, 0), (1, 2), (1, 0) \mid (0, 0), (0, 1), (1, 0), (1, 1) \mid \\
& (1, 2), (1, 0), (0, 2), (0, 0) \mid (1, 0), (1, 1), (0, 0), (0, 1)\} = A_0 \\
& B_7B_8 = \{(0, 1), (0, 2), (1, 1), (1, 2)\} \cdot \{(0, 0), (0, 1), (1, 0), (1, 1)\} = \\
& \{(0, 1), (0, 2), (1, 1), (1, 2) \mid (0, 2), (0, 0), (1, 2), (1, 0) \mid \\
& (1, 1), (1, 2), (0, 1), (0, 2) \mid (1, 2), (1, 0), (0, 2), (0, 0)\} = A_0 \\
& B_8B_8 = \{(0, 0), (0, 1), (1, 0), (1, 1)\} \cdot \{(0, 0), (0, 1), (1, 0), (1, 1)\} = \\
& \{(0, 0), (0, 1), (1, 0), (1, 1) \mid (0, 1), (0, 2), (1, 1), (1, 2) \mid \\
& (1, 0), (1, 1), (0, 0), (0, 1) \mid (1, 1), (1, 2), (0, 1), (0, 2)\} = A_0
\end{aligned}$$

B^*	B_1	B_2	B_5	B_0	B_3	B_4	B_6	B_7	B_8	A_0
B_1	B_1	B_1	B_5	A_0						
B_2		B_1	B_5	B_6	B_8	B_7	A_0	A_0	A_0	A_0
B_5			B_1	A_0						
B_0				B_0	B_3	B_4	B_6	B_7	B_8	A_0
B_3					B_4	B_0	B_8	B_6	B_7	A_0
B_4						B_3	B_7	B_8	B_6	A_0
B_6							A_0	A_0	A_0	A_0
B_7								A_0	A_0	A_0
B_8									A_0	A_0
A_0										A_0

We shall consider now the hypergroupoid B^* on $\{B_i \mid 0 \leq i \leq 8\} \cup \{A_0\}$ and we shall study the corresponding hypergroupoid on

$$H^* = \{(0,0), (0,2), (1,0), (1,2), (1,1), (0,1)\}.$$

From B^* we obtain the following H^* , which we divide into two tables, H^{*1} and H^{*2} .

We first do the following notations:

denote the elements $(0,0), (0,1), (0,2)$ of B_1 by $00_{B_1}, 01_{B_1}, 02_{B_1}$, respectively;

denote the elements $(0,0), (0,2)$ of B_2 by $00_{B_2}, 02_{B_2}$, respectively;

denote the elements $(1,0), (1,1), (1,2)$ of B_5 by $10_{B_5}, 11_{B_5}, 12_{B_5}$, respectively;

denote the elements $(0,0), (1,0)$ of B_0 by $00_{B_0}, 10_{B_0}$, respectively;

denote the elements $(0,1), (1,1)$ of B_3 by $01_{B_3}, 11_{B_3}$, respectively;

denote the elements $(0,2), (1,2)$ of B_4 by $02_{B_4}, 12_{B_4}$, respectively;

denote the elements $(0,0), (1,0), (0,2), (1,2)$ of B_6 by $00_{B_6}, 10_{B_6}, 02_{B_6}, 12_{B_6}$, respectively;

denote the elements $(0,1), (0,2), (1,1), (1,2)$ of B_7 by $01_{B_7}, 02_{B_7}, 11_{B_7}, 12_{B_7}$,

respectively and

H^{*2}	00_{B_6}	10_{B_6}	02_{B_6}	12_{B_6}	01_{B_7}	02_{B_7}	11_{B_7}	12_{B_7}	00_{B_8}	01_{B_8}	10_{B_8}	11_{B_8}	A_0
00_{B_1}	A_0	A_0											
01_{B_1}	A_0	A_0											
02_{B_1}	A_0	A_0											
00_{B_2}	A_0	A_0											
02_{B_2}	A_0	A_0											
10_{B_5}	A_0	A_0											
11_{B_5}	A_0	A_0											
12_{B_5}	A_0	A_0											
00_{B_0}	B_6	B_6	B_6	B_6	B_7	B_7	B_7	B_7	B_8	B_8	B_8	B_8	A_0
10_{B_0}	B_6	B_6	B_6	B_6	B_7	B_7	B_7	B_7	B_8	B_8	B_8	B_8	A_0
01_{B_3}	B_8	B_8	B_8	B_8	B_6	B_6	B_6	B_6	B_7	B_7	B_7	B_7	A_0
11_{B_3}	B_8	B_8	B_8	B_8	B_6	B_6	B_6	B_6	B_7	B_7	B_7	B_7	A_0
02_{B_4}	B_7	B_7	B_7	B_7	B_8	B_8	B_8	B_8	B_6	B_6	B_6	B_6	A_0
12_{B_4}	B_7	B_7	B_7	B_7	B_8	B_8	B_8	B_8	B_6	B_6	B_6	B_6	A_0
00_{B_6}	A_0	A_0											
10_{B_6}	A_0	A_0											
02_{B_6}	A_0	A_0											
12_{B_6}	A_0	A_0											
01_{B_7}	A_0	A_0											
02_{B_7}	A_0	A_0											
11_{B_7}	A_0	A_0											
12_{B_7}	A_0	A_0											
00_{B_8}	A_0	A_0											
01_{B_8}	A_0	A_0											
10_{B_8}	A_0	A_0											
11_{B_8}	A_0	A_0											
A_0	A_0												

Let us calculate now the $\mu_1(i, j)$ for $i \in \{0, 1\}$, $j \in \{0, 1, 2\}$.

We have

$$A_1(0, 1) = 25/3 + 18 \cdot 3 \cdot 2/6 + 2 \cdot 4/4 + 2 \cdot 4/4 + 12 \cdot 2 \cdot 2/6 + 9/3 + 18 \cdot 3 \cdot 2/6 + 2 \cdot 8/4 + 2 \cdot 8/4 + 2 \cdot 8/4 + 4/2 + 2 \cdot 8/4 + 2 \cdot 8/4 + 144/6 = 101 + 25/3 = 328/3,$$

which corresponds to the sets:

- $B_1 B_1, B_2 B_2, B_1 B_2$ divided by $|B_1|$;
- $B_1(B_0, B_3, B_4, B_6, B_7, B_8)$ divided by $|A_0|$;
- $B_2 B_3$ divided by $|B_8|$;
- $B_2 B_4$ divided by $|B_7|$;
- $B_2(B_6, B_7, B_8)$ divided by $|A_0|$;
- $B_5 B_5$ divided by $|B_1|$;
- $B_1(B_0, B_3, B_4, B_5, B_6, B_7, B_8)$ divided by $|A_0|$;
- $B_0 B_7$ divided by $|B_7|$;
- $B_0 B_8$ divided by $|B_8|$;
- $B_4 B_6$ divided by $|B_7|$;
- $B_4 B_7$ divided by $|B_8|$;

B_4B_4 divided by $|B_3|$;

B_3B_6 divided by $|B_8|$;

B_3B_8 divided by $|B_7|$;

and 144 divided by $|A_0|$, respectively. So, $A_1(0, 1) = 328/3$.

$$q_1(0, 1) = 25 + 108 + 8 + 8 + 48 + 9 + 108 + 16 + 16 + 16 + 16 + 16 + 16 + 144 = \\ = 558;$$

$$\mu_1(0, 1) = 328/(3 \cdot 558) = 0.1959.$$

$$A_1(1, 1) = 2 \cdot 9/3 + 18 \cdot 3 \cdot 2/6 + 2 \cdot 2 \cdot 3/3 + 2 \cdot 4/4 + 2 \cdot 4/4 + 12 \cdot 2 \cdot 2/6 + 9/3 + \\ 18 \cdot 3 \cdot 2/6 + 4/2 + 2 \cdot 8/4 + 144/6 = \\ 6 + 4 + 18 + 2 + 2 + 8 + 3 + 18 + 4 + 4 + 4 + 2 + 4 + 4 + 24 = 111, \text{ which} \\ \text{corresponds to the sets:}$$

B_1B_5 divided by $|B_5|$;

$B_1(B_0, B_3, B_4, B_6, B_7, B_8)$ divided by $|A_0|$;

B_2B_5 divided by $|B_5|$;

B_2B_3 divided by $|B_8|$;

B_2B_4 divided by $|B_7|$;

$B_2(B_6, B_7, B_8)$ divided by $|A_0|$;

B_5B_5 divided by $|B_1|$;

$B_1(B_0, B_3, B_4, B_6, B_7, B_8)$ divided by $|A_0|$;

B_4B_4 divided by $|B_3|$;

B_3B_6 divided by $|B_8|$;

B_3B_8 divided by $|B_7|$;

B_4B_6 divided by $|B_7|$;

B_4B_7 divided by $|B_8|$;

B_0B_7 divided by $|B_7|$;

B_0B_8 divided by $|B_8|$;

and 144 divided by $|A_0|$, respectively. So, $A_1(1, 1) = 111$.

$$q(1, 1) = 18 + 12 + 108 + 8 + 8 + 48 + 9 + 108 + 16 + 16 + 16 + 16 + 4 + 16 + 16 + 144 = \\ 563$$

$$\mu_1(1, 1) = 111/563 = 0.19716.$$

$$A_1(1, 0) = 2 \cdot 9/3 + 18 \cdot 3 \cdot 2/6 + 2 \cdot 3 \cdot 2/3 + 2 \cdot 4/4 + 2 \cdot 4/4 + 12 \cdot 2 \cdot 2/6 + 18 \cdot 3 \cdot 2/6 + \\ 4/2 + 2 \cdot 4/4 + 2 \cdot 8/4 + 2 \cdot 8/4 + 2 \cdot 4/2 + 2 \cdot 8/4 + 2 \cdot 8/4 + 2 \cdot 8/4 + 144/6 = 114,$$

which corresponds to the sets:

B_1B_5 divided by $|B_5|$;

$B_1(B_0, B_3, B_4, B_6, B_7, B_8)$ divided by $|A_0|$;

B_2B_5 divided by $|B_5|$;

B_2B_0 divided by $|B_6|$;

B_2B_3 divided by $|B_8|$;

$B_2(B_6, B_7, B_8)$ divided by $|A_0|$;

$B_5(B_0, B_3, B_4, B_6, B_7, B_8)$ divided by $|A_0|$;

B_0B_0 divided by $|B_0|$;

B_0B_4 divided by $|B_6|$;
 B_0B_6 divided by $|B_6|$;
 B_0B_8 divided by $|B_8|$;
 B_3B_4 divided by $|B_0|$;
 B_3B_6 divided by $|B_8|$;
 B_3B_7 divided by $|B_6|$;
 B_4B_7 divided by $|B_8|$;
 B_4B_8 divided by $|B_6|$

and 144 divided by $|A_0|$, respectively. So, $A_1(1, 0) = 114$.

$$q_1(1, 0) = 18 + 108 + 12 + 8 + 8 + 48 + 108 + 4 + 8 + 16 + 16 + 16 + 8 + 16 + 16 + 16 + 144 = 570;$$

$$\mu_1(1, 0) = 114/570 = 0.2.$$

$$A_1(1, 2) = 2 \cdot 9/3 + 18 \cdot 3 \cdot 2/6 + 2 \cdot 3 \cdot 2/3 + 2 \cdot 4/4 + 2 \cdot 4/4 + 12 \cdot 2 \cdot 2/6 + 18 \cdot 3 \cdot 2/6 + 2 \cdot 4/2 + 2 \cdot 4/4 + 2 \cdot 8/4 + 2 \cdot 8/4 + 2 \cdot 8/4 + 2 \cdot 8/4 + 4/4 + 4/2 + 144/6 = 114,$$

which corresponds to the sets:

B_1B_5 divided by $|B_5|$;
 $B_1(B_0, B_3, B_4, B_6, B_7, B_8)$ divided by $|A_0|$;
 B_2B_5 divided by $|B_5|$;
 B_2B_0 divided by $|B_6|$;
 B_2B_4 divided by $|B_7|$;
 $B_2(B_6, B_7, B_8)$ divided by $|A_0|$;
 $B_5(B_0, B_3, B_4, B_6, B_7, B_8)$ divided by $|A_0|$;
 B_0B_3 divided by $|B_4|$;
 B_0B_4 divided by $|B_6|$;
 B_0B_6 divided by $|B_6|$;
 B_0B_7 divided by $|B_7|$;
 B_4B_6 divided by $|B_7|$;
 B_4B_8 divided by $|B_6|$;
 B_3B_7 divided by $|B_6|$;
 B_3B_8 divided by $|B_7|$;
 B_3B_3 divided by $|B_4|$

and 144 divided by $|A_0|$, respectively. So, $A_1(1, 2) = 114$.

$$q_1(1, 2) = 18 + 108 + 12 + 8 + 8 + 48 + 108 + 8 + 8 + 16 + 16 + 16 + 16 + 16 + 4 + 144 = 570;$$

$$\mu_1(1, 2) = 114/570 = 0.2.$$

$$A_1(0, 0) = 25/3 + 18 \cdot 2 \cdot 3/6 + 4 \cdot 2/4 + 4 \cdot 2/4 + 12 \cdot 2 \cdot 2/6 + 9/3 + 18 \cdot 3 \cdot 2/6 + 4/2 + 2 \cdot 4/4 + 2 \cdot 8/4 + 2 \cdot 8/4 + 2 \cdot 4/2 + 2 \cdot 8/4 + 2 \cdot 8/4 + 2 \cdot 8/4 + 144/6 = 346/3,$$

$$q_1(0, 0) = 25 + 108 + 8 + 8 + 48 + 9 + 108 + 4 + 8 + 16 + 16 + 8 + 16 + 16 + 16 + 16 + 144 = 574;$$

$$\mu_1(0, 0) = 346/(3 \cdot 574) = 0.2009.$$

$A_1(0, 2) = 25/3 + 18 \cdot 2 \cdot 3/6 + 4 \cdot 2/4 + 4 \cdot 2/4 + 12 \cdot 2 \cdot 2/6 + 9/3 + 18 \cdot 3 \cdot 2/6 + 2 \cdot 4/4 + 2 \cdot 4/2 + 2 \cdot 8/4 + 2 \cdot 8/4 + 2 \cdot 8/2 + 2 \cdot 8/4 + 2 \cdot 8/4 + 2 \cdot 8/4 + 144/6 = 346/3$, which corresponds to the sets:

$(B_1 + B_2) \cdot (B_1 + B_2)$ divided by $|B_1|$;
 $B_1(B_0, B_3, B_4, B_6, B_7, B_8)$ divided by $|A_0|$;
 B_2B_0 divided by $|B_6|$;
 B_2B_4 divided by $|B_7|$;
 $B_2(B_6, B_7, B_8)$ divided by $|A_0|$;
 B_5, B_5 divided by $|B_1|$;
 $B_5(B_0, B_3, B_4, B_6, B_7, B_8)$ divided by $|A_0|$;
 B_0B_4 divided by $|B_6|$;
 B_0B_3 divided by $|B_4|$;
 B_0B_6 divided by $|B_6|$;
 B_0B_7 divided by $|B_7|$;
 B_3B_7 divided by $|B_6|$;
 B_3B_8 divided by $|B_7|$;
 B_4B_6 divided by $|B_7|$;
 B_4B_8 divided by $|B_6|$;
 B_3B_3 divided by $|B_4|$

and 144 divided by $|A_0|$ respectively. So, $A_1(0, 2) = 346/3$.

$q_1(0, 2) = 25 + 108 + 8 + 8 + 48 + 9 + 108 + 8 + 8 + 16 + 16 + 16 + 16 + 16 + 4 + 144 = 574$;

$$\mu_1(0, 2) = \mu_1(0, 0) = 346/(3 \cdot 574) = 0.200929.$$

So, we have

$$\begin{aligned} \mu_1(0, 1) &= 0.1959 \\ \mu_1(1, 1) &= 0.19716 \\ \mu_1(1, 0) &= \mu_1(1, 2) = 0.2 \\ \mu_1(0, 0) &= \mu_1(0, 2) = 0.200929. \end{aligned}$$

Let us denote

$$\begin{aligned} (0, 1) &\text{ by } 1, & (1, 1) &\text{ by } 2, \\ (1, 0) &\text{ by } 3, & (1, 2) &\text{ by } 4, \\ (0, 0) &\text{ by } 5, & (0, 2) &\text{ by } 6. \end{aligned}$$

Let H be the set $\{1, 2, 3, 4, 5, 6\}$.

Hence we obtain the following structure:

H_1^*	1	2	3	4	5	6
1	1	1, 2	1, 2, 3, 4	1, 2, 3, 4	H	H
2		2	2, 3, 4	2, 3, 4	2, 3, 4, 5, 6	2, 3, 4, 5, 6
3			3, 4	3, 4	3, 4, 5, 6	3, 4, 5, 6
4				3, 4	3, 4, 5, 6	3, 4, 5, 6
5					5, 6	5, 6
6						5, 6

From H_1^* we obtain

$$A_2(2) = 1 + 2/2 + 4/4 + 4/3 + 4/5 + 4/6 = 29/5.$$

$$q_2(2) = 2 + 1 + 4 + 4 + 4 + 4 = 19$$

$$\mu_2(2) = 29/(5 \cdot 19) = 0.305263.$$

$$A_2(3) = A_2(4) = 4/2 + 4/4 + 4/3 + 4/5 + 8/4 + 4/6 = 39/5.$$

$$q_2(3) = q_2(4) = 4 + 4 + 4 + 4 + 4 + 8 = 28$$

$$\mu_2(3) = \mu_2(4) = 39/(5 \cdot 28) = 0.27857.$$

$$A_2(5) = A_2(6) = 4/2 + 8/3 + 4/5 + 4/6 = 82/15.$$

$$q_2(5) = q_2(6) = 4 + 8 + 4 + 4 = 20$$

$$\mu_2(5) = \mu_2(6) = 82/(15 \cdot 20) = 0.27333.$$

$$A_2(1) = 1 + 2/2 + 4/4 + 4/6 = 11/3$$

$$q_2(1) = 11, \quad \mu_2(1) = 11/(11 \cdot 3) = 0.3333$$

Notice that

$$\mu_2(1) = 0.3333 > \mu_2(2) = 0.305263 > \mu_2(3) = \mu_2(4) = 0.27857 > \mu_2(5) = \mu_2(6) = 0.27333.$$

Thus, $H_2 \simeq H_1$, whence $\partial H_1^* = 1$.

3 Conclusion

Clearly, H_2 and H_1 are isomorphic and by consequence, the fuzzy grade of H^* is one.

References

- [1] R. Ameri, M.M. Zahedi, *Hypergroup and join space induced by a fuzzy subset*, PU.M.A., vol. 8, (2-3-4) (1997).
- [2] P. Corsini, *Prolegomena of Hypergroup Theory*, Aviani Editore, (1993).
- [3] P. Corsini, *On Chinese hyperstructures*, J. of Discrete Math. Sci and Cryptography, Vol 6 (2-3) (2003).

- [4] P. Corsini, *Join Spaces, Power Sets, Fuzzy Sets*, Proc. Fifth International Congress on A.H.A., 1993, Iasi, Romania
- [5] P. Corsini, *A new connection between hypergroups and fuzzy sets*, Southeast Bulletin of Math., 27 (2003)
- [6] P. Corsini, *Hyperstructures associated with ordered sets*, Bulletin of the Greek Mathematical Society, vol. 48, (2003)
- [7] P. Corsini, *Join Spaces, multivalued functions and soft sets*, Proceedings of the International Conference on Algebra 2010, (ICA 2010), Universitas Gadjah Mada and the Southeast Asian Math.
- [8] P. Corsini, *HX -groups and Hypergroup*, Analele Univ. "Ovidius", Math. Series n. 3, (2016)
- [9] P. Corsini, I. Cristea, *Fuzzy grade of i.p.s. hypergroups of order less or equal to 6*, PU.M.A., vol. 14, n. 4, (2003)
- [10] P. Corsini, I. Cristea, *Fuzzy grade of i.p.s. hypergroups of order 7*, Iran J. of Fuzzy Systems, 1 (2004)
- [11] P. Corsini, I. Cristea, *Fuzzy sets and non complete 1-hypergroups*, An. St. Univ. Ovidius Constanta, 13 (1) (2005)
- [12] P. Corsini, B. Davvaz, *New connections among multivalued functions, hyperstructures and fuzzy sets*, Jordan Journal of Mathematics and Statistics, (JJMS) 3 (3) (2010)
- [13] P. Corsini, V. Leoreanu-Fotea, *Applications of Hyperstructure Theory*, Advances in Mathematics, Kluwer Academic Publishers, (2003).
- [14] P. Corsini, V. Leoreanu-Fotea, *Join Spaces associated with Fuzzy Sets*, J. of Combinatorics, Information and System Sciences, vol. 20, n. 1 (1995)
- [15] P. Corsini, V. Leoreanu-Fotea, *On the grade of a sequence of fuzzy sets and join spaces determined by a hypergraph*, Southeast Asian Bulletin of Mathematics, 34 (2010)
- [16] P. Corsini, V. Leoreanu-Fotea, A. Iranmanesh, *On the sequence of join spaces and membership functions determined by a hypergraph*, Journal of Multivalued-Logic and Soft Computing, vol. 14, issue 6, (2008)
- [17] P. Corsini, R. Mahjoob, *Multivalued functions, fuzzy subsets and join spaces*, Ratio Mathematica, 20 (2010)
- [18] I. Cristea, *A property of the connection between fuzzy sets and hypergroupoids*, Italian Journal of Pure and Applied Mathematics, 21 (2007).
- [19] S. Hoskova, P. Chvalina, C. Rackova, *Noncommutative join spaces of integral operators and related hyperstructures*, Advances MT, 1 (2000).

- [20] L. Hongxing, D. Qinzhi, W. Peizhuang, *Hypergroup* (I). Busefal, Vol. 23, (1985).
- [21] L. Hongxing, W. Peizhuang, *Hypergroup* (II). Busefal, vol. 25, (1986).
- [22] L. Hongxing, *HX -Groups*, Busefal, Vol. 33, (1987).
- [23] M. Honghai, *Uniform HX -groups*, Busefal Vol. 47, (1991).
- [24] A. Maturo, I. Tofan, *Iperstrutture, strutture fuzzy ed applicazioni*, monografia di 168 pagine, pubblicazione finanziata con i fondi del progetto internazionale Socrates-Erasmus 2000/2001, Italia-Romania, Dierre Edizioni San Salvo, agosto.
- [25] W. Prenowitz, J. Jantosciak, *Join Geometries*, Springer-Verlag UTM, (1979).
- [26] S. Rasovic, *Hyperrings constructed by multiendomorphisms of hypergroups*, Proceedings of the 10th International Congress on AHA, Brno, (2008).
- [27] K. Serafimidis, A. Kehagias, M. Konstantinidou, *The L -fuzzy Corsini join hyperoperation*, Italian Journal of Pure and Applied Mathematics, 12 (2003).
- [28] S. Spartalis, *The hyperoperation relation and the Corsini's partial or not partial hypergroupoid*, Italian Journal of Pure and Applied Mathematics, 24 (2008).
- [29] M. Stefanescu, I. Cristea, *On the fuzzy grade of hypergroups*, Fuzzy Sets and Systems, 159 (2008).
- [30] T. Vougiouklis, *Hyperstructures and their representations*, Hadronic Press Inc. (1994).
- [31] M. Yavari, *Corsini's method and construction of join spaces*, Italian Journal of Pure and Applied Mathematics, 23 (2008).
- [32] F. Yuming, *Algebraic hyperstructures obtained from algebraic structures with binary fuzzy binary relations*, Italian Journal of Pure and Applied Mathematics, 25 (2009).
- [33] Z. Zhenliang, *The properties of HX -Groups*, Italian Journal of Pure and Applied Mathematics, Vol. 2, (1997).
- [34] Z. Zhenliang, *Classifications of HX -Groups and their chains of normal subgroups*, Italian Journal of Pure and Applied Mathematics, Vol. 5, (1999).

Piergiulio CORSINI,
 Department of Polytechnical Engineering,
 University of Udine,
 Via delle Scienze 206, 33100 Udine, Italy.
 Email: piergiuliocorsini@gmail.com