



Generalized Horadam Polynomials and Numbers

Snezana S. Djordjevic and Gospava B. Djordjevic

Abstract

We consider the polynomials $h_{n,m}(x)$ ($m \geq 2$) and the numbers $h_{n,m}$ ($x = 1$), which are the generalized Horadam polynomials and the generalized Horadam numbers, respectively. We also consider the polynomials $h_{n,m}^{(s)}(x)$ - convolutions of the polynomials $h_{n,m}(x)$, and the sequence of numbers $h_{n,m}^{(s)}$ - convolutions of the numbers $h_{n,m}$, where $s \geq 0$.

1 Introduction

In the paper [8] authors considered Horadam polynomials $h_n(x)$, which are given by the following recurrence relation

$$h_n(x) = pxh_{n-1}(x) + qh_{n-2}(x), \quad n > 2, \quad (1)$$

with $h_1(x) = a$, $h_2(x) = bx$, (a, b, p, q are some real constants).

We emphasize some particular cases of the polynomials $h_n(x)$:

- 1° For $a = b = p = q = 1$, we get the Fibonacci polynomials $F_n(x)$;
- 2° For $a = 2$, $b = p = q = 1$, we get the Lucas polynomials $L_n(x)$;
- 3° If $a = q = 1$, $b = p = 2$, then we get the Pell polynomials $P_n(x)$;
- 4° If $a = 1$, $b = p = 2$, $q = -1$, then we get the Chebyshev polynomials of the second kind $U_n(x)$.

Key Words: Generalized Horadam polynomials and numbers, Convolution, Recurrence relation; Explicit representation, Generating function.

2010 Mathematics Subject Classification: Primary 11B83; Secondary 11B37, 11B39.

Received: 28.03.2017

Revised: 02.09.2017

Accepted: 05.09.2017

2 Generalized polynomials

In this section we introduce the polynomials $h_{n,m}(x)$ ($m \geq 2$), the generalized Horadam polynomials, by

$$h_{n,m}(x) = pxh_{n-1,m}(x) + qh_{n-m,m}(x), \quad n > m, \quad (2)$$

with $h_{1,m}(x) = a$, $h_{n,m}(x) = bp^{n-2}x^{n-1}$, for $n = 2, \dots, m$.

For $x = 1$ in (2), we obtain the generalized Horadam numbers $h_{n,m}$:

$$h_{n,m} = ph_{n-1,m} + qh_{n-m,m}, \quad n > m, \quad (3)$$

with $h_{1,m} = a$, $h_{n,m} = bp^{n-2}$, for $n = 2, \dots, m$.

Remark 1. For $m = 2$ in (2) and (3), we get Horadam polynomials $h_n(x)$ and Horadam numbers h_n , respectively (see [8]).

Now, using the standard method, starting with the recurrence relation (2), we find that the function

$$F(x, t) = \frac{a + xt(b - ap)}{1 - pxt - qt^m} = \sum_{n=1}^{\infty} h_{n,m}(x)t^{n-1} \quad (4)$$

is the generating function of the polynomials $h_{n,m}(x)$.

Remark 2. For $m = 2$, the relation (4) yields the generating function $g(x, t)$ of the Horadam polynomials $h_n(x)$ (see [8], (13)):

$$g(x, t) = \frac{a + xt(b - ap)}{1 - pxt - qt^2} = \sum_{n=1}^{\infty} h_n(x)t^{n-1}.$$

Using the development of the function $F(x, t)$, given by (4), into the series on t , then comparing the corresponding coefficients to t^n , we obtain the explicit formula for the polynomials $h_{n,m}(x)$

$$\begin{aligned} h_{n,m}(x) = & a \sum_{k=0}^{[(n-1)/m]} \binom{n-1-(m-1)k}{k} (px)^{n-1-mk} q^k \\ & + \left(\frac{b}{p} - a \right) \sum_{k=0}^{[(n-2)/m]} \binom{n-2-(m-1)k}{k} (px)^{n-1-mk} q^k. \end{aligned} \quad (2.4)$$

Remark 3. For $m = 2$ the formula (2.4) yields the explicit formula for Horadam polynomials $h_n(x)$ (see [8], (16)).

Taking $x = 1$ in (2.4), we get the explicit formula for the generalized Horadam numbers $h_{n,m}$:

$$h_{n,m} = ap^{n-1} \sum_{k=0}^{[(n-1)/m]} \binom{n-1-(m-1)k}{k} \left(\frac{q}{p^m}\right)^k + p^{n-2}(b-ap) \sum_{k=0}^{[(n-2)/m]} \binom{n-2-(m-1)k}{k} \left(\frac{q}{p^m}\right)^k. \quad (2.5)$$

Some particular cases of the polynomials $h_{n,m}(x)$ (see [1, 3, 4, 5, 6]) are:

$$F_{n+1,m}(x) = \sum_{k=0}^{[n/m]} \binom{n-(m-1)k}{k} x^{n-mk} - \text{Fibonacci polynomials};$$

$$P_{n+1,m}(x) = \sum_{k=0}^{[n/m]} \binom{n-(m-1)k}{k} (2x)^{n-mk} - \text{Pell polynomials};$$

$$J_{n+1,m}(y) = \sum_{k=0}^{[n/m]} \binom{n-(m-1)k}{k} (2y)^k - \text{Jacobsthal polynomials};$$

$$U_{n,m}(x) = \sum_{k=0}^{[n/m]} (-1)^k \binom{n-(m-1)k}{k} (2x)^{n-mk} - \text{Chebyshev polynomials}.$$

3 Some properties

Theorem 1. The polynomials $h_{n,m}(x)$ satisfy the following relation

$$\sum_{k=1}^{n-1} h_{k,m}(x) = \frac{h_{n,m}(x) + q \sum_{i=1}^{m-1} h_{n-i,m}(x) - a - x(b-ap)}{px + q - 1}. \quad (5)$$

Proof. Starting from the recurrence relation (2) and by the corresponding characteristic equation

$$\lambda^m - px\lambda^{m-1} - q = 0, \quad (6)$$

where $\lambda_i, i = 1, \dots, m$ are the solutions of the equation (3.2), we have:

$$\begin{aligned} \sum_{k=1}^{n-1} h_{k,m}(x) &= \sum_{k=1}^{n-1} (A_1\lambda_1^{k-1} + A_2\lambda_2^{k-1} + \dots + A_m\lambda_m^{k-1}) \\ &= A_1(1 + \lambda_1 + \lambda_1^2 + \dots + \lambda_1^{n-2}) \\ &\quad + A_2(1 + \lambda_2 + \lambda_2^2 + \dots + \lambda_2^{n-2}) + \dots + \\ &\quad + A_m(1 + \lambda_m + \lambda_m^2 + \dots + \lambda_m^{n-2}) \\ &= A_1 \frac{1 - \lambda_1^{n-1}}{1 - \lambda_1} + A_2 \frac{1 - \lambda_2^{n-1}}{1 - \lambda_2} + \dots + A_m \frac{1 - \lambda_m^{n-1}}{1 - \lambda_m} \\ &= \frac{1}{(1 - \lambda_1)(1 - \lambda_2) \cdots (1 - \lambda_m)} \times \\ &\quad \left(A_1(1 - \lambda_1^{n-1} \prod_{i=2}^m (1 - \lambda_i)) + A_2(1 - \lambda_2^{n-1} \prod_{i=1, i \neq 2}^m (1 - \lambda_i)) + \dots + A_m(1 - \lambda_m^{n-1} \prod_{i=1}^{m-1} (1 - \lambda_i)) \right). \end{aligned}$$

Using the relations (by (3.2))

$$\lambda_1 + \lambda_2 + \dots + \lambda_m = px, \quad \lambda_1\lambda_2 + \lambda_1\lambda_3 + \dots + \lambda_{m-1}\lambda_m = 0, \dots, (-1)^m\lambda_1\lambda_2 \cdots \lambda_m = -q,$$

we get

$$\begin{aligned} &\left(\sum_{i=1}^m A_i(1 - \lambda_i^{n-1}) \prod_{j=1, j \neq i}^m (1 - \lambda_j) \right) \cdot ((1 - \lambda_1)(1 - \lambda_2) \cdots (1 - \lambda_m))^{-1} \\ &= \frac{1}{1 - px - q} \times \\ &\quad \sum_{i=1}^m A_i(1 - \lambda_i^{n-1}) (1 - \lambda_i(px - \lambda_i) - \lambda_i^2(px - \lambda_i) + \dots + (-1)^{m-1} \lambda_i^{m-1}(px - \lambda_i)) \\ &= \frac{1}{1 - px - q} \left(a - apx + bx - h_{n,m}(x) - q \sum_{i=1}^{m-1} h_{n-i,m}(x) \right). \end{aligned}$$

□

Remark 4. For $m = 2$ the formula (5) yields (see [8], (18))

$$\sum_{k=1}^{n-1} h_k(x) = \frac{h_n(x) + qh_{n-1}(x) - a - x(b - ap)}{px + q - 1}.$$

Some special cases of the formula (5) are (see [5]):

$$\sum_{k=1}^{n-1} F_{k,m}(x) = \frac{F_{n,m}(x) + \sum_{i=1}^{m-1} F_{n-i,m}(x) - 1}{x},$$

$F_{n,m}(x)$ – the generalized Fibonacci polynomials;

$$\sum_{k=1}^{n-1} P_{k,m}(x) = \frac{P_{n,m}(x) + \sum_{i=1}^{m-1} P_{n-i,m}(x) - 1}{2x},$$

$P_{n,m}(x)$ – the generalized Pell polynomials;

$$\sum_{k=1}^{n-1} U_{k,m}(x) = \frac{U_{n,m}(x) - \sum_{i=1}^{m-1} U_{n-i,m}(x) - 1}{2x - 2},$$

$U_{n,m}(x)$ – the generalized Chebyshev polynomials of the second kind.

4 Convolutions of the generalized Horadam polynomials

In this section we introduce the polynomials $h_{n,m}^{(s)}(x)$, the convolutions of the polynomials $h_{n,m}(x)$, by

$$G(x, t) = \left(\frac{a + xt(b - ap)}{1 - pxt - qt^m} \right)^{s+1} = \sum_{n=1}^{\infty} h_{n,m}^{(s)}(x)t^{n-1}, \quad (7)$$

where $n, m \in \mathbb{N}$, $s \in \mathbb{N} \cup \{0\}$, $n \geq m$.

Starting from (7), we get the following representation of the polynomials $h_{n,m}^{(s)}(x)$:

$$\begin{aligned} h_{n,m}^{(s)}(x) &= \sum_{i=0}^{s+1} \binom{s+1}{i} a^{s+1-i} \left(\frac{b}{p} - a \right)^i \\ &\times \sum_{k=0}^{[(n-1)/m]} \frac{(s+1)_{n-1-i-(m-1)k}}{k!(n-1-i-mk)!} (px)^{n-1-mk} q^k. \end{aligned} \quad (4.2)$$

where $\binom{n}{k} = 0$ for $n < k$.

Some particular cases of the polynomials $h_{n,m}^{(s)}(x)$ are:

1° For $a = b = p = q = 1$, we have $F_{n,m}^{(s)}(x)$, the convolutions of the generalized Fibonacci polynomials and (4.2) becomes

$$F_{n,m}^{(s)}(x) = \sum_{k=0}^{[(n-1)/m]} \frac{(s+1)_{n-1-(m-1)k}}{k!(n-1-mk)!} x^{n-1-mk}. \quad (4.3)$$

If we use the known relations ([7], [4]):

$$(\alpha)_{n+k} = (\alpha)_n (\alpha+n)_k; \quad \frac{(-1)^k}{(n-k)!} = \frac{(-n)_k}{n!}, \quad (\alpha)_{n-k} = \frac{(-1)^k (\alpha)_n}{(1-\alpha-n)_k},$$

then (4.3) takes the following hypergeometric representation

$$F_{n+1,m}^{(s)}(x) = \frac{x^n (s+1)^n}{n!} {}_m F_{m-1} \left[\begin{matrix} \frac{-n}{m}, \frac{1-n}{m}, \dots, \frac{m-1-n}{m}; & -x^{-m} \\ \frac{-s-n}{m-1}, \frac{1-s-n}{m-1}, \dots, \frac{m-2-s-n}{m-1} \end{matrix} \right].$$

2° For $a = q = 1$, $b = p = 2$, we obtain $P_{n,m}^{(s)}(x)$, the convolutions of the generalized Pell polynomials

$$P_{n+1,m}^{(s)}(x) = \sum_{k=0}^{[n/m]} \frac{(s+1)_{n-(m-1)k}}{k!(n-mk)!} (2x)^{n-mk},$$

or

$$P_{n+1,m}^{(s)}(x) = \frac{(2x)^n (s+1)_n}{n!} {}_m F_{m-1} \left[\begin{matrix} \frac{-n}{m}, \frac{1-n}{m}, \dots, \frac{m-1-n}{m}; & (2x)^{-m} \\ \frac{-s-n}{m-1}, \frac{1-s-n}{m-1}, \dots, \frac{m-2-s-n}{m-1} \end{matrix} \right].$$

3° For $a = 1$, $b = p = 2$, $q = -1$, in (4.2), we get $U_{n,m}^{(s)}(x)$, the convolutions of the generalized Chebyshev polynomials

$$U_{n,m}^{(s)}(x) = \frac{(2x)^n (s+1)_n}{n!} {}_m F_{m-1} \left[\begin{matrix} \frac{-n}{m}, \frac{1-n}{m}, \dots, \frac{m-1-n}{m}; & (2x)^{-m} \\ \frac{-s-n}{m-1}, \frac{1-s-n}{m-1}, \dots, \frac{m-2-s-n}{m-1} \end{matrix} \right].$$

Next, for $b = ap$ in (4), we get

$$F(x, t) = a(1 - pt - qt^m)^{-1} = \sum_{n=1}^{\infty} h_{n,m}(x) t^{n-1}. \quad (4.3)$$

Differentiating both sides of (4.3) to x , one by one, s -times, we find that

$$h_{n-s,m}^{(s)}(x) = \frac{1}{ap^s s!} D^s h_{n,m}(x), \quad \text{where } h_{n,m}^s(x) \equiv \frac{\partial^s h_{n,m}(x)}{\partial x^s}. \quad (4.4)$$

Namely, using the formula (4.4), we easily calculate the convolutions of the polynomials $h_{n,m}^{(s)}(x)$. Next, we give the examples for $m = 3$ and $s = 0, 1, 2, 3$, and for $m = 4$ and $s = 0, 1, 2, 3$.

Table 1: $h_{n,3}^{(s)}(x)$

n	$s = 0$	$s = 1$
1	a	1
2	apx	$2px$
3	$a(px)^2$	$3(px)^2$
4	$a(px)^3 + aq$	$4(px)^3 + 2q$
5	$a(px)^4 + 2apqx$	$5(px)^4 + 6pqx$
6	$a(px)^5 + 3aq(px)^2$	$6(px)^5 + 12q(px)^2$
7	$a(px)^6 + 4aq(px)^3 + aq^2$	$7(px)^6 + 20q(px)^3 + 3q^2$
8	$a(px)^7 + 5aq(px)^4 + 3aq^2px$	$8(px)^7 + 30q(px)^4 + 12q^2px$
9	$a(px)^8 + 6aq(px)^5 + 6aq^2(px)^2$	$9(px)^8 + 42q(px)^5 + 30q^2(px)^2$

Table 2: $h_{n,3}^{(s)}(x)$

n	$s = 2$	$s = 3$
1	1	1
2	$3px$	$4px$
3	$6(px)^2$	$10(px)^2$
4	$10(px)^3 + 3q$	$20(px)^3 + 4q$
5	$15(px)^4 + 12pqx$	$35(px)^4 + 20pqx$
6	$21(px)^5 + 30q(px)^2$	$56(px)^5 + 60q(px)^2$
7	$28(px)^6 + 60q(px)^3 + 6q^2$	$84(px)^6 + 140q(px)^3 + 10q^2$
8	$36(px)^7 + 105q(px)^4 + 30q^2px$	$120(px)^7 + 280q(px)^4 + 60q^2px$
9	$45(px)^8 + 168q(px)^5 + 90q^2(px)^2$	$165(px)^8 + 504q(px)^5 + 210q^2(px)^2$

Table 3: $h_{n,4}^{(s)}(x)$

n	$s = 0$	$s = 1$
1	a	1
2	apx	$2px$
3	$a(px)^2$	$3(px)^2$
4	$a(px)^3$	$4(px)^3$
5	$a(px)^4 + aq$	$5(px)^4 + 2q$
6	$a(px)^5 + 2aqpx$	$6(px)^5 + 6qpx$
7	$a(px)^6 + 3aq(px)^2$	$7(px)^6 + 12q(px)^2$
8	$a(px)^7 + 4aq(px)^3$	$8(px)^7 + 20q(px)^3$
9	$a(px)^8 + 5aq(px)^4 + aq^2$	$9(px)^8 + 30q(px)^4 + 3q^2$

Table 4: $h_{n,4}^{(s)}(x)$

n	$s = 2$	$s = 3$
1	1	1
2	$3px$	$4px$
3	$6(px)^2$	$10(px)^2$
4	$10(px)^3$	$20(px)^3$
5	$15(px)^4 + 3q$	$35(px)^4 + 4q$
6	$21(px)^5 + 12qpx$	$56(px)^5 + 20qpx$
7	$28(px)^6 + 30q(px)^2$	$84(px)^6 + 60q(px)^2$
8	$36(px)^7 + 60q(px)^3$	$120(px)^7 + 140q(px)^3$
9	$45(px)^8 + 105q(px)^4 + 6q^2$	$165(px)^8 + 56q(px)^4 + 10q^2$

Table 5: $P_{n,4}^{(s)}(x)$ -Pell polynomials

n	$s = 0$	$s = 1$
1	1	1
2	$2x$	$2(2x)$
3	$(2x)^2$	$3(2x)^2$
4	$(2x)^3$	$4(2x)^3$
5	$(2x)^4 + 1$	$5(2x)^4 + 2$
6	$(2x)^5 + 2(2x)$	$6(2x)^5 + 6(2x)$
7	$(2x)^6 + 3(2x)^2$	$7(2x)^6 + 12(2x)^2$
8	$(2x)^7 + 4(2x)^3$	$8(2x)^7 + 20(2x)^3$
9	$(2x)^8 + 5(2x)^4 + 1$	$9(2x)^8 + 30(2x)^4 + 3$

Table 6: $P_{n,4}^{(s)}(x)$

n	$s = 2$	$s = 3$
1	1	1
2	$3(2x)$	$4(2x)$
3	$6(2x)^2$	$10(2x)^2$
4	$10(2x)^3$	$20(2x)^3$
5	$15(2x)^4 + 3$	$35(2x)^4 + 4$
6	$21(2x)^5 + 12(2x)$	$56(2x)^5 + 20(2x)$
7	$28(2x)^6 + 30(2x)^2$	$84(2x)^6 + 60(2x)^2$
8	$36(2x)^7 + 60(2x)^3$	$120(2x)^7 + 140(2x)^3$
9	$45(2x)^8 + 105(2x)^4 + 6$	$165(2x)^8 + 56(2x)^4 + 10$

Table 7: $P_{n,4}^{(s)}$ -Pell numbers

n	$s = 0$	$s = 1$	$s = 2$	$s = 3$
1	1	1	1	1
2	2	4	6	8
3	4	12	24	40
4	8	32	80	160
5	17	82	243	564
6	36	204	696	1832
7	76	7496	1912	5616
8	160	1184	4608	16480
9	337	2787	13206	43146

References

- [1] G. B. Djordjevic, Some generalization of the Jacobsthal numbers, Filomat, 24(2)(2010), 143-151.
- [2] G. B. Djordjevic, Generalization of the Fibonacci and Lucas polynomials, Filomat, 23(3)(2009), 291-301.
- [3] G. B. Djordjevic, H. M. Srivastava, Some generalization of the incomplete Fibonacci and incomplete Lucas polynomials, Adv. Stud. Contemp. Math. 11(2005), 11-32.
- [4] G. B. Djordjevic, Convolutions of the generalized Pell and Pell-Lucas numbers, Filomat, 30(1)(2016), 105-112.
- [5] G. B. Djordjevic, Polynomials related to generalized Chebyshev polynomials, Filomat, 23(3)(2009), 279-290.
- [6] A. F. Horadam, Convolution summations for Pell and Pell-Lucas numbers, The Fibonacci Quart. 38(5)(2000), 451-457.
- [7] E. D. Rainville, Special functions, The Macmillan Company, New York, 1960.
- [8] Tugba Horcum, E. Gokcen Kocer, On some properties of Horadam polynomials, Int. Math. Forum, 4(25)(2009), 1243-1252.

Snezana S. Djordjevic,
Faculty of technology, Department of Mathematics,
University of Nis,
Bulevar Oslobođenja 124, 16000 Leskovac, Serbia.
Email: snezanadjordjevic1971@gmail.com

Gospava B. Djordjevic,
Faculty of technology, Department of Mathematics,
University of Nis,
Bulevar Oslobođenja 124, 16000 Leskovac, Serbia.
Email: gospava48@ptt.rs