

Antioxidant activity as indicator of UV radiation and other abiotic stress factors on *Agaricus bisporus* (Lange/Imbach) and *Sedum hybridum* (L.)

A. Szabó

e-mail: anna.szabo@uni-corvinus.hu

A. Geösel

e-mail: andras.geosel@uni-corvinus.hu

Corvinus University of Budapest, Faculty of Food Science,
Department of Vegetable and Mushroom Growing,
H-1118 Budapest, Villányi St 29–43., Hungary

Z. Kókai

e-mail: zoltan.kokai@uni-corvinus.hu

Corvinus University of Budapest, Faculty of Food Science,
Department of Postharvest Science and Sensory Evaluation,
H-1118 Budapest, Villányi St 29–43., Hungary

Cs. Orbán

e-mail: orban.csaba@se-etk.hu

K. Tőreki

e-mail: toreki.kristof@gmail.com

Semmelweis University, Faculty of Health Sciences,
Department of Dietetics and Nutrition Sciences,
H-1088 Budapest, Vas St 17., Hungary

A. Szőke

e-mail: szoke.andrea83@gmail.com

Corvinus University of Budapest, Faculty of Horticultural Sciences,
Department of Floriculture and Dendrology,
H-1118 Budapest, Villányi St 29–43., Hungary

Keywords and phrases: free radicals, redox adaptation, button mushroom, extensive green roof

Abstract. Investigation of stress level might be facilitated also in plant and horticultural sciences, but currently mainly morphological parameters are in use. Antioxidant activity routinely measured in food-oriented researches and several studies indirectly indicated that stress factors can influence this parameter. Our aim was to assess the potential direct indicator role of antioxidant activity in stress conditions. We measured the effects of UVB and soil-delivered stress on *Agaricus bisporus* and *Sedum hybridum*. Our results indicate that UVB slightly decreases, while the inadequate soil conditions increase antioxidant activity; hence these measurements are suitable for determining the level of stress in different living samples.

1 Introduction

Antioxidant activity is a parameter that describes the capacity of a sample to neutralize the free radicals. It has got a crucial role in human health, as many diseases (e.g. malignant tumours, non-alcoholic fat liver, autoimmune diseases etc.) (Felton *et al.*, 1994; Geösel *et al.*, 2011; Ozzard *et al.*, 2008) can be developed by the damages in DNA and the structural and functional proteins evoked by these reactive substances (Bagchi *et al.*, 2000). Because of this, many studies investigated how efficient the different foods are (e.g. fruits and vegetables) in delivering protective antioxidants to the human body.

In plants, the role of the antioxidant type of characteristic substances is the same as in the human body: protect the structure and function of plant cell organelles from free-radical-caused damages, and thus maintain homeostasis.

In mammalian, plant, and fungal cells, the antioxidant capacity mainly comes from two systems: the first one is the non-enzymatic, which utilizes organic substances (e.g. ascorbic acid, lycopene, α -tocopherol, glutathione) to quench the Reactive Oxygen Species (ROS), while the enzymatic antioxidant system contains enzyme proteins, such as Glutathion-S-transferase, Ascorbate-peroxidase, to facilitate electron or hydrogen atom transfer (Edwards *et al.*, 2000; Venisse, 2003).

Most of the non-enzymatic participants are represented in a relatively small amount in the different samples. Increasing of their levels can be influenced by many factors, e.g. the concentration of lycopene levels depends on temperature (Helyes *et al.*, 2007). Polyphenols are expanded by the photo-oxidative UV-stress in many fruits, e.g. *Sambucus sp.*, *Fragaria sp.* (Asami *et al.*, 2003; Murugesan *et al.*, 2012). The joint feature of these induction factors is that they both mean stress to the plants.

Many horticultural products have been tested for this phenomenon with different methods (*Csambalik et al.*, 2014; *Geösel et al.*, 2011; *Sipos et al.*, 2013). Most of the studies investigate the effect of environmental factors on the health-promoting molecule contents. Some studies suggested that stress factors indicate the elevated synthesis of carotenoids (*Kim et al.*, 2012). Others show that polyphenol content increases during UV-stress to protect the plant tissues from damages (*Ozzard et al.*, 2008). Similar results were obtained from measurements performed on fruits exposed to insect invasions (*Felton et al.*, 1994; *Hemingway & Laks*, 1992). Other abiotic stress factors, such as water shortage or chilling injury, also alter the bioactive compound levels (*Esteban et al.*, 2001). The availability of mineral elements determine the redox status as well (*Tewari et al.*, 2006). Competition with weed also means relevant stress to the plants.

The antioxidant-producing ability of the different species describes the capacity of the plants to adapt to these stress factors and hence contribute to the viability characteristics. As measurements of antioxidant activity can be performed by rapid and economical methods, another paradigm also arises (*Huang et al.*, 2005). It is hard to deceive the effect of stress factors to the investigated species. Most of the agricultural and food engineers assess the optimal condition by the quality parameters e.g. size, yield, growing rate and dynamics, ornamental value (*VanWoert et al.*, 2005). These are indirect parameters.

As antioxidant capacity measures the total ROS elimination capacity of the processed tissue, and these radicals formed during the exposition to the stress factors, it is possible that the antioxidant capacity indicates the intensity of stress itself.

Because of the aforementioned points, we came up with a novel approach to determine the connection of antioxidant activity with the different stress factors in mushrooms, as a relevant part of human nutrition, and in *Sedum*-s as an ornamentally interesting species.

UVB-treated white button mushrooms were chosen for the purpose of this study since nowadays they have become one of the novelty mushroom products with their increased vitamin D content, representing a natural source of this vitamin of high importance for the human nutrition (*Ozzard et al.*, 2008). The other object is *Sedum hybridum*, a commonly applied species on green roofs, especially extensive green roofs for its exceptional tolerance for climatic conditions and high decorative value (*Szöke et al.*, 2012).

2 Materials and methods

2.1 Investigated samples

Ultraviolet-B (UVB)-treated *Agaricus bisporus* samples were provided by the Department of Vegetable and Mushroom Growing. The biologically active, still developing fruitbodies were treated by a 15W output 290–315 nm wavelength VL-115 M lamp. 0, 5, 15, 20, 25, and 30 minutes of UVB radiation were applied on three consecutive days (a total of: 0, 15, 30, 45, 60, 75, 90 minutes). Samples were taken on 15th January 2015.

Sedum hybridum plants were propagated vegetatively. The samples were grown on a 3-year extensive green roof in four different substrate mixtures (S1: soil-zeolite-sand-based substrate, S2: riolite-based substrate, S3: brick-fracture- and ytong-based substrate and S4: soil) and in a 10-cm layer, respectively, as potential sources of substrate-associated stress factors. The control plants were grown on the ground in the original soil of the plot under the same ambient conditions. Samples were collected at the same phenological phase on the 10th of April 2015.

2.2 Sample preparation

Samples were homogenized with distilled water in a 1:1 ratio at 24000 min⁻¹ RPM by a teflon homogenizer. This homogenate was kept in ultrasonic water bath for 15 min, and then spanned at 2000 g for 15 min. The supernatant was used for all of the measurements.

2.3 Antioxidant activity measurements

Ferric reduction antioxidant power (FRAP) was determined according to *Benzie & Strain* (1996). 100 µl of sample was added to pH = 3.6, 300 mM acetate buffer – 10 mmol/litre TPTZ (2,4,6-tripyridyl-s-triazine) in 40 mM HCl – 20 mmol/litre FeCl₃ · 6H₂O, and then, after 5 minutes, absorbance was read at $\lambda = 593$ nm. Results were generated in ascorbic acid equivalence (*Benzie & Strain*, 1996).

DPPH (2,2-diphenyl-1-picrylhydrazyl) elimination assay was performed as described by *Brand-Williams et al.* (1995). 100 µl of sample was added to 3.9 ml of $6 \cdot 10^{-5}$ M methanolic DPPH solution, and then incubated for 20 minutes in dark. Absorbance readings were performed at $\lambda = 517$ nm, and inhibition % was calculated (*Brand-Williams et al.*, 1995).

CUPRAC assay was measured by the method of *Apak et al.* (2008). 1 ml

10^{-2} M CuCl_2 , 1 ml $7.5 \cdot 10^{-3}$ M neocuproine solution, 1 ml pH = 7.4, 1 M NH_4Ac buffer, 100 μl sample, and 1 ml distilled water was mixed and incubated for 30 minutes in dark at room temperature. After the absorbance reading at $\lambda = 450$ nm, values were calculated to trolox equivalents (*Apak et al.*, 2008).

ABTS-assay was measured as described by *Huang et al.* (2005). Reaction mixture contained 10 μl of sample; 20 μl of 3.50 mg/ml myoglobin in 50 mM, pH = 7.4, 9% NaCl and 1% glucose containing potassium-phosphate buffer; 150 μl of 1 mg ABTS and 25 μl 3% H_2O_2 in 0.1 M pH5 citrate buffer. This mixture was shaken for 5 minutes at 37 °C, then alkaline stop solution was added and measured at $\lambda = 405$ nm against trolox calibration curve (*Huang et al.*, 2005).

Total Phenolic Compound (TPC) was recorded as described by *Singleton and Rossi* (1965). 1250 μl of 10-fold diluted Folin-Cioalteaou reagent, 240 μl methanol, 10 μl of sample, and 1 ml of 0.7 M NaCO_3 was mixed, and then kept at 50 °C for 5 minutes, then measured at $\lambda = 765$ nm to calibration curve set-up with gallic acid (*Singleton & Rossi*, 1965). All of the measurements were carried out in five replicates.

2.4 Statistical analysis

Raw data were processed during statistical analysis. Since the resulting dataset is considered to be low from statistical point of view ($n < 13$), the conditions for parametric tests were not met. That is why the Kruskal-Wallis nonparametric test (with the exact p-value calculation) was chosen (equivalent to ANOVA parametric test), which is not sensitive either to the normality of our data distribution or to the heterogeneity of its deviation (*Conover*, 1980).

Multiple pairwise comparisons using Dunn's procedure – as a suitable method for detecting significant differences with Bonferroni correction – was applied. The analyses were implemented by the XLStat-Sensory solution software, version 2013.1.01 (Addinsoft, 28 West 27th Street, Suite 503, New York, NY 10001, USA).

To visualize and compare data in graphs, a 0 to 100 scaling method was applied to normalize the data.

3 Results and discussions

Our results (*Table 1*) indicate that in the case of UVB exposition the antioxidant activity does not alter. None of the applied methods show either increasing or decreasing patterns via photo-oxidative stress.

Table 1: Antioxidant activity and phenolic compound alteration patterns via different stress factors

Species	Stressor	DPPH		CUPRAC		ABTS		FRAP		TPC	
		mean±SD		mean±SD		mean±SD		mean±SD		mean±SD	
Agaricus bisporus	UVB-0	61.76±5.34 ^a		100.00±0.80 ^b		100.00±5.30 ^c		99.53±2.86 ^b		100.00±1.54 ^b	
	UVB-15	96.48±2.61 ^b		81.66±1.76 ^b		48.09±0.38 ^{ab}		68.56±4.16 ^{ab}		85.88±1.34 ^{ab}	
	UVB-30	79.50±0.00 ^{ab}		66.50±0.69 ^{ab}		47.96±0.29 ^{ab}		66.06±0.00 ^{ab}		74.92±0.00 ^a	
	UVB-45	97.98±2.33 ^b		77.45±1.59 ^{ab}		47.99±0.67 ^a		75.13±0.25 ^{ab}		81.47±1.56 ^{ab}	
	UVB-60	82.19±1.38 ^{ab}		50.19±0.00 ^{ab}		53.85±0.03 ^{abc}		100.00±4.79 ^b		84.81±4.58 ^{ab}	
	UVB-75	98.67±2.70 ^b		65.55±1.11 ^{ab}		54.17±0.17 ^{abc}		95.36±0.11 ^b		95.66±5.87 ^b	
	UVB-90	100.00±6.86 ^b		50.94±0.61 ^a		54.59±0.00 ^{bc}		18.90±3.29 ^a		71.09±2.46 ^a	
	Range	38.24		49.81		52.04		81.10		28.91	
Sedum hybridum	Control	9.29±0.00 ^a		69.65±0.26 ^{ab}		100.00±0.00 ^{ab}		61.20±4.72 ^{ab}		29.34±0.88 ^{ab}	
	S1	85.76±5.90 ^{ab}		69.70±1.51 ^{ab}		55.15±13.86 ^a		61.48±0.00 ^a		27.65±1.33 ^a	
	S2	85.91±0.43 ^{ab}		68.61±0.00 ^a		97.53±11.69 ^b		65.62±0.88 ^{ab}		28.45±0.86 ^a	
	S3	79.50±0.78 ^a		100.00±0.13 ^b		52.28±23.39 ^a		100.00±3.75 ^b		100.00±22.27 ^b	
	S4	100.00±0.92 ^b		80.82±1.71 ^{ab}		68.03±21.32 ^{ab}		72.01±7.00 ^{ab}		32.87±0.00 ^{ab}	
	Range	90.71		31.39		47.72		38.80		72.35	

Data is given in mean ± SD format obtained from five parallel measurements. **a**, **b**, and **c** indicate homogeneous groups, **ab** and **abc** indicate heterogeneous groups. Homogeneity and heterogeneity are indicated by methods and species, respectively.

Although the size of changes is not meaningful, there was a slight loss of radical-eliminating molecules during the treatments. The highest values were observed in the control samples. The total phenolic compound changes also demonstrate the decreasing pattern observed in the case of antioxidant activity (Figure 1).




Figure 1: Antioxidant activity and phenolic compound characteristics of UVB-treated white button mushroom samples (mean, SD, n=5)

In the case of soil-derived stress, we observed the organic-poor substrate-derived stress, as the highest increase in the antioxidant activity was manifested in the S3 growth sample. Values of high-soil-content S1 and S4 samples show similar patterns in the case of each of the methods (Figure 2).




Figure 2: Antioxidant activity and phenolic compound characteristics of *Sedum hybridum* samples (mean, SD, n=5)

Range values show that the sensitivity of the applied 5 methods is different. In the case of *Agaricus*, the increasing order of ranking is TPC<DPPH<CUPRAC<ABTS<FRAP. In the case of *Sedum*, the following ranking applies: CUPRAC<FRAP<ABTS<TPC<DPPH.

4 Conclusion

In this study, we assessed the potential role of antioxidant-activity assays in the monitoring of photo-oxidative UVB- and substrate-associated stress factors on a nutritionally relevant edible mushroom and on an ornamentally valuable plant. Phenolic compound levels were also determined. During the measurements, we utilized the most common methods that can be easily adapted by any laboratory without the need of expensive instrumentation (*Huang et al.*, 2005).

Our results indicated that UVB stress did not cause an increasing pattern in mushroom samples, which is similar to what was observed in shiitake mushrooms by *Jiang et al.* (2010). It is maybe due to the fact that mushrooms do not possess a photosynthetic apparatus. There is evidence that the thylakoid membrane-bound enzymes of the photosynthetic system are the most sensitive proteins in the plants to the ROS damages. Avoiding the function loss can be achieved by the enhanced antioxidant capacity (enzymatic and non-enzymatic as well). The difference between *Sedum* and *Agaricus sp.*s may come from the lack of chloroplast, whence the necessity of fast-reacting antioxidant systems (*Halliwell*, 1989). The absence of phenolic content changes is due to the fact that mushrooms are not considered as a good source for these molecules (*Wong et al.*, 2013).

The results also indicated that alteration in antioxidant activity is a hallmark of the response to the related stress, as in the case of *Sedum hybridum* species the soils with the hypothesized different adequacy levels. The highest values measured in S3 substrates verified this assumption. This result is in line with others, who also highlighted the relevance of adequate salinity or other soil-delivered necessary compounds (*Esteban & Villanueva*, 2001; *Sairam et al.*, 2005).

Alteration in phenolic compound levels was also significant, which is in agreement with the results obtained from test performed on berries (*Asami et al.*, 2003; *Murugesan et al.*, 2012).

Our different range/rank/order results between the *Sedum* and *Agaricus sp.*s are also in line with others' conclusion that the simultaneous utilization of

different antioxidant activity methods is necessary as their sensitivity differs due to their different chemical background (*Huang et al.*, 2005).

On the basis of our results, we can conclude that antioxidant activity is a suitable derived parameter for the assessment of stress level.

Acknowledgements

The results presented here are an output of the research funds TÁMOP-4.2.1./B-09/1/KMR 2010-0005 and VITADFUN (TECH_08/2008) supported by NKTH.

References

- [1] R. Apak, K. Güclü, M. Özyürek, S. E. Celik, Mechanism of antioxidant capacity assays and the CUPRAC (cupric ion reducing antioxidant capacity) assay. *Microchim. Acta*, 160. (2008) 413–419.
- [2] D. K. Asami, Y. J. Hong, D. M. Barrett, A. E. Mitchell, Comparison of the total phenolic and ascorbic acid content of freeze-dried and air-dried marionberry, strawberry, and corn grown using conventional, organic, and sustainable agricultural practices. *Agric. Food Chem.*, 51. (2003) 1237–1241.
- [3] D. Bagchi, M. Bagchi, S. J. Stohs, D. K. Das, S. D. Ray, C. A. Kuszynski, Free radicals and grape seed proanthocyanidin extract: importance in human health and disease prevention. *Toxicology*, 148. (2000) 187–197.
- [4] F. Benzie, J. J. Strain, The ferric reducing ability of plasma (FRAP) as a measure of “antioxidant power”: the FRAP assay. *Anal. Biochem.*, 239. (1996) 70–76.
- [5] W. Brand Williams, M. E. Cuvelier, C. Berset, Use of a free radical method to evaluate antioxidant activity. *Lebensm.-Wiss. u. Technol.*, 28. (1995) 25–30.
- [6] W. J. Conover, Practical nonparametric statistics. John Wiley and Sons, New York. (1980) 229–238.
- [7] L. Csambalik, A. Divéky-Ertsey, Z. Pap, Cs. Orbán, M. Stégerné Máté, A. Gere, É. Stefanovits-Bányai, L. Sipos, Coherences of instrumental

- and sensory characteristics: case study on cherry tomatoes. *J. of Food Sci.*, 79. (2014) 2192–2202.
- [8] R. Edwards, D. P. Dixon, V. Walbot, Plant glutathione S-transferases: enzymes with multiple functions in sickness and in health. *Trends Plant. Sci.*, 5. (2000) 193–198.
- [9] M. A. Esteban, M. J. Villanueva, J. R. Lissarrague, Effect of irrigation on changes in the anthocyanin composition of the skin of cv Tempranillo (*Vitis vinifera* L.) grape berries during ripening. *J. Sci. Food Agr.*, 81. (2001) 409–420.
- [10] G. W. Felton, C. B. Summers, A. J. Mueller, Oxidative responses in soybean foliage to herbivory by bean leaf beetle and three-cornered alfalfa hopper. *J. of Chem. Ecology*, 20. (1994) 639–650.
- [11] A. Geösel, L. Sipos, É. Stefanovitsné-Bányai, Z. Kókai, J. Györfi, Antioxidant, polyphenol and sensory analysis of *Agaricus bisporus* and *Agaricus subrufescens* varieties. *Acta Alimentaria*. 40. (2011) 33–40.
- [12] B. Halliwell, Free radicals, reactive oxygen species and human disease: a critical evaluation with special reference to atherosclerosis. *Br. J. Exp. Pathol.*, 70. (1989) 737–757.
- [13] L. Helyes, A. Lugasi, Z. Pék, Effect of natural light on surface temperature and lycopene content of vine ripened tomato fruit. *Canadian J. of Plant Sci.*, 87. (2007) 927–929.
- [14] R. W. Hemingway, P. E. Laks, Plant Polyphenols: Synthesis, properties, significance. Springer Science & Business Media. (1992) 1053.
- [15] D. Huang, B. Ou, R. L. Prior, The chemistry behind antioxidant capacity assays. *J. Agric. Food Chem.*, 53. (2005) 1841–1856.
- [16] T. Jiang, M. M. Jahangir, Z. Jiang, X. Lu, T. Ying, Influence of UV-C treatment on antioxidant capacity, antioxidant enzyme activity and texture of postharvest shiitake (*Lentinus edodes*) mushrooms during storage. *Postharvest Biol. and Techn.*, 56. (2010) 209–215.
- [17] S. H. Kim, Y. H. Kim, O. A. Young, A. Mi-Jeong, J. C. Jeong, H. S. Lee, S. S. Kwak, Downregulation of the lycopene ϵ -cyclase gene increases carotenoid synthesis via the β -branch-specific pathway and en-

- hances salt-stress tolerance in sweetpotato transgenic calli. Article first published online: 1 OCT 2012 DOI: 10.1111/j.1399-3054.2012.01688.
- [18] R. Murugesan, V. Orsat, M. Lefsrud, Effect of pulsed ultraviolet light on the total phenol content of elderberry (*Sambucus nigra*). *Fruit Food and Nutr. Sci.*, 3. (2012) 774–783.
- [19] A. Ozzard, G. Morrison, M. Hoskin, Vitamin D deficiency treated by consuming UVB-irradiated mushrooms. *Br. J. Gen. Pract.*, 58. (2008) 644–645.
- [20] R. K. Sairam, G. C. Srivastava, S. Agarwal, R. C. Meena, Differences in antioxidant activity in response to salinity stress in tolerant and susceptible wheat genotypes. *Biol. Plantarum*, 49. (2005) 85–91.
- [21] V. L. Singleton, J. A. Rossi, Colorimetry of total phenolics with posphomolibdic phosphotungstic acid reagents. *Am. J. of Enology and Viti-culture*, 161. (1965) 144–158.
- [22] L. Sipos, G. Ficzek, Z. Kókai, M. Tóth, New multiresistant apple cultivars – complex assessment of sensory and some instrumental attributes. *Acta Alimentaria*, 42. (2013) 132–142.
- [23] Szőke, A., Gerzson, L., Forró, E., Extenzív zöldtetők termesztő közegeinek kritikus paraméterei. Talajtani Vándorgyűlés Miskolc. (2012) 23–25.
- [24] R. K. Tewari, P. Kumar, P. N. Sharma, Magnesium deficiency induced oxidative stress and antioxidant responses in mulberry plants. *Scientia Horticulturae*, 108. (2006) 7–14.
- [25] N. D. VanWoert, D. B. Rowe, J. A. Andresen, C. L. Rugh, L. Xiao, Watering regime and green roof substrate design affect sedum plant growth. *HortScience*, 40. (2005) 659–664.
- [26] J. S. Venisse, Involvement of three pathogenicity factors of *Erwinia amylovora* in the oxidative stress associated with compatible interaction in pear. *FEBS Letters*, 537. (2003) 198–202.
- [27] F. C. Wong, T. T. Chai, S. L. Tan, A. L. Yong, Evaluation of bioactivities and phenolic content of selected edible mushrooms in Malaysia. *Tropical J. of Pharmaceutical Research*, 12. (2013). ISSN (electronic) 1596-9827.