

# On the k-Fibonacci words

José L. RAMÍREZ

Instituto de Matemáticas y sus  
Aplicaciones  
Universidad Sergio Arboleda, Colombia  
email:  
josel.ramirez@ima.usergioarboleda.edu.co

Gustavo N. RUBIANO

Departamento de Matemáticas  
Universidad Nacional de Colombia,  
Colombia  
email: gnrubiano@unal.edu.co

**Abstract.** In this paper we define the k-Fibonacci words in analogy with the definition of the k-Fibonacci numbers. We study their properties and we associate to this family of words a family of curves with interesting patterns.

## 1 Introduction

Fibonacci numbers and their generalizations have many interesting properties and applications to almost every fields of science and arts (e.g. see [13]). The Fibonacci numbers  $F_n$  are the terms of the sequence  $0, 1, 1, 2, 3, 5, \dots$  wherein each term is the sum of the two previous terms, beginning with the values  $F_0 = 0$ , and  $F_1 = 1$ .

Besides the usual Fibonacci numbers many kinds of generalizations of these numbers have been presented in the literature. In particular, a generalization is the k-Fibonacci numbers [11].

For any positive real number  $k$ , the k-Fibonacci sequence, say  $\{F_{k,n}\}_{n \in \mathbb{N}}$  is defined recurrently by

$$F_{k,0} = 0, F_{k,1} = 1 \quad \text{and} \quad F_{k,n+1} = kF_{k,n} + F_{k,n-1}, \quad n \geq 1. \quad (1)$$

---

**Computing Classification System 1998:** I.3.7, G.2.0

**Mathematics Subject Classification 2010:** 11B39, 05A05, 68R15

**Key words and phrases:** Fibonacci word, k-Fibonacci numbers, k-Fibonacci words, k-Fibonacci curves

In [11],  $k$ -Fibonacci numbers were found by studying the recursive application of two geometrical transformations used in the four-triangle longest-edge (4TLE) partition. These numbers have been studied in several papers, see [5, 10, 11, 12, 18, 19, 23].

The characteristic equation associated to the recurrence relation (1) is  $x^2 = kx + 1$ . The roots of this equation are

$$r_{k,1} = \frac{k + \sqrt{k^2 + 4}}{2}, \quad \text{and} \quad r_{k,2} = \frac{k - \sqrt{k^2 + 4}}{2}.$$

Some of the properties that the  $k$ -Fibonacci numbers verify are (see [11, 12] for the proofs).

- Binet Formula:  $F_{k,n} = \frac{r_{k,1}^n - r_{k,2}^n}{r_{k,1} - r_{k,2}}$ .
- Combinatorial Formula:  $F_{k,n} = \sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n-1-i}{i} k^{n-1-2i}$ .
- $\lim_{n \rightarrow \infty} \frac{F_{k,n}}{F_{k,n-1}} = r_{k,1}$ .

On the other hand, there is a word-combinatorial interpretation of the Fibonacci sequence. Fibonacci words are words over  $\{0, 1\}$  defined recursively as follows:

$$f_0 = 1, \quad f_1 = 0, \quad f_n = f_{n-1}f_{n-2}, \quad n \geq 2.$$

The words  $f_n$  are referred to as the finite Fibonacci words and it is clear that  $|f_n| = F_{n+1}$ . The limit

$$\mathbf{f} = \lim_{n \rightarrow \infty} f_n = 0100101001001010010100100101 \dots$$

is called the Fibonacci word. This word is certainly one of the most studied words in the combinatorics on words, (see, e.g., [2, 6, 7, 9, 15, 22]). It is the archetype of a Sturmian word [14]. This word can be associated with a curve, which has fractal properties obtained from combinatorial properties of  $\mathbf{f}$  [3, 16, 21].

In this paper we introduce a family of words  $\mathbf{f}_k$  that generalize the Fibonacci word. Specifically, the  $k$ -Fibonacci words are words over  $\{0, 1\}$  defined inductively as follows

$$f_{k,0} = 0, \quad f_{k,1} = 0^{k-1}1, \quad f_{k,n} = f_{k,n-1}^k f_{k,n-2},$$

for all  $n \geq 2$  and  $k \geq 1$ . Then it is clear that  $|f_{k,n}| = F_{k,n+1}$ . The infinite word

$$f_k := \lim_{n \rightarrow \infty} f_{k,n}$$

is called the  $k$ -Fibonacci word. In connection with this definition, we investigate some new combinatorial properties and we associate a family of curves with interesting patterns.

## 2 Definitions and notation

The terminology and notations are mainly those of Lothaire [14] and Allouche and Shallit [1].

Let  $\Sigma$  be a finite alphabet, whose elements are called symbols. A word over  $\Sigma$  is a finite sequence of symbols from  $\Sigma$ . The set of all words over  $\Sigma$ , i.e., the free monoid generated by  $\Sigma$ , is denoted by  $\Sigma^*$ . The identity element  $\epsilon$  of  $\Sigma^*$  is called the empty word and  $\Sigma^+ = \Sigma^* \setminus \{\epsilon\}$ . For any word  $w \in \Sigma^*$ ,  $|w|$  denotes its length, i.e., the number of symbols occurring in  $w$ . The length of  $\epsilon$  is taken to be equal to 0. If  $a \in \Sigma$  and  $w \in \Sigma^*$ , then  $|w|_a$  denotes the number of occurrences of  $a$  in  $w$ .

For two words  $u = a_1 a_2 \cdots a_k$  and  $v = b_1 b_2 \cdots b_s$  in  $\Sigma^*$  we denote by  $uv$  the concatenation of the two words, that is,  $uv = a_1 a_2 \cdots a_k b_1 b_2 \cdots b_s$ . If  $v = \epsilon$  then  $u\epsilon = \epsilon u = u$ , moreover, by  $u^n$  we denote the word  $uu \cdots u$  ( $n$  times). A word  $v$  is a subword (or factor) of  $u$  if there exist  $x, y \in \Sigma^*$  such that  $u = xvy$ . If  $x = \epsilon$  ( $y = \epsilon$ ), then  $v$  is called prefix (suffix) of  $u$ .

The reversal of a word  $u = a_1 a_2 \cdots a_n$  is the word  $u^R = a_n \cdots a_2 a_1$  and  $\epsilon^R = \epsilon$ . A word  $u$  is a palindrome if  $u^R = u$ .

An infinite word over  $\Sigma$  is a map  $u : \mathbb{N} \rightarrow \Sigma$ . It is written  $u = a_1 a_2 a_3 \cdots$ . The set of all infinite words over  $\Sigma$  is denoted by  $\Sigma^\omega$ .

**Example 1** The word  $p = (p_n)_{n \geq 1} = 0110101000101 \cdots$ , where  $p_n = 1$  if  $n$  is a prime number and  $p_n = 0$  otherwise, is an example of an infinite word.  $p$  is called the characteristic sequence of the prime numbers.

**Definition 2** Let  $\Sigma$  and  $\Delta$  be alphabets. A morphism is a map  $h : \Sigma^* \rightarrow \Delta^*$  such that  $h(xy) = h(x)h(y)$  for all  $x, y \in \Sigma^*$ . It is clear that  $h(\epsilon) = \epsilon$ .

There is a special class of words, with many remarkable properties, the so-called Sturmian words. These words admit several equivalent definitions (see, e.g. [1] or [14]).

**Definition 3** Let  $\mathbf{w} \in \Sigma^\omega$ . We define  $P(\mathbf{w}, n)$ , the complexity function of  $\mathbf{w}$ , to be the map that counts, for all integer  $n \geq 0$ , the number of subwords of length  $n$  in  $\mathbf{w}$ . An infinite word  $\mathbf{w}$  is a Sturmian word if  $P(\mathbf{w}, n) = n + 1$  for all integer  $n \geq 0$ .

Since for any Sturmian word  $P(\mathbf{w}, 1) = 2$ , then Sturmian words are over two symbols. The word  $\mathbf{p}$ , in Example 1, is not a Sturmian word because  $P(\mathbf{p}, 2) = 4$ .

Given two real numbers  $\alpha, \beta \in \mathbb{R}$  with  $\alpha$  irrational and  $0 < \alpha < 1, 0 \leq \beta < 1$ , we define the infinite word  $\mathbf{w} = w_1 w_2 w_3 \cdots$  as

$$w_n = \lfloor (n+1)\alpha + \beta \rfloor - \lfloor n\alpha + \beta \rfloor.$$

The numbers  $\alpha$  and  $\beta$  are the slope and the intercept, respectively. This word is called mechanical. The mechanical words are equivalent to Sturmian words [14]. As special case, when  $\beta = 0$ , we obtain the characteristic words.

**Definition 4** Let  $\alpha$  be an irrational number with  $0 < \alpha < 1$ . For  $n \geq 1$ , define

$$w_\alpha(n) := \lfloor (n+1)\alpha \rfloor - \lfloor n\alpha \rfloor,$$

and

$$\mathbf{w}(\alpha) := w_\alpha(1)w_\alpha(2)w_\alpha(3)\cdots,$$

then  $\mathbf{w}(\alpha)$  is called a characteristic word with slope  $\alpha$ .

On the other hand, note that every irrational  $\alpha \in (0, 1)$  has a unique continued fraction expansion

$$\alpha = [0, a_1, a_2, a_3, \dots] = \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}},$$

where each  $a_i$  is a positive integer. Let  $\alpha = [0, 1 + d_1, d_2, \dots]$  be an irrational number with  $d_1 \geq 0$  and  $d_n > 0$  for  $n > 1$ . To the directive sequence  $(d_1, d_2, \dots, d_n, \dots)$ , we associate a sequence  $(s_n)_{n \geq -1}$  of words defined by

$$s_{-1} = 1, \quad s_0 = 0, \quad s_n = s_{n-1}^{d_n} s_{n-2}, \quad (n \geq 1).$$

Such a sequence of words is called a standard sequence. This sequence is related to characteristic words in the following way. Observe that, for any  $n \geq 0$ ,  $s_n$  is a prefix of  $s_{n+1}$ , which gives meaning to  $\lim_{n \rightarrow \infty} s_n$  as an infinite word. In fact, one can prove [14] that each  $s_n$  is a prefix of  $\mathbf{w}(\alpha)$  for all  $n \geq 0$  and

$$\mathbf{w}(\alpha) = \lim_{n \rightarrow \infty} s_n. \quad (2)$$

**Example 5** *The infinite Fibonacci word  $\mathbf{f} = 0100101001001010 \cdots$  is a Sturmian word [14], exactly  $\mathbf{f} = \mathbf{w}\left(\frac{1}{\phi^2}\right)$  where  $\phi = \frac{1+\sqrt{5}}{2}$  is the golden ratio.*

**Definition 6** *The Fibonacci morphism  $\sigma : \{0, 1\} \rightarrow \{0, 1\}$  is defined by  $\sigma(0) = 01$  and  $\sigma(1) = 0$ .*

The Fibonacci word  $\mathbf{f}$  satisfies that  $\lim_{n \rightarrow \infty} \sigma^n(1) = \mathbf{f}[1]$ .

### 3 The $k$ -Fibonacci words

**Definition 7** *The  $n$ th  $k$ -Fibonacci words are words over  $\{0, 1\}$  defined inductively as follows*

$$f_{k,0} = 0, \quad f_{k,1} = 0^{k-1}1, \quad f_{k,n} = f_{k,n-1}^k f_{k,n-2},$$

for all  $n \geq 2$  and  $k \geq 1$ . The infinite word

$$\mathbf{f}_k := \lim_{n \rightarrow \infty} f_{k,n}$$

is called the  $k$ -Fibonacci word.

It is clear that  $|f_{k,n}| = F_{k,n+1}$ . For  $k = 1$  we have the word  $\bar{\mathbf{f}} = 1011010110110 \dots$ , where  $\bar{\mathbf{a}}$  is a morphism, with  $\mathbf{a} \in \{0, 1\}$ , defined by  $\bar{0} = 1, \bar{1} = 0$ .

**Example 8** *The first  $k$ -Fibonacci words are*

$$\begin{aligned} f_1 &= 1011010110110 \cdots = \bar{\mathbf{f}}, & f_2 &= 0101001010010 \cdots, & f_3 &= 0010010010001 \cdots, \\ f_4 &= 0001000100010 \cdots, & f_5 &= 0000100001000 \cdots, & f_6 &= 0000010000010 \cdots. \end{aligned}$$

**Definition 9** *The  $k$ -Fibonacci morphism  $\sigma_k : \{0, 1\} \rightarrow \{0, 1\}$  is defined by  $\sigma_k(0) = 0^{k-1}1$  and  $\sigma_k(1) = 0^{k-1}10$ .*

**Theorem 10** *For all  $n \geq 0$ ,  $\sigma_k^n(0) = f_{k,n}$  and  $\sigma_k^{n+1}(1) = f_{k,n+1}f_{k,n}$ . Hence, the  $k$ -Fibonacci word  $\mathbf{f}_k$  satisfies that  $\lim_{n \rightarrow \infty} \sigma_k^n(0) = \mathbf{f}_k$ .*

**Proof.** We prove the two assertions about  $\sigma_k^n$  by induction on  $n$ . They are clearly true for  $n = 0, 1$ . Assume for all  $j < n$ ; we prove them for  $n$ :

$$\begin{aligned}\sigma_k^{n+1}(0) &= \sigma_k^n(0^{k-1}1) = (\sigma_k^n(0))^{k-1}\sigma_k^n(1) = f_{k,n}^{k-1}f_{k,n}f_{k,n-1} = f_{k,n}^kf_{k,n-1} = f_{k,n+1}. \\ \sigma_k^{n+2}(1) &= \sigma_k^{n+1}(0^{k-1}10) = (\sigma_k^{n+1}(0))^{k-1}\sigma_k^{n+1}(1)\sigma_k^{n+1}(0) = f_{k,n+1}^{k-1}f_{k,n+1}f_{k,n}f_{k,n+1} \\ &= f_{k,n+1}^kf_{k,n}f_{k,n+1} = f_{k,n+2}f_{k,n+1}. \quad \square\end{aligned}$$

### Proposition 11

1.  $|f_{k,n}|_1 = F_{k,n}$  and  $|f_{k,n+1}|_0 = F_{k,n+1} + F_{k,n}$  for all  $n \geq 0$ .
2.  $\lim_{n \rightarrow \infty} \frac{|f_{k,n}|_0}{|f_{k,n}|_1} = \frac{r_{k,1}^2}{1 + r_{k,1}}.$
3.  $\lim_{n \rightarrow \infty} \frac{|f_{k,n}|_0}{|f_{k,n}|_1} = r_{k,1}.$
4.  $\lim_{n \rightarrow \infty} \frac{|f_{k,n}|_0}{|f_{k,n}|_1} = 1 + \frac{1}{r_{k,1}}.$

**Proof.**

1. It is clear by induction on  $n$ .

$$\begin{aligned}2. \quad \lim_{n \rightarrow \infty} \frac{|f_{k,n}|_0}{|f_{k,n}|_1} &= \lim_{n \rightarrow \infty} \frac{F_{k,n+1}}{F_{k,n} + F_{k,n-1}} = \lim_{n \rightarrow \infty} \frac{\frac{F_{k,n+1}}{F_{k,n}}}{1 + \frac{F_{k,n-1}}{F_{k,n}}} = \frac{r_{k,1}^2}{1 + r_{k,1}}. \\ 3. \quad \lim_{n \rightarrow \infty} \frac{|f_{k,n}|_0}{|f_{k,n}|_1} &= \lim_{n \rightarrow \infty} \frac{F_{k,n+1}}{F_{k,n}} = r_{k,1}. \\ 4. \quad \lim_{n \rightarrow \infty} \frac{|f_{k,n}|_0}{|f_{k,n}|_1} &= \lim_{n \rightarrow \infty} \frac{F_{k,n} + F_{k,n-1}}{F_{k,n}} = 1 + \frac{1}{r_{k,1}}. \quad \square\end{aligned}$$

**Proposition 12** *The  $k$ -Fibonacci word and the  $n$ th  $k$ -Fibonacci word satisfy that*

1. *The word 11 is not a subword of the  $k$ -Fibonacci word,  $k \geq 2$ .*
2. *Let  $ab$  be the last two symbols of  $f_{k,n}$ . For  $n \geq 1$ , we have  $ab = 10$  if  $n$  is even and  $ab = 01$  if  $n$  is odd,  $k \geq 2$ .*
3. *The concatenation of two successive  $k$ -Fibonacci words is “almost commutative”, i.e.,  $f_{k,n-1}f_{k,n-2}$  and  $f_{k,n-2}f_{k,n-1}$  have a common prefix the length  $F_{k,n} + F_{k,n-1} - 2$  for all  $n \geq 2$ .*

**Proof.**

1. It suffices to prove that 11 is not a subword of  $f_{k,n}$ , for all  $n \geq 0$ . By induction on  $n$ . For  $n = 0, 1$  it is clear. Assume for all  $j < n$ ; we prove it for  $n$ . We know that  $f_{k,n} = f_{k,n-1}^k f_{k,n-2}$  so by the induction hypothesis we have that 11 is not a subword of  $f_{k,n-1}$  and  $f_{k,n-2}$ . Therefore, the only possibility is that 1 is a suffix and prefix of  $f_{k,n-1}$  or 1 is a suffix of  $f_{k,n-1}$  and a prefix of  $f_{k,n-2}$ , both there are impossible.
2. By induction on  $n$ . For  $n = 1, 2$  it is clear. Assume for all  $j < n$ ; we prove it for  $n$ . We know that  $f_{k,n+1} = f_{k,n}^k f_{k,n-1}$ , if  $n+1$  is even then by the induction hypothesis the last two symbols of  $f_{k,n-1}$  are 10, therefore the last two symbols of  $f_{k,n+1}$  are 10. Analogously, if  $n+1$  is odd.
3. By induction on  $n$ . For  $n = 1, 2$  it is clear. Assume for all  $j < n$ ; we prove it for  $n$ . By definition of  $f_{k,n}$ , we have

$$\begin{aligned} f_{k,n-1} f_{k,n-2} &= f_{k,n-2}^k f_{k,n-3} \cdot f_{k,n-3}^k f_{k,n-4} \\ &= (f_{k,n-3}^k f_{k,n-4})^k \cdot f_{k,n-3}^k f_{k,n-3} f_{k,n-4}, \end{aligned}$$

and

$$\begin{aligned} f_{k,n-2} f_{k,n-1} &= f_{k,n-3}^k f_{k,n-4} \cdot f_{k,n-2}^k f_{k,n-3} \\ &= f_{k,n-3}^k f_{k,n-4} \cdot (f_{k,n-3}^k f_{k,n-4})^k \cdot f_{k,n-3} \\ &= (f_{k,n-3}^k f_{k,n-4})^k f_{k,n-3}^k f_{k,n-4} f_{k,n-3}. \end{aligned}$$

Hence the words have a common prefix of length  $k(kF_{k,n-2} + F_{k,n-3}) + kF_{k,n-2} = k(F_{k,n-1} + F_{k,n-2})$ . By the induction hypothesis  $f_{k,n-3} f_{k,n-4}$  and  $f_{k,n-4} f_{k,n-3}$  have a common prefix of length  $F_{k,n-2} + F_{k,n-3} - 2$ . Therefore the words have a common prefix of length

$$k(F_{k,n-1} + F_{k,n-2}) + F_{k,n-2} + F_{k,n-3} - 2 = F_{k,n} + F_{k,n-1} - 2. \quad \square$$

**Definition 13** Let  $\Phi : \{0, 1\}^* \rightarrow \{0, 1\}^*$  be a map such that  $\Phi$  deletes the last two symbols, i.e.,  $\Phi(a_1 a_2 \cdots a_n) = a_1 a_2 \cdots a_{n-2}$  ( $n \geq 2$ ).

**Corollary 14** The  $n$ th  $k$ -Fibonacci word, satisfy for all  $n \geq 2$  that

1.  $\Phi(f_{k,n-1} f_{k,n-2}) = \Phi(f_{k,n-2} f_{k,n-1})$ .
2.  $\Phi(f_{k,n-1} f_{k,n-2}) = f_{k,n-2} \Phi(f_{k,n-1}) = f_{k,n-1} \Phi(f_{k,n-2})$ .
3. If  $f_{k,n} = \Phi(f_{k,n}) ab$ , then  $\Phi(f_{k,n-2}) ab \Phi(f_{k,n-1}) = f_{k,n-1} \Phi(f_{k,n-2})$ .

4. If  $f_{k,n} = \Phi(f_{k,n})ab$ , then  $\Phi(f_{k,n-2})(ab\Phi(f_{k,n-1}))^k = \Phi(f_{k,n})$ .

**Proof.** Parts (a) and (b) follow immediately from Proposition 12-(3) and because of  $|f_{k,n}| \geq 2$  for all  $n \geq 2$ . (c) In fact, if  $f_{k,n} = \Phi(f_{k,n})ab$  then from Proposition 12-(2) we have  $f_{k,n-2} = \Phi(f_{k,n-2})ab$ . Hence  $\Phi(f_{k,n-2})ab\Phi(f_{k,n-1}) = f_{k,n-2}\Phi(f_{k,n-1}) = f_{k,n-1}\Phi(f_{k,n-2})$ . (d) It is clear from (c) and definition of  $f_{k,n}$ .  $\square$

**Theorem 15**  $\Phi(f_{k,n})$  is a palindrome for all  $n \geq 1$  and  $k \geq 1$ .

**Proof.** By induction on  $n$ . If  $n = 2$  then  $\Phi(f_{k,2}) = (0^{k-1}1)^{k-1}0^{k-1}$  is a palindrome. Now suppose that the result is true for all  $j < n$ ; we prove it for  $n$ .

$$\begin{aligned} (\Phi(f_{k,n}))^R &= (\Phi(f_{k,n-1}^k f_{k,n-2}))^R = (f_{k,n-1}^k \Phi(f_{k,n-2}))^R = \Phi(f_{k,n-2})^R (f_{k,n-1}^k)^R \\ &= \Phi(f_{k,n-2})(f_{k,n-1}^R)^k. \end{aligned}$$

If  $n$  is even then  $f_{k,n} = \Phi(f_{k,n})10$  and from Corollary 14-(4), we have that

$$\begin{aligned} (\Phi(f_{k,n}))^R &= \Phi(f_{k,n-2})((\Phi(f_{k,n-1})01)^R)^k = \Phi(f_{k,n-2})(10(\Phi(f_{k,n-1}))^R)^k \\ &= \Phi(f_{k,n-2})(10\Phi(f_{k,n-1}))^k = \Phi(f_{k,n}). \end{aligned}$$

If  $n$  is odd, the proof is analogous.  $\square$

**Corollary 16** 1. If  $f_{k,n} = \Phi(f_{k,n})ab$  then  $ba\Phi(f_{k,n})ab$  is a palindrome.  
2. If  $u$  is a subword of the  $k$ -Fibonacci word, then so is its reversal,  $u^R$ .

**Theorem 17** Let  $\alpha = [0, \bar{k}]$  be an irrational number, with  $k$  a positive integer, then

$$w(\alpha) = f_k.$$

**Proof.** Let  $\alpha = [0, \bar{k}]$  an irrational number, then its associated standard sequence is

$$s_{-1} = 1, \quad s_0 = 0, \quad s_1 = s_0^{k-1}s_{-1} = 0^{k-1}1 \text{ and } s_n = s_{n-1}^k s_{n-2}, \quad n \geq 2.$$

Hence  $\{s_n\}_{n \geq 0} = \{f_{k,n}\}_{n \geq 0}$  and from equation (2), we have

$$w(\alpha) = \lim_{n \rightarrow \infty} s_n = f_k. \quad \square$$



**Remark.** Note that

$$[0, \overline{k}] = \frac{1}{k + \frac{1}{k + \frac{1}{k + \frac{1}{\ddots}}}} = \frac{-k + \sqrt{k^2 + 4}}{2} = -r_{k,2}$$

From the above theorem, we conclude that  $k$ -Fibonacci words are Sturmian words.

A fractional power is a word of the form  $z = x^n y$ , where  $n \in \mathbb{Z}^+$  and  $x \in \Sigma^+$ , and  $y$  is power prefix of  $x$ . If  $|z| = p$  and  $|x| = q$ , we say that  $z$  is a  $p/q$ -power, or  $z = x^{p/q}$ . In the expression  $x^{p/q}$ , the number  $p/q$  is the power's exponent. For example, 01201201 is a  $8/3$ -power,  $01201201 = (012)^{8/3}$ . The index of an infinite word  $w \in \Sigma^\omega$  is defined by

$$\text{Ind}(w) := \sup\{r \in \mathbb{Q}_{\geq 1} : w \text{ contains an } r\text{-power}\}$$

For example  $\text{Ind}(\mathbf{f}) > 3$  because the cube  $(010)^3$  occurs in  $\mathbf{f}$  at position 6. In [15] the authors proof that  $\text{Ind}(\mathbf{f}) = 2 + \phi \approx 3.618$ . A general formula for the index of a Sturmian word was given in [8].

**Theorem 18** *If  $\mathbf{u}$  is a Sturmian word of slope  $\alpha = [0, a_1, a_2, a_3, \dots]$ , then*

$$\text{Ind}(w) = \sup_{n \geq 0} \left\{ 2 + a_{n+1} + \frac{q_{n-1} - 2}{q_n} \right\},$$

where  $q_n$  is the denominator of  $\alpha = [0, a_1, a_2, a_3, \dots, a_n]$  and satisfies  $q_{-1} = 0, q_0 = 1, q_{n+1} = a_{n+1}q_n + q_{n-1}$ .

**Corollary 19** *The index of  $k$ -Fibonacci words is  $\text{Ind}(\mathbf{f}_k) = 2 + k + \frac{1}{r_{k,1}}$ .*

**Proof.**  $\mathbf{f}_k$  is a Sturmian word of slope  $\alpha = [0, \overline{k}]$ , then it is clear that  $q_n = F_{k,n+1}$ , and from above theorem

$$\text{Ind}(\mathbf{f}_k) = \sup_{n \geq 0} \left\{ 2 + k + \frac{F_{k,n} - 2}{F_{k,n+1}} \right\} = 2 + k + \frac{1}{r_{k,1}}. \quad \square$$

## 4 The $k$ -Fibonacci Word Curve

The Fibonacci word can be associated to a curve from a drawing rule. We must travel the word in a particular way, depending on the symbol read a particular action is produced, this idea is the same as that used in the L-Systems [17]. In this case, the drawing rule is called “odd-even drawing rule” [16], this is defined as shown in the following table.

Symbol	Action
1	Draw a line forward.
0	Draw a line forward and if the symbol 0 is in a position even then turn $\theta$ degree and if 0 is in a position odd then turn $-\theta$ degrees.

**Definition 20** The  $n$ th-curve of Fibonacci, denoted by  $\mathcal{F}_n$ , is obtained by applying the odd-even drawing rule to the word  $f_n$ . The Fibonacci word fractal  $\mathcal{F}$ , is defined as

$$\mathcal{F} := \lim_{n \rightarrow \infty} \mathcal{F}_n.$$

**Example 21** In Figure 1 we show the curve  $\mathcal{F}_{10}$  and  $\mathcal{F}_{17}$ . The graphics in this paper were generated using the software **Mathematica 9.0**, [20].





Figure 1: Fibonacci curves  $\mathcal{F}_{10}$  and  $\mathcal{F}_{17}$  corresponding to the words  $f_{10}$  and  $f_{17}$

Properties of Fibonacci Word Fractal can be found in [3, 4, 16].

**Definition 22** The  $n$ th  $k$ -curve of Fibonacci, denoted by  $\mathcal{F}_{k,n}$ , is obtained by applying the odd-even drawing rule to the word  $f_{k,n}$ . The  $k$ -Fibonacci word curve  $\mathcal{F}_k$  is defined as

$$\mathcal{F}_k := \lim_{n \rightarrow \infty} \mathcal{F}_{k,n}.$$

Table 1: Some curves  $\mathcal{F}_{k,n}$  with  $\theta = 90^\circ$ 

In Table 1, we show some curves  $\mathcal{F}_{k,n}$  with an angle  $\theta = 90^\circ$ .


In Table 2, we show some curves  $\mathcal{F}_{k,n}$  with an angle  $\theta = 60^\circ$ . In general these curves have a lot of patterns because the index is large, see Corollary 19.

**Proposition 23** *The  $k$ -Fibonacci word curve and the curve  $\mathcal{F}_{k,n}$  have the following properties:*

1. The  $k$ -Fibonacci curve  $\mathcal{F}_k$  is composed only of segments of length 1 or 2.
2. The  $\mathcal{F}_{k,n}$  is symmetric.
3. The number of turns in the curve  $\mathcal{F}_{k,n}$  is  $F_{k,n} + F_{k,n-1}$ .
4. If  $n$  is even then the  $\mathcal{F}_{k,n}$  curve is similar to the curve  $\mathcal{F}_{k,n-2}$  and if  $n$  is odd then the  $\mathcal{F}_{k,n}$  curve is similar to the curve  $\mathcal{F}_{k,n-3}$ .

**Proof.**

1. It is clear from Proposition 12-1, because 110 and 111 are not subwords of  $\mathbf{f}_k$ .
2. It is clear from Theorem 15, because  $\mathbf{f}_{k,n} = \Phi(\mathbf{f}_{k,n})\mathbf{ab}$ , where  $\Phi(\mathbf{f}_{k,n})$  is a palindrome.
3. It is clear from definition of odd-even drawn rule and because  $|\mathbf{f}_{k,n+1}|_0 = F_{k,n+1} + F_{k,n}$ .

Table 2: Some curves  $\mathcal{F}_{k,n}$  with  $\theta = 60^\circ$ 

4. If  $n$  is even. It is clear that  $\sigma_k^2(f_{k,n-2}) = f_{k,n}$ . We are going to prove that  $\sigma_k^2$  guaranties the odd-even alternation required by the odd-even drawing rule. In fact,  $\sigma_k^2(0) = \sigma_k(0^{k-1}1) = (0^{k-1}1)^k 0$  and  $\sigma_k^2(1) = (0^{k-1}1)^k 0^k 1$ . As  $k$  is even, then  $|\sigma_k^2(0)|$  and  $|\sigma_k^2(1)|$  are odd. Hence if  $|w|$  is even (odd) then  $|\sigma_k^2(w)|$  is even (odd). Since  $\sigma_k^2$  preserves the parity of length then any subword in the  $k$ -Fibonacci word preserves the parity of position.

Finally, we have to proof that the resulting angle of a pattern must be preserved or inverted by  $\sigma_k^2$ . Let  $\alpha(w)$  be the function that gives the resulting angle of a word  $w$  through the odd-even drawing rule of angle

$\theta$ . Note that  $a(00) = 0^\circ$ ,  $a(01) = -\theta^\circ$  and  $a(10) = \theta^\circ$ . Therefore

$$\begin{aligned}
 a(\sigma_k^2(00)) &= a((0^{k-1}1)^k 0 (0^{k-1}1)^k 0) \\
 &= a((0^{k-1}1)^k) a(00^{k-1}) a(1(0^{k-1}1)^{k-1}0) \\
 &= -k\theta^\circ + 0^\circ + k\theta^\circ = 0^\circ \\
 a(\sigma_k^2(01)) &= a((0^{k-1}1)^k 0 (0^{k-1}1)^k 0^k 1) \\
 &= a((0^{k-1}1)^k) a(00^{k-1}) a(1(0^{k-1}1)^{k-1}0) a(0^{k-1}1) \\
 &= -k\theta^\circ + 0^\circ + k\theta^\circ - \theta^\circ = -\theta^\circ \\
 a(\sigma_k^2(10)) &= a((0^{k-1}1)^k 0^k 1 (0^{k-1}1)^k 0) \\
 &= a((0^{k-1}1)^k) a(0^k) a(1(0^{k-1}1)^k 0) \\
 &= -k\theta^\circ + 0^\circ + (k+1)\theta^\circ = \theta^\circ
 \end{aligned}$$

Then  $\sigma_k^2$  inverts the resulting angle, i.e.,  $a(w) = -a(\sigma_k^2(w))$  for any word  $w$ . Therefore the image of a pattern by  $\sigma_k^2$  is the rotation of this pattern by a rotation of  $-\theta^\circ$ . Since  $\sigma_k^2(f_{k,n-2}) = f_{k,n}$ , then the curve  $\mathcal{F}_{k,n}$  is similar to the curve  $\mathcal{F}_{k,n-2}$ .

If  $n$  is odd the proof is similar, but using  $\sigma_k^3$ . □

**Example 24** In Figure 2  $\mathcal{F}_{4,4}$  looks similar to  $\mathcal{F}_{4,6}, \mathcal{F}_{4,8}$  and so on.




Figure 2: Curves  $\mathcal{F}_{4,4}, \mathcal{F}_{4,6}, \mathcal{F}_{4,8}$  with  $\theta = 60^\circ$

In Figure 3  $\mathcal{F}_{5,3}$  looks similar to  $\mathcal{F}_{5,6}$ .

## Acknowledgements

The first author was partially supported by Universidad Sergio Arboleda under grant number USA-II-2012-14.




Figure 3: Curves  $\mathcal{F}_{5,3}, \mathcal{F}_{5,6}$  with  $\theta = 60^\circ$

## References

- [1] J. Allouche, J. Shallit, *Automatic Sequences*, Cambridge University Press, 2003.  $\Rightarrow$  214, 216
- [2] J. Berstel, Fibonacci words—a survey, in: G. Rosenberg, A. Salomaa (Eds.), *The Book of L*, Springer, Berlin, 1986, pp. 11–26.  $\Rightarrow$  213
- [3] A. Blondin-Mass, S. Brlek, A. Garon, S. Labb, Two infinite families of polyominoes that tile the plane by translation in two distinct ways, *Theoret. Comput. Sci.* **412** (2011) 4778–4786.  $\Rightarrow$  213, 221
- [4] A. Blondin-Mass, S. Brlek, S. Labb, M. Mends France, Complexity of the Fibonacci snowflake, *Fractals* **20** (2012) 257–260.  $\Rightarrow$  221
- [5] C. Bolat, H. Kse, On the properties of  $k$ -Fibonacci numbers, *Int. J. Contemp. Math. Sciences*, **22**, 5 (2010) 1097–1105.  $\Rightarrow$  213
- [6] J. Cassaigne, On extremal properties of the Fibonacci word, *RAIRO - Theor. Inf. Appl.* **42**, (4) (2008) 701–715.  $\Rightarrow$  213
- [7] W. Chuan, Fibonacci words, *Fibonacci Quart.*, **30**, 1 (1992) 68–76.  $\Rightarrow$  213
- [8] D. Damanik, D. Lenz, The index of Sturmian sequences, *European J. Combin.*, **23** (2002) 23–29.  $\Rightarrow$  220
- [9] A. de Luca, Sturmian words: structure, combinatorics, and their arithmetics, *Theor. Comput. Sci.* **183**, 1 (1997) 45–82.  $\Rightarrow$  213
- [10] S. Falcon, A. Plaza, On  $k$ -Fibonacci sequences and polynomials and their derivatives, *Chaos Solitons Fractals* **39**, 3 (2009) 1005–1019.  $\Rightarrow$  213
- [11] S. Falcon, A. Plaza, On the Fibonacci  $k$ -numbers, *Chaos Solitons Fractals* **32**, 5 (2007) 1615–1624.  $\Rightarrow$  212, 213
- [12] S. Falcon, A. Plaza, The  $k$ -Fibonacci sequence and the Pascal 2-triangle, *Chaos Solitons Fractals* **33**, 1 (2007) 38–49.  $\Rightarrow$  213
- [13] T. Koshy, *Fibonacci and Lucas numbers with Applications*, Wiley-Interscience, 2001.  $\Rightarrow$  212
- [14] M. Lothaire, *Algebraic Combinatorics on Words*, Encyclopedia of Mathematics and its Applications, Cambridge University Press, 2002.  $\Rightarrow$  213, 214, 215, 216

- [15] F. Mignosi, G. Pirillo, Repetitions in the Fibonacci infinite word, *RAIRO – Theor. Inf. Appl.* **26** (1992) 199–204.  $\Rightarrow$  213, 220
- [16] A. Monnerot, The Fibonacci word fractal, preprint, 2009.  $\Rightarrow$  213, 221
- [17] P. Prusinkiewicz, A. Lindenmayer, *The Algorithmic Beauty of Plants*, Springer-Verlag, Nueva York, 2004.  $\Rightarrow$  221
- [18] J. Ramírez, Incomplete k-Fibonacci and k-Lucas numbers, *Chinese Journal of Mathematics* (2013).  $\Rightarrow$  213
- [19] J. Ramírez. Some properties of convolved k-Fibonacci numbers. *ISRN Combinatorics* (2013) ID759641, 5pp.  $\Rightarrow$  213
- [20] J. Ramírez, G. Rubiano, Generating fractals curves from homomorphisms between languages [with *Mathematica*<sup>®</sup>] (Spanish), *Rev. Integr. Temas Mat.* **30**, 2 (2012) 129–150.  $\Rightarrow$  221
- [21] J. Ramírez, G. Rubiano, R. de Castro, A generalization of the Fibonacci word fractal and the Fibonacci snowflake, preprint arXiv:1212.1368, 2013.  $\Rightarrow$  213
- [22] W. Rytter, The structure of subword graphs and suffix trees of Fibonacci words, *Theoret. Comput. Sci.* **363**, 2 (2006) 211–223.  $\Rightarrow$  213
- [23] A. Salas. About k-Fibonacci numbers and their associated numbers, *Int. Math. Forum.* **50**, 6 (2011) 2473–2479.  $\Rightarrow$  213

*Received: June 12, 2013 • Revised: October 25, 2013*