

Spread of *Eragrostis albensis* (Poaceae) and *Dittrichia graveolens* (Asteraceae) in the southern Poland

Anna Wróbel & Marcin Nobis

Spread of *Eragrostis albensis* (Poaceae) and *Dittrichia graveolens* (Asteraceae) in the southern Poland.
– Acta Mus. Siles. Sci. Natur., 66: 117-120, 2017.

Abstract: New localities of *Eragrostis albensis* H. Scholz and *Dittrichia graveolens* (L.) Greuter have been found in the southern Poland. The former taxon is currently considered a kenophyte (epicophyte and holoagriophyte) in the country. It occurs on sandy alluvia along Vistula, Oder and San River Valleys as well as on anthropogenic sites mainly in the eastern and south-eastern Poland. The latter species is a recent newcomer regarded as an ephemerophyte, which so far has been reported from only one locality in Śląskie Province. In 2017 we discovered 16 new localities of *E. albensis* and five of *D. graveolens* on the territory of the southern Poland. Populations of both species consisted of few to several dozen individuals which grew within anthropogenic habitats, mainly roadsides. Distribution maps of both species in the southern Poland were presented.

Key words: new locality, expansion, spreading species, distribution, Poland

Introduction

It is commonly observed that anthropogenic habitats serve as perfect migration corridors for plants. Thus, numerous species propagate along transport networks where effective long-distance seed dispersal may occur (Clifford 1959; Tikka et al. 2001; Rauschert et al. 2017). In addition, many alien species spread rapidly through human-modified landscape and especially in this way invade new areas beyond their native distribution (Vakhlamova et al. 2016; Benedetti & Morelli 2017). In Central Europe, two species, i.e. *Eragrostis albensis* H. Scholz and *Dittrichia graveolens* (L.) Greuter, may serve as good examples of plants which follow such propagation strategy and still expand their non-native range limits (Guzik & Sudnik-Wójcikowska 2005; Michalewska & Nobis 2005; Kocián 2015).

Eragrostis albensis H. Scholz is a terophyte recently distinguished as a new species to science (Scholz 1996). It belongs to *Eragrostis pectinacea-pilosa* complex which is regarded as taxonomically problematic group within large *Eragrostis* genus (Scholz 1996; Špryňar & Kubát 2004). According to Scholz (1996), *E. albensis* should be considered Central European endemic taxon which could have recently originated after rapid speciation. However, soon after it was described in Germany, *E. albensis* was also noted at numerous localities in: the Netherlands, Austria, the Czech Republic, Slovakia, Poland, Ukraine, Belarus and eastern part of Russia (Scholz 1996; Špryňar & Kubát 2004; Guzik & Sudnik-Wójcikowska 2005; Hohla 2006; Hohla & Kleesadle 2006; Scholz 2010).

The revision of herbaria materials from Poland indicated that almost all specimens previously identified as *E. pilosa* (L.) P. Beauv. should be classified as *E. albensis* (Guzik & Sudnik-Wójcikowska 2005; Michalewska & Nobis 2005). Moreover, the oldest individuals were in fact not collected in Germany; it turned out that the species was recorded much earlier in Poland, the Czech Republic and Slovakia (Guzik & Sudnik-Wójcikowska 2005; Špryňar & Kubát 2004). As a consequence, Špryňar & Kubát (2004) suggested that *E. albensis* should be treated rather as a neophyte in Central Europe, which might have probably immigrated from

the east Eurasia. In our opinion, such hypothesis seems to be much more convincing than possibility of recent speciation postulated by Scholz (1996).

Eragrostis albensis occupies natural habitats such as sandy river banks, but it also grows on anthropogenic sites, especially along roads and railway tracks (Guzik & Sudnik-Wójcikowska 1996, 2005; Michalewska & Nobis 2005; Wrzesień 2005; Kącki & Szczęśniak 2009; Nobis & Nobis 2015). Still, it is difficult to conclude whether spread of *E. albensis* began in one type of habitat or probably, in both simultaneously. Possible hypotheses were thoroughly discussed by Guzik & Sudnik-Wójcikowska 1996, 2005 and Michalewska & Nobis 2005.

In Poland, *E. albensis* is an alien species, regarded as an holoagriophyte or epecophyte. So far, it has been observed on sandy alluvia along Vistula, Oder and San River valleys (Guzik & Sudnik-Wójcikowska 1996, 2005; Kącki & Szczęśniak 2009). It also occupies many localities within anthropogenic sites mainly in eastern and south-eastern Poland (Michalewska & Nobis 2005; Wrzesień 2005; Nobis & Nobis 2006, 2010).

Another alien species to Central Europe, *Dittrichia graveolens*, is an annual plant characteristic for the flora of the Mediterranean region (Ball 1976; Brullo & de Marco 2000). Currently, its spread is observed in Western Atlantic European coast, Middle East (Brullo & de Marco 2000), Central Europe (Frank 2006; Frajman & Kaligarič 2009; Király et al. 2014; Kocián 2015) as well as in two other continents where it was introduced, i.e. North America and Australia (Kloot 1987; Brownsey et al. 2013). In Central Europe, the species' rapid expansion has been noted along roads, particularly motorways (Frajman & Kaligarič 2009; Király et al. 2014; Kocián 2015). Its presence has been recently also confirmed in Poland (Kocián 2015). In 2013 *D. graveolens* was found in Śląskie Province near the S1 expressway (DF 90 ATPOL cartogram unit); hence, the species was regarded as an ephemerophyte in the country. Although *D. graveolens* has been so far observed in Poland on only this single locality, its further spread has been suggested as very probable (Kocián 2015).

New localities of *Eragrostis albensis*

During field studies in 2017 the Authors found 16 new localities of *E. albensis* on the territory of Małopolskie Province (Fig. 1). In accordance with ATPOL cartogram method (Zajac 1978) these sites are located within 9 ATPOL grid units (squares 10 km x 10 km):

DF 99 – Myślenice, Kazimierza Wielkiego Street; Stróża; **DG 09** – Krzczonów; **EF 81** – Marszowice near Gdów; **EG 00** – Mszana Dolna, Krakowska Street; Kasinka Mała; Lubień; **EG 11** – Mszana Górna; **EG 12** – Szczawa; Kamienica; **EG 23** – Maszkowice near Jazowsko; Tylmanowa; Zabrzeż; **EG 24** – Naszacowice; **EG 33** – Kłodne near Tylmanowa; Krościenko nad Dunajcem, Sobieskiego Street and Jagiellońska Street.

The distribution map of *E. albensis* in the southern Poland (Fig. 1) was based on Nobis & Nobis 2010, 2015 and it was supplemented with newly found localities. Some of them are currently the southernmost records of *E. albensis* in Poland (Fig. 1). In Małopolskie Province the species spreads within anthropogenic habitats. It grows in the crevices between roads and curbs or in pavement slabs. In all localities from few to several dozen individuals were noted. It is highly probable that further spread of *E. albensis* will be observed in southern Poland in the nearest future.

New localities of *Dittrichia graveolens*

In 2017 the species was found on roadsides, on few localities along A4 motorway between Kraków and Katowice as well as on western ring road of Kraków (Fig. 2). In accordance with ATPOL cartogram method (Zajac 1978) these sites are located within four ATPOL grid units (squares 10 km x 10 km): **DF 55** – Wygoda; **DF 56** – Chrzanów; Bołęcin; **DF 66** – Regulice; **DF 69** – Kryspinów / Kraków Bielany. In all localities from few to several dozen individuals




Fig 1: Distribution of *Eragrostis albensis* H. Scholz in the southern Poland: ● – known localities (according to Nobis & Nobis 2010, 2015); ▲ – new localities.




Fig 2: Distribution of *Dittrichia graveolens* (L.) Greuter in the southern Poland: ● – known locality (according to Kocián 2015); ▲ – new localities.

of the species were noted. Our findings proved that *D. graveolens* quickly extending its range in southern Poland, and currently the species should be regarded as an epiphyte, established in anthropogenic habitats. It is very likely that further localities of this potentially invasive species will soon be found also in other parts of the country.

References

Ball P.W. (1976): *Dittrichia* W. Greuter. In: Tutin T.G., Heywood V.H., Burges N.A., Valentine D.H., Walters S.M. & Webb D.A. [eds.]: *Flora Europaea* 4, pp. 136–137. Cambridge University Press, Cambridge.

- Benedetti Y. & Morelli F. (2017): Spatial mismatch analysis among hotspots of alien plant species, road and railway networks in Germany and Austria. – PLoS ONE 12(8): 1–13.
- Brownsey R., Kyser G.B. & DiTomaso J.M. (2013): Stinkwort is rapidly expanding its range in California. – California Agriculture 67: 110–115.
- Brullo S. & de Marco G. (2000): Taxonomical revision of the genus *Dittrichia* (Asteraceae). – Portugaliae Acta Biologica 19: 341–354.
- Clifford H. T. (1959): Seed dispersal by motor vehicles. – Journal of Ecology 47(2): 311–315.
- Frajman B. & Kaligarič M. (2009): *Dittrichia graveolens*, nova tujerodna vrsta slovenske flore. – Hladnikia 24: 35–43.
- Frank C. (2006): Beobachtungen zur Einbürgerung neuer Arten in Sachsen-Anhalt. – Mitt. florist. Kart. Sachsen-Anhalt (Halle) 11: 81–90.
- Guzik J. & Sudnik-Wójcikowska B. (1996): The spread and habitats of *Eragrostis pilosa* (Poaceae) in the Vistula valley. – Fragmenta Floristica et Geobotanica 41(2): 753–769.
- Guzik J. & Sudnik-Wójcikowska B. (2005): Critical review of species of the genus *Eragrostis* in Poland. In: Frey L. (ed.): Biology of grasses, pp. 45–58. Instytut Botaniki im. W. Szafera, Polska Akademia Nauk, Kraków.
- Hohla M. (2006): Neues über die Verbreitung von *Eragrostis albensis*, *Eragrostis multicaulis* und *Eragrostis pilosa* in Österreich. – Linzer Biologische Beiträge 38(2): 1233–1253.
- Hohla M. & Kleesadle G. (2006): *Eragrostis albensis* – neu für Österreich – und weitere bemerkenswerte Funde zur Flora von Oberösterreich. – Beiträge zur Naturkunde Oberösterreichs 16: 197–202.
- Kącki Z. & Szczęśniak E. (2009): Gatunki z rodzaju miłka *Eragrostis* spp. In: Dajdok Z. & Pawlaczyk P. (eds.): Inwazyjne gatunki roślin ekosystemów mokradłowych Polski, pp. 77–79. Wydawnictwo Klubu Przyrodników, Świebodzin.
- Király G., Eliáš P. jun. & Ditě D. (2014): Two thermophilic alien species new to the flora of Slovakia. – Thaiszia Journal of Botany 24: 125–134.
- Kloot P.M. (1987): The naturalised flora of South Australia 1. The documentation of its development. – Journal of the Adelaide Botanic Gardens 10(1): 81–90.
- Kocián P. (2015): *Dittrichia graveolens* (L.) Greuter – a new alien species in Poland. – Acta Musei Silesiae, Scientiae Naturales 64: 193–197.
- Michalewska A. & Nobis M. (2005): Ekspansja *Eragrostis albensis* (Poaceae) na antropogenicznych siedliskach w południowo-wschodniej Polsce. – Fragmenta Floristica et Geobotanica Polonica 12(1): 45–55.
- Nobis M. & Nobis A. (2006): Interesujące, rzadkie i rozprzestrzeniające się gatunki roślin naczyniowych notowane na terenach kolejowych w południowo-wschodniej Polsce. – Fragmenta Floristica et Geobotanica Polonica 13(2): 301–308.
- Nobis M. & Nobis A. (2010): Rzadkie i rozprzestrzeniające się gatunki roślin naczyniowych na terenach kolejowych wschodniej części polskich Karpat i ich przedpola. – Fragmenta Floristica et Geobotanica Polonica 17(2): 275–284.
- Nobis M. & Nobis A. (2015): *Eragrostis albensis* H. Scholz. In: Zajac A. & Zajac M. (eds.): Rozmieszczenie kenofitów w Karpatach polskich i na ich przedpolu, pp. 83–84. Instytut Botaniki, Uniwersytet Jagielloński, Kraków.
- Rauschert E.S.J., Mortensen D.A. & Bloser S.M. (2017): Human-mediated dispersal via rural road maintenance can move invasive propagules. – Biological Invasions 19: 2047–2058.
- Scholz H. (1996): *Eragrostis albensis* (Gramineae), das Elb-Liebesgras – ein neuer Neo-Endemit Mitteleuropas. – Verhandlungen des Botanischen Vereins für die Provinz Brandenburg 128(1995): 73–82.
- Scholz H. (2010): *Eragrostis* N.M. Wolf. In: Sukhorukov A.P. (ed.): The identification manual of vascular plants of the Tambov region, pp. 84–85. Grif and K, Tula.
- Špryňar P. & Kubát K. (2004): *Eragrostis albensis* and *Eragrostis pectinacea*, two new alien grass species for the flora of the Czech Republic (Poaceae). – Zprávy České Botanické Společnosti 39: 1–24.
- Tikka P.M., Högmänder H. & Koski P.S. (2001): Road and railway verges serve as dispersal corridors for grassland plants. – Landscape Ecology 16: 659–666.
- Vakhlamova T., Rusterholz H.P., Kanibolotskaya Y., Baur B. (2016): Effects of road type and urbanization on the diversity and abundance of alien species in roadside verges in Western Siberia. – Plant Ecology 217: 241–252.
- Wrzesień M. (2005): Alien species of grasses in the flora of the railway areas of the central-eastern Poland. In: Frey L. (ed.): Biology of grasses, pp. 139–150. Instytut Botaniki im. W. Szafera, Polska Akademia Nauk, Kraków.
- Zajac A. (1978): Atlas of distribution of vascular plants in Poland (ATPOL). – Taxon 27(5/6): 481–484.

Authors' addresses: Anna Wróbel & Marcin Nobis, Department of Taxonomy, Phytogeography and Paleobotany of Institute of Botany, Jagiellonian University, Kopernika 27, 31-501 Kraków, Poland.
E-mail of corresponding author: m.nobis@uj.edu.pl