

# Optimization of *in vitro* asymbiotic seed germination protocol for *Serapias vomeracea*

Arda Acemi\* and Fazıl Özen

## Abstract

*Serapias vomeracea* is an economically important orchid species which is over-collected from nature, because of its glucomannan-rich tubers. Thus, optimization of *in vitro* culture methodology in this species is required to meet industrial needs and to secure its populations in nature. This study aimed to optimize the surface sterilization protocol for *S. vomeracea* seeds and to select the optimal seed germination medium by comparing the commonly used media in *in vitro* orchid culture. During seed surface sterilization, ethyl alcohol (EtOH) pre-treatment prior to sodium hypochlorite (NaOCl) treatment increased the disinfection success and viable seed yield when examined using the triphenyl tetrazolium chloride (TTC) seed viability test. Also, low-g force centrifugation as an additional step in the surface sterilization method separated the seeds without embryo from the viable seeds and thereby decreased potential counting errors after incubation. Comparison of media showed that solid Knudson C (KN) medium induced the highest number of germinated seeds. However, seed germination success of Lindemann (LN) and Vacin & Went (VW) media was found to be higher when the media was used in liquid form. Half-strength liquid VW was the only medium that induced higher germination success than the other full-strength media. The highest number of ungerminated seeds was found when using KN medium whereas liquid VW medium gave the lowest number. In general, protocorm formation was triggered when the media were used in liquid form. However, rhizoid elongation was suppressed in liquid media. These findings suggest that this optimized seed surface sterilization method offers a simple and effective alternative to classical methods. Additionally, solid KN medium may be considered as a cost-effective and reliable alternative to other commonly-used complex media in *S. vomeracea* cultures.

**Keywords:** Medium selection, method optimization, orchid seed germination, seed viability, surface sterilization

## Introduction


*Serapias vomeracea* is a wide-spread, perennial orchid species that can be found in arid meadows, on abandoned agricultural land and heathlands up to a height of 1000 m. The plant's distribution extends over Mediterranean-Atlantic countries (Portugal, Spain, France, Italy, the Balkans, Greece, and Turkey). However, it is also found in North Africa (Morocco and Tunisia) where it is under high pressure of inbreeding depression (1). This species is one of the most widespread tuberous orchids in Turkey. Thus it is easily accessible in many cities and regions. Ease of accessibility in the environment and the economic value of its tubers make *S. vomeracea* one of the most collected orchid species from its native habitats (2). The component that makes the tubers of orchids so valuable is the polysaccharide glucomannan. Glucomannan is obtained from the tubers of orchids and the Konjac plant (*Amorphophallus konjac*). Glucomannan is currently used as a raw material for Salep production which is a traditional hot beverage in Turkey and is also used in food supplements, especially in weight loss capsules and other nutritional supplements, due to its hydrophilic properties (2). Also, the polysaccharide glucomannan is used as a thickener in the food industry and has been used to

Department of Biology, Faculty of Arts and Sciences, Kocaeli University, İzmit, Kocaeli, Turkey

\*Corresponding author: A. Acemi  
E-mail: arda.acemi@kocaeli.edu.tr

DOI: 10.2478/ebtj-2019-0017

© 2019 Authors. This work was licensed under the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 License.



**Figure 1.** Stereomicroscope view of *Serapias vomeracea* seeds. A: 16x magnification, the scale bar represents 0.5 mm, B: 80x magnification, the scale bar represents 0.1 mm.

treat high cholesterol (3, 4). Even though tuberous orchids are not yet extinct, there is evidence of population decline due to over-collection because of increasing commercial value and rising demand from industry. Therefore, an effective strategy should be designed, not only to conserve them in nature but also to take advantage of their economic potential through scientific investigation and development of commercially viable farming. One element of this strategy would be to develop and standardize both the optimum seed surface sterilization method and an efficient *in vitro* seed germination protocol that can be applied to economically valuable orchids including exotic hybrid ones, on a commercial scale.

As orchids can produce countless, dust-like seeds in their ovaries, the seeds have been mostly preferred as starting materials in tissue culture studies on terrestrial orchids (5). However, orchids are slow-growing plants in nature, and their seeds require a mycorrhizal relationship to germinate (6). Thus their seed germination rates are very low in nature. In contrast, *in vitro* cultures can offer an alternative for faster and more efficient production of orchids, and similar plants, that are difficult to germinate and develop in nature (7). Examination of various culture media, especially those developed mainly for orchids, can offer a detailed understanding of their nutritional requirements for *in vitro* germination.

In surface sterilization experiments of many *in vitro* germination studies, disinfectants such as ethyl alcohol (EtOH), sodium hypochlorite (NaOCl), calcium hypochlorite (CaOCl), hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), silver nitrate (AgNO<sub>3</sub>) and mercuric chloride (HgCl<sub>2</sub>) have been commonly used (8). However, less toxic chemical compounds should be used in the surface sterilization treatment of orchid seeds since they have thin seed coats which make the embryo particularly vulnerable to toxic chemicals such as disinfectants (9). In this context, in addition to disinfectant selection, treatment duration is also important.

Another major factor is the method of surface sterilization used for orchid seeds. Orchid seeds are generally weighed and put into a pouch made from filter paper before they are

dipped into disinfectants for surface sterilization following which they are sown using a scalpel or inoculation loop (10, 11). However, because the seeds are so small and difficult to handle a large number of seeds remain in the paper pouches.

As can be seen, there is scope for a range of improvements in the successful culturing of orchids including a selection of the optimal sterilizing agent, improved seed recovery following sterilization and selection of the optimal culture medium. Also, by investigating the optimal *in vitro* seed germination medium for *Serapias vomeracea* a better understanding of the plant's developmental physiology would be forthcoming, which would be of benefit in the development of controlled farming of this glucomannan-rich species for industrial purposes. Therefore, this study aimed to optimize the surface sterilization protocol for *S. vomeracea* seeds and to select the optimal seed germination medium by comparing the commonly used media in *in vitro* orchid culture.

## Materials and Methods

### Seed source, and storage

Seeds of *Serapias vomeracea* (Burm.f.) Briq. were used as starting material for *in vitro* cultures (Fig. 1). These seeds were obtained from the Directorate of Aegean Agricultural Research Institute (Menemen, İzmir, Turkey). The orchid seeds were kept dry and in the dark at 4 °C until they were used.

### Development of an efficient surface sterilization protocol

The seeds were weighed to 1 mg and then placed in round-bottom centrifuge tubes. In order to understand whether a pretreatment procedure was necessary for surface sterilization and to develop a more effective disinfection protocol, the seeds were divided into four groups and subjected to different sterilization treatments as shown in Table 1. During surface sterilization treatment, centrifuge tubes were stirred on a vortex shaker (Biosan Multi-Vortex V-32, Riga, Latvia). To reduce surface tension, 0.5 ml of Tween-20 was added to the sterilization stock solutions. After surface sterilization procedures, the viable embryos were separated from empty seed coats by low g

**Table 1.** Seed surface sterilization treatments tested on *Serapias vomeracea* seeds

Treatment	Concentration (%)	Time (min)
NaOCl	0.5	8
NaOCl	1.0	8
EtOH + NaOCl	70 + 0.5	0.5 + 8
EtOH + NaOCl	70 + 1.0	0.5 + 8

centrifugation using the weight difference between viable seeds and non-viable seeds. The seed/sterilization mixtures were centrifuged at 2000 g for 1 min (CAPP CappRondo Micro-centrifuge, Nordhausen, Germany). The non-viable seed coats were then removed with the supernatant using a micropipette whereas the seeds with embryo formed a loose pellet at the bottom of the tube. Finally, viable seeds were rinsed with sterile H<sub>2</sub>O to remove all traces of remaining NaOCl. The used H<sub>2</sub>O was finally pipetted out from the tubes. To assess the success of the various surface sterilization treatments, the remaining seeds were sown onto Knudson C (KN) medium (12) and the cultures were observed for 30 days.

#### Determination of seed viability

The tetrazolium (2,3,5-Triphenyltetrazolium chloride; TTC) test, which is a reliable and validated method for various plant seeds, was used for determination of seed viability (5). TTC solution 1% (*m:v*) was freshly prepared before the test. TTC (0.2 g) was dissolved in 20 ml of distilled water, pH was adjusted to 6.95, the solution was sterilized through a syringe filter, and then it was stored in an amber bottle at 4 °C until use. The seeds were treated with 70% EtOH for 2 min followed by 1% (*m:v*) NaOCl for 10 min. The seeds were divided into three groups prior to conducting the TTC test of seed viability after exposure to surface sterilization and H<sub>2</sub>O treatments. The first group of seeds was subjected to the TTC test without any prior treatment. The second group was left in the dark at 30 °C for 24 h in distilled water (dH<sub>2</sub>O), and then the TTC test was performed. The last group was subjected directly to the above-mentioned disinfection processes before the TTC test was applied. Each group of seeds was examined under a stereomicroscope (Olympus SZ61, Tokyo, Japan) following the TTC test and the stained seeds were counted.

#### Preparation of seed germination media, and seed sowing

In the experiments, KN, Vacin and Went (VW), and Lindemann (LN) media (13, 14) were used in full and half strength (½), both in solid (S) and liquid (L) forms. These formulations of the commercially available media are given in Table 2. All media were supplemented with 20 g L<sup>-1</sup> sucrose and, if solidified, with 3.5 g L<sup>-1</sup> phytagel (Duchefa, Haarlem, The Netherlands). The pH of the media was set to 5.6 using 1 N NaOH or 1 N HCl before autoclaving at 121°C under pressure of 118 kPa for 20 min (Nüve NC-40M, Ankara, Turkey). To transfer all the sur-

**Table 2.** Media compositions tested in *in vitro* germination of *Serapias vomeracea* seeds

Content	Quantity (mg L <sup>-1</sup> )		
Macroelements	KN	VW	LN
KNO <sub>3</sub>		525.000	
NH <sub>4</sub> NO <sub>3</sub>	500.000		
NH <sub>4</sub> H <sub>2</sub> PO <sub>4</sub>	500.000		
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>		500.000	1000.000
Ca <sub>3</sub> (PO <sub>4</sub> ) <sub>2</sub>		200.000	
Ca(NO <sub>3</sub> ) <sub>2</sub>	347.200		347.200
MgSO <sub>4</sub>	122.125	122.100	58.620
KH <sub>2</sub> PO <sub>4</sub>	250.000	250.000	135.000
KCl	250.000		1050.000
Microelements			
AlCl <sub>3</sub> ·6H <sub>2</sub> O			0.056
NiCl <sub>2</sub> ·6H <sub>2</sub> O			0.031
MnSO <sub>4</sub> ·H <sub>2</sub> O	5.682	5.600	0.052
KI			0.099
H <sub>3</sub> BO <sub>3</sub>			1.014
ZnSO <sub>4</sub> ·7H <sub>2</sub> O			0.565
CuSO <sub>4</sub> ·5H <sub>2</sub> O			0.019
Iron source			
C <sub>6</sub> H <sub>5</sub> FeO <sub>7</sub>			4.400
FeSO <sub>4</sub> ·7H <sub>2</sub> O	25.000	27.800	
Complexing agent			
Na <sub>2</sub> EDTA·2H <sub>2</sub> O		37.260	
Vitamins			
Nicotinic acid			1.000
Pyridoxine-HCl			1.000
Thiamin-HCl		0.400	10.000
myo-Inositol			100.000
Aminoacids			
Glycine			2.000
Carbon source			
Sucrose	20000.000	20000.000	20000.000

face-sterilized seeds to the germination medium, 200 µl dH<sub>2</sub>O was added into the centrifuge tube after rinse water was carefully removed. The seeds were pipetted several times to ensure all the remaining seeds were dispersed in water. Finally, the seeds were

**Table 3.** Stages of orchid seed germination according to Yamazaki and Miyoshi (2006)

Germination stage	Indicators
Stage 0	'No germination' stage. No growth of embryo occurs.
Stage 1	'Pre-germination' stage. Embryo swells to fill the seed coat.
Stage 2	'Germination' stage. Embryo emerges from the seed coat.
Stage 3	'Protocorm' stage. Embryo is completely discharged from the seed coat.
Stage 4	'Rhizoid' stage. Rhizoids are formed on the protocorm surface.
Stage 5	'Shoot' stage. Shoot is differentiated from the protocorm.

transferred in sterile water and placed onto the desired medium using a micropipette. To disperse the seeds onto the medium, the Petri dishes were gently shaken after seed sowing.

#### Culture conditions

The solid cultures were incubated at  $23 \pm 1$  °C in a plant growth chamber (Sanyo MLR-352H, Osaka, Japan) while a refrigerated and illuminated incubator shaker (New Brunswick Innova 42R, CT, USA) was used for the liquid cultures. The cultures were maintained under illumination of  $60 \mu\text{mol m}^{-2} \text{s}^{-1}$  photo-synthetic photon flux density with a 16-h photoperiod. In order to reduce evaporation from the cultures, the relative humidity of the growth chamber was adjusted to 50% during the germination experiments.

#### Seed count and statistical analysis

The seeds were observed and counted by stereo microscope under 20-fold magnification at the end of a 45-day incubation period. The Petri dishes were divided into squares of  $2 \times 2$  cm in size for easier counting. Germination stages of the seeds were evaluated according to Yamazaki and Miyoshi (15). These stages are given in Table 3. The germination percentages were calculated using the following formula for each medium (1).

(1) Germination percentage (%) =  $\Sigma$  Seed number (Stage 2–5)  $\times$  100 /  $\Sigma$  Seed number (Stage 0–5)

One milligram of seed ( $\approx$ 180–200 seeds) per Petri dish was sown on each medium for seed surface sterilization and germination experiment. Each experiment was performed in

triplicate. Surface sterilization success of the treatments was evaluated based on the observed bacterial and/or fungal contamination over 30 Petri dishes.

Seed germination and viability data were given as mean  $\pm$  standard deviation (SD). Means were compared using Duncan's Multiple Range Test (DMRT) at a significance level of  $p < 0.05$ . IBM SPSS Statistics software, version 22, was used for statistical analysis (IBM SPSS Statistics, Chicago, IL, USA).

## Results

#### Efficacy of surface sterilization method

The combination of EtOH application for 30 seconds prior to 1.0% NaOCl application for eight minutes fully inhibited contamination in the medium. Thus, this combination gave the most efficient surface sterilization result, and it was subsequently used in further experiments. However, a decrease in the concentration of NaOCl (0.5%) used in this combination resulted in the contamination of  $23.33 \pm 5.77\%$  of the cultures. When EtOH pretreatment was omitted, the contamination rate increased proportionally to the decrease in the concentration of NaOCl. In the absence of EtOH pretreatment, treatment with 1.0% NaOCl resulted in the contamination of  $46.66 \pm 5.77\%$  of the cultures while the highest contamination rate of  $66.66 \pm 11.55\%$  was observed in cultures exposed to 0.5% NaOCl.

#### Seed viability

The seeds which were kept in distilled water for one day showed a higher rate of viability on TTC test than those which

**Table 4.** Effects of different solid media on *in vitro* germination of *Serapias vomeracea* seeds

Medium	Seed germination stages (%)				
	Stage 0	Stage 1	Stage 2	Stage 3	Stage 4
KN	$15.04 \pm 2.68^a$	$26.19 \pm 5.32^d$	$58.77 \pm 5.51^a$	$0.00 \pm 0.00^d$	$0.00 \pm 0.00^b$
½KN	$13.04 \pm 2.16^{ab}$	$35.46 \pm 4.40^c$	$51.51 \pm 3.13^a$	$0.00 \pm 0.00^d$	$0.00 \pm 0.00^b$
VW	$9.75 \pm 0.98^{bc}$	$46.00 \pm 5.16^b$	$40.82 \pm 1.92^b$	$2.83 \pm 0.30^a$	$0.79 \pm 0.10^a$
½VW	$7.45 \pm 0.47^c$	$58.71 \pm 7.69^a$	$33.71 \pm 7.11^{bc}$	$0.13 \pm 0.05^d$	$0.00 \pm 0.00^b$
LN	$13.86 \pm 3.25^a$	$57.64 \pm 2.16^a$	$27.47 \pm 4.28^{cd}$	$1.03 \pm 0.16^c$	$0.00 \pm 0.00^b$
½LN	$13.01 \pm 1.59^{ab}$	$61.22 \pm 4.95^a$	$23.55 \pm 5.23^d$	$2.21 \pm 0.30^b$	$0.00 \pm 0.00^b$

Data represent mean  $\pm$  SD. Means having the same superscript letters in a column were not significantly different by Duncan's multiple range test ( $p < 0.05$ )



**Table 5.** Effects of different liquid media on *in vitro* germination of *Serapias vomeracea* seeds


Medium	Seed germination stages (%)				
	Stage 0	Stage 1	Stage 2	Stage 3	Stage 4
<b>KN</b>	11.85 ± 1.36 <sup>a</sup>	34.35 ± 3.55 <sup>c</sup>	44.61 ± 1.50 <sup>b</sup>	8.46 ± 0.88 <sup>a</sup>	0.72 ± 0.14 <sup>a</sup>
<b>½KN</b>	9.43 ± 1.80 <sup>abc</sup>	41.38 ± 3.14 <sup>b</sup>	46.75 ± 1.31 <sup>b</sup>	2.44 ± 0.92 <sup>b</sup>	0.00 ± 0.00 <sup>b</sup>
<b>VW</b>	6.29 ± 1.37 <sup>c</sup>	41.91 ± 0.10 <sup>b</sup>	51.62 ± 1.12 <sup>a</sup>	0.17 ± 0.03 <sup>c</sup>	0.00 ± 0.00 <sup>b</sup>
<b>½VW</b>	7.94 ± 1.52 <sup>bc</sup>	36.91 ± 3.42 <sup>bc</sup>	52.41 ± 2.34 <sup>a</sup>	2.74 ± 0.25 <sup>b</sup>	0.00 ± 0.00 <sup>b</sup>
<b>LN</b>	10.98 ± 2.07 <sup>ab</sup>	48.19 ± 1.91 <sup>a</sup>	40.24 ± 1.88 <sup>c</sup>	0.60 ± 0.06 <sup>c</sup>	0.00 ± 0.00 <sup>b</sup>
<b>½LN</b>	10.55 ± 2.26 <sup>ab</sup>	50.60 ± 2.57 <sup>a</sup>	38.66 ± 2.77 <sup>c</sup>	0.19 ± 0.00 <sup>c</sup>	0.00 ± 0.00 <sup>b</sup>

Data represent mean ± SD. Means having the same superscript letters in a column were not significantly different by Duncan's multiple range test ( $p < 0.05$ )

were not, after surface sterilization. The viability percentage of the seeds kept in dH<sub>2</sub>O before surface sterilization was found to be 54.67±4.16 while viability percentage was found to be 43.67±3.06 in response to direct surface sterilization without prior dH<sub>2</sub>O treatment. Direct application of TTC test without any surface sterilization treatment on the seeds resulted in 41.66±2.89% viability. Consequently, sterilization treatment slightly increased seed viability when calculated by the TTC test, but dH<sub>2</sub>O pretreatment made the most significant contribution to seed viability.


#### Effect of type and physical state of media on germination of the seeds

The detailed germination data are given in Table 4 and 5 for solid and liquid media, respectively. The highest number of stage 0 seeds was found from KN medium both in solid and liquid phases. In general, liquid media resulted in fewer stage 0 seeds than solid media. The lowest number of stage 0 seeds was found when liquid VW medium was used. All half-strength solid media resulted in fewer stage 0 seeds than full-strength solid media. Only half-strength VW medium gave a slightly higher number of stage 0 seeds than its full-strength variant. Solid LN and KN media gave statistically the same stage 0 seed percentages. Stage 1 seeds were observed in higher numbers from both solid and liquid LN medium while KN medium induced the lowest number of stage 1 seeds (Fig. 2A). Among the tested liquid media, full- and half-strength LN media gave statistically the same result for stage 1 seed percentages while a similar statistical result was obtained using half-strength VW, full strength LN and half-strength LN. All half-strength solid media induced a higher percentage of stage 1 seeds than full-strength solid media. Solid KN medium gave the highest proportion of stage 2 seeds than other media (Fig. 2B). However, the results from full- and half-strength KN medium both in solid and liquid phases were statistically the same. Using liquid media, more stage 2 seeds were counted when cultured in full- and half-strength VW medium. The results obtained from these media were statistically the same. The lowest number of stage 2 seeds was counted when cultured in half-strength LN medium both in liquid and solid phases. For stage 3 seed induction the most successful medium was solid VW whereas no



**Figure 2.** Germination stages of *Serapias vomeracea* seeds. A: Stage 1 seed in solid KN medium, B: Stage 2 seed at the beginning of germination in solid KN medium, C: Stage 3 seed from solid VW medium, D: General view of stage 3 seeds in liquid KN medium, E: Stunted rhizoids of the seed in liquid medium, F: Elongated rhizoids on solid VW medium.

stage 3 seeds were observed from solid KN, regardless of the medium strength. All other solid media induced stage 3 seed development (Fig. 2C). However, liquid KN medium gave the highest percentage of stage 3 seeds while the lowest percentage was found when using liquid VW medium. It was observed that liquid media triggered the formation of the protocorm



**Figure 3.** Comparison of *in vitro* germination rates of *Serapias vomeracea* seeds on solid and liquid media. Data represent mean  $\pm$  SD. Means having the same superscript letters in a column were not significantly different by Duncan's multiple range test ( $p < 0.05$ ).

(stage 3 germination) more successfully than solid media (Fig. 2D). Only culturing with solid VW and liquid KN medium resulted in stage 4 seed germination. The solid VW medium gave slightly better results than liquid KN medium with the rhizoids of stage 4 seeds on the solid medium being longer than that of liquid media-germinated seeds (Figs. 2E and 2F). No stage 5 seeds were observed in any of the media tested regardless of strength or form. Considering the germination parameters for orchid seeds, all seeds found at stage 2 or later were considered to have germinated. When the data were analyzed in this way, solid KN medium was found to be the best germination medium for *S. vomeracea* seeds ( $58.77 \pm 5.51\%$ ), whereas semi-strength solid LN medium gave the lowest seed germination percentage ( $25.76 \pm 4.93\%$ ) (Fig. 3). In addition, germinated seeds on KN medium successfully progressed towards subsequent developmental stages with prolonged incubation.

## Discussions

When using seeds as starting materials in tissue culture studies the most important measure of success is the percentage of the seeds which germinate. Successful germination will be due to the fact that the seeds cultured on the medium receive water and swell and take macro- and microelements from the medium (16). Therefore, it is necessary to eliminate the waxy compounds that may prevent the water from entering through the seed coat and thus to convey a hydrophilic nature to the seeds (17). It has been shown that ethyl alcohol provides a hydrophilic structure to the coats of orchid seeds by eliminating pectin, suberin and other waxy compounds present on the coat, thereby ensuring effective contact with NaOCl during surface sterilization (18). In addition, the same researchers reported that pretreatment with EtOH significantly increases germination percentages. Although the composition of hydrophobic compounds on the seed coat varies from species to species,

the improved results with EtOH showed that the hydrophobic compounds on the seed coat of *S. vomeracea* might be too dense to be eliminated by NaOCl sterilization alone. Ponert *et al.* (18) also recommended placing the seeds in a syringe during surface sterilization treatment in order to prevent seed loss during sowing. In addition, it should be noted that since the microbial load of orchid seeds may differ, the durations of the surface sterilization applications may vary for each study. The seed quantity is also another important factor affecting the success and the length of the surface sterilization process. For example, Ercole *et al.* (19) used 25 mg of *S. vomeracea* seeds per centrifuge tube but subjected them to 1 ml of 1% NaOCl solution for 20 min. In that study, no seed viability test was employed, but the authors reported higher germination rates of up to 95% in symbiotic cultures on solid oat medium. However, the authors did not assess the developmental stages of the seeds. Furthermore, considering that 1 mg of *S. vomeracea* seeds would be between 180-200 individual seeds, counting 25 mg of seeds under the microscope at the beginning, middle and the end of the incubation period would be much more time-consuming than our method, and may result in counting inaccuracy due to human error.

In this study, the effectiveness of the TTC test was increased by keeping the seeds in  $\text{dH}_2\text{O}$  for 24 hours. The reason for this is likely to be due to the onset and/or acceleration of metabolic activities in the seeds receiving water. An additional effect may be because of the increased permeability of the seed coat due to the oxidation of lignin (20). In a study on *Orchis coriophora*, Bektas *et al.* (21) incubated the seeds in 5% sucrose with several drops of commercial bleach solution for 12 hours before the TTC test. Seed viability was found to be  $54 \pm 2.1\%$ . However, in a similar study on *S. vomeracea*, the result of seed viability was not indicated (22). Plant seeds need a carbon and energy source until they germinate and become autotrophic when cultured *in*

*vitro* (23). This is more evident in non-endosperm-containing orchid seeds. Therefore, sucrose application may increase seed viability results on the TTC test by also accelerating metabolic activities.

The seeds of epiphytic and terrestrial orchids have been reported to germinate faster and contain more water when they are grown symbiotically. This is because mycorrhizal fungi increase the seed germination rate by facilitating water, carbon and nutrients intake (24). Thus, the length of germination time in asymbiotic cultures may be related to the slower water, carbon and nutrients intake (25). In this study, although solid KN medium was found to be the best overall inducer of germination, higher germination rates were generally observed from liquid media. This finding suggests that liquid media facilitates easier intake of water and other required nutrients compared with solid media. In this context, epiphytic *Doritaenopsis* orchid cultures have been reported to germinate faster in liquid medium than solid medium, but the overall germination rate remained the same at the end of the incubation period (26). A general observation from our study was that at the end of extended incubation periods the rhizoids of the protocorms were much shorter in liquid media, but in the solid media, the rhizoids developed normally. This may be related to the pressure and/or surface tension of the liquid culture on the protocorms and/or to easier access of the protocorms to nutrients. Rhizoids are primitive root precursors and conduits for nutrients to the protocorms. Rhizoids also play a role as carbohydrate depositories for subsequent stages of growth (27). Since the seeds float in liquid media and are not present on a single solid surface as with solid media, there are disadvantageous in using liquid media during the seed counting step. Also, it is necessary to agitate the cultures during incubation to prevent elements and the seeds precipitating in liquid media. For all these reasons, it is recommended to use solid media for culturing *S. vomeracea*.

Compared to LN medium used in the experiments, KN and VW media are the simplest ones in terms of microelements, vitamins, and amino acids. However, both media gave the highest seed germination results among the liquid cultures. Macroelements cover the largest portion in culture medium composition. Therefore, this difference in germination success can be attributed to the differences between the macroelement compositions of the media. LN medium contains 46% less of  $\text{KH}_2\text{PO}_4$  than KN and VW medium. VW medium contains  $\text{Ca}_3(\text{PO}_4)_2$  while KN medium contains  $\text{NH}_4\text{H}_2\text{PO}_4$  which act as additional phosphate sources. Therefore, the success of germination of these liquid culture media may be due to increased availability of phosphate to the seeds. Rich phosphate content in Mishra medium was found to be responsible for increased seed germination rates in *Cymbidium eburneum* (28). In nature, enhanced plant development due to increased phosphate uptake as a result of mycorrhizal symbiosis was reported for *Goodyera repens* (29). Rodrigues *et al.* (30) stated that the initial phosphorus (P) demand in young orchid plants may be greater than that of adult plants. Therefore, phosphate-rich media should be considered in asymbiotic germination of *S. vomer-*

*acea* seeds. However, KN medium which had the highest seed germination rate, is also distinguished from the other media by the higher total nitrogen ratio, mainly coming from  $\text{NH}_4\text{NO}_3$  and  $\text{Ca}(\text{NO}_3)_2$ . Thus, the success of germination in solid media may also be related to the presence of nitrate, which is present in much greater amounts than phosphate. Furthermore, in addition to organic carbon, nitrogen (N) is probably a major nutrient transferred to the plant because orchid tissues are highly nitrogen-enriched (31). Similar to the findings obtained in this study, Gümüş and Ellialtıoğlu (11) reported that the highest germination rate of *S. vomeracea* seeds (76.85%) was found when using KN medium, whereas seed germination decreased down to 47.51% after a reduction of inorganic nitrogen sources from the medium. They supplemented the KC medium with 20 g L<sup>-1</sup> sucrose but collected the germination data after 90 d of incubation in the dark. Bektaş and Sökmen (22) stated that the germination rate of *S. vomeracea* seeds in KN medium without sucrose was only 27.95±2.4% after 60 days of incubation. Furthermore, at the same incubation period, the authors reached to 73.74±1.6% germination ratio after testing Orchimax medium which is enriched by trypton to provide an additional source of reduced organic nitrogen, vitamins and nutritional agents. The medium also includes activated charcoal and MES 2-(N-Morpholino)ethanesulfonic acid buffer. However, the authors showed that absence of activated charcoal in Orchimax medium reduced the germination ratio down to 43.11±1.5%. Therefore, the main effective component is activated charcoal in this complex medium. In our study, we showed that there is no need to use such a complex medium, but simple Knudson C medium can offer an easy-to-prepare and cost-effective alternative to other complex media. In the present study, 58.77±5.51% of the seeds germinated on sucrose-enriched KN medium after 45 days of incubation. It is evident that the germination percentage of the seeds is closely related to the extent of the incubation period. If the incubation period is extended, more germinated seeds will be counted. KN medium showed high germination rates also for *Calopogon tuberosus* var. *tuberosus* (32), *Dendrobium aphyllum*, *Rhyncostlis retusa*, *Cymbidium longifolium* (33) and *C. aloifolium* cultures (34). Our results also showed that glycine amino acid and vitamins such as nicotinic acid, pyridoxine, thiamin, and *myo*-inositol which are completely absent from KN media and largely absent from VW media are not required for the germination of *S. vomeracea* seeds. However, at subsequent developmental stages, the nutritional needs of the plant would differ.

## Conclusion

The additional low-g centrifugation treatment in our seed surface sterilization method successfully separated empty seed coats (with no embryo) from the seeds with embryos using their mass difference. This method is ideal for eliminate counting errors, especially for stage 0 and 1 seeds, since it removes the non-viable seed coats before sowing. Here, a rapid pre-treatment with EtOH is also advised to enhance the effectiveness of the TTC test to achieve more reliable results. Our study showed



that the use of solid KN medium appears to be the optimum choice for *S. vomeracea* seed culture germination. The medium is known for its simplicity and ease of preparation while offering various modification possibilities that can enhance germination success. Based on the KN medium, the addition of several growth regulators such as auxin or cytokinin derivatives, or promoters such as oligomeric or polymeric chitosans, natural carbon sources such as banana homogenate or coconut water may enhance the effectiveness of KN medium in germinating *S. vomeracea* seeds, but further study would be required to confirm this. The outcome of this study suggests that our improved surface sterilization method and use of a solid KN medium may be a simple, cost-effective and reliable alternative to commonly used sterilization methods and use of other more complex media for *S. vomeracea* cultures.

### Acknowledgments

The authors would like to thank Mr. Jeremy Jones, of the Kocaeli University Academic Writing Department, for his help with the English used in this paper.

### Conflict of Interest

The authors declare that they have no conflicts of interest.

### Ethical Compliance

This article does not contain any studies involving human participants or animals performed by any of the authors.

### References

- Bellusci F, Pellegrino G, Musacchio A. Different levels of inbreeding depression between outcrossing and selfing *Serapias* species (Orchidaceae). *Biol Plant* 2009; 53: 175-178.
- Acemi A, Çobanoğlu Ö, Türker-Kaya S. FTIR-based comparative analysis of glucomannan contents in some tuberous orchids, and effects of pre-processing on glucomannan measurement. *J Sci Food Agric* 2019; 99: 3681-3686.
- Arvill A, Bodin L. Effect of short-term ingestion of konjac glucomannan on serum cholesterol in healthy men. *Am J Clin Nutr* 1995; 61: 585-589.
- Sood N, Baker LW, Coleman CI. Effect of glucomannan on plasma lipid and glucose concentrations, body weight, and blood pressure: Systematic review and meta-analysis. *Am J Clin Nutr* 2008; 88: 1167-1175.
- Arditti J. *Micropropagation of orchids 2<sup>nd</sup> ed.*, Blackwell Publishing, UK, 2008.
- Rasmussen HN, Dixon KW, Jersáková J, Těšitelová T. Germination and seedling establishment in orchids: a complex of requirements. *Ann Bot* 2015; 116: 391-402.
- Paul S, Kumaria S, Tandon P. An effective nutrient medium for asymbiotic seed germination and large-scale *in vitro* regeneration of *Dendrobium hookerianum*, a threatened orchid of northeast India, *AoB Plants* 2012; plr032.
- Yeung EC, Park J, Harry IS. Orchid seed germination and micropropagation I: Background information and related protocols; pp. 101-125 in Lee YI; Yeung ET (Eds) *Orchid Propagation: From laboratories to greenhouses—Methods and protocols*. New York, Springer Protocols Handbooks, Humana Press, 2018.
- Katsalirou E, Gerakis A, Haldas X, Deconinck G. Optimal disinfection times for seeds of Mediterranean orchids propagated on nutrient media. *Eur J Environ Sci* 2017; 7: 119-124.
- Özkoç I, Dalcı M. Germination of the seeds of *Orchis laxiflora* Lam. (Orchidaceae) through asymbiotic culture techniques. *Turk J Bot* 1994; 18: 461-464.
- Gümüş C, Ellialtıoğlu Ş. Seed germination and development of *Serapias vomeracea* (Burm. fil.) Briq. ssp. *orientalis* Greuter in tissue culture. *Res J Biotechnol* 2012; 7: 4-8.
- Knudson L. A new nutrient solution for germination of orchid seed. *Amer Orchid Soc Bull* 1946; 15: 214-217.
- Vacin E, Went FW. Some pH changes in nutrient solutions. *Bot Gaz* 1949; 110: 605-613.
- Lindemann EG, Gunkel JE, Davidson OW. Meristem culture of *Cattleya*, *Amer Orchid Soc Bull* 1970; 39: 1002-1004.
- Yamazaki J, Miyoshi K. *In vitro* asymbiotic germination of immature seed and formation of protocorm by *Cephalanthera falcata* (Orchidaceae), *Ann Bot* 2006; 98: 1197-1206.
- Yeung EC. A perspective on orchid seed and protocorm development. *Bot Stud* 2017; 58: 33.
- Kauth PJ, Dutra D, Johnson TR, Stewart SL, Kane ME, Vendrame W. Techniques and applications of *in vitro* orchid seed germination; pp. 375-391 in Teixeira da Silva, J.A. (Ed.) *Floriculture, Ornamental and Plant Biotechnology: Advances and Topical Issues*. Isleworth, UK, Global Science Book, 2008.
- Ponert J, Vosolsobě S, Kmecová K, Lipavská H. European orchid cultivation – from seed to mature plant. *Eur J Environ Sci* 2011; 1: 95-107.
- Ercole E, Rodda M, Girlanda M, Perotto S. Establishment of a symbiotic *in vitro* system between a green meadow orchid and a *Rhizoctonia*-like fungus. *Bio-protocol* 2015; 5(10): e1482.
- Barsberg S, Rasmussen HN, Kodahl N. Composition of *Cypripedium calceolus* (Orchidaceae) seeds analyzed by attenuated total reflectance infrared spectroscopy: In search of understanding longevity in the ground. *Am J Bot* 2013; 100(10): 2066-2073.
- Bektaş E, Cüce M, Sökmen A. *In vitro* germination, protocorm formation and plantlet development of *Orchis coriophora* (Orchidaceae), a naturally growing orchid species in Turkey. *Turk J Bot* 2013; 37: 336-342.
- Bektaş E, Sökmen A. *In vitro* seed germination, plantlet growth, tuberization, and synthetic seed production of *Serapias vomeracea* (Burm.f.) Briq. *Turk J Bot* 2016; 40: 584-594.
- Zahara M, Datta A, Boonkorkaew P, Mishra A. The effects of Different media, sucrose concentrations and natural additives on plantlet growth of *Phalaenopsis* hybrid 'Pink'. *Braz Arch Biol Technol* 2017; 60: e17160149.
- De Long JR, Swarts ND, Dixon KW, Egerton-Warburton LM. Mycorrhizal preference promotes habitat invasion by a native Australian orchid: *Microtis media*. *Ann Bot* 2012; 111: 409-18.
- Yoder JA, Zettler LW, Stewart SL. Water requirements of terrestrial and epiphytic orchid seeds and seedlings, and evidence for water uptake by means of mycotrophy. *Plant Sci* 2000; 156: 145-150.
- Tsai W-T, Chu C-Y. Static liquid culture of *Doritaenopsis* seedlings. *Hortscience* 2008; 43: 206-210.
- Edwin FG. *Plant propagation by tissue culture*. Exegetics Ltd., Edington Wilts, UK, 1993.
- Gogoi K, Kumaria S, Tandon P. *Ex situ* conservation of *Cymbidium eburneum* Lindl.: a threatened and vulnerable orchid, by asymbiotic seed germination. *3 Biotech* 2012; 2: 337.
- Alexander C, Alexander IJ, Hadley G. Phosphate uptake by *Goodyera repens* in relation to mycorrhizal infection. *New Phytol* 1984; 97: 401-411.
- Rodrigues DT, Novais RF, Alvarez VH, Dias JMM, Villani EMA. Orchid growth and nutrition in response to mineral and organic fertilizers. *Rev Bras Cienc Solo* 2010; 34: 1609-1616.
- Fochi V, Chitarra V, Kohler A, Voyron S, Singan VR, Lindquist EA, Barry KW, Girlanda M, Grigoriev IV, Martin F, Balestrini R, Perotto S. Fungal and plant gene expression in the *Tulasnella calospora*



- *Serapias vomeracea* symbiosis provides clues about nitrogen pathways in orchid mycorrhizas. *New Phytol* 2017; 213: 365-379.
32. Kauth PJ, Vendrame W, Kane ME. *In vitro* seed culture and seedling development of *Calopogon tuberosus*. *Plant Cell Tissue Organ Cult* 2006; 85: 95-102.
33. Das AK, Das J, Gogoi HK, Srivastava RB. Mass propagation of orchids through *in vitro* seed culture technology. *J Cell Tissue Cult Res* 2008; 8: 1585-1588.
34. Philip RJ, Kakati JP, Sebestinraj J, Suriya K. *In vitro* seed germination of *Cymbidium aloifolium* (L.) Sw., a potential medicinal orchid from Eastern Ghats of Tamil Nadu, India. *J Plant Biotechnol* 2017; 44: 343-348.