

Succession of ground beetle (Coleoptera: Carabidae) communities after windthrow disturbance in a montane Norway spruce forest in the Hrubý Jeseník Mts. (Czech Republic)

Josef Kašák^{1*}, Jiří Foit¹, Marek Hučín²

¹Mendel University in Brno, Faculty of Forestry and Wood Technology, Zemědělská 3, CZ – 613 00 Brno, Czech Republic

²Gymnasium Třebíč, square Masarykovo náměstí 9/116, CZ – 674 01 Třebíč, Czech Republic

Abstract

Wind disturbances are a key factor that is significantly involved in the life cycle of natural boreomontane coniferous forests. As most of these forests are currently intensively managed, we have limited knowledge on succession following natural disturbance. Succession in a Norway spruce stand after a windthrow event was studied using ground beetles as model bioindication taxa in the Hrubý Jeseník Mts. (Czech Republic). The study documented that the composition of ground beetle communities was significantly associated with stand age and its microclimatic parameters (particularly the minimal temperature and average and minimal humidity). Forest species including prey specialists, hygrophilous species, as well as habitat generalists were the most abundant in the mature stand, where the forest had the highest humidity and the least profound minimal temperatures. In contrast, open-habitat species, including euryoecious species and relict species of higher elevations, reached their highest abundances in clearing shortly after the windthrow. In clearing the humidity was lower and the temperature fluctuated significantly (the lowest minimal temperatures). Ground beetles, including forest species, were the least abundant in young stands (10 and 20 years after windthrow). We conclude that old stands are of particular importance because they harbour the highest abundance and diversity of ground beetles with various ecological requirements. Natural wind disturbances are important as well since they increase diversity by enabling the occurrence of many non-forest species. Hence, a mosaic of stands of different ages with a sufficient proportion of old stands should be maintained when managing montane coniferous forests.

Key words: forest stand development; microclimate; multivariate analysis, windstorm

Editor: Tomáš Hlásny

1. Introduction

Natural disturbances, such as windthrows, are key factors in the process of regeneration in continuity of coniferous temperate and boreal forests (Peterken 1996). As trees are broken or uprooted and open tree-fall gaps arise, the stable conditions of shady forest ecosystems are disrupted. Tree-fall gaps are characterised by bare patches of mineral soil, high amounts of dead wood, and increased sunlight reaching the forest floor. Thus, windthrows are followed by important changes in microclimate, ground structural heterogeneity and vegetation. These changes significantly affect the invertebrate fauna (Bouget & Duelli 2004).

Ground beetles (Carabidae) as a model bioindication group (Rainio & Niemelä 2003) are among the most studied invertebrates, and thus the succession of their

communities within the traditional forest management cycle from clear-cutting (i.e., artificial disturbance) to mature stands is also well studied (e.g., Butterfield 1997; Koivula et al. 2002; Magura et al. 2003; Purchart et al. 2013). In general, changes in ground beetle communities after natural disturbances (e.g., windthrows) are insufficiently studied (Skłodowski & Garbalińska 2007). However, some studies dealing with this topic have been conducted in the Alps (Duelli et al. 2002; Winter et al. 2015), northern Poland (Skłodowski & Garbalińska 2007), the Tatra Mountains (Urbanovičová et al. 2010; Šustek & Vido 2013) and North America (Gandhi et al. 2008). The results of these studies can be summarised into three main outcomes: i) windthrows are followed by decreased abundances of forest species, ii) tree-fall gaps are characterised by the occurrence of species specific to open habitats, and iii) changes in temperature

*Corresponding author. Josef Kašák, e-mail: Abovic@seznam.cz, phone: +420 606 278 554

and humidity are considered to be the most significant factors driving ground beetle community succession in these studies.

Although climatic factors (above all, temperature and humidity) are considered to substantially affect the succession of ground beetle communities after windthrows (see above), the exact relationships between changes in climatic factors and beetle succession were not addressed in the aforementioned studies. In addition, a majority of these studies were carried out in mountains, where climatic factors largely limit insect survival (Mani 1968). Therefore, the present study investigates the effects of selected forest stand characteristics and climatic factors on communities of ground beetles in a montane coniferous forest after windthrow. The following hypotheses were tested: i) ground beetle species composition and abundance are affected by forest disturbance by windthrow and its following regeneration and ii) ground beetle species composition and abundance are affected by habitat temperature and humidity. Both hypotheses were tested at the community, ecological guild and species levels. Based on our obtained results, we suggest implications for forestry management.

2. Methods

2.1. Study site

The study area is located in northern Moravia (the Czech Republic) in the Jeseníky Protected Landscape Area, and is 2 km east of Praděd Mt. (1492 m a. s. l.), near the Kamzičí chata cottage, at an elevation of 1200 – 1220 m a. s. l. (50.0820747N, 17.2619078E). The locality is situated in a seminatural montane spruce forest. The climate is cold, with a mean temperature of 1.9 °C, the long-term annual average rainfall is 1100 – 1200 mm, and snow cover lasts approximately 180 days, typically from November to early May (Quitt 1971).

We sampled four stands of different ages with areas ranging 2 – 5 ha, each stand representing a different stage of succession of spruce forest after windthrow of a mature stand: i) clearing 3 years after the event (hereafter “3”), ii) stand approximately 10 years after the event, young growth (“10”), iii) stand approximately 20 years after the event, young stand (“20”), iv) stand approximately 90 years old, mature stand (“90”). Following windthrow, the surviving trees (a few trees per ha) were kept in situ, but a majority of the dead wood caused by the windthrows or bark beetles was removed from the stands. As a result, a low amount of coarse dead wood was present in the studied stands. The studied stands were established by natural regeneration following the windstorm events, which was supported by artificial regeneration. The study stands were dominated by Norway spruce (*Picea abies* [L.] H. Karst.) (90%) with dispersed individuals of rowan (*Sorbus* sp.), European beech (*Fagus sylvatica* L.) and birch (*Betula pendula* Roth.).

2.2. Field sampling and variable measurement

Ground beetles were sampled using pitfall traps (8 cm in diameter, 15 cm deep, one-third filled with a 4% formaldehyde solution), a standard method used to sample epigeic invertebrates (Spence & Niemelä 1994). In each study stand, 8 traps were installed in a line at a spacing of 10 m. The trap lines were located in the centres of the study stands, and no trap was closer than 10 m from the stand edge. In total, 32 traps were used. The pitfall traps were exposed from 30. 5. to 16. 10. 2011 and were emptied at approximately three-week interval (21. 6., 9. 7., 27. 7., 14. 8., 1. 9., 19. 9. and 16. 10. 2011). Each sample was preserved in 95% ethanol. All ground beetles were identified to the species level using keys (Hůrka 1996). The used nomenclature follows de Jong (2012).

Additionally, one datalogger (Votcraft DL-120TH) was installed in the middle of each study stand and was active during the whole period of beetle trapping. Dataloggers were mounted 10 cm above the ground in a shaded location. The dataloggers recorded the following climatic characteristics: minimal daily temperature (hereafter “Tmin”), average daily temperature (“Tave”), maximal daily temperature (“Tmax”), minimal daily humidity (“Hmin”) and average daily humidity (“Have”).

2.3. Data analysis

Prior to data analysis, the number of individuals of each species per sample was converted to the number of individuals per sampling day (relative abundance) to remove possible bias caused by the slightly uneven lengths of the trap exposure intervals. For each interval of trap exposure, the mean values of the explanatory climatic variables (Tmin, Tave, Tmax, Hmin, Have) were calculated and coded as continuous variables.

To relate gradients in species composition and abundance to the studied predictors and to test the importance of these predictors, a canonical correspondence analysis (CCA) was used (ter Braak & Šmilauer 2002), which is appropriate for data with a larger gradient length (the gradient length was previously examined by detrended correspondence analysis (DCA)). The data were standardised and centred by sample, and a Monte Carlo permutation test was used (2000 permutations). All the explanatory variables (i.e., climatic variables and stand age) were treated as continuous variables. The four sampled stands (i.e., 3, 10, 20, 90) were treated as supplementary variables in the analysis. A forward selection procedure was used to test environmental variables. Since ground beetle communities are strongly affected by phenology (Rainio & Niemelä 2003), we treated the date of sample collection (coded as an ordinal week of the year) as a covariate.

Subsequently, we used generalised linear models (GLMs) with a Poisson distribution and log link function to separately assess the impact of selected single fac-

3.2. Responses of specific groups of ground beetles to stand age

The results of the K–W tests revealed that the abundance of ground beetle guilds differed significantly among stands of different ages (Fig. 2). The biggest differences were found in the occurrence of the guild of forest species, which reached its highest abundance by far in the mature 90-year-old stand. Open-habitat ground beetles were the most abundant in the youngest 3-year-old stand, less abundant in the mature 90-year-old stand and almost absent in the 10- and 20-year-old stands. Habitat generalists were the most abundant in the 90-year-old stand.




Fig. 2. Differences in abundance of three ecological guilds of ground beetles among stands in different stages of forest succession following windthrow in montane Norway spruce stand of Hrubý Jeseník Mts. On the x axis is age of stand in years. The box and whisker plots are composed of outliers (asterisks), non-outlier ranges (whiskers), lower and upper quartiles (boxes) and medians (middle lines). Results of Kruskal–Wallis tests are shown as well.

3.3. Responses of ground beetle species to stand characteristics

Using GLM analyses, the occurrences of 19 species of ground beetles were found to be significantly affected by at least one of the tested habitat factors (Table 2). Specifically, 18 species responded significantly to stand age (12 increased in abundance, 6 decreased), 12 were affected by the minimal temperature (all increased), and 12 varied in abundance according to the average humidity (9 increased, 3 decreased).

Thus, typical forest ground beetles increased in abundance in stands of higher age, which was particularly remarkable in the case of species of the genus *Carabus* and food specialists that prey on molluscs, such as *Cychrus caraboides* and *C. attenuates* (Table 2). In contrast, the abundance of open-habitat species (*Amara* spp. and *Carabus sylvestris*) and some euryoecious species (e.g., *Leistus ferrugineus* and *Notiophilus palustris*) that are associated with disturbed open habitats, such

Table 2. Results of generalised linear model analyses for beetles and their responses to different features of spruce stands.

Species	N	Age			Tmin			Have		
		Intercept	Slope	AIC	Intercept	Slope	AIC	Intercept	Slope	AIC
<i>Amara lumicollis</i>	6	-14	-0.06	1.45	1.12	0.14	2.43	-13.29	0.61	2.05
<i>Calathus micropterus</i>	90	1.8	1.05	11.55	2.96	0.86	13.96	9.96	0.82	14.51
<i>Carabus auronitens</i>	123	1.3	1.69	15.91	4.09	1.01	16.87	10.72	1	18.16
<i>Carabus glabratus</i>	6	12.44	0.09	1.63	2.59	0.22	2.51	22.94	0.34	2.36
<i>Carabus linnaei</i>	792	1.94	5.18	76.12	5.17	2.74	96.42	10.02	2.45	109.24
<i>Carabus sylvestris</i>	312	-0.34	-0.87	35.46	1.46	1.26	33.98	-3.43	-0.94	35.41
<i>Carabus violaceus</i>	164	0.52	0.95	18.62	2.67	1.11	17.11	2.43	0.38	19.47
<i>Cychrus attenuatus</i>	7	12.66	0.09	1.75	19.77	0.44	2.26	7.44	0.19	2.81
<i>Cychrus caraboides</i>	15	1.41	0.63	3.78	5.96	0.39	3.8	15.63	0.43	3.96
<i>Leistus ferrugineus</i>	13	-2.17	-0.74	3.71	-0.91	-0.35	4.43	-14.3	-1	3.55
<i>Leistus piceus</i>	31	-0.98	-0.74	7.41	0.14	0.06	8.05	-5.65	-0.54	7.79
<i>Notiophilus biguttatus</i>	10	-13.48	-0.1	1.66	-0.17	-0.05	3.27	-12.68	-0.77	2.67
<i>Notiophilus palustris</i>	10	-1.68	-0.57	2.96	0.27	0.06	3.44	-8.72	-0.5	3.24
<i>Patrobis atrovirens</i>	17	3.52	0.75	3.07	6.57	0.41	4.84	16.87	0.49	5.07
<i>Pterostichus aethiops</i>	5	-0.48	-0.15	1.77	1.16	0.13	1.77	-6.12	-0.22	1.75
<i>Pterostichus diligens</i>	13	0.04	0.02	4.68	1.14	0.40 ^{ns}	4.56	6.8	0.23	4.59
<i>Pterostichus oblongopunctatus</i>	80	0.18 ^{ns}	0.15	13.57	0.64	1.81	13.43	5.27	0.49	13.29
<i>Pterostichus rufitarsis</i>	17	0.03 ^{ns}	0.05	4.72	0.26	0.07	4.71	-4.19	-0.27	4.64
<i>Pterostichus unctulatus</i>	615	0.73	2.39	60.34	2.29	2.04	59.41	4.97	1.28	64.86
<i>Trechus splendens</i>	91	1.53	1.58	15.11	2.79	0.85	17.28	14.6	1.02	17.13
<i>Trechus striatulus</i>	236	2.23	2.89	28.32	2.64	1.34	40.04	18.06	1.84	37.77
<i>Trichotichnus laevicollis</i>	37	0.35	0.3	8.38	2.29	0.52	7.95	-2.77	-0.26	8.35

^{ns} not significant; * p < 0.05; ** p < 0.01; *** p < 0.001; p-values in bold are significant at α = 0.05; positive responses of species to the tested environmental variables are highlighted by a grey background.

as windthrow gaps, decreased in stands of higher age. Increasing humidity was associated with higher abundances of hygrophilous species of the genus *Trechus* and some forest species of the genus *Carabus*.

4. Discussion

Our results confirm that windstorm events in Norway spruce mountain forest and subsequent forest regeneration and succession are associated with dramatic changes in ground beetle communities. Among the tested factors, stand age (i.e., time elapsed after windthrow event), minimal temperature and average humidity affected ground beetles the most significantly (Table 1). In general, the abundance of forest species increased with an increase in the age of the stand. However, this increase is likely not linear but instead commences later in succession because the abundances of forest species were even slightly lower in 10- and 20-year-old stands (Fig. 2). Thus, forest ground beetles, as well as ground beetles in total, reached maximal abundances in the mature 90-year-old stand, which was characterised by the highest values of minimal temperature and average humidity. In contrast, open-habitat species and euryoecious species occurred most numerous in the clearing 3 years after windthrow. These results are in accordance with previous studies on carabid succession after wind disturbances (Šustek & Vido 2013; Winter et al. 2015) or during the standard forestry cycle (Magura et al. 2003).

A decrease in the abundance of forest ground beetles following windthrow or felling is explained mainly by dramatic changes in abiotic conditions (Koivula et al. 2002; Sklodowski & Garbalinska 2007). It is believed that forest ground beetles require a relatively stable microclimate (Thiele 1977). Such conditions are better maintained in mature (or undisturbed) forest stands, which protect the ground against direct sun and wind exposure due to the higher canopy closure (Sklodowski 2017). In contrast, the temperature fluctuates strongly in windthrow gaps or clear-cut areas (Chen et al. 1999), where high temperatures might be reached during a sunny day but temperatures decrease substantially at night or during a windy and cloudy day (Warren & Key 1991). In accordance, the highest minimal temperatures, reflecting the most stable microclimate, were maintained in the mature stands in the present study, where the highest abundance of forest ground beetles was also found (Fig. 1 and 2). Furthermore, our study documented the minimal temperature to be a more important climatic factor than average temperature (Table 1). Correspondingly, the survival and reproduction of animals are mostly affected by extreme values of habitat characteristics rather than by average values (Begon et al. 2006).

The aforementioned increased sun and wind exposure of the ground after windthrow or felling lead to lowered humidity (regardless of a higher total amount

of rainfall) (Ishizuka et al. 2002). Subsequently, as the canopy densifies during succession in a new forest stand, the humidity rises again, as was documented in the present study (Fig. 1), leading to increased abundances of hygrophilous forest ground beetles, such as *Trechus* spp., which were more numerous in the older stands (90 years old) in the present study (Fig. 1, Table 2). Similarly, in the present study, species of the genus *Cychrus* that feed upon molluscs (Hůrka 1996) also experienced an increase in abundance in older stands. The molluscs were previously reported to diminish after windthrow events due to lower humidity (Duelli et al. 2002). In the present study, the mature forest stand (90 years old), with the most stable climatic conditions (including a smaller range of extreme values), hosted the highest numbers of forest ground beetles as well as all ground beetles in total. Ground beetles may also prefer old stands because woody debris and litter provide shelter and preserve microhabitat conditions (Sklodowski 2016). Similar results were reported by Urbanovičová et al. (2010) in the Tatra Mts., showing the highest abundance of ground beetles in undisturbed forest stands.

Despite the substantial decrease in the abundance of forest ground beetles after windthrow, the abundance of forest ground beetles remained higher in the windthrow clearing (3 years after the event) than in slightly older (10- and 20-year-old) forest stands. This phenomenon has also been documented in other similar studies (Magura et al. 2003; Purchart et al. 2013; Sklodowski 2017), and it can be explained by the following: a) some ground beetle species (members of the genus *Carabus*) can survive several years in suboptimal habitats (Koivula et al. 2002) because their development takes several years (Hůrka 1996), and adults can live for a few years as well; b) a windthrow area might temporarily offer high numbers of prey because an open habitat with a substantial amount of freshly dead wood and an array of herbs is characterised by a high abundance of saproxylic and herbivorous invertebrates (Winter et al. 2015; Sklodowski 2017) c) temporarily increased amount of dead wood provides shelter for carabids (Sklodowski 2016; Sklodowski 2017).

Overall, the lowest abundances of forest ground beetles were recorded in 10- and 20-year-old stands in the present study (Fig. 2). Similarly, a decrease in abundance of forest ground beetles during the first half of the forest life cycle has been observed in several studies (Butterfield 1997; Magura et al. 2003; Purchart et al. 2013). This might be attributed to the fact that forest stands in this stage are usually very shady with a negligible amount of dead wood and a poorly developed herbaceous understorey (Purchart et al. 2013), so the supply of prey and shelters in such habitats is strongly limited (Taboada et al. 2008).

As expected, open-habitat ground beetle species reached their highest abundance in the clearing 3 years after windthrow, which corresponds to the results of other studies (Sklodowski & Garbalinska 2007; Gandhi

et al. 2008; Šustek & Vido 2013; Sklodowski 2017). The second highest abundance of open-habitat species was surprisingly found in the mature 90-year-old stand. This corresponds to the fact that mature montane spruce forests usually exhibit a partly open structure because their canopies are already disturbed by dieback or falling/felling of individual trees (Winter et al. 2015).

The clearing (3 years after windthrow) hosted the highest abundance of open-habitat species (*Amara* spp. and *Harpalus* spp.) and euryoecious ground beetles typical of disturbed habitats (*Bembidion lampros*, *B. quadrimaculatum*, *Leistus ferrugineus* and *Notiophilus palustris*). Several important bioindicator species, such as the boreomontane species *Amara erratica* and *Carabus sylvestris*, which are typically distributed in alpine meadows (Stanovský & Pulpán 2006), occurred in the clearing as well. Clearings after windthrows represent temporarily open habitats that can enable the reproduction of these species. Additionally, these species could easily colonise the clearing at our study site because of its proximity to alpine meadows in the Hrubý Jeseník Mts., where they are quite numerous (cf. Kašák et al. 2015).

5. Conclusion

Based on the results of the present, as well as several other studies (Magura et al. 2003; Niemelä et al. 2007), we can assume that the presence of old forest stands at a particular location is a fundamental prerequisite for the long-term survival of populations of specialised forest ground beetles. At the same time, newly regenerated forest stands after a windthrow event or felling do not become optimal for forest ground beetles until they are 60–90 years old. Therefore, montane forests should be managed to ensure that a sufficient area of mature forest stands is maintained near any clearing until the clearing becomes a middle-age-class stand. On the other hand, disturbed areas, such as windthrow clearings, support the occurrence of many non-forest species (e.g., open-habitat, euryoecious species) and thereby increase the overall diversity of ground beetles in these ecosystems. Hence, a fine mosaic of forest stands of various ages can enable the contemporary survival of many different ground beetle ecological groups and species.

Acknowledgements

We thank Tomáš Kuras (Palacký University in Olomouc) for technical support and advice regarding the study design. We would like to thank Jiří Stanovský (Ostrava) for the determination and revision of some problematic ground beetle individuals. Employees of PLAČKO Jeseníky are acknowledged for allowing us to carry out this study at the selected location. Last but not least we thank two anonymous reviewers for their valuable comments and suggestions that improved the manuscript.

References

- Begon, M., Townsed, C. R., Harper, J. L., 2006: Ecology. From individuals to ecosystems. Blackwell Publishing, Oxford, 738 p.
- Bouget, Ch., Duelli P., 2004: The effects of windthrow on forest insect communities: a literature review. *Biological Conservation*, 118:281–299.
- Butterfield, J., 1997: Carabid community succession during the forestry cycle in conifer plantations. *Ecography*, 20:614–625.
- de Jong, Y. S. D. M. (ed.), 2012: Fauna Europaea version 2.6.
- Duelli, P., Obrist, M., Wermelinger, B., 2002: Windthrow-induced changes in faunistic biodiversity in alpine spruce forests. *Forest Snow and Landscape Research*, 77:117–131.
- Gandhi, K. J. K., Gilmore, D. W., Katovich, S. A., Matson, W. J., Zasada, J. C., Seybold, S., 2008: Catastrophic windstorm and fuel-reduction treatments alter ground beetle (Coleoptera: Carabidae) assemblages in a North American sub-boreal forest. *Forest Ecology and Management*, 256:1104–1123.
- Hollander, M., Wolfe, D. A., 1999: Nonparametric statistical methods, 2nd edition. Wiley, New York, 816 p.
- Hůrka, 1996: Carabidae of the Czech and Slovak Republics—illustrated key. Kabourek, Zlín, 565 p.
- Hůrka, K., Veselý, P., Farkač, J., 1996: Využití střevlíkovitých (Coleoptera: Carabidae) k indikaci kvality prostředí. *Klapalekiana*, 32:15–26.
- Chen, J. Q., Saunders, S. C., Crow, T. R., Naiman, R. J., Brosnoff, K. D., Mroz, G. D. et al., 1999: Microclimate in forest ecosystem and landscape ecology – Variation in local climate can be used to monitor and compare the effects of different management regimes. *Bioscience*, 49:288–297.
- Ishizuka, M., Ochiai, Y., Utsugi, H., 2002: Microenvironments and growth in gaps. In: Nakashizuka, M. (ed.): Diversity and interaction in a temperate forest community: Ogawa forest reserve of Japan. Springer, Tokyo, p. 229–244.
- Kašák, J., Mazalová, M., Šipoš, J., Kuras, T., 2015: Dwarf pine – invasive plant threatens biodiversity of alpine beetles. *Biodiversity and Conservation*, 10:2399–2415.
- Koivula, M., Kukkonen, Niemelä, J., 2002: Boreal carabid-beetle (Coleoptera, Carabidae) assemblages along the clear-cut originated succession gradient. *Biodiversity and Conservation*, 11:1269–1288.
- Magura, T., Tóthmérész, B., Elek, Z., 2003: Diversity and composition of carabids during a forestry cycle. *Biodiversity and Conservation*, 12:73–85.
- Mani, M. S., 1968: Ecology and biogeography of high altitude insects. *Series Entomologica*, 4:1–528.
- Niemelä, J., Koivula, M., Kotze, D. J., 2007: The effects of forestry on carabid beetles (Coleoptera: Carabidae) in boreal forests. *Journal of Insect Conservation*, 11:5–18.

- Peterken, G. F., 1996: Natural woodland. Ecology and conservation in northern temperate regions. Cambridge University Press, Cambridge, 540 p.
- Purchart, L., Tuf, I. H., Hula, V., Suchomel, J., 2013: Arthropod assemblages in Norway spruce monocultures during a forest cycle – A multi-taxa approach. *Forest Ecology and Management*, 306:42–51.
- Quitt, E., 1971: Klimatické oblasti Československa. Geografický ústav ČSAV, Brno, 73 p.
- Rainio, J., Niemelä, J., 2003: Ground beetles (Coleoptera: Carabidae) as bioindicators. *Biodiversity and Conservation*, 12:487–506.
- Sklodowski, J., Garbalińska, P., 2007: Ground beetle assemblages (Coleoptera, Carabidae) in the third year of regeneration after hurricane in the Puszcza Piska pine forests. *Baltic Journal of Coleopterologist*, 7:17–35.
- Sklodowski, J., 2016: Manual soil preparation and piles of branches can support ground beetles (Coleoptera, carabidae) better than four different mechanical soil treatments in clear-cut area of a closed-canopy pine forest in northern Poland. *Scandinavian Journal of Forest Research*, 32:123–133.
- Sklodowski, J., 2017: Three phases of changes in carabid assemblages during secondary succession in a pine forest disturbed by windthrow – results from first 10 years of observations. *Insect Conservation and Diversity*, doi: 10.1111/icad.12237
- Spence, J. R., Niemelä, J. K., 1994: Sampling ground beetle assemblages with pitfall traps: the madness and the method. *The Canadian Entomologist*, 126:881–894.
- StatSoft Inc. 2013: Electronic statistics textbook. OK, Tulsa. [www document]. URL: <http://www.statsoft.com/textbook/>. (Site visited on 25 October 2016).
- Stanovský, J., Pulpán, J., 2006: Střevlíkovití brouci Slezska (severovýchodní Moravy). Muzeum Beskyd, Frýdek-Místek, 159 p.
- Šustek, Z., Vido, J., 2013: Vegetation state and extreme drought as factors determining differentiation and succession of Carabidae communities in forests damaged by a windstorm in the High Tatra Mts. *Biologia*, 68:1198–1210.
- Taboada, A., Kotze, D. J., Tarrega, J., Salgado, J. M., 2008: Carabids of differently aged reforested pine-woods and a natural pine forest in a historically modified landscape. *Basic and Applied Ecology*, 9:161–171.
- ter Braak, C. J. F., Šmilauer, P., 2002: CANOCO reference manual and CanoDraw for Windows user's guide: software for canonical community ordination (version 4.5). Microcomputer Power, Ithaca, NY, USA, 500 p.
- Thiele, H. U., 1977: Carabid beetles in their environments. Springer, Berlin, 366 p.
- Urbanovičová, V., Kováč, L., Miklisová, D., 2010: Epigeic arthropod communities of spruce forest stands in the High Tatra Mts. (Slovakia) with special reference to Collembola – first year after windthrow. *Acta Societatis Zoologicae Bohemicae*, 74:141–152.
- Warren, M. S., Key, R. S., 1991: Woodlands: past, present and potential for insects. In: Collins, N. M., Thomas, J. A. (eds.): *The conservation of insects and their habitats*. Academic Press, London, p. 144–212.
- Winter, M-B., Ammaer, Ch., Baier, R., Donato, D. C., Seibold, S., Müller, J., 2015: Multi-taxon alpha diversity following bark beetle disturbance: Evaluation multi-decade persistence of a diverse early-seral phase. *Forest Ecology and Management*, 338:32–45.

Appendix 1. Abundances of the ground beetles sampled in the Hrubý Jeseník Mts. and their ecological characteristics.

Species	H ^a	R ^b	Age of stand				Total
			3	10	20	90	
<i>Amara erratica</i> (Duftschmid, 1812)	O	R	2	0	0	0	2
<i>Amara lunicollis</i> Schiöde, 1837	O	A	6	0	0	0	6
<i>Bembidion lampros</i> (Herbst, 1784)	O	E	1	0	0	0	1
<i>Bembidion quadrimaculatum</i> (Linnaeus, 1761)	O	E	1	0	0	0	1
<i>Calathus micropterus</i> (Duftschmid, 1812)	F	A	10	4	1	75	90
<i>Carabus arvensis</i> (Herbst, 1784)	G	A	2	0	0	1	3
<i>Carabus auronitens</i> Fabricius, 1792	F	A	10	14	20	79	123
<i>Carabus glabratus</i> Paykull, 1790	F	A	0	0	0	6	6
<i>Carabus intricatus</i> Linnaeus, 1761	F	A	1	0	0	0	1
<i>Carabus linnaei</i> Panzer, 1810	F	A	89	5	17	681	792
<i>Carabus sylvestris</i> Panzer, 1793	O	A	157	16	19	120	312
<i>Carabus violaceus</i> Linnaeus, 1758	G	A	33	32	19	80	164
<i>Cychrus attenuatus</i> (Fabricius, 1792)	F	R	0	0	0	7	7
<i>Cychrus caraboides</i> (Linnaeus, 1758)	F	A	1	1	3	10	15
<i>Harpalus latus</i> (Linnaeus, 1758)	F	A	0	1	0	0	1
<i>Leistus ferrugineus</i> (Linnaeus, 1758)	G	E	11	0	1	1	13
<i>Leistus piceus</i> Frölich, 1799	F	A	18	6	1	6	31
<i>Notiophilus biguttatus</i> (Fabricius, 1779)	F	A	10	0	0	0	10
<i>Notiophilus palustris</i> (Duftschmid, 1812)	G	E	6	3	1	0	10
<i>Patrobus atrorufus</i> (Stroem, 1768)	G	A	0	1	0	16	17
<i>Pseudoophonus rufipes</i> (De Geer, 1774)	O	E	0	1	0	0	1
<i>Pterostichus aethiops</i> (Panzer, 1797)	F	A	2	1	1	1	5
<i>Pterostichus diligens</i> (Sturm, 1824)	G	A	0	4	8	1	13
<i>Pterostichus niger</i> (Schaller, 1783)	F	E	0	1	1	0	2
<i>Pterostichus oblongopunctatus</i> (Fabricius, 1787)	F	A	8	21	39	12	80
<i>Pterostichus rufitarsis cordatus</i> Letzner, 1842	F	R	6	3	2	6	17
<i>Pterostichus unctulatus</i> (Duftschmid, 1812)	F	A	101	106	73	335	615
<i>Trechus pilisensis sudeticus</i> Pawlowski, 1975	F	A	0	0	2	0	2
<i>Trechus splendens</i> Gemminger et Harold, 1868	F	A	2	15	13	61	91
<i>Trechus striatulus</i> Putzeys, 1847	G	A	8	24	6	198	236
<i>Trichotichnus laevicollis</i> (Duftschmid, 1812)	F	A	13	2	1	21	37
Number of species			23	20	19	20	31
Number of individuals			498	261	228	1717	2704

Beetles were divided into three groups either ^a according to their habitat association: generalist (G), forest (F) and open-habitat (O) species, or ^b according to their ability to cope with environmental changes following Hürka et al. (1996) as either relicts (R), adaptive (A) or eurytopic (E).