

# The Product Space of Real Normed Spaces and its Properties

Noboru Endou  
 Gifu National College of Technology  
 Japan

Yasunari Shidama  
 Shinshu University  
 Nagano, Japan

Keiichi Miyajima  
 Ibaraki University  
 Hitachi, Japan

**Summary.** In this article, we define the product space of real linear spaces and real normed spaces. We also describe properties of these spaces.

MML identifier: PRVECT\_2, version: 7.8.05 4.84.971

The terminology and notation used here are introduced in the following articles: [20], [9], [22], [2], [1], [19], [5], [23], [7], [10], [8], [4], [13], [12], [21], [14], [3], [6], [16], [11], [15], [17], and [18].

## 1. THE PRODUCT SPACE OF REAL LINEAR SPACES

The following propositions are true:

- (1) Let  $s, t$  be sequences of real numbers and  $g$  be a real number. Suppose that for every element  $n$  of  $\mathbb{N}$  holds  $t(n) = |s(n) - g|$ . Then  $s$  is convergent and  $\lim s = g$  if and only if  $t$  is convergent and  $\lim t = 0$ .
- (2) Let  $x, y$  be finite sequences of elements of  $\mathbb{R}$ . Suppose  $\text{len } x = \text{len } y$  and for every element  $i$  of  $\mathbb{N}$  such that  $i \in \text{Seg len } x$  holds  $0 \leq x(i)$  and  $x(i) \leq y(i)$ . Then  $|x| \leq |y|$ .
- (3) Let  $F$  be a finite sequence of elements of  $\mathbb{R}$ . If for every element  $i$  of  $\mathbb{N}$  such that  $i \in \text{dom } F$  holds  $F(i) = 0$ , then  $\sum F = 0$ .

Let  $f$  be a function and let  $X$  be a set. A function is called a multi-operation of  $X$  and  $f$  if:

- (Def. 1)  $\text{dom } f = \text{dom } f$  and for every set  $i$  such that  $i \in \text{dom } f$  holds  $f(i)$  is a function from  $\{X, f(i)\}$  into  $f(i)$ .

Let  $F$  be a sequence of non empty sets and let  $X$  be a set. Observe that every multi-operation of  $X$  and  $F$  is finite sequence-like.

We now state the proposition

- (4) Let  $X$  be a set,  $F$  be a sequence of non empty sets, and  $p$  be a finite sequence. Then  $p$  is a multi-operation of  $X$  and  $F$  if and only if  $\text{len } p = \text{len } F$  and for every set  $i$  such that  $i \in \text{dom } F$  holds  $p(i)$  is a function from  $\{X, F(i)\}$  into  $F(i)$ .

Let  $F$  be a sequence of non empty sets, let  $X$  be a set, let  $p$  be a multi-operation of  $X$  and  $F$ , and let  $i$  be an element of  $\text{dom } F$ . Then  $p(i)$  is a function from  $\{X, F(i)\}$  into  $F(i)$ .

Next we state the proposition

- (5) Let  $X$  be a non empty set,  $F$  be a sequence of non empty sets, and  $f, g$  be functions from  $\{X, \prod F\}$  into  $\prod F$ . Suppose that for every element  $x$  of  $X$  and for every element  $d$  of  $\prod F$  and for every element  $i$  of  $\text{dom } F$  holds  $f(x, d)(i) = g(x, d)(i)$ . Then  $f = g$ .

Let  $F$  be a sequence of non empty sets, let  $X$  be a non empty set, and let  $p$  be a multi-operation of  $X$  and  $F$ . The functor  $\prod^\circ p$  yielding a function from  $\{X, \prod F\}$  into  $\prod F$  is defined as follows:

- (Def. 2) For every element  $x$  of  $X$  and for every element  $d$  of  $\prod F$  and for every element  $i$  of  $\text{dom } F$  holds  $(\prod^\circ p)(x, d)(i) = p(i)(x, d(i))$ .

Let  $R$  be a binary relation. We say that  $R$  is real-linear-space-yielding if and only if:

- (Def. 3) For every set  $S$  such that  $S \in \text{rng } R$  holds  $S$  is a real linear space.

Let us note that there exists a finite sequence which is non empty and real-linear-space-yielding.

A real linear space-sequence is a non empty real-linear-space-yielding finite sequence.

Let  $G$  be a real linear space-sequence and let  $j$  be an element of  $\text{dom } G$ . Then  $G(j)$  is a real linear space.

Let  $G$  be a real linear space-sequence. The functor  $\overline{G}$  yielding a sequence of non empty sets is defined by:

- (Def. 4)  $\text{len } \overline{G} = \text{len } G$  and for every element  $j$  of  $\text{dom } G$  holds  $\overline{G}(j) = \text{the carrier of } G(j)$ .

Let  $G$  be a real linear space-sequence and let  $j$  be an element of  $\text{dom } \overline{G}$ . Then  $G(j)$  is a real linear space.

Let  $G$  be a real linear space-sequence. The functor  $\langle +_{G_i} \rangle_i$  yielding a family of binary operations of  $\overline{G}$  is defined as follows:

(Def. 5)  $\text{len}(\langle +_{G_i} \rangle_i) = \text{len } \overline{G}$  and for every element  $j$  of  $\text{dom } \overline{G}$  holds  $\langle +_{G_i} \rangle_i(j) =$  the addition of  $G(j)$ .

The functor  $\langle -_{G_i} \rangle_i$  yields a family of unary operations of  $\overline{G}$  and is defined as follows:

(Def. 6)  $\text{len}(\langle -_{G_i} \rangle_i) = \text{len } \overline{G}$  and for every element  $j$  of  $\text{dom } \overline{G}$  holds  $\langle -_{G_i} \rangle_i(j) =$   $\text{comp } G(j)$ .

The functor  $\langle 0_{G_i} \rangle_i$  yielding an element of  $\prod \overline{G}$  is defined by:

(Def. 7) For every element  $j$  of  $\text{dom } \overline{G}$  holds  $\langle 0_{G_i} \rangle_i(j) =$  the zero of  $G(j)$ .

The functor  $\text{multop } G$  yields a multi-operation of  $\mathbb{R}$  and  $\overline{G}$  and is defined by:

(Def. 8)  $\text{len } \text{multop } G = \text{len } \overline{G}$  and for every element  $j$  of  $\text{dom } \overline{G}$  holds  $(\text{multop } G)(j) =$  the external multiplication of  $G(j)$ .

Let  $G$  be a real linear space-sequence. The functor  $\prod G$  yielding a strict non empty RLS structure is defined by:

(Def. 9)  $\prod G = \langle \prod \overline{G}, \langle 0_{G_i} \rangle_i, \prod^\circ(\langle +_{G_i} \rangle_i), \prod^\circ \text{multop } G \rangle$ .

Let  $G$  be a real linear space-sequence. One can check that  $\prod G$  is Abelian, add-associative, right zeroed, right complementable, and real linear space-like.

## 2. THE PRODUCT SPACE OF REAL NORMED SPACES

Let  $R$  be a binary relation. We say that  $R$  is real-norm-space-yielding if and only if:

(Def. 10) For every set  $x$  such that  $x \in \text{rng } R$  holds  $x$  is a real normed space.

One can check that there exists a finite sequence which is non empty and real-norm-space-yielding.

A real norm space-sequence is a non empty real-norm-space-yielding finite sequence.

Let  $G$  be a real norm space-sequence and let  $j$  be an element of  $\text{dom } G$ . Then  $G(j)$  is a real normed space.

Let us note that every finite sequence which is real-norm-space-yielding is also real-linear-space-yielding.

Let  $G$  be a real norm space-sequence and let  $x$  be an element of  $\prod \overline{G}$ . The functor  $\text{normsequence}(G, x)$  yields an element of  $\mathcal{R}^{\text{len } G}$  and is defined as follows:

(Def. 11)  $\text{len } \text{normsequence}(G, x) = \text{len } G$  and for every element  $j$  of  $\text{dom } G$  holds  $(\text{normsequence}(G, x))(j) = (\text{the norm of } G(j))(x(j))$ .

Let  $G$  be a real norm space-sequence. The functor  $\text{productnorm } G$  yields a function from  $\prod (\overline{G} \text{ qua real linear space-sequence})$  into  $\mathbb{R}$  and is defined by:

(Def. 12) For every element  $x$  of  $\prod \overline{G}$  holds  $(\text{productnorm } G)(x) = |\text{normsequence}(G, x)|$ .

Let  $G$  be a real norm space-sequence. The functor  $\prod G$  yielding a strict non empty normed structure is defined as follows:

(Def. 13) The RLS structure of  $\prod G = \prod(G \text{ qua real linear space-sequence})$  and the norm of  $\prod G = \text{productnorm } G$ .

In the sequel  $G$  is a real norm space-sequence.

We now state four propositions:

- (6)  $\prod G = \langle \prod \overline{G}, \langle 0_{G_i} \rangle_i, \prod^\circ(\langle +_{G_i} \rangle_i), \prod^\circ \text{multop } G, \text{productnorm } G \rangle$ .
- (7) For every vector  $x$  of  $\prod G$  and for every element  $y$  of  $\prod \overline{G}$  such that  $x = y$  holds  $\|x\| = |\text{normsequence}(G, y)|$ .
- (8) For all elements  $x, y, z$  of  $\prod \overline{G}$  and for every element  $i$  of  $\mathbb{N}$  such that  $i \in \text{dom } x$  and  $z = (\prod^\circ(\langle +_{G_i} \rangle_i))(x, y)$  holds  $(\text{normsequence}(G, z))(i) \leq (\text{normsequence}(G, x) + \text{normsequence}(G, y))(i)$ .
- (9) For every element  $x$  of  $\prod \overline{G}$  and for every element  $i$  of  $\mathbb{N}$  such that  $i \in \text{dom } x$  holds  $0 \leq (\text{normsequence}(G, x))(i)$ .

Let  $G$  be a real norm space-sequence. Observe that  $\prod G$  is real normed space-like, real linear space-like, Abelian, add-associative, right zeroed, and right complementable.

One can prove the following propositions:

- (10) Let  $G$  be a real norm space-sequence,  $i$  be an element of  $\text{dom } G$ ,  $x$  be a point of  $\prod G$ ,  $y$  be an element of  $\prod \overline{G}$ , and  $x_1$  be a point of  $G(i)$ . If  $y = x$  and  $x_1 = y(i)$ , then  $\|x_1\| \leq \|x\|$ .
- (11) Let  $G$  be a real norm space-sequence,  $i$  be an element of  $\text{dom } G$ ,  $x, y$  be points of  $\prod G$ ,  $x_1, y_1$  be points of  $G(i)$ , and  $z_1, z_2$  be elements of  $\prod \overline{G}$ . If  $x_1 = z_1(i)$  and  $z_1 = x$  and  $y_1 = z_2(i)$  and  $z_2 = y$ , then  $\|y_1 - x_1\| \leq \|y - x\|$ .
- (12) Let  $G$  be a real norm space-sequence,  $s_1$  be a sequence of  $\prod G$ ,  $x_0$  be a point of  $\prod G$ , and  $y_0$  be an element of  $\prod \overline{G}$ . Suppose  $x_0 = y_0$  and  $s_1$  is convergent and  $\lim s_1 = x_0$ . Let  $i$  be an element of  $\text{dom } G$ . Then there exists a sequence  $s_2$  of  $G(i)$  such that  $s_2$  is convergent and  $y_0(i) = \lim s_2$  and for every element  $m$  of  $\mathbb{N}$  there exists an element  $s_3$  of  $\prod \overline{G}$  such that  $s_3 = s_1(m)$  and  $s_2(m) = s_3(i)$ .
- (13) Let  $G$  be a real norm space-sequence,  $s_1$  be a sequence of  $\prod G$ ,  $x_0$  be a point of  $\prod G$ , and  $y_0$  be an element of  $\prod \overline{G}$ . Suppose that
  - (i)  $x_0 = y_0$ , and
  - (ii) for every element  $i$  of  $\text{dom } G$  there exists a sequence  $s_2$  of  $G(i)$  such that  $s_2$  is convergent and  $y_0(i) = \lim s_2$  and for every element  $m$  of  $\mathbb{N}$  there exists an element  $s_3$  of  $\prod \overline{G}$  such that  $s_3 = s_1(m)$  and  $s_2(m) = s_3(i)$ .
 Then  $s_1$  is convergent and  $\lim s_1 = x_0$ .
- (14) For every real norm space-sequence  $G$  such that for every element  $i$  of  $\text{dom } G$  holds  $G(i)$  is complete holds  $\prod G$  is complete.

## REFERENCES

- [1] Grzegorz Bancerek. König's theorem. *Formalized Mathematics*, 1(3):589–593, 1990.
- [2] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [4] Czesław Byliński. Binary operations. *Formalized Mathematics*, 1(1):175–180, 1990.
- [5] Czesław Byliński. The complex numbers. *Formalized Mathematics*, 1(3):507–513, 1990.
- [6] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(3):529–536, 1990.
- [7] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [8] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [9] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [10] Czesław Byliński. The sum and product of finite sequences of real numbers. *Formalized Mathematics*, 1(4):661–668, 1990.
- [11] Agata Darmochwał. The Euclidean space. *Formalized Mathematics*, 2(4):599–603, 1991.
- [12] Jarosław Kotowicz. Convergent sequences and the limit of sequences. *Formalized Mathematics*, 1(2):273–275, 1990.
- [13] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(2):269–272, 1990.
- [14] Eugeniusz Kusak, Wojciech Leńczuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Formalized Mathematics*, 1(2):335–342, 1990.
- [15] Anna Lango and Grzegorz Bancerek. Product of families of groups and vector spaces. *Formalized Mathematics*, 3(2):235–240, 1992.
- [16] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(1):223–230, 1990.
- [17] Jan Popiołek. Real normed space. *Formalized Mathematics*, 2(1):111–115, 1991.
- [18] Yasunari Shidama. Banach space of bounded linear operators. *Formalized Mathematics*, 12(1):39–48, 2004.
- [19] Andrzej Trybulec. Subsets of complex numbers. *To appear in Formalized Mathematics*.
- [20] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [21] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(2):291–296, 1990.
- [22] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [23] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.

Received July 9, 2007

---