The First Mean Value Theorem for Integrals

Keiko Narita Hirosaki-city Aomori, Japan Noboru Endou Gifu National College of Technology Japan

Yasunari Shidama Shinshu University Nagano, Japan

Summary. In this article, we prove the first mean value theorem for integrals [16]. The formalization of various theorems about the properties of the Lebesgue integral is also presented.

MML identifier: MESFUNC7, version: 7.8.09 4.97.1001

The notation and terminology used in this paper are introduced in the following articles: [20], [2], [17], [6], [1], [4], [21], [22], [11], [3], [9], [8], [10], [18], [19], [5], [13], [12], [14], [15], and [7].

1. Lemmas for Extended Real Valued Functions

For simplicity, we use the following convention: X is a non empty set, S is a σ -field of subsets of X, M is a σ -measure on S, f, g are partial functions from X to $\overline{\mathbb{R}}$, and E is an element of S.

One can prove the following three propositions:

- (1) If for every element x of X such that $x \in \text{dom } f$ holds $f(x) \leq g(x)$, then g f is non-negative.
- (2) For every set Y and for every partial function f from X to $\overline{\mathbb{R}}$ and for every real number r holds $(r f) \upharpoonright Y = r (f \upharpoonright Y)$.
- (3) Suppose f is integrable on M and g is integrable on M and g f is non-negative. Then there exists an element E of S such that $E = \operatorname{dom} f \cap \operatorname{dom} g$ and $\int f \upharpoonright E \, \mathrm{d}M \leq \int g \upharpoonright E \, \mathrm{d}M$.

KEIKO NARITA et al.

2. σ -Finite Sets

Let us consider X. One can verify that there exists a partial function from X to $\overline{\mathbb{R}}$ which is non-negative.

Let us consider X, f. Then |f| is a non-negative partial function from X to $\overline{\mathbb{R}}$.

Next we state the proposition

- (4) Suppose f is integrable on M. Then there exists a function F from \mathbb{N} into S such that
- (i) for every element n of N holds $F(n) = \text{dom } f \cap \text{GTE-dom}(|f|, \overline{\mathbb{R}}(\frac{1}{n+1})),$
- (ii) dom $f \setminus \text{EQ-dom}(f, 0_{\overline{\mathbb{R}}}) = \bigcup \text{rng } F$, and
- (iii) for every element n of N holds $F(n) \in S$ and $M(F(n)) < +\infty$.

3. The First Mean Value Theorem for Integrals

Let F be a binary relation. We introduce F is extreal-yielding as a synonym of F is extended real-valued.

Let k be a natural number and let x be an element of $\overline{\mathbb{R}}$. Then $k \mapsto x$ is a finite sequence of elements of $\overline{\mathbb{R}}$.

Let us note that there exists a finite sequence which is extreal-yielding.

The binary operation $\cdot_{\overline{\mathbb{R}}}$ on $\overline{\mathbb{R}}$ is defined by:

 $(\text{Def. 2})^1 \ \ \text{For all elements } x, \, y \text{ of } \overline{\mathbb{R}} \text{ holds } \cdot_{\overline{\mathbb{R}}}(x, \, y) = x \cdot y.$

One can check that $\cdot_{\overline{\mathbb{R}}}$ is commutative and associative.

One can prove the following proposition

(5) $\mathbf{1}_{\cdot_{\overline{\mathbb{R}}}} = 1.$

One can check that $\cdot_{\overline{\mathbb{R}}}$ is unital.

Let F be an extreal-yielding finite sequence. The functor $\prod F$ yields an element of $\overline{\mathbb{R}}$ and is defined by:

(Def. 3) There exists a finite sequence f of elements of $\overline{\mathbb{R}}$ such that f = F and $\prod F = \bigcup_{\overline{\mathbb{R}}} \circledast f$.

Let x be an element of $\overline{\mathbb{R}}$ and let n be a natural number. Note that $n \mapsto x$ is extreal-yielding.

Let x be an element of $\overline{\mathbb{R}}$ and let k be a natural number. The functor x^k is defined by:

(Def. 4) $x^k = \prod (k \mapsto x).$

Let x be an element of $\overline{\mathbb{R}}$ and let k be a natural number. Then x^k is an extended real number.

Let us note that $\varepsilon_{\overline{\mathbb{R}}}$ is extreal-yielding.

¹The definition (Def. 1) has been removed.

Let r be an element of $\overline{\mathbb{R}}$. Note that $\langle r \rangle$ is extreal-yielding.

We now state two propositions:

- (6) $\prod(\varepsilon_{\overline{\mathbb{R}}}) = 1.$
- (7) For every element r of $\overline{\mathbb{R}}$ holds $\prod \langle r \rangle = r$.

Let f, g be extreal-yielding finite sequences. Observe that $f \cap g$ is extreal-yielding.

We now state three propositions:

- (8) For every extreal-yielding finite sequence F and for every element r of $\overline{\mathbb{R}}$ holds $\prod (F \cap \langle r \rangle) = \prod F \cdot r$.
- (9) For every element x of $\overline{\mathbb{R}}$ holds $x^1 = x$.
- (10) For every element x of $\overline{\mathbb{R}}$ and for every natural number k holds $x^{k+1} = x^k \cdot x$.

Let k be a natural number and let us consider X, f. The functor f^k yields a partial function from X to $\overline{\mathbb{R}}$ and is defined by:

(Def. 5) $\operatorname{dom}(f^k) = \operatorname{dom} f$ and for every element x of X such that $x \in \operatorname{dom}(f^k)$ holds $f^k(x) = f(x)^k$.

Next we state several propositions:

- (11) For every element x of $\overline{\mathbb{R}}$ and for every real number y and for every natural number k such that x = y holds $x^k = y^k$.
- (12) For every element x of $\overline{\mathbb{R}}$ and for every natural number k such that $0 \leq x$ holds $0 \leq x^k$.
- (13) For every natural number k such that $1 \le k$ holds $+\infty^k = +\infty$.
- (14) Let k be a natural number and given X, S, f, E. If $E \subseteq \text{dom } f$ and f is measurable on E, then $|f|^k$ is measurable on E.
- (15) Suppose dom $f \cap \text{dom } g = E$ and f is finite and g is finite and f is measurable on E and g is measurable on E. Then f g is measurable on E.
- (16) If $\operatorname{rng} f$ is bounded, then f is finite.
- (17) Let M be a σ -measure on S, f, g be partial functions from X to $\overline{\mathbb{R}}$, E be an element of S, and F be a non empty subset of $\overline{\mathbb{R}}$. Suppose dom $f \cap$ dom g = E and rng f = F and g is finite and f is measurable on E and rng f is bounded and g is integrable on M. Then $(f g) \upharpoonright E$ is integrable on M and there exists an element c of \mathbb{R} such that $c \ge \inf F$ and $c \le \sup F$ and $\int (f |g|) \upharpoonright E \, \mathrm{d}M = \overline{\mathbb{R}}(c) \cdot \int |g| \upharpoonright E \, \mathrm{d}M$.

4. Selected Properties of Integrals

We use the following convention: E_1 , E_2 denote elements of S, x, A denote sets, and a, b denote real numbers.

The following propositions are true:

- (18) $|f| \upharpoonright A = |f \upharpoonright A|.$
- (19) $\operatorname{dom}(|f| + |g|) = \operatorname{dom} f \cap \operatorname{dom} g$ and $\operatorname{dom} |f + g| \subseteq \operatorname{dom} |f|$.
- (20) $|f| \mid \operatorname{dom} |f+g| + |g| \mid \operatorname{dom} |f+g| = (|f|+|g|) \mid \operatorname{dom} |f+g|.$
- (21) If $x \in \text{dom} |f + g|$, then $|f + g|(x) \le (|f| + |g|)(x)$.
- (22) Suppose f is integrable on M and g is integrable on M. Then there exists an element E of S such that $E = \operatorname{dom}(f+g)$ and $\int |f+g| \upharpoonright E \, \mathrm{d}M \leq \int |f| \upharpoonright E \, \mathrm{d}M + \int |g| \upharpoonright E \, \mathrm{d}M$.
- (23) $\max_{+}(\chi_{A,X}) = \chi_{A,X}.$
- (24) If $M(E) < +\infty$, then $\chi_{E,X}$ is integrable on M and $\int \chi_{E,X} dM = M(E)$ and $\int \chi_{E,X} \upharpoonright E dM = M(E)$.
- (25) If $M(E_1 \cap E_2) < +\infty$, then $\int \chi_{(E_1),X} \upharpoonright E_2 \, \mathrm{d}M = M(E_1 \cap E_2)$.
- (26) Suppose f is integrable on M and $E \subseteq \text{dom } f$ and $M(E) < +\infty$ and for every element x of X such that $x \in E$ holds $a \leq f(x) \leq b$. Then $\overline{\mathbb{R}}(a) \cdot M(E) \leq \int f \upharpoonright E \, \mathrm{d}M \leq \overline{\mathbb{R}}(b) \cdot M(E).$

References

- Grzegorz Bancerek. The fundamental properties of natural numbers. Formalized Mathematics, 1(1):41-46, 1990.
- [2] Grzegorz Bancerek. The ordinal numbers. Formalized Mathematics, 1(1):91–96, 1990.
- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. Formalized Mathematics, 1(1):107–114, 1990.
- [4] Józef Białas. Infimum and supremum of the set of real numbers. Measure theory. Formalized Mathematics, 2(1):163-171, 1991.
- [5] Józef Białas. Series of positive real numbers. Measure theory. Formalized Mathematics, 2(1):173–183, 1991.
- [6] Józef Białas. The σ -additive measure theory. Formalized Mathematics, 2(2):263–270, 1991.
- [7] Józef Białas. Some properties of the intervals. Formalized Mathematics, 5(1):21–26, 1996.
- [8] Czesław Byliński. Binary operations. Formalized Mathematics, 1(1):175–180, 1990.
- Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. Formalized Mathematics, 1(3):529–536, 1990.
- [10] Czesław Byliński. Functions from a set to a set. Formalized Mathematics, 1(1):153–164, 1990;
- [11] Čzesław Byliński. Partial functions. Formalized Mathematics, 1(2):357–367, 1990.
- [12] Noboru Endou and Yasunari Shidama. Integral of measurable function. Formalized Mathematics, 14(2):53–70, 2006.
- [13] Noboru Endou, Katsumi Wasaki, and Yasunari Shidama. Basic properties of extended real numbers. Formalized Mathematics, 9(3):491–494, 2001.
- [14] Noboru Endou, Katsumi Wasaki, and Yasunari Shidama. Definitions and basic properties of measurable functions. *Formalized Mathematics*, 9(3):495–500, 2001.
- [15] Noboru Endou, Katsumi Wasaki, and Yasunari Shidama. The measurability of extended real valued functions. *Formalized Mathematics*, 9(3):525–529, 2001.
- [16] P. R. Halmos. *Measure Theory*. Springer-Verlag, 1987.
- [17] Krzysztof Hryniewiecki. Basic properties of real numbers. Formalized Mathematics, 1(1):35–40, 1990.
- [18] Rafał Kwiatek. Factorial and Newton coefficients. *Formalized Mathematics*, 1(5):887–890, 1990.
- [19] Andrzej Nędzusiak. σ -fields and probability. Formalized Mathematics, 1(2):401–407, 1990.
- [20] Zinaida Trybulec. Properties of subsets. Formalized Mathematics, 1(1):67–71, 1990.
- [21] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.

[22] Edmund Woronowicz. Relations defined on sets. Formalized Mathematics, 1(1):181–186, 1990.

Received October 30, 2007