

# Ramsey's Theorem

Marco Riccardi  
 Casella Postale 49  
 54038 Montignoso, Italy

**Summary.** The goal of this article is to formalize two versions of Ramsey's theorem. The theorems are not phrased in the usually pictorial representation of a coloured graph but use a set-theoretic terminology. After some useful lemma, the second section presents a generalization of Ramsey's theorem on infinite set closely following the book [9]. The last section includes the formalization of the theorem in a more known version (see [1]).

MML identifier: RAMSEY\_1, version: 7.9.01 4.101.1015

The notation and terminology used here are introduced in the following papers: [15], [16], [17], [4], [3], [6], [12], [7], [2], [5], [8], [14], [13], [10], and [11].

## 1. PRELIMINARIES

For simplicity, we adopt the following convention:  $n, m, k$  are natural numbers,  $X, Y, Z$  are sets,  $f$  is a function from  $X$  into  $Y$ , and  $H$  is a subset of  $X$ .

Let us consider  $X, Y, H$  and let  $P$  be a partition of  $[X]^Y$ . We say that  $H$  is homogeneous for  $P$  if and only if:

(Def. 1) There exists an element  $p$  of  $P$  such that  $[H]^Y \subseteq p$ .

Let us consider  $n$  and let  $X$  be an infinite set. One can check that  $[X]^n$  is non empty.

Let us consider  $n, X, Y, f$ . Let us assume that  $f$  is one-to-one and  $\overline{n} \subseteq \overline{X}$  and  $X$  is non empty and  $Y$  is non empty. The functor  $f||^n$  yields a function from  $[X]^n$  into  $[Y]^n$  and is defined by:

(Def. 2) For every element  $x$  of  $[X]^n$  holds  $(f||^n)(x) = f \circ x$ .

Next we state four propositions:

- (1) If  $f$  is one-to-one and  $\overline{n} \subseteq \overline{X}$  and  $X$  is non empty and  $Y$  is non empty, then  $[f \circ H]^n = (f||^n)^\circ([H]^n)$ .
- (2) If  $X$  is infinite and  $X \subseteq \omega$ , then  $\overline{X} = \omega$ .
- (3) If  $X$  is infinite, then  $X \cup Y$  is infinite.
- (4) If  $X$  is infinite and  $Y$  is finite, then  $X \setminus Y$  is infinite.

Let  $X$  be an infinite set and let  $Y$  be a set. Note that  $X \cup Y$  is infinite.

Let  $X$  be an infinite set and let  $Y$  be a finite set. One can verify that  $X \setminus Y$  is infinite.

The following propositions are true:

- (5)  $[X]^0 = \{0\}$ .
- (6) For every finite set  $X$  such that  $\text{card } X < n$  holds  $[X]^n$  is empty.
- (7) If  $X \subseteq Y$ , then  $[X]^Z \subseteq [Y]^Z$ .
- (8) If  $X$  is finite and  $Y$  is finite and  $\overline{Y} = X$ , then  $[Y]^X = \{Y\}$ .
- (9) If  $X$  is non empty and  $Y$  is non empty, then  $f$  is constant iff there exists an element  $y$  of  $Y$  such that  $\text{rng } f = \{y\}$ .
- (10) For every finite set  $X$  such that  $k \leq \text{card } X$  there exists a subset  $Y$  of  $X$  such that  $\text{card } Y = k$ .
- (11) If  $m \geq 1$ , then  $n + 1 \leq \binom{n+m}{m}$ .
- (12) If  $m \geq 1$  and  $n \geq 1$ , then  $m + 1 \leq \binom{n+m}{m}$ .
- (13) Let  $X$  be a non empty set,  $p_1, p_2$  be elements of  $X$ ,  $P$  be a partition of  $X$ , and  $A$  be an element of  $P$ . Suppose  $p_1 \in A$  and (the projection onto  $P$ )( $p_1$ ) = (the projection onto  $P$ )( $p_2$ ). Then  $p_2 \in A$ .

## 2. INFINITE RAMSEY THEOREM

We now state two propositions:

- (14) Let  $F$  be a function from  $[X]^n$  into  $k$ . Suppose  $k \neq 0$  and  $X$  is infinite. Then there exists  $H$  such that  $H$  is infinite and  $F|[H]^n$  is constant.
- (15) Let  $X$  be an infinite set and  $P$  be a partition of  $[X]^n$ . If  $\overline{P} = k$ , then there exists a subset of  $X$  which is infinite and homogeneous for  $P$ .

## 3. RAMSEY'S THEOREM

The scheme *BinInd2* concerns a binary predicate  $\mathcal{P}$ , and states that:

$$\mathcal{P}[m, n]$$

provided the following conditions are satisfied:

- $\mathcal{P}[0, n]$  and  $\mathcal{P}[n, 0]$ , and
- If  $\mathcal{P}[m + 1, n]$  and  $\mathcal{P}[m, n + 1]$ , then  $\mathcal{P}[m + 1, n + 1]$ .

We now state two propositions:

- (16) Suppose  $m \geq 2$  and  $n \geq 2$ . Then there exists a natural number  $r$  such that
- (i)  $r \leq \binom{m+n-2}{m-1}$ ,
  - (ii)  $r \geq 2$ , and
  - (iii) for every finite set  $X$  and for every function  $F$  from  $[X]^2$  into  $\text{Seg } 2$  such that  $\text{card } X \geq r$  there exists a subset  $S$  of  $X$  such that  $\text{card } S \geq m$  and  $\text{rng}(F \upharpoonright [S]^2) = \{1\}$  or  $\text{card } S \geq n$  and  $\text{rng}(F \upharpoonright [S]^2) = \{2\}$ .
- (17) Let  $m$  be a natural number. Then there exists a natural number  $r$  such that for every finite set  $X$  and for every partition  $P$  of  $[X]^2$  if  $\text{card } X \geq r$  and  $\overline{P} = 2$ , then there exists a subset  $S$  of  $X$  such that  $\text{card } S \geq m$  and  $S$  is homogeneous for  $P$ .

## REFERENCES

- [1] M. Aigner and G. M. Ziegler. *Proofs from THE BOOK*. Springer-Verlag, Berlin Heidelberg New York, 2004.
- [2] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(2):377–382, 1990.
- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [4] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [5] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [6] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [7] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(1):165–167, 1990.
- [8] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [9] T. J. Jech. *Set Theory*. Springer-Verlag, Berlin Heidelberg New York, 2002.
- [10] Rafał Kwiatek. Factorial and Newton coefficients. *Formalized Mathematics*, 1(5):887–890, 1990.
- [11] Takaya Nishiyama and Yasuho Mizuhara. Binary arithmetics. *Formalized Mathematics*, 4(1):83–86, 1993.
- [12] Konrad Raczkowski and Paweł Sadowski. Equivalence relations and classes of abstraction. *Formalized Mathematics*, 1(3):441–444, 1990.
- [13] Marco Riccardi. The sylow theorems. *Formalized Mathematics*, 15(3):159–165, 2007.
- [14] Andrzej Trybulec. A Borsuk theorem on homotopy types. *Formalized Mathematics*, 2(4):535–545, 1991.
- [15] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [16] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [17] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.

Received April 18, 2008

---