

# Extended Riemann Integral of Functions of Real Variable and One-sided Laplace Transform<sup>1</sup>

Masahiko Yamazaki  
 Shinshu University  
 Nagano, Japan

Hiroshi Yamazaki  
 Shinshu University  
 Nagano, Japan

Yasunari Shidama  
 Shinshu University  
 Nagano, Japan

**Summary.** In this article, we defined a variety of extended Riemann integrals and proved that such integration is linear. Furthermore, we defined the one-sided Laplace transform and proved the linearity of that operator.

MML identifier: INTEGR10, version: 7.9.01 4.101.1015

The papers [11], [1], [5], [12], [10], [2], [7], [6], [8], [9], [3], [4], and [13] provide the terminology and notation for this paper.

## 1. PRELIMINARIES

In this paper  $a, b, r$  are elements of  $\mathbb{R}$ .

We now state three propositions:

- (1) For all real numbers  $a, b, g_1, M$  such that  $a < b$  and  $0 < g_1$  and  $0 < M$  there exists  $r$  such that  $a < r < b$  and  $(b - r) \cdot M < g_1$ .
- (2) For all real numbers  $a, b, g_1, M$  such that  $a < b$  and  $0 < g_1$  and  $0 < M$  there exists  $r$  such that  $a < r < b$  and  $(r - a) \cdot M < g_1$ .
- (3)  $\exp b - \exp a = \int_a^b (\text{the function } \exp)(x)dx.$

---

<sup>1</sup>This work has been partially supported by the MEXT grant Grant-in-Aid for Young Scientists (B)16700156.

## 2. THE DEFINITION OF EXTENDED RIEMANN INTEGRAL

Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$  and let  $a, b$  be real numbers. We say that  $f$  is right extended Riemann integrable on  $a, b$  if and only if the conditions (Def. 1) are satisfied.

- (Def. 1)(i) For every real number  $d$  such that  $a \leq d < b$  holds  $f$  is integrable on  $[a, d]$  and  $f|_{[a, d]}$  is bounded, and  
(ii) there exists a partial function  $\mathcal{I}$  from  $\mathbb{R}$  to  $\mathbb{R}$  such that  $\text{dom } \mathcal{I} = [a, b[$  and for every real number  $x$  such that  $x \in \text{dom } \mathcal{I}$  holds  $\mathcal{I}(x) = \int_a^x f(x)dx$  and  $\mathcal{I}$  is left convergent in  $b$ .

Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$  and let  $a, b$  be real numbers. We say that  $f$  is left extended Riemann integrable on  $a, b$  if and only if the conditions (Def. 2) are satisfied.

- (Def. 2)(i) For every real number  $d$  such that  $a < d \leq b$  holds  $f$  is integrable on  $[d, b]$  and  $f|_{[d, b]}$  is bounded, and  
(ii) there exists a partial function  $\mathcal{I}$  from  $\mathbb{R}$  to  $\mathbb{R}$  such that  $\text{dom } \mathcal{I} = ]a, b]$  and for every real number  $x$  such that  $x \in \text{dom } \mathcal{I}$  holds  $\mathcal{I}(x) = \int_x^b f(x)dx$  and  $\mathcal{I}$  is right convergent in  $a$ .

Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$  and let  $a, b$  be real numbers. Let us assume that  $f$  is right extended Riemann integrable on  $a, b$ . The functor

$(R^>) \int_a^b f(x)dx$  yielding a real number is defined by the condition (Def. 3).

- (Def. 3) There exists a partial function  $\mathcal{I}$  from  $\mathbb{R}$  to  $\mathbb{R}$  such that  $\text{dom } \mathcal{I} = [a, b[$  and for every real number  $x$  such that  $x \in \text{dom } \mathcal{I}$  holds  $\mathcal{I}(x) = \int_a^x f(x)dx$  and  $\mathcal{I}$  is left convergent in  $b$  and  $(R^>) \int_a^b f(x)dx = \lim_{b^-} \mathcal{I}$ .

Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$  and let  $a, b$  be real numbers. Let us assume that  $f$  is left extended Riemann integrable on  $a, b$ . The functor

$(R^<) \int_a^b f(x)dx$  yielding a real number is defined by the condition (Def. 4).

- (Def. 4) There exists a partial function  $\mathcal{I}$  from  $\mathbb{R}$  to  $\mathbb{R}$  such that  $\text{dom } \mathcal{I} = ]a, b]$  and for every real number  $x$  such that  $x \in \text{dom } \mathcal{I}$  holds  $\mathcal{I}(x) = \int_x^b f(x)dx$

and  $\mathcal{I}$  is right convergent in  $a$  and  $(R^<) \int_a^b f(x)dx = \lim_{a^+} \mathcal{I}$ .

Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$  and let  $a$  be a real number. We say that  $f$  is extended Riemann integrable on  $a, +\infty$  if and only if the conditions (Def. 5) are satisfied.

- (Def. 5)(i) For every real number  $b$  such that  $a \leq b$  holds  $f$  is integrable on  $[a, b]$  and  $f|_{[a, b]}$  is bounded, and  
(ii) there exists a partial function  $\mathcal{I}$  from  $\mathbb{R}$  to  $\mathbb{R}$  such that  $\text{dom } \mathcal{I} = [a, +\infty[$  and for every real number  $x$  such that  $x \in \text{dom } \mathcal{I}$  holds  $\mathcal{I}(x) = \int_a^x f(x)dx$  and  $\mathcal{I}$  is convergent in  $+\infty$ .

Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$  and let  $b$  be a real number. We say that  $f$  is extended Riemann integrable on  $-\infty, b$  if and only if the conditions (Def. 6) are satisfied.

- (Def. 6)(i) For every real number  $a$  such that  $a \leq b$  holds  $f$  is integrable on  $[a, b]$  and  $f|_{[a, b]}$  is bounded, and  
(ii) there exists a partial function  $\mathcal{I}$  from  $\mathbb{R}$  to  $\mathbb{R}$  such that  $\text{dom } \mathcal{I} = ]-\infty, b]$  and for every real number  $x$  such that  $x \in \text{dom } \mathcal{I}$  holds  $\mathcal{I}(x) = \int_x^b f(x)dx$  and  $\mathcal{I}$  is convergent in  $-\infty$ .

Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$  and let  $a$  be a real number. Let us assume that  $f$  is extended Riemann integrable on  $a, +\infty$ . The functor

$(R^>) \int_a^{+\infty} f(x)dx$  yielding a real number is defined by the condition (Def. 7).

- (Def. 7) There exists a partial function  $\mathcal{I}$  from  $\mathbb{R}$  to  $\mathbb{R}$  such that  $\text{dom } \mathcal{I} = [a, +\infty[$  and for every real number  $x$  such that  $x \in \text{dom } \mathcal{I}$  holds  $\mathcal{I}(x) = \int_a^x f(x)dx$  and  $\mathcal{I}$  is convergent in  $+\infty$  and  $(R^>) \int_a^{+\infty} f(x)dx = \lim_{+\infty} \mathcal{I}$ .

Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$  and let  $b$  be a real number. Let us assume that  $f$  is extended Riemann integrable on  $-\infty, b$ . The functor

$(R^<) \int_{-\infty}^b f(x)dx$  yields a real number and is defined by the condition (Def. 8).

- (Def. 8) There exists a partial function  $\mathcal{I}$  from  $\mathbb{R}$  to  $\mathbb{R}$  such that  $\text{dom } \mathcal{I} = ]-\infty, b]$  and for every real number  $x$  such that  $x \in \text{dom } \mathcal{I}$  holds  $\mathcal{I}(x) = \int_x^b f(x)dx$

and  $\mathcal{I}$  is convergent in  $-\infty$  and  $(R^<) \int_{-\infty}^b f(x)dx = \lim_{-\infty} \mathcal{I}$ .

Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$ . We say that  $f$  is  $\infty$ -extended Riemann integrable if and only if:

(Def. 9)  $f$  is extended Riemann integrable on  $0, +\infty$  and extended Riemann integrable on  $-\infty, 0$ .

Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$ . The functor  $(R) \int_{-\infty}^{+\infty} f(x)dx$  yields a real number and is defined by:

$$(Def. 10) \quad (R) \int_{-\infty}^{+\infty} f(x)dx = (R^>) \int_0^{+\infty} f(x)dx + (R^<) \int_{-\infty}^0 f(x)dx.$$

### 3. LINEARITY OF EXTENDED RIEMANN INTEGRAL

One can prove the following propositions:

(4) Let  $f, g$  be partial functions from  $\mathbb{R}$  to  $\mathbb{R}$  and  $a, b$  be real numbers. Suppose that

- (i)  $a < b$ ,
- (ii)  $[a, b] \subseteq \text{dom } f$ ,
- (iii)  $[a, b] \subseteq \text{dom } g$ ,
- (iv)  $f$  is right extended Riemann integrable on  $a, b$ , and
- (v)  $g$  is right extended Riemann integrable on  $a, b$ .

Then  $f + g$  is right extended Riemann integrable on  $a, b$  and

$$(R^>) \int_a^b (f + g)(x)dx = (R^>) \int_a^b f(x)dx + (R^>) \int_a^b g(x)dx.$$

(5) Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$  and  $a, b$  be real numbers. Suppose  $a < b$  and  $[a, b] \subseteq \text{dom } f$  and  $f$  is right extended Riemann integrable on  $a, b$ . Let  $r$  be a real number. Then  $rf$  is right extended Riemann integrable

$$\text{on } a, b \text{ and } (R^>) \int_a^b (rf)(x)dx = r \cdot (R^>) \int_a^b f(x)dx.$$

(6) Let  $f, g$  be partial functions from  $\mathbb{R}$  to  $\mathbb{R}$  and  $a, b$  be real numbers. Suppose that

- (i)  $a < b$ ,
- (ii)  $[a, b] \subseteq \text{dom } f$ ,
- (iii)  $[a, b] \subseteq \text{dom } g$ ,
- (iv)  $f$  is left extended Riemann integrable on  $a, b$ , and
- (v)  $g$  is left extended Riemann integrable on  $a, b$ .

Then  $f + g$  is left extended Riemann integrable on  $a, b$  and

$$(R^<) \int_a^b (f + g)(x) dx = (R^<) \int_a^b f(x) dx + (R^<) \int_a^b g(x) dx.$$

- (7) Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$  and  $a, b$  be real numbers. Suppose  $a < b$  and  $[a, b] \subseteq \text{dom } f$  and  $f$  is left extended Riemann integrable on  $a, b$ . Let  $r$  be a real number. Then  $rf$  is left extended Riemann integrable

$$\text{on } a, b \text{ and } (R^<) \int_a^b (rf)(x) dx = r \cdot (R^<) \int_a^b f(x) dx.$$

- (8) Let  $f, g$  be partial functions from  $\mathbb{R}$  to  $\mathbb{R}$  and  $a$  be a real number. Suppose that

- (i)  $[a, +\infty[ \subseteq \text{dom } f$ ,
- (ii)  $[a, +\infty[ \subseteq \text{dom } g$ ,
- (iii)  $f$  is extended Riemann integrable on  $a, +\infty$ , and
- (iv)  $g$  is extended Riemann integrable on  $a, +\infty$ .

Then  $f + g$  is extended Riemann integrable on  $a, +\infty$  and

$$(R^>) \int_a^{+\infty} (f + g)(x) dx = (R^>) \int_a^{+\infty} f(x) dx + (R^>) \int_a^{+\infty} g(x) dx.$$

- (9) Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$  and  $a$  be a real number. Suppose  $[a, +\infty[ \subseteq \text{dom } f$  and  $f$  is extended Riemann integrable on  $a, +\infty$ . Let  $r$  be a real number. Then  $rf$  is extended Riemann integrable on  $a, +\infty$  and

$$(R^>) \int_a^{+\infty} (rf)(x) dx = r \cdot (R^>) \int_a^{+\infty} f(x) dx.$$

- (10) Let  $f, g$  be partial functions from  $\mathbb{R}$  to  $\mathbb{R}$  and  $b$  be a real number. Suppose that

- (i)  $] -\infty, b] \subseteq \text{dom } f$ ,
- (ii)  $] -\infty, b] \subseteq \text{dom } g$ ,
- (iii)  $f$  is extended Riemann integrable on  $-\infty, b$ , and
- (iv)  $g$  is extended Riemann integrable on  $-\infty, b$ .

Then  $f + g$  is extended Riemann integrable on  $-\infty, b$  and

$$(R^<) \int_{-\infty}^b (f + g)(x) dx = (R^<) \int_{-\infty}^b f(x) dx + (R^<) \int_{-\infty}^b g(x) dx.$$

- (11) Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$  and  $b$  be a real number. Suppose  $] -\infty, b] \subseteq \text{dom } f$  and  $f$  is extended Riemann integrable on  $-\infty, b$ . Let  $r$  be a real number. Then  $rf$  is extended Riemann integrable on  $-\infty, b$  and

$$(R^<) \int_{-\infty}^b (rf)(x) dx = r \cdot (R^<) \int_{-\infty}^b f(x) dx.$$

- (12) Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$  and  $a, b$  be real numbers.

Suppose  $a < b$  and  $[a, b] \subseteq \text{dom } f$  and  $f$  is integrable on  $[a, b]$  and  $f|_{[a, b]}$  is bounded. Then  $(R^>) \int_a^b f(x)dx = \int_a^b f(x)dx$ .

- (13) Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$  and  $a, b$  be real numbers. Suppose  $a < b$  and  $[a, b] \subseteq \text{dom } f$  and  $f$  is integrable on  $[a, b]$  and  $f|_{[a, b]}$  is bounded. Then  $(R^<) \int_a^b f(x)dx = \int_a^b f(x)dx$ .

#### 4. THE DEFINITION OF ONE-SIDED LAPLACE TRANSFORM

Let  $s$  be a real number. The functor  $e^{-s \cdot \square}$  yielding a function from  $\mathbb{R}$  into  $\mathbb{R}$  is defined by:

- (Def. 11) For every real number  $t$  holds  $e^{-s \cdot \square}(t) = (\text{the function exp})(-s \cdot t)$ .

Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$ . The one-sided Laplace transform of  $f$  yielding a partial function from  $\mathbb{R}$  to  $\mathbb{R}$  is defined by the conditions (Def. 12).

- (Def. 12)(i)  $\text{dom}(\text{the one-sided Laplace transform of } f) = ]0, +\infty[$ , and  
(ii) for every real number  $s$  such that  $s \in \text{dom}(\text{the one-sided Laplace transform of } f)$  holds  $(\text{the one-sided Laplace transform of } f)(s) = (R^>) \int_0^{+\infty} (f e^{-s \cdot \square})(x)dx$ .

#### 5. LINEARITY OF ONE-SIDED LAPLACE TRANSFORM

Next we state two propositions:

- (14) Let  $f, g$  be partial functions from  $\mathbb{R}$  to  $\mathbb{R}$ . Suppose that  
(i)  $\text{dom } f = [0, +\infty[$ ,  
(ii)  $\text{dom } g = [0, +\infty[$ ,  
(iii) for every real number  $s$  such that  $s \in ]0, +\infty[$  holds  $f e^{-s \cdot \square}$  is extended Riemann integrable on  $0, +\infty$ , and  
(iv) for every real number  $s$  such that  $s \in ]0, +\infty[$  holds  $g e^{-s \cdot \square}$  is extended Riemann integrable on  $0, +\infty$ .

Then

- (v) for every real number  $s$  such that  $s \in ]0, +\infty[$  holds  $(f + g) e^{-s \cdot \square}$  is extended Riemann integrable on  $0, +\infty$ , and  
(vi) the one-sided Laplace transform of  $f + g = (\text{the one-sided Laplace transform of } f) + (\text{the one-sided Laplace transform of } g)$ .  
(15) Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$  and  $a$  be a real number. Suppose  $\text{dom } f = [0, +\infty[$  and for every real number  $s$  such that  $s \in ]0, +\infty[$  holds  $f e^{-s \cdot \square}$  is extended Riemann integrable on  $0, +\infty$ . Then

- (i) for every real number  $s$  such that  $s \in ]0, +\infty[$  holds  $a f e^{-s \cdot \square}$  is extended Riemann integrable on  $0, +\infty$ , and
- (ii) the one-sided Laplace transform of  $a f = a$  the one-sided Laplace transform of  $f$ .

## REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [2] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [3] Noboru Endou and Artur Korniłowicz. The definition of the Riemann definite integral and some related lemmas. *Formalized Mathematics*, 8(1):93–102, 1999.
- [4] Noboru Endou, Katsumi Wasaki, and Yasunari Shidama. Definition of integrability for partial functions from  $\mathbb{R}$  to  $\mathbb{R}$  and integrability for continuous functions. *Formalized Mathematics*, 9(2):281–284, 2001.
- [5] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [6] Jarosław Kotowicz. Convergent sequences and the limit of sequences. *Formalized Mathematics*, 1(2):273–275, 1990.
- [7] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(2):269–272, 1990.
- [8] Jarosław Kotowicz. The limit of a real function at infinity. *Formalized Mathematics*, 2(1):17–28, 1991.
- [9] Jarosław Kotowicz. One-side limits of a real function at a point. *Formalized Mathematics*, 2(1):29–40, 1991.
- [10] Yatsuka Nakamura. Half open intervals in real numbers. *Formalized Mathematics*, 10(1):21–22, 2002.
- [11] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [12] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.
- [13] Yuguang Yang and Yasunari Shidama. Trigonometric functions and existence of circle ratio. *Formalized Mathematics*, 7(2):255–263, 1998.

Received July 22, 2008

---