

# Arithmetic Operations on Functions from Sets into Functional Sets

Artur Kornilowicz<sup>1</sup>  
 Institute of Computer Science  
 University of Białystok  
 Sosnowa 64, 15-887 Białystok  
 Poland

**Summary.** In this paper we introduce sets containing number-valued functions. Different arithmetic operations on maps between any set and such functional sets are later defined.

MML identifier: VALUED\_2, version: 7.11.01 4.117.1046

The notation and terminology used here are introduced in the following papers: [4], [9], [10], [2], [11], [6], [3], [1], [8], [5], and [7].

## 1. FUNCTIONAL SETS

In this paper  $x$ ,  $X$ ,  $X_1$ ,  $X_2$  are sets.

Let  $Y$  be a functional set. The functor  $\text{DOMS}(Y)$  is defined by:

(Def. 1)  $\text{DOMS}(Y) = \bigcup \{\text{dom } f : f \text{ ranges over elements of } Y\}$ .

Let us consider  $X$ . We say that  $X$  is complex-functions-membered if and only if:

(Def. 2) If  $x \in X$ , then  $x$  is a complex-valued function.

Let us consider  $X$ . We say that  $X$  is extended-real-functions-membered if and only if:

(Def. 3) If  $x \in X$ , then  $x$  is an extended real-valued function.

Let us consider  $X$ . We say that  $X$  is real-functions-membered if and only if:

---

<sup>1</sup>The article was written while the author visited Shinshu University, Nagano, Japan.

(Def. 4) If  $x \in X$ , then  $x$  is a real-valued function.

Let us consider  $X$ . We say that  $X$  is rational-functions-membered if and only if:

(Def. 5) If  $x \in X$ , then  $x$  is a rational-valued function.

Let us consider  $X$ . We say that  $X$  is integer-functions-membered if and only if:

(Def. 6) If  $x \in X$ , then  $x$  is an integer-valued function.

Let us consider  $X$ . We say that  $X$  is natural-functions-membered if and only if:

(Def. 7) If  $x \in X$ , then  $x$  is a natural-valued function.

One can check the following observations:

- \* every set which is natural-functions-membered is also integer-functions-membered,
- \* every set which is integer-functions-membered is also rational-functions-membered,
- \* every set which is rational-functions-membered is also real-functions-membered,
- \* every set which is real-functions-membered is also complex-functions-membered, and
- \* every set which is real-functions-membered is also extended-real-functions-membered.

Let us mention that every set which is empty is also natural-functions-membered.

Let  $f$  be a complex-valued function. Observe that  $\{f\}$  is complex-functions-membered.

One can verify that every set which is complex-functions-membered is also functional and every set which is extended-real-functions-membered is also functional.

One can verify that there exists a set which is natural-functions-membered and non empty.

Let  $X$  be a complex-functions-membered set. One can verify that every subset of  $X$  is complex-functions-membered.

Let  $X$  be an extended-real-functions-membered set. Note that every subset of  $X$  is extended-real-functions-membered.

Let  $X$  be a real-functions-membered set. Note that every subset of  $X$  is real-functions-membered.

Let  $X$  be a rational-functions-membered set. Observe that every subset of  $X$  is rational-functions-membered.

Let  $X$  be an integer-functions-membered set. Note that every subset of  $X$  is integer-functions-membered.

Let  $X$  be a natural-functions-membered set. Observe that every subset of  $X$  is natural-functions-membered.

Let  $D$  be a set. The functor  $\mathbb{C}\text{-PFuncs } D$  yields a set and is defined by:

(Def. 8) For every set  $f$  holds  $f \in \mathbb{C}\text{-PFuncs } D$  iff  $f$  is a partial function from  $D$  to  $\mathbb{C}$ .

Let  $D$  be a set. The functor  $\mathbb{C}\text{-Funcs } D$  yielding a set is defined by:

(Def. 9) For every set  $f$  holds  $f \in \mathbb{C}\text{-Funcs } D$  iff  $f$  is a function from  $D$  into  $\mathbb{C}$ .

Let  $D$  be a set. The functor  $\overline{\mathbb{R}}\text{-PFuncs } D$  yields a set and is defined by:

(Def. 10) For every set  $f$  holds  $f \in \overline{\mathbb{R}}\text{-PFuncs } D$  iff  $f$  is a partial function from  $D$  to  $\overline{\mathbb{R}}$ .

Let  $D$  be a set. The functor  $\overline{\mathbb{R}}\text{-Funcs } D$  yields a set and is defined as follows:

(Def. 11) For every set  $f$  holds  $f \in \overline{\mathbb{R}}\text{-Funcs } D$  iff  $f$  is a function from  $D$  into  $\overline{\mathbb{R}}$ .

Let  $D$  be a set. The functor  $\mathbb{R}\text{-PFuncs } D$  yielding a set is defined by:

(Def. 12) For every set  $f$  holds  $f \in \mathbb{R}\text{-PFuncs } D$  iff  $f$  is a partial function from  $D$  to  $\mathbb{R}$ .

Let  $D$  be a set. The functor  $\mathbb{R}\text{-Funcs } D$  yielding a set is defined by:

(Def. 13) For every set  $f$  holds  $f \in \mathbb{R}\text{-Funcs } D$  iff  $f$  is a function from  $D$  into  $\mathbb{R}$ .

Let  $D$  be a set. The functor  $\mathbb{Q}\text{-PFuncs } D$  yields a set and is defined as follows:

(Def. 14) For every set  $f$  holds  $f \in \mathbb{Q}\text{-PFuncs } D$  iff  $f$  is a partial function from  $D$  to  $\mathbb{Q}$ .

Let  $D$  be a set. The functor  $\mathbb{Q}\text{-Funcs } D$  yields a set and is defined by:

(Def. 15) For every set  $f$  holds  $f \in \mathbb{Q}\text{-Funcs } D$  iff  $f$  is a function from  $D$  into  $\mathbb{Q}$ .

Let  $D$  be a set. The functor  $\mathbb{Z}\text{-PFuncs } D$  yielding a set is defined by:

(Def. 16) For every set  $f$  holds  $f \in \mathbb{Z}\text{-PFuncs } D$  iff  $f$  is a partial function from  $D$  to  $\mathbb{Z}$ .

Let  $D$  be a set. The functor  $\mathbb{Z}\text{-Funcs } D$  yields a set and is defined as follows:

(Def. 17) For every set  $f$  holds  $f \in \mathbb{Z}\text{-Funcs } D$  iff  $f$  is a function from  $D$  into  $\mathbb{Z}$ .

Let  $D$  be a set. The functor  $\mathbb{N}\text{-PFuncs } D$  yields a set and is defined by:

(Def. 18) For every set  $f$  holds  $f \in \mathbb{N}\text{-PFuncs } D$  iff  $f$  is a partial function from  $D$  to  $\mathbb{N}$ .

Let  $D$  be a set. The functor  $\mathbb{N}\text{-Funcs } D$  yielding a set is defined by:

(Def. 19) For every set  $f$  holds  $f \in \mathbb{N}\text{-Funcs } D$  iff  $f$  is a function from  $D$  into  $\mathbb{N}$ .

The following propositions are true:

- (1)  $\mathbb{C}\text{-Funcs } X$  is a subset of  $\mathbb{C}\text{-PFuncs } X$ .
- (2)  $\overline{\mathbb{R}}\text{-Funcs } X$  is a subset of  $\overline{\mathbb{R}}\text{-PFuncs } X$ .
- (3)  $\mathbb{R}\text{-Funcs } X$  is a subset of  $\mathbb{R}\text{-PFuncs } X$ .
- (4)  $\mathbb{Q}\text{-Funcs } X$  is a subset of  $\mathbb{Q}\text{-PFuncs } X$ .
- (5)  $\mathbb{Z}\text{-Funcs } X$  is a subset of  $\mathbb{Z}\text{-PFuncs } X$ .

(6)  $\mathbb{N}$ -Funcs  $X$  is a subset of  $\mathbb{N}$ -PFuncs  $X$ .

Let us consider  $X$ . One can verify the following observations:

- \*  $\mathbb{C}$ -PFuncs  $X$  is complex-functions-membered,
- \*  $\mathbb{C}$ -Funcs  $X$  is complex-functions-membered,
- \*  $\overline{\mathbb{R}}$ -PFuncs  $X$  is extended-real-functions-membered,
- \*  $\overline{\mathbb{R}}$ -Funcs  $X$  is extended-real-functions-membered,
- \*  $\mathbb{R}$ -PFuncs  $X$  is real-functions-membered,
- \*  $\mathbb{R}$ -Funcs  $X$  is real-functions-membered,
- \*  $\mathbb{Q}$ -PFuncs  $X$  is rational-functions-membered,
- \*  $\mathbb{Q}$ -Funcs  $X$  is rational-functions-membered,
- \*  $\mathbb{Z}$ -PFuncs  $X$  is integer-functions-membered,
- \*  $\mathbb{Z}$ -Funcs  $X$  is integer-functions-membered,
- \*  $\mathbb{N}$ -PFuncs  $X$  is natural-functions-membered, and
- \*  $\mathbb{N}$ -Funcs  $X$  is natural-functions-membered.

Let  $X$  be a complex-functions-membered set. Observe that every element of  $X$  is complex-valued.

Let  $X$  be an extended-real-functions-membered set. One can check that every element of  $X$  is extended real-valued.

Let  $X$  be a real-functions-membered set. One can check that every element of  $X$  is real-valued.

Let  $X$  be a rational-functions-membered set. One can check that every element of  $X$  is rational-valued.

Let  $X$  be an integer-functions-membered set. Observe that every element of  $X$  is integer-valued.

Let  $X$  be a natural-functions-membered set. Observe that every element of  $X$  is natural-valued.

Let  $X, x$  be sets, let  $Y$  be a complex-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . Observe that  $f(x)$  is function-like and relation-like.

Let  $X, x$  be sets, let  $Y$  be an extended-real-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . Observe that  $f(x)$  is function-like and relation-like.

Let us consider  $X, x$ , let  $Y$  be a complex-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . One can check that  $f(x)$  is complex-valued.

Let us consider  $X, x$ , let  $Y$  be an extended-real-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . One can verify that  $f(x)$  is extended real-valued.

Let us consider  $X, x$ , let  $Y$  be a real-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . Note that  $f(x)$  is real-valued.

Let us consider  $X$ ,  $x$ , let  $Y$  be a rational-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . Note that  $f(x)$  is rational-valued.

Let us consider  $X$ ,  $x$ , let  $Y$  be an integer-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . Note that  $f(x)$  is integer-valued.

Let us consider  $X$ ,  $x$ , let  $Y$  be a natural-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . One can check that  $f(x)$  is natural-valued.

Let us consider  $X$  and let  $Y$  be a complex-membered set. One can check that  $X \rightarrowtail Y$  is complex-functions-membered.

Let us consider  $X$  and let  $Y$  be an extended real-membered set. Observe that  $X \rightarrowtail Y$  is extended-real-functions-membered.

Let us consider  $X$  and let  $Y$  be a real-membered set. Observe that  $X \rightarrowtail Y$  is real-functions-membered.

Let us consider  $X$  and let  $Y$  be a rational-membered set. Observe that  $X \rightarrowtail Y$  is rational-functions-membered.

Let us consider  $X$  and let  $Y$  be an integer-membered set. Observe that  $X \rightarrowtail Y$  is integer-functions-membered.

Let us consider  $X$  and let  $Y$  be a natural-membered set. One can verify that  $X \rightarrowtail Y$  is natural-functions-membered.

Let us consider  $X$  and let  $Y$  be a complex-membered set. Note that  $Y^X$  is complex-functions-membered.

Let us consider  $X$  and let  $Y$  be an extended real-membered set. Note that  $Y^X$  is extended-real-functions-membered.

Let us consider  $X$  and let  $Y$  be a real-membered set. Note that  $Y^X$  is real-functions-membered.

Let us consider  $X$  and let  $Y$  be a rational-membered set. Note that  $Y^X$  is rational-functions-membered.

Let us consider  $X$  and let  $Y$  be an integer-membered set. Note that  $Y^X$  is integer-functions-membered.

Let us consider  $X$  and let  $Y$  be a natural-membered set. One can check that  $Y^X$  is natural-functions-membered.

Let  $R$  be a binary relation. We say that  $R$  is complex-functions-valued if and only if:

(Def. 20)  $\text{rng } R$  is complex-functions-membered.

We say that  $R$  is extended-real-functions-valued if and only if:

(Def. 21)  $\text{rng } R$  is extended-real-functions-membered.

We say that  $R$  is real-functions-valued if and only if:

(Def. 22)  $\text{rng } R$  is real-functions-membered.

We say that  $R$  is rational-functions-valued if and only if:

(Def. 23)  $\text{rng } R$  is rational-functions-membered.

We say that  $R$  is integer-functions-valued if and only if:

(Def. 24)  $\text{rng } R$  is integer-functions-membered.

We say that  $R$  is natural-functions-valued if and only if:

(Def. 25)  $\text{rng } R$  is natural-functions-membered.

Let  $f$  be a function. Let us observe that  $f$  is complex-functions-valued if and only if:

(Def. 26) For every set  $x$  such that  $x \in \text{dom } f$  holds  $f(x)$  is a complex-valued function.

Let us observe that  $f$  is extended-real-functions-valued if and only if:

(Def. 27) For every set  $x$  such that  $x \in \text{dom } f$  holds  $f(x)$  is an extended real-valued function.

Let us observe that  $f$  is real-functions-valued if and only if:

(Def. 28) For every set  $x$  such that  $x \in \text{dom } f$  holds  $f(x)$  is a real-valued function.

Let us observe that  $f$  is rational-functions-valued if and only if:

(Def. 29) For every set  $x$  such that  $x \in \text{dom } f$  holds  $f(x)$  is a rational-valued function.

Let us observe that  $f$  is integer-functions-valued if and only if:

(Def. 30) For every set  $x$  such that  $x \in \text{dom } f$  holds  $f(x)$  is an integer-valued function.

Let us observe that  $f$  is natural-functions-valued if and only if:

(Def. 31) For every set  $x$  such that  $x \in \text{dom } f$  holds  $f(x)$  is a natural-valued function.

One can verify the following observations:

- \* every binary relation which is natural-functions-valued is also integer-functions-valued,
- \* every binary relation which is integer-functions-valued is also rational-functions-valued,
- \* every binary relation which is rational-functions-valued is also real-functions-valued,
- \* every binary relation which is real-functions-valued is also extended-real-functions-valued, and
- \* every binary relation which is real-functions-valued is also complex-functions-valued.

Let us note that every binary relation which is empty is also natural-functions-valued.

Let us mention that there exists a function which is natural-functions-valued.

Let  $R$  be a complex-functions-valued binary relation. Note that  $\text{rng } R$  is complex-functions-membered.

Let  $R$  be an extended-real-functions-valued binary relation. Observe that  $\text{rng } R$  is extended-real-functions-membered.

Let  $R$  be a real-functions-valued binary relation. Note that  $\text{rng } R$  is real-functions-membered.

Let  $R$  be a rational-functions-valued binary relation. Observe that  $\text{rng } R$  is rational-functions-membered.

Let  $R$  be an integer-functions-valued binary relation. One can verify that  $\text{rng } R$  is integer-functions-membered.

Let  $R$  be a natural-functions-valued binary relation. One can check that  $\text{rng } R$  is natural-functions-membered.

Let us consider  $X$  and let  $Y$  be a complex-functions-membered set. Observe that every partial function from  $X$  to  $Y$  is complex-functions-valued.

Let us consider  $X$  and let  $Y$  be an extended-real-functions-membered set. One can check that every partial function from  $X$  to  $Y$  is extended-real-functions-valued.

Let us consider  $X$  and let  $Y$  be a real-functions-membered set. One can check that every partial function from  $X$  to  $Y$  is real-functions-valued.

Let us consider  $X$  and let  $Y$  be a rational-functions-membered set. Observe that every partial function from  $X$  to  $Y$  is rational-functions-valued.

Let us consider  $X$  and let  $Y$  be an integer-functions-membered set. Observe that every partial function from  $X$  to  $Y$  is integer-functions-valued.

Let us consider  $X$  and let  $Y$  be a natural-functions-membered set. Note that every partial function from  $X$  to  $Y$  is natural-functions-valued.

Let  $f$  be a complex-functions-valued function and let us consider  $x$ . Note that  $f(x)$  is function-like and relation-like.

Let  $f$  be an extended-real-functions-valued function and let us consider  $x$ . Observe that  $f(x)$  is function-like and relation-like.

Let  $f$  be a complex-functions-valued function and let us consider  $x$ . One can verify that  $f(x)$  is complex-valued.

Let  $f$  be an extended-real-functions-valued function and let us consider  $x$ . Note that  $f(x)$  is extended real-valued.

Let  $f$  be a real-functions-valued function and let us consider  $x$ . One can verify that  $f(x)$  is real-valued.

Let  $f$  be a rational-functions-valued function and let us consider  $x$ . Observe that  $f(x)$  is rational-valued.

Let  $f$  be an integer-functions-valued function and let us consider  $x$ . Note that  $f(x)$  is integer-valued.

Let  $f$  be a natural-functions-valued function and let us consider  $x$ . One can check that  $f(x)$  is natural-valued.

## 2. OPERATIONS

For simplicity, we adopt the following rules:  $Y, Y_1, Y_2$  are complex-functions-membered sets,  $c, c_1, c_2$  are complex numbers,  $f$  is a partial function from  $X$

to  $Y$ ,  $f_1$  is a partial function from  $X_1$  to  $Y_1$ ,  $f_2$  is a partial function from  $X_2$  to  $Y_2$ , and  $g, h, k$  are complex-valued functions.

We now state a number of propositions:

- (7) If  $g \neq \emptyset$  and  $g + c_1 = g + c_2$ , then  $c_1 = c_2$ .
- (8) If  $g \neq \emptyset$  and  $g - c_1 = g - c_2$ , then  $c_1 = c_2$ .
- (9) If  $g \neq \emptyset$  and  $g$  is non-empty and  $g c_1 = g c_2$ , then  $c_1 = c_2$ .
- (10)  $-(g + c) = -g - c$ .
- (11)  $-(g - c) = -g + c$ .
- (12)  $(g + c_1) + c_2 = g + (c_1 + c_2)$ .
- (13)  $(g + c_1) - c_2 = g + (c_1 - c_2)$ .
- (14)  $(g - c_1) + c_2 = g - (c_1 - c_2)$ .
- (15)  $g - c_1 - c_2 = g - (c_1 + c_2)$ .
- (16)  $g c_1 c_2 = g (c_1 \cdot c_2)$ .
- (17)  $-(g + h) = -g - h$ .
- (18)  $g - h = -(h - g)$ .
- (19)  $(g h)/k = g (h/k)$ .
- (20)  $(g/h) k = (g k)/h$ .
- (21)  $g/h/k = g/(h k)$ .
- (22)  $c - g = (-c) g$ .
- (23)  $c - g = -c g$ .
- (24)  $(-c) g = -c g$ .
- (25)  $-g h = (-g) h$ .
- (26)  $-g/h = (-g)/h$ .
- (27)  $-g/h = g/-h$ .

Let  $f$  be a complex-valued function and let  $c$  be a complex number. The functor  $f/c$  yields a function and is defined as follows:

(Def. 32)  $f/c = \frac{1}{c} f$ .

Let  $f$  be a complex-valued function and let  $c$  be a complex number. Note that  $f/c$  is complex-valued.

Let  $f$  be a real-valued function and let  $r$  be a real number. Note that  $f/r$  is real-valued.

Let  $f$  be a rational-valued function and let  $r$  be a rational number. One can check that  $f/r$  is rational-valued.

Let  $f$  be a complex-valued finite sequence and let  $c$  be a complex number. One can check that  $f/c$  is finite sequence-like.

The following propositions are true:

- (28)  $\text{dom}(g/c) = \text{dom } g$ .
- (29)  $(g/c)(x) = \frac{g(x)}{c}$ .



- (30)  $(-g)/c = -g/c.$
- (31)  $g/-c = -g/c.$
- (32)  $g/-c = (-g)/c.$
- (33) If  $g \neq \emptyset$  and  $g$  is non-empty and  $g/c_1 = g/c_2$ , then  $c_1 = c_2$ .
- (34)  $(g c_1)/c_2 = g \frac{c_1}{c_2}.$
- (35)  $(g/c_1) c_2 = (g c_2)/c_1.$
- (36)  $g/c_1/c_2 = g/(c_1 \cdot c_2).$
- (37)  $(g + h)/c = g/c + h/c.$
- (38)  $(g - h)/c = g/c - h/c.$
- (39)  $(g h)/c = g (h/c).$
- (40)  $(g/h)/c = g/(h c).$

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . The functor  $-f$  yields a function and is defined by:

(Def. 33)  $\text{dom}(-f) = \text{dom } f$  and for every set  $x$  such that  $x \in \text{dom}(-f)$  holds  $(-f)(x) = -f(x).$

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . Then  $-f$  is a partial function from  $X$  to  $\mathbb{C}\text{-PFuns DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a real-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . Then  $-f$  is a partial function from  $X$  to  $\mathbb{R}\text{-PFuns DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a rational-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . Then  $-f$  is a partial function from  $X$  to  $\mathbb{Q}\text{-PFuns DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be an integer-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . Then  $-f$  is a partial function from  $X$  to  $\mathbb{Z}\text{-PFuns DOMS}(Y)$ .

Let  $Y$  be a complex-functions-membered set and let  $f$  be a finite sequence of elements of  $Y$ . One can check that  $-f$  is finite sequence-like.

We now state two propositions:

- (41)  $--f = f.$
- (42) If  $-f_1 = -f_2$ , then  $f_1 = f_2.$

Let  $X$  be a complex-functions-membered set, let  $Y$  be a set, and let  $f$  be a partial function from  $X$  to  $Y$ . The functor  $f \circ -$  yielding a function is defined as follows:

(Def. 34)  $\text{dom}(f \circ -) = \text{dom } f$  and for every complex-valued function  $x$  such that  $x \in \text{dom}(f \circ -)$  holds  $(f \circ -)(x) = f(-x).$

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . The functor  $^1/f$  yields a function and is defined as follows:

(Def. 35)  $\text{dom } ^1/f = \text{dom } f$  and for every set  $x$  such that  $x \in \text{dom } ^1/f$  holds  $(^1/f)(x) = f(x)^{-1}$ .

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . Then  $^1/f$  is a partial function from  $X$  to  $\mathbb{C}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a real-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . Then  $^1/f$  is a partial function from  $X$  to  $\mathbb{R}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a rational-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . Then  $^1/f$  is a partial function from  $X$  to  $\mathbb{Q}\text{-PFuncs DOMS}(Y)$ .

Let  $Y$  be a complex-functions-membered set and let  $f$  be a finite sequence of elements of  $Y$ . Note that  $^1/f$  is finite sequence-like.

The following proposition is true

$$(43) \quad ^1/^1/f = f.$$

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . The functor  $|f|$  yields a function and is defined by:

(Def. 36)  $\text{dom } |f| = \text{dom } f$  and for every set  $x$  such that  $x \in \text{dom } |f|$  holds  $|f|(x) = |f(x)|$ .

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . Then  $|f|$  is a partial function from  $X$  to  $\mathbb{C}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a real-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . Then  $|f|$  is a partial function from  $X$  to  $\mathbb{R}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a rational-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . Then  $|f|$  is a partial function from  $X$  to  $\mathbb{Q}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be an integer-functions-membered set, and let  $f$  be a partial function from  $X$  to  $Y$ . Then  $|f|$  is a partial function from  $X$  to  $\mathbb{N}\text{-PFuncs DOMS}(Y)$ .

Let  $Y$  be a complex-functions-membered set and let  $f$  be a finite sequence of elements of  $Y$ . Note that  $|f|$  is finite sequence-like.

We now state the proposition

$$(44) \quad ||f|| = |f|.$$

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be a complex number. The functor  $f + c$

yields a function and is defined by:

(Def. 37)  $\text{dom}(f + c) = \text{dom } f$  and for every set  $x$  such that  $x \in \text{dom}(f + c)$  holds  $(f + c)(x) = c + f(x)$ .

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be a complex number. Then  $f + c$  is a partial function from  $X$  to  $\mathbb{C}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a real-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be a real number. Then  $f + c$  is a partial function from  $X$  to  $\mathbb{R}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a rational-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be a rational number. Then  $f + c$  is a partial function from  $X$  to  $\mathbb{Q}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be an integer-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be an integer number. Then  $f + c$  is a partial function from  $X$  to  $\mathbb{Z}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a natural-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be a natural number. Then  $f + c$  is a partial function from  $X$  to  $\mathbb{N}\text{-PFuncs DOMS}(Y)$ .

One can prove the following propositions:

$$(45) \quad f + c_1 + c_2 = f + (c_1 + c_2).$$

$$(46) \quad \text{If } f \neq \emptyset \text{ and } f \text{ is non-empty and } f + c_1 = f + c_2, \text{ then } c_1 = c_2.$$

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be a complex number. The functor  $f - c$  yields a function and is defined as follows:

(Def. 38)  $f - c = f + -c$ .

We now state two propositions:

$$(47) \quad \text{dom}(f - c) = \text{dom } f.$$

$$(48) \quad \text{If } x \in \text{dom}(f - c), \text{ then } (f - c)(x) = f(x) - c.$$

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be a complex number. Then  $f - c$  is a partial function from  $X$  to  $\mathbb{C}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a real-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be a real number. Then  $f - c$  is a partial function from  $X$  to  $\mathbb{R}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a rational-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be a rational number. Then  $f - c$  is a partial function from  $X$  to  $\mathbb{Q}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be an integer-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be an integer number. Then  $f - c$  is a partial function from  $X$  to  $\mathbb{Z}\text{-PFuncs DOMS}(Y)$ .

We now state four propositions:

(49) If  $f \neq \emptyset$  and  $f$  is non-empty and  $f - c_1 = f - c_2$ , then  $c_1 = c_2$ .

(50)  $(f + c_1) - c_2 = f + (c_1 - c_2)$ .

(51)  $(f - c_1) + c_2 = f - (c_1 - c_2)$ .

(52)  $f - c_1 - c_2 = f - (c_1 + c_2)$ .

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be a complex number. The functor  $f \cdot c$  yielding a function is defined as follows:

(Def. 39)  $\text{dom}(f \cdot c) = \text{dom } f$  and for every set  $x$  such that  $x \in \text{dom}(f \cdot c)$  holds  $(f \cdot c)(x) = c f(x)$ .

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be a complex number. Then  $f \cdot c$  is a partial function from  $X$  to  $\mathbb{C}\text{-PFuns DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a real-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be a real number. Then  $f \cdot c$  is a partial function from  $X$  to  $\mathbb{R}\text{-PFuns DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a rational-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be a rational number. Then  $f \cdot c$  is a partial function from  $X$  to  $\mathbb{Q}\text{-PFuns DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be an integer-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be an integer number. Then  $f \cdot c$  is a partial function from  $X$  to  $\mathbb{Z}\text{-PFuns DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a natural-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be a natural number. Then  $f \cdot c$  is a partial function from  $X$  to  $\mathbb{N}\text{-PFuns DOMS}(Y)$ .

The following two propositions are true:

(53)  $f \cdot c_1 \cdot c_2 = f \cdot (c_1 \cdot c_2)$ .

(54) If  $f \neq \emptyset$  and  $f$  is non-empty and for every  $x$  such that  $x \in \text{dom } f$  holds  $f(x)$  is non-empty and  $f \cdot c_1 = f \cdot c_2$ , then  $c_1 = c_2$ .

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be a complex number. The functor  $f/c$  yielding a function is defined as follows:

(Def. 40)  $f/c = f \cdot c^{-1}$ .

One can prove the following propositions:

(55)  $\text{dom}(f/c) = \text{dom } f$ .

(56) If  $x \in \text{dom}(f/c)$ , then  $(f/c)(x) = c^{-1} f(x)$ .

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be a complex number. Then  $f/c$  is a partial function from  $X$  to  $\mathbb{C}\text{-PFuns DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a real-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be a real number. Then  $f/c$  is a partial function from  $X$  to  $\mathbb{R}\text{-PFunCS DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a rational-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $c$  be a rational number. Then  $f/c$  is a partial function from  $X$  to  $\mathbb{Q}\text{-PFunCS DOMS}(Y)$ .

The following propositions are true:

$$(57) \quad f/c_1/c_2 = f/(c_1 \cdot c_2).$$

$$(58) \quad \text{If } f \neq \emptyset \text{ and } f \text{ is non-empty and for every } x \text{ such that } x \in \text{dom } f \text{ holds } f(x) \text{ is non-empty and } f/c_1 = f/c_2, \text{ then } c_1 = c_2.$$

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be a complex-valued function. The functor  $f + g$  yielding a function is defined as follows:

$$(\text{Def. 41}) \quad \text{dom}(f+g) = \text{dom } f \cap \text{dom } g \text{ and for every set } x \text{ such that } x \in \text{dom}(f+g) \text{ holds } (f+g)(x) = f(x) + g(x).$$

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be a complex-valued function. Then  $f + g$  is a partial function from  $X \cap \text{dom } g$  to  $\mathbb{C}\text{-PFunCS DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a real-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be a real-valued function. Then  $f + g$  is a partial function from  $X \cap \text{dom } g$  to  $\mathbb{R}\text{-PFunCS DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a rational-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be a rational-valued function. Then  $f + g$  is a partial function from  $X \cap \text{dom } g$  to  $\mathbb{Q}\text{-PFunCS DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be an integer-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be an integer-valued function. Then  $f + g$  is a partial function from  $X \cap \text{dom } g$  to  $\mathbb{Z}\text{-PFunCS DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a natural-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be a natural-valued function. Then  $f + g$  is a partial function from  $X \cap \text{dom } g$  to  $\mathbb{N}\text{-PFunCS DOMS}(Y)$ .

Next we state two propositions:

$$(59) \quad f + g + h = f + (g + h).$$

$$(60) \quad -(f + g) = (-f) + -g.$$

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be a complex-valued function. The functor  $f - g$  yields a function and is defined by:

$$(\text{Def. 42}) \quad f - g = f + -g.$$

We now state two propositions:

$$(61) \quad \text{dom}(f - g) = \text{dom } f \cap \text{dom } g.$$

$$(62) \quad \text{If } x \in \text{dom}(f - g), \text{ then } (f - g)(x) = f(x) - g(x).$$

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be a complex-valued function. Then  $f - g$  is a partial function from  $X \cap \text{dom } g$  to  $\mathbb{C}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a real-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be a real-valued function. Then  $f - g$  is a partial function from  $X \cap \text{dom } g$  to  $\mathbb{R}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a rational-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be a rational-valued function. Then  $f - g$  is a partial function from  $X \cap \text{dom } g$  to  $\mathbb{Q}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be an integer-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be an integer-valued function. Then  $f - g$  is a partial function from  $X \cap \text{dom } g$  to  $\mathbb{Z}\text{-PFuncs DOMS}(Y)$ .

The following propositions are true:

$$(63) \quad f - -g = f + g.$$

$$(64) \quad -(f - g) = (-f) + g.$$

$$(65) \quad (f + g) - h = f + (g - h).$$

$$(66) \quad (f - g) + h = f - (g - h).$$

$$(67) \quad f - g - h = f - (g + h).$$

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be a complex-valued function. The functor  $f \cdot g$  yielding a function is defined by:

(Def. 43)  $\text{dom}(f \cdot g) = \text{dom } f \cap \text{dom } g$  and for every set  $x$  such that  $x \in \text{dom}(f \cdot g)$  holds  $(f \cdot g)(x) = f(x) g(x)$ .

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be a complex-valued function. Then  $f \cdot g$  is a partial function from  $X \cap \text{dom } g$  to  $\mathbb{C}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a real-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be a real-valued function. Then  $f \cdot g$  is a partial function from  $X \cap \text{dom } g$  to  $\mathbb{R}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a rational-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be a rational-valued function. Then  $f \cdot g$  is a partial function from  $X \cap \text{dom } g$  to  $\mathbb{Q}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be an integer-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be an integer-valued function. Then  $f \cdot g$  is a partial function from  $X \cap \text{dom } g$  to  $\mathbb{Z}\text{-PFuncs DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a natural-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be a natural-valued function. Then  $f \cdot g$  is a partial function from  $X \cap \text{dom } g$  to  $\mathbb{N}\text{-PFuncs DOMS}(Y)$ .

Next we state three propositions:

$$(68) \quad f \cdot -g = (-f) \cdot g.$$

$$(69) \quad f \cdot -g = -f \cdot g.$$

$$(70) \quad f \cdot g \cdot h = f \cdot (gh).$$

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be a complex-valued function. The functor  $f/g$  yields a function and is defined by:

$$(\text{Def. 44}) \quad f/g = f \cdot g^{-1}.$$

Next we state two propositions:

$$(71) \quad \text{dom}(f/g) = \text{dom } f \cap \text{dom } g.$$

$$(72) \quad \text{If } x \in \text{dom}(f/g), \text{ then } (f/g)(x) = f(x)/g(x).$$

Let us consider  $X$ , let  $Y$  be a complex-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be a complex-valued function. Then  $f/g$  is a partial function from  $X \cap \text{dom } g$  to  $\mathbb{C}\text{-PFuns DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a real-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be a real-valued function. Then  $f/g$  is a partial function from  $X \cap \text{dom } g$  to  $\mathbb{R}\text{-PFuns DOMS}(Y)$ .

Let us consider  $X$ , let  $Y$  be a rational-functions-membered set, let  $f$  be a partial function from  $X$  to  $Y$ , and let  $g$  be a rational-valued function. Then  $f/g$  is a partial function from  $X \cap \text{dom } g$  to  $\mathbb{Q}\text{-PFuns DOMS}(Y)$ .

Next we state the proposition

$$(73) \quad (f \cdot g)/h = f \cdot (g/h).$$

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be complex-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ . The functor  $f + g$  yielding a function is defined as follows:

$$(\text{Def. 45}) \quad \text{dom}(f+g) = \text{dom } f \cap \text{dom } g \text{ and for every set } x \text{ such that } x \in \text{dom}(f+g) \\ \text{holds } (f+g)(x) = f(x) + g(x).$$

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be complex-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ . Then  $f + g$  is a partial function from  $X_1 \cap X_2$  to  $\mathbb{C}\text{-PFuns}(\text{DOMS}(Y_1) \cap \text{DOMS}(Y_2))$ .

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be real-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ . Then  $f + g$  is a partial function from  $X_1 \cap X_2$  to  $\mathbb{R}\text{-PFuns}(\text{DOMS}(Y_1) \cap \text{DOMS}(Y_2))$ .

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be rational-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ . Then  $f + g$  is a partial function from  $X_1 \cap X_2$  to  $\mathbb{Q}\text{-PFuns}(\text{DOMS}(Y_1) \cap \text{DOMS}(Y_2))$ .

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be integer-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to

$Y_2$ . Then  $f + g$  is a partial function from  $X_1 \cap X_2$  to  $\mathbb{Z}\text{-PFuncs}(\text{DOMS}(Y_1) \cap \text{DOMS}(Y_2))$ .

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be natural-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ . Then  $f + g$  is a partial function from  $X_1 \cap X_2$  to  $\mathbb{N}\text{-PFuncs}(\text{DOMS}(Y_1) \cap \text{DOMS}(Y_2))$ .

We now state three propositions:

$$(74) \quad f_1 + f_2 = f_2 + f_1.$$

$$(75) \quad (f + f_1) + f_2 = f + (f_1 + f_2).$$

$$(76) \quad -(f_1 + f_2) = (-f_1) + -f_2.$$

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be complex-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ . The functor  $f - g$  yields a function and is defined by:

(Def. 46)  $\text{dom}(f - g) = \text{dom } f \cap \text{dom } g$  and for every set  $x$  such that  $x \in \text{dom}(f - g)$  holds  $(f - g)(x) = f(x) - g(x)$ .

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be complex-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ . Then  $f - g$  is a partial function from  $X_1 \cap X_2$  to  $\mathbb{C}\text{-PFuncs}(\text{DOMS}(Y_1) \cap \text{DOMS}(Y_2))$ .

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be real-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ . Then  $f - g$  is a partial function from  $X_1 \cap X_2$  to  $\mathbb{R}\text{-PFuncs}(\text{DOMS}(Y_1) \cap \text{DOMS}(Y_2))$ .

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be rational-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ . Then  $f - g$  is a partial function from  $X_1 \cap X_2$  to  $\mathbb{Q}\text{-PFuncs}(\text{DOMS}(Y_1) \cap \text{DOMS}(Y_2))$ .

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be integer-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ . Then  $f - g$  is a partial function from  $X_1 \cap X_2$  to  $\mathbb{Z}\text{-PFuncs}(\text{DOMS}(Y_1) \cap \text{DOMS}(Y_2))$ .

One can prove the following propositions:

$$(77) \quad f_1 - f_2 = -(f_2 - f_1).$$

$$(78) \quad -(f_1 - f_2) = (-f_1) + f_2.$$

$$(79) \quad (f + f_1) - f_2 = f + (f_1 - f_2).$$

$$(80) \quad (f - f_1) + f_2 = f - (f_1 - f_2).$$

$$(81) \quad f - f_1 - f_2 = f - (f_1 + f_2).$$

$$(82) \quad f - f_1 - f_2 = f - f_2 - f_1.$$

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be complex-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ .



The functor  $f \cdot g$  yields a function and is defined by:

(Def. 47)  $\text{dom}(f \cdot g) = \text{dom } f \cap \text{dom } g$  and for every set  $x$  such that  $x \in \text{dom}(f \cdot g)$  holds  $(f \cdot g)(x) = f(x) g(x)$ .

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be complex-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ . Then  $f \cdot g$  is a partial function from  $X_1 \cap X_2$  to  $\mathbb{C}\text{-PFuncs}(\text{DOMS}(Y_1) \cap \text{DOMS}(Y_2))$ .

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be real-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ . Then  $f \cdot g$  is a partial function from  $X_1 \cap X_2$  to  $\mathbb{R}\text{-PFuncs}(\text{DOMS}(Y_1) \cap \text{DOMS}(Y_2))$ .

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be rational-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ . Then  $f \cdot g$  is a partial function from  $X_1 \cap X_2$  to  $\mathbb{Q}\text{-PFuncs}(\text{DOMS}(Y_1) \cap \text{DOMS}(Y_2))$ .

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be integer-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ . Then  $f \cdot g$  is a partial function from  $X_1 \cap X_2$  to  $\mathbb{Z}\text{-PFuncs}(\text{DOMS}(Y_1) \cap \text{DOMS}(Y_2))$ .

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be natural-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ . Then  $f \cdot g$  is a partial function from  $X_1 \cap X_2$  to  $\mathbb{N}\text{-PFuncs}(\text{DOMS}(Y_1) \cap \text{DOMS}(Y_2))$ .

We now state several propositions:

$$(83) \quad f_1 \cdot f_2 = f_2 \cdot f_1.$$

$$(84) \quad (f \cdot f_1) \cdot f_2 = f \cdot (f_1 \cdot f_2).$$

$$(85) \quad (-f_1) \cdot f_2 = -f_1 \cdot f_2.$$

$$(86) \quad f_1 \cdot -f_2 = -f_1 \cdot f_2.$$

$$(87) \quad f \cdot (f_1 + f_2) = f \cdot f_1 + f \cdot f_2.$$

$$(88) \quad (f_1 + f_2) \cdot f = f_1 \cdot f + f_2 \cdot f.$$

$$(89) \quad f \cdot (f_1 - f_2) = f \cdot f_1 - f \cdot f_2.$$

$$(90) \quad (f_1 - f_2) \cdot f = f_1 \cdot f - f_2 \cdot f.$$

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be complex-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ . The functor  $f/g$  yields a function and is defined by:

(Def. 48)  $\text{dom}(f/g) = \text{dom } f \cap \text{dom } g$  and for every set  $x$  such that  $x \in \text{dom}(f/g)$  holds  $(f/g)(x) = f(x)/g(x)$ .

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be complex-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$

to  $Y_2$ . Then  $f/g$  is a partial function from  $X_1 \cap X_2$  to  $\mathbb{C}\text{-PFuncs}(\text{DOMS}(Y_1) \cap \text{DOMS}(Y_2))$ .

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be real-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ . Then  $f/g$  is a partial function from  $X_1 \cap X_2$  to  $\mathbb{R}\text{-PFuncs}(\text{DOMS}(Y_1) \cap \text{DOMS}(Y_2))$ .

Let  $X_1, X_2$  be sets, let  $Y_1, Y_2$  be rational-functions-membered sets, let  $f$  be a partial function from  $X_1$  to  $Y_1$ , and let  $g$  be a partial function from  $X_2$  to  $Y_2$ . Then  $f/g$  is a partial function from  $X_1 \cap X_2$  to  $\mathbb{Q}\text{-PFuncs}(\text{DOMS}(Y_1) \cap \text{DOMS}(Y_2))$ .

One can prove the following propositions:

- (91)  $(-f_1)/f_2 = -f_1/f_2$ .
- (92)  $f_1/-f_2 = -f_1/f_2$ .
- (93)  $(f \cdot f_1)/f_2 = f \cdot (f_1/f_2)$ .
- (94)  $(f/f_1) \cdot f_2 = (f \cdot f_2)/f_1$ .
- (95)  $f/f_1/f_2 = f/(f_1 \cdot f_2)$ .
- (96)  $(f_1 + f_2)/f = f_1/f + f_2/f$ .
- (97)  $(f_1 - f_2)/f = f_1/f - f_2/f$ .

#### REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [2] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [3] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [4] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [5] Andrzej Kondracki. Basic properties of rational numbers. *Formalized Mathematics*, 1(5):841–845, 1990.
- [6] Andrzej Trybulec. Tuples, projections and Cartesian products. *Formalized Mathematics*, 1(1):97–105, 1990.
- [7] Andrzej Trybulec. On the sets inhabited by numbers. *Formalized Mathematics*, 11(4):341–347, 2003.
- [8] Michał J. Trybulec. Integers. *Formalized Mathematics*, 1(3):501–505, 1990.
- [9] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [10] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [11] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.

Received October 15, 2008

---