

The Definition of Topological Manifolds

Marco Riccardi
 Via del Pero 102
 54038 Montignoso, Italy

Summary. This article introduces the definition of n -locally Euclidean topological spaces and topological manifolds [13].

MML identifier: MFOLD_1, version: 7.11.07 4.156.1112

The papers [8], [1], [6], [15], [7], [18], [3], [4], [17], [2], [16], [9], [19], [20], [11], [12], [10], [14], and [5] provide the terminology and notation for this paper.

1. PRELIMINARIES

Let x, y be sets. Observe that $\{\langle x, y \rangle\}$ is one-to-one.

In the sequel n denotes a natural number.

One can prove the following two propositions:

- (1) For every non empty topological space T holds T and $T|\Omega_T$ are homeomorphic.
- (2) Let X be a non empty subspace of \mathcal{E}_T^n and f be a function from X into \mathbb{R}^1 . Suppose f is continuous. Then there exists a function g from X into \mathcal{E}_T^n such that
 - (i) for every point a of X and for every point b of \mathcal{E}_T^n and for every real number r such that $a = b$ and $f(a) = r$ holds $g(b) = r \cdot b$, and
 - (ii) g is continuous.

Let us consider n and let S be a subset of \mathcal{E}_T^n . We say that S is ball if and only if:

- (Def. 1) There exists a point p of \mathcal{E}_T^n and there exists a real number r such that $S = \text{Ball}(p, r)$.

Let us consider n . Observe that there exists a subset of \mathcal{E}_T^n which is ball and every subset of \mathcal{E}_T^n which is ball is also open.

Let us consider n . One can verify that there exists a subset of \mathcal{E}_T^n which is non empty and ball.

In the sequel p denotes a point of \mathcal{E}_T^n and r denotes a real number.

The following proposition is true

- (3) For every open subset S of \mathcal{E}_T^n such that $p \in S$ there exists ball subset B of \mathcal{E}_T^n such that $B \subseteq S$ and $p \in B$.

Let us consider n, p, r . The functor $\mathbb{B}_r(p)$ yields a subspace of \mathcal{E}_T^n and is defined as follows:

(Def. 2) $\mathbb{B}_r(p) = \mathcal{E}_T^n \upharpoonright \text{Ball}(p, r)$.

Let us consider n . The functor \mathbb{B}^n yields a subspace of \mathcal{E}_T^n and is defined as follows:

(Def. 3) $\mathbb{B}^n = \mathbb{B}_1(0_{\mathcal{E}_T^n})$.

Let us consider n . One can verify that \mathbb{B}^n is non empty. Let us consider p and let s be a positive real number. Observe that $\mathbb{B}_s(p)$ is non empty.

The following propositions are true:

- (4) The carrier of $\mathbb{B}_r(p) = \text{Ball}(p, r)$.
- (5) If $n \neq 0$ and p is a point of \mathbb{B}^n , then $|p| < 1$.
- (6) Let f be a function from \mathbb{B}^n into \mathcal{E}_T^n . Suppose $n \neq 0$ and for every point a of \mathbb{B}^n and for every point b of \mathcal{E}_T^n such that $a = b$ holds $f(a) = \frac{1}{1-|b| \cdot |b|} \cdot b$. Then f is homeomorphism.
- (7) Let r be a positive real number and f be a function from \mathbb{B}^n into $\mathbb{B}_r(p)$. Suppose $n \neq 0$ and for every point a of \mathbb{B}^n and for every point b of \mathcal{E}_T^n such that $a = b$ holds $f(a) = r \cdot b + p$. Then f is homeomorphism.
- (8) \mathbb{B}^n and \mathcal{E}_T^n are homeomorphic.

In the sequel q denotes a point of \mathcal{E}_T^n .

We now state three propositions:

- (9) For all positive real numbers r, s holds $\mathbb{B}_r(p)$ and $\mathbb{B}_s(q)$ are homeomorphic.
- (10) For every non empty ball subset B of \mathcal{E}_T^n holds B and $\Omega_{\mathcal{E}_T^n}$ are homeomorphic.
- (11) Let M, N be non empty topological spaces, p be a point of M , U be a neighbourhood of p , and B be an open subset of N . Suppose U and B are homeomorphic. Then there exists an open subset V of M and there exists an open subset S of N such that $V \subseteq U$ and $p \in V$ and V and S are homeomorphic.

2. MANIFOLD

In the sequel M is a non empty topological space.

Let us consider n , M . We say that M is n -locally Euclidean if and only if the condition (Def. 4) is satisfied.

(Def. 4) Let p be a point of M . Then there exists a neighbourhood U of p and there exists an open subset S of \mathcal{E}_T^n such that U and S are homeomorphic.

Let us consider n . Observe that \mathcal{E}_T^n is n -locally Euclidean.

Let us consider n . Observe that there exists a non empty topological space which is n -locally Euclidean.

We now state two propositions:

(12) M is n -locally Euclidean if and only if for every point p of M there exists a neighbourhood U of p and there exists ball subset B of \mathcal{E}_T^n such that U and B are homeomorphic.

(13) M is n -locally Euclidean if and only if for every point p of M there exists a neighbourhood U of p such that U and $\Omega_{\mathcal{E}_T^n}$ are homeomorphic.

Let us consider n . Observe that every non empty topological space which is n -locally Euclidean is also first-countable.

Let us note that every non empty topological space which is 0-locally Euclidean is also discrete and every non empty topological space which is discrete is also 0-locally Euclidean.

Let us consider n . One can verify that \mathcal{E}_T^n is second-countable.

Let us consider n . Note that there exists a non empty topological space which is second-countable, Hausdorff, and n -locally Euclidean.

Let us consider n , M . We say that M is n -manifold if and only if:

(Def. 5) M is second-countable, Hausdorff, and n -locally Euclidean.

Let us consider M . We say that M is manifold-like if and only if:

(Def. 6) There exists n such that M is n -manifold.

Let us consider n . Observe that there exists a non empty topological space which is n -manifold.

Let us consider n . One can check the following observations:

- * every non empty topological space which is n -manifold is also second-countable, Hausdorff, and n -locally Euclidean,
- * every non empty topological space which is second-countable, Hausdorff, and n -locally Euclidean is also n -manifold, and
- * every non empty topological space which is n -manifold is also manifold-like.

Let us note that every non empty topological space which is second-countable and discrete is also 0-manifold.

Let us consider n and let M be an n -manifold non empty topological space. One can verify that every non empty subspace of M which is open is also n -manifold.

Let us note that there exists a non empty topological space which is manifold-like.

A manifold is a manifold-like non empty topological space.

REFERENCES

- [1] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(2):377–382, 1990.
- [2] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [3] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [4] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [5] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [6] Agata Darmochwał. Compact spaces. *Formalized Mathematics*, 1(2):383–386, 1990.
- [7] Agata Darmochwał. The Euclidean space. *Formalized Mathematics*, 2(4):599–603, 1991.
- [8] Adam Grabowski. Properties of the product of compact topological spaces. *Formalized Mathematics*, 8(1):55–59, 1999.
- [9] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [10] Zbigniew Karno. Separated and weakly separated subspaces of topological spaces. *Formalized Mathematics*, 2(5):665–674, 1991.
- [11] Zbigniew Karno. The lattice of domains of an extremally disconnected space. *Formalized Mathematics*, 3(2):143–149, 1992.
- [12] Artur Kornilowicz and Yasunari Shidama. Intersections of intervals and balls in \mathcal{E}_T^n . *Formalized Mathematics*, 12(3):301–306, 2004.
- [13] John M. Lee. *Introduction to Topological Manifolds*. Springer-Verlag, New York Berlin Heidelberg, 2000.
- [14] Robert Milewski. Bases of continuous lattices. *Formalized Mathematics*, 7(2):285–294, 1998.
- [15] Beata Padlewska. Locally connected spaces. *Formalized Mathematics*, 2(1):93–96, 1991.
- [16] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(1):223–230, 1990.
- [17] Karol Pąk. Basic properties of metrizable topological spaces. *Formalized Mathematics*, 17(3):201–205, 2009, doi: 10.2478/v10037-009-0024-8.
- [18] Bartłomiej Skorulski. First-countable, sequential, and Frechet spaces. *Formalized Mathematics*, 7(1):81–86, 1998.
- [19] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(2):291–296, 1990.
- [20] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.

Received August 17, 2010
