Linear Transformations of Euclidean Topological Spaces

Karol Pąk Institute of Informatics University of Białystok Poland

Summary. We introduce linear transformations of Euclidean topological spaces given by a transformation matrix. Next, we prove selected properties and basic arithmetic operations on these linear transformations. Finally, we show that a linear transformation given by an invertible matrix is a homeomorphism.

MML identifier: MATRTOP1, version: 7.11.07 4.156.1112

The papers [2], [12], [6], [26], [7], [8], [30], [21], [22], [23], [15], [31], [29], [19], [24], [3], [4], [9], [16], [5], [20], [18], [1], [14], [28], [13], [10], [25], [27], [11], and [17] provide the notation and terminology for this paper.

1. Preliminaries

For simplicity, we adopt the following rules: X, Y denote sets, n, m, k, i denote natural numbers, r denotes a real number, R denotes an element of \mathbb{R}_{F} , K denotes a field, f, f_1, f_2, g_1, g_2 denote finite sequences, r_1, r_2, r_3 denote real-valued finite sequences, c_1, c_2 denote complex-valued finite sequences, and F denotes a function.

Let us consider X, Y and let F be a positive yielding partial function from X to \mathbb{R} . One can check that $F \upharpoonright Y$ is positive yielding.

Let us consider X, Y and let F be a negative yielding partial function from X to \mathbb{R} . One can verify that $F \upharpoonright Y$ is negative yielding.

Let us consider X, Y and let F be a non-positive yielding partial function from X to \mathbb{R} . Note that $F \upharpoonright Y$ is non-positive yielding.

C 2011 University of Białystok ISSN 1426-2630(p), 1898-9934(e) Let us consider X, Y and let F be a non-negative yielding partial function from X to \mathbb{R} . Note that $F \upharpoonright Y$ is non-negative yielding.

Let us consider r_1 . One can check that $\sqrt{r_1}$ is finite sequence-like.

Let us consider r_1 . The functor [@] r_1 yielding a finite sequence of elements of \mathbb{R}_F is defined by:

(Def. 1) $^{@}r_1 = r_1.$

Let p be a finite sequence of elements of \mathbb{R}_{F} . The functor [@]p yields a finite sequence of elements of \mathbb{R} and is defined as follows:

(Def. 2) $^{@}p = p$.

We now state several propositions:

(1) $({}^{@}r_2) + {}^{@}r_3 = r_2 + r_3.$

(2)
$$\sqrt{r_2} \cap r_3 = \sqrt{r_2} \cap \sqrt{r_3}.$$

- (3) $\sqrt{\langle r \rangle} = \langle \sqrt{r} \rangle.$
- (4) $\sqrt{r_1^2} = |r_1|.$
- (5) If r_1 is non-negative yielding, then $\sqrt{\sum r_1} \leq \sum \sqrt{r_1}$.
- (6) There exists X such that $X \subseteq \text{dom } F$ and $\text{rng } F = \text{rng}(F \upharpoonright X)$ and $F \upharpoonright X$ is one-to-one.

Let us consider c_1 , X. Observe that $c_1 - X$ is complex-valued.

Let us consider r_1 , X. Observe that $r_1 - X$ is real-valued.

Let c_1 be a complex-valued finite subsequence. Note that Seq c_1 is complex-valued.

Let r_1 be a real-valued finite subsequence. Observe that Seq r_1 is real-valued. One can prove the following propositions:

- (7) For every permutation P of dom f such that $f_1 = f \cdot P$ there exists a permutation Q of dom(f X) such that $f_1 X = (f X) \cdot Q$.
- (8) For every permutation P of dom c_1 such that $c_2 = c_1 \cdot P$ holds $\sum (c_2 X) = \sum (c_1 X)$.
- (9) Let f, f_1 be finite subsequences and P be a permutation of dom f. If $f_1 = f \cdot P$, then there exists a permutation Q of dom Seq $(f_1 \upharpoonright P^{-1}(X))$ such that Seq $(f \upharpoonright X) =$ Seq $(f_1 \upharpoonright P^{-1}(X)) \cdot Q$.
- (10) Let c_1, c_2 be complex-valued finite subsequences and P be a permutation of dom c_1 . If $c_2 = c_1 \cdot P$, then $\sum \text{Seq}(c_1 \upharpoonright X) = \sum \text{Seq}(c_2 \upharpoonright P^{-1}(X))$.
- (11) Let f be a finite subsequence and n be an element of \mathbb{N} . If for every i holds $i + n \in X$ iff $i \in Y$, then $\operatorname{Shift}^n f \upharpoonright X = \operatorname{Shift}^n(f \upharpoonright Y)$.
- (12) There exists a subset Y of N such that $\text{Seq}((f_1 \cap f_2) \upharpoonright X) = (\text{Seq}(f_1 \upharpoonright X)) \cap \text{Seq}(f_2 \upharpoonright Y)$ and for every n such that n > 0 holds $n \in Y$ iff $n + \text{len } f_1 \in X \cap \text{dom}(f_1 \cap f_2)$.
- (13) If len $g_1 = \text{len } f_1$ and len $g_2 \leq \text{len } f_2$, then $\text{Seq}((f_1 \cap f_2) \upharpoonright (g_1 \cap g_2)^{-1}(X)) = (\text{Seq}(f_1 \upharpoonright g_1^{-1}(X))) \cap \text{Seq}(f_2 \upharpoonright g_2^{-1}(X)).$

- (14) Let D be a non empty set and M be a matrix over D of dimension $n \times m$. Then M X is a matrix over D of dimension $n \frac{1}{M^{-1}(X)} \times m$.
- (15) Let F be a function from Seg n into Seg n, D be a non empty set, M be a matrix over D of dimension $n \times m$, and given i. If $i \in$ Seg width M, then $(M \cdot F)_{\Box,i} = M_{\Box,i} \cdot F$.
- (16) Let A be a matrix over K of dimension $n \times m$. Suppose $\operatorname{rk}(A) = n$. Then there exists a matrix B over K of dimension $m - n \times m$ such that $\operatorname{rk}(A \cap B) = m$.
- (17) Let A be a matrix over K of dimension $n \times m$. Suppose $\operatorname{rk}(A) = m$. Then there exists a matrix B over K of dimension $n \times n - m'$ such that $\operatorname{rk}(A \cap B) = n$.

For simplicity, we adopt the following convention: f, f_1 , f_2 denote n-element real-valued finite sequences, p, p_1 , p_2 denote points of $\mathcal{E}^n_{\mathrm{T}}$, M, M_1 , M_2 denote matrices over \mathbb{R}_{F} of dimension $n \times m$, and A, B denote square matrices over \mathbb{R}_{F} of dimension n.

2. LINEAR TRANSFORMATIONS OF EUCLIDEAN TOPOLOGICAL SPACES GIVEN BY A TRANSFORMATION MATRIX

Let us consider n, m, M. The functor Mx2Tran M yielding a function from $\mathcal{E}^n_{\mathrm{T}}$ into $\mathcal{E}^m_{\mathrm{T}}$ is defined by:

(Def. 3)(i) $(Mx2Tran M)(f) = Line(LineVec2Mx(^{@}f) \cdot M, 1)$ if $n \neq 0$,

(ii) $(Mx2Tran M)(f) = 0_{\mathcal{E}_T^m}$, otherwise.

Let us consider n, m, M and let x be a set. One can check that (Mx2Tran M)(x) is function-like and relation-like and (Mx2Tran M)(x) is real-valued and finite sequence-like.

Let us consider n, m, M, f. One can check that (Mx2Tran M)(f) is *m*-element.

One can prove the following propositions:

(18) If $1 \le i \le m$ and $n \ne 0$, then $(Mx2Tran M)(f)(i) = (^{@}f) \cdot M_{\Box i}$.

- (19) len MX2FinS $(I_K^{n \times n}) = n$.
- (20) Let B_1 be an ordered basis of the *n*-dimension vector space over \mathbb{R}_F and B_2 be an ordered basis of the *m*-dimension vector space over \mathbb{R}_F . Suppose $B_1 = \text{MX2FinS}(I_{\mathbb{R}_F}^{n \times n})$ and $B_2 = \text{MX2FinS}(I_{\mathbb{R}_F}^{m \times m})$. Let M_1 be a matrix over \mathbb{R}_F of dimension len $B_1 \times \text{len } B_2$. If $M_1 = M$, then Mx2Tran $M = \text{Mx2Tran}(M_1, B_1, B_2)$.
- (21) For every permutation P of $\operatorname{Seg} n$ holds $(\operatorname{Mx2Tran} M)(f) = (\operatorname{Mx2Tran}(M \cdot P))(f \cdot P)$ and $f \cdot P$ is an *n*-element finite sequence of elements of \mathbb{R} .
- (22) $(Mx2Tran M)(f_1 + f_2) = (Mx2Tran M)(f_1) + (Mx2Tran M)(f_2).$

- (23) $(Mx2Tran M)(r \cdot f) = r \cdot (Mx2Tran M)(f).$
- (24) $(Mx2Tran M)(f_1 f_2) = (Mx2Tran M)(f_1) (Mx2Tran M)(f_2).$
- (25) $(Mx2Tran(M_1 + M_2))(f) = (Mx2Tran M_1)(f) + (Mx2Tran M_2)(f).$
- (26) $(R) \cdot (Mx2Tran M)(f) = (Mx2Tran(R \cdot M))(f).$
- (27) $(Mx2Tran M)(p_1 + p_2) = (Mx2Tran M)(p_1) + (Mx2Tran M)(p_2).$
- (28) $(Mx2Tran M)(p_1 p_2) = (Mx2Tran M)(p_1) (Mx2Tran M)(p_2).$
- (29) $(\operatorname{Mx2Tran} M)(0_{\mathcal{E}_{T}^{n}}) = 0_{\mathcal{E}_{T}^{m}}.$
- (30) For every subset A of $\mathcal{E}_{\mathrm{T}}^{n}$ holds $(\mathrm{Mx}2\mathrm{Tran}\,M)^{\circ}(p + A) = (\mathrm{Mx}2\mathrm{Tran}\,M)(p) + (\mathrm{Mx}2\mathrm{Tran}\,M)^{\circ}A.$
- (31) For every subset A of $\mathcal{E}_{\mathrm{T}}^{m}$ holds $(\mathrm{Mx}2\mathrm{Tran}\,M)^{-1}((\mathrm{Mx}2\mathrm{Tran}\,M)(p)+A) = p + (\mathrm{Mx}2\mathrm{Tran}\,M)^{-1}(A).$
- (32) Let A be a matrix over \mathbb{R}_{F} of dimension $n \times m$ and B be a matrix over \mathbb{R}_{F} of dimension width $A \times k$. If if n = 0, then m = 0 and if m = 0, then k = 0, then Mx2Tran $B \cdot \mathrm{Mx2Tran} A = \mathrm{Mx2Tran}(A \cdot B)$.
- (33) Mx2Tran $(I_{\mathbb{R}_{\mathrm{F}}}^{n \times n}) = \mathrm{id}_{\mathcal{E}_{\mathrm{T}}^{n}}.$
- (34) If Mx2Tran $M_1 = Mx2Tran M_2$, then $M_1 = M_2$.
- (35) Let A be a matrix over \mathbb{R}_{F} of dimension $n \times m$ and B be a matrix over \mathbb{R}_{F} of dimension $k \times m$. Then $(\mathrm{Mx2Tran}(A \cap B))(f \cap (k \mapsto 0)) =$ $(\mathrm{Mx2Tran} A)(f)$ and $(\mathrm{Mx2Tran}(B \cap A))((k \mapsto 0) \cap f) = (\mathrm{Mx2Tran} A)(f)$.
- (36) Let A be a matrix over \mathbb{R}_{F} of dimension $n \times m$, B be a matrix over \mathbb{R}_{F} of dimension $k \times m$, and g be a k-element real-valued finite sequence. Then $(\mathrm{Mx}2\mathrm{Tran}(A \cap B))(f \cap g) = (\mathrm{Mx}2\mathrm{Tran} A)(f) + (\mathrm{Mx}2\mathrm{Tran} B)(g)$.
- (37) Let A be a matrix over \mathbb{R}_{F} of dimension $n \times k$ and B be a matrix over \mathbb{R}_{F} of dimension $n \times m$ such that if n = 0, then k + m = 0. Then $(\mathrm{Mx2Tran}(A \cap B))(f) = (\mathrm{Mx2Tran} A)(f) \cap (\mathrm{Mx2Tran} B)(f)$.
- (38) $(Mx2Tran(I_{\mathbb{R}_{F}}^{m \times m} \restriction n))(f) \restriction n = f.$

3. Selected Properties of the Mx2Tran Operator

Next we state several propositions:

- (39) Mx2Tran M is one-to-one iff rk(M) = n.
- (40) Mx2Tran A is one-to-one iff $\operatorname{Det} A \neq 0_{\mathbb{R}_{\mathrm{F}}}$.
- (41) Mx2Tran M is onto iff rk(M) = m.
- (42) Mx2Tran A is onto iff Det $A \neq 0_{\mathbb{R}_{\mathrm{F}}}$.
- (43) For all A, B such that $\text{Det } A \neq 0_{\mathbb{R}_{\mathrm{F}}}$ holds $(\text{Mx}2\text{Tran } A)^{-1} = \text{Mx}2\text{Tran } B$ iff $A^{\sim} = B$.
- (44) There exists an *m*-element finite sequence *L* of elements of \mathbb{R} such that for every *i* such that $i \in \text{dom } L$ holds $L(i) = |^{@}(M_{\Box,i})|$ and for every *f* holds $|(\text{Mx2Tran } M)(f)| \leq \sum L \cdot |f|.$

- (45) There exists a real number L such that L > 0 and for every f holds $|(Mx2Tran M)(f)| \le L \cdot |f|.$
- (46) If $\operatorname{rk}(M) = n$, then there exists a real number L such that L > 0 and for every f holds $|f| \leq L \cdot |(\operatorname{Mx2Tran} M)(f)|$.
- (47) Mx2Tran M is continuous.

Let us consider n, K. One can check that there exists a square matrix over K of dimension n which is invertible.

Let us consider n and let A be an invertible square matrix over \mathbb{R}_{F} of dimension n. Note that Mx2Tran A is homeomorphism.

References

- Jesse Alama. The rank+nullity theorem. Formalized Mathematics, 15(3):137–142, 2007, doi:10.2478/v10037-007-0015-6.
- [2] Grzegorz Bancerek. Cardinal numbers. Formalized Mathematics, 1(2):377–382, 1990.
- [3] Grzegorz Bancerek. The fundamental properties of natural numbers. Formalized Mathematics, 1(1):41-46, 1990.
- [4] Grzegorz Bancerek. The ordinal numbers. Formalized Mathematics, 1(1):91–96, 1990.
- [5] Grzegorz Bancerek, Mitsuru Aoki, Akio Matsumoto, and Yasunari Shidama. Processes in Petri nets. Formalized Mathematics, 11(1):125–132, 2003.
- [6] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. Formalized Mathematics, 1(1):107–114, 1990.
- [7] Czesław Byliński. Functions and their basic properties. Formalized Mathematics, 1(1):55– 65, 1990.
- [8] Czesław Byliński. Functions from a set to a set. Formalized Mathematics, 1(1):153-164, 1990.
- [9] Czesław Byliński. Partial functions. Formalized Mathematics, 1(2):357–367, 1990.
- [10] Czesław Byliński. The sum and product of finite sequences of real numbers. Formalized Mathematics, 1(4):661–668, 1990.
- [11] Agata Darmochwał. Families of subsets, subspaces and mappings in topological spaces. *Formalized Mathematics*, 1(2):257-261, 1990.
- [12] Agata Darmochwał. The Euclidean space. Formalized Mathematics, 2(4):599–603, 1991.
- [13] Noboru Endou, Takashi Mitsuishi, and Yasunari Shidama. Dimension of real unitary space. Formalized Mathematics, 11(1):23–28, 2003.
- [14] Krzysztof Hryniewiecki. Basic properties of real numbers. Formalized Mathematics, 1(1):35-40, 1990.
- [15] Katarzyna Jankowska. Matrices. Abelian group of matrices. Formalized Mathematics, 2(4):475–480, 1991.
- [16] Artur Korniłowicz. On the real valued functions. Formalized Mathematics, 13(1):181–187, 2005.
- [17] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Formalized Mathematics*, 1(2):335–342, 1990.
- [18] Anna Lango and Grzegorz Bancerek. Product of families of groups and vector spaces. Formalized Mathematics, 3(2):235–240, 1992.
- [19] Robert Milewski. Associated matrix of linear map. Formalized Mathematics, 5(3):339– 345, 1996.
- [20] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. Formalized Mathematics, 1(1):223–230, 1990.
- [21] Karol Pak. Basic properties of determinants of square matrices over a field. Formalized Mathematics, 15(1):17–25, 2007, doi:10.2478/v10037-007-0003-x.
- [22] Karol Pąk. Basic properties of the rank of matrices over a field. Formalized Mathematics, 15(4):199–211, 2007, doi:10.2478/v10037-007-0024-5.
- [23] Karol Pąk. Solutions of linear equations. Formalized Mathematics, 16(1):81–90, 2008, doi:10.2478/v10037-008-0012-4.

KAROL PĄK

- [24] Karol Pak. Linear map of matrices. Formalized Mathematics, 16(3):269–275, 2008, doi:10.2478/v10037-008-0032-0.
- [25] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers. Formalized Mathematics, 1(3):445–449, 1990.
- [26] Wojciech A. Trybulec. Non-contiguous substrings and one-to-one finite sequences. Formalized Mathematics, 1(3):569–573, 1990.
- [27] Zinaida Trybulec. Properties of subsets. Formalized Mathematics, 1(1):67–71, 1990.
- [28] Edmund Woronowicz. Relations and their basic properties. Formalized Mathematics, 1(1):73–83, 1990.
- [29] Xiaopeng Yue, Xiquan Liang, and Zhongpin Sun. Some properties of some special matrices. Formalized Mathematics, 13(4):541–547, 2005.
- [30] Katarzyna Zawadzka. The sum and product of finite sequences of elements of a field. Formalized Mathematics, 3(2):205-211, 1992.
- [31] Katarzyna Zawadzka. The product and the determinant of matrices with entries in a field. Formalized Mathematics, 4(1):1–8, 1993.

Received October 26, 2010