

Leibniz Series for π^1

Karol Pąk
Institute of Informatics
University of Białystok
Ciołkowskiego 1M, 15-245 Białystok
Poland

Summary. In this article we prove the Leibniz series for π which states that

$$\frac{\pi}{4} = \sum_{n=0}^{\infty} \frac{(-1)^n}{2 \cdot n + 1}.$$

The formalization follows K. Knopp [8], [1] and [6]. *Leibniz's Series for Pi* is item #26 from the “Formalizing 100 Theorems” list maintained by Freek Wiedijk at <http://www.cs.ru.nl/F.Wiedijk/100/>.

MSC: 40G99 03B35

Keywords: π approximation; Leibniz theorem; Leibniz series

MML identifier: LEIBNIZ1, version: 8.1.05 5.39.1282

1. PRELIMINARIES

From now on i, n, m denote natural numbers, r, s denote real numbers, and A denotes a non empty, closed interval subset of \mathbb{R} .

Now we state the proposition:

$$(1) \quad \text{rng}((\text{the function } \tan)|] - \frac{\pi}{2}, \frac{\pi}{2}[) = \mathbb{R}.$$

PROOF: Set $P = \frac{\pi}{2}$. Set $I =] - P, P[$. $\mathbb{R} \subseteq \text{rng}((\text{the function } \tan)|I)$ by [4, (50)], [20, (30)], [14, (15)], [16, (1)]. \square

¹This work has been financed by the resources of the Polish National Science Centre granted by decision no. DEC-2012/07/N/ST6/02147.

One can verify that the function \arctan is total and the function \arctan is differentiable.

Now we state the propositions:

$$(2) \quad (\text{The function } \arctan)'(r) = \frac{1}{1+r^2}.$$

(3) Let us consider an open subset Z of \mathbb{R} . Then

(i) the function \arctan is differentiable on Z , and

(ii) for every r such that $r \in Z$ holds $(\text{the function } \arctan)'|_Z(r) = \frac{1}{1+r^2}$.

The theorem is a consequence of (2).

Let us consider n . One can verify that \square^n is continuous.

Now we state the propositions:

$$(4) \quad (i) \quad \text{dom}(\frac{\square^n}{\square^0 + \square^2}) = \mathbb{R}, \text{ and}$$

(ii) $\frac{\square^n}{\square^0 + \square^2}$ is continuous, and

$$(iii) \quad (\frac{\square^n}{\square^0 + \square^2})(r) = \frac{r^n}{1+r^2}.$$

$$(5) \quad \int_A (\frac{\square^0}{\square^0 + \square^2})(x) dx =$$

$(\text{the function } \arctan)(\sup A) - (\text{the function } \arctan)(\inf A).$

PROOF: Set $Z_0 = \square^0$. Set $Z_2 = \square^2$. Set $f = \frac{Z_0}{Z_0 + Z_2}$. $\text{dom } f = \mathbb{R}$. f is continuous. If $r \in \mathbb{R}$, then $f(r) = \frac{1}{1+r^2}$ by [13, (4)], (4). For every element x of \mathbb{R} such that $x \in \text{dom}(\text{the function } \arctan)'|_{\mathbb{R}}$ holds $(\text{the function } \arctan)'|_{\mathbb{R}}(x) = f(x)$. \square

$$(6) \quad \int_A ((-1)^i \cdot (\frac{\square^{2 \cdot n}}{\square^0 + \square^2}))(x) dx = (-1)^i \cdot ((\frac{1}{2 \cdot n + 1}) \cdot (\sup A)^{2 \cdot n + 1} - (\frac{1}{2 \cdot n + 1}) \cdot (\inf A)^{2 \cdot n + 1}) + \int_A ((-1)^{i+1} \cdot (\frac{\square^{2 \cdot (n+1)}}{\square^0 + \square^2}))(x) dx.$$

PROOF: Set $I_1 = (-1)^i$. Set $i_1 = i + 1$. Set $n_1 = n + 1$. Set $I_2 = (-1)^{i_1}$. Set $Z_0 = \square^0$. Set $Z_2 = \square^2$. Set $Z_{2n} = \square^{2 \cdot n}$. Set $f = I_1 \cdot Z_{2n}$. Set $g = I_2 \cdot (\frac{\square^{2 \cdot n_1}}{Z_0 + Z_2})$. $\text{dom } g = \mathbb{R}$. For every element x of \mathbb{R} , $(I_1 \cdot (\frac{Z_{2n}}{Z_0 + Z_2}))(x) = (f + g)(x)$ by [13, (6)], [17, (36)], (4). $f + g = I_1 \cdot (\frac{Z_{2n}}{Z_0 + Z_2})$. $\frac{\square^{2 \cdot n_1}}{Z_0 + Z_2}$ is continuous. \square

$$(7) \quad \text{Suppose } A = [0, r] \text{ and } r \geq 0. \text{ Then } |\int_A (\frac{\square^{2 \cdot n}}{\square^0 + \square^2})(x) dx| \leq (\frac{1}{2 \cdot n + 1}) \cdot r^{2 \cdot n + 1}.$$

PROOF: Set $Z_0 = \square^0$. Set $Z_2 = \square^2$. Set $N = 2 \cdot n$. Set $Z_n = \square^N$. Set $f = \frac{Z_n}{Z_0 + Z_2}$. f is continuous and $\text{dom } f = \mathbb{R}$. Reconsider $f_1 = f \upharpoonright A$ as a function from A into \mathbb{R} . Reconsider $Z_1 = Z_n \upharpoonright A$ as a function from A into \mathbb{R} . For every r such that $r \in A$ holds $f_1(r) \leq Z_1(r)$ by [4, (49)], [17,

(36)], [18, (3)], (4). For every object x such that $x \in \mathbb{R}$ holds $f(x) = |f|(x)$ by [13, (8)], (4). \square

2. EULER TRANSFORMATION

Let a be a sequence of real numbers. The alternating series of a yielding a sequence of real numbers is defined by

(Def. 1) $it(i) = (-1)^i \cdot a(i)$.

Now we state the proposition:

- (8) Let us consider a sequence a of real numbers. Suppose a is non-negative yielding, non-increasing, and convergent and $\lim a = 0$. Then
- (i) the alternating series of a is summable, and
 - (ii) for every n , $(\sum_{\alpha=0}^{\kappa} (\text{the alternating series of } a)(\alpha))_{\kappa \in \mathbb{N}}(2 \cdot n) \geq \sum(\text{the alternating series of } a) \geq (\sum_{\alpha=0}^{\kappa} (\text{the alternating series of } a)(\alpha))_{\kappa \in \mathbb{N}}(2 \cdot n + 1)$.

PROOF: Set $A =$ the alternating series of a . Set $P = (\sum_{\alpha=0}^{\kappa} A(\alpha))_{\kappa \in \mathbb{N}}$. Define $\mathcal{T}[\text{natural number, object}] \equiv \$_2 = P(2 \cdot \$_1)$. Define $\mathcal{S}[\text{natural number, object}] \equiv \$_2 = P(2 \cdot \$_1 + 1)$. Consider T being a function from \mathbb{N} into \mathbb{R} such that for every element x of \mathbb{N} , $\mathcal{T}[x, T(x)]$ from [5, Sch. 3].

Consider S being a function from \mathbb{N} into \mathbb{R} such that for every element x of \mathbb{N} , $\mathcal{S}[x, S(x)]$ from [5, Sch. 3]. For every natural number n , $S(n) \leq S(n+1)$. For every natural number n , $T(n) \geq T(n+1)$. For every natural number n , $T(n) \geq S(n)$. For every natural number n , $T(n) > S(0) - 1$ by [10, (6)]. For every natural number n , $S(n) < T(0) + 1$ by [10, (8)].

Define $\mathcal{D}(\text{natural number}) = 2 \cdot \$_1 + 1$. Consider D being a function from \mathbb{N} into \mathbb{N} such that for every element x of \mathbb{N} , $D(x) = \mathcal{D}(x)$ from [5, Sch. 8]. Reconsider $D_1 = D$ as a many sorted set indexed by \mathbb{N} . For every natural number n , $D(n) < D(n+1)$ by [2, (13)]. Reconsider $a_2 = a \cdot D_1$ as a sequence of real numbers.

For every object x such that $x \in \mathbb{N}$ holds $a_2(x) = (T - S)(x)$ by [4, (12)]. For every real number p such that $0 < p$ there exists a natural number n such that for every natural number m such that $n \leq m$ holds $|P(m) - \lim T| < p$ by [19, (9)]. \square

3. MAIN THEOREM

Let us consider r . The Leibniz series of r yielding a sequence of real numbers is defined by

$$(\text{Def. 2}) \quad it(n) = \frac{(-1)^n \cdot r^{2 \cdot n + 1}}{2 \cdot n + 1}.$$

The Leibniz series yielding a sequence of real numbers is defined by the term

$$(\text{Def. 3}) \quad \text{the Leibniz series of 1.}$$

Now we state the propositions:

(9) Suppose $r \in [-1, 1]$. Then

(i) |the Leibniz series of r | is non-negative yielding, non-increasing, and convergent, and

(ii) $\lim | \text{the Leibniz series of } r | = 0$.

PROOF: Set $r_1 = \text{the Leibniz series of } r$. Set $A = |r_1|$. $A(n) = \frac{|r|^{2 \cdot n + 1}}{2 \cdot n + 1}$ by [15, (1)], [3, (67), (65)]. $A(n) \geq A(n + 1)$ by [3, (46)], [15, (1)], [13, (6)], [2, (13)]. Set $C = \{0\}_{n \in \mathbb{N}}$. Define $\mathcal{F}(\text{natural number}) = \frac{\frac{1}{2}}{\$1 + \frac{1}{2}}$. Consider f being a sequence of real numbers such that $f(n) = \mathcal{F}(n)$ from [11, Sch. 1]. $C(n) \leq A(n) \leq f(n)$ by [11, (57)], [3, (46)], [13, (11)], [2, (11)]. \square

(10) (i) if $r \geq 0$, then the alternating series of |the Leibniz series of r | = the Leibniz series of r , and

(ii) if $r < 0$, then $(-1) \cdot (\text{the alternating series of |the Leibniz series of } r|) = \text{the Leibniz series of } r$.

PROOF: Set $r_1 = \text{the Leibniz series of } r$. Set $A = |r_1|$. Set $a_1 = \text{the alternating series of } A$. $a_1(n) = (-1)^n \cdot (\frac{|r|^{2 \cdot n + 1}}{2 \cdot n + 1})$ by [15, (1)], [3, (67), (65)]. If $r \geq 0$, then $a_1 = r_1$. \square

(11) If $r \in [-1, 1]$, then the Leibniz series of r is summable. The theorem is a consequence of (9), (8), and (10).

(12) Suppose $A = [0, r]$ and $r \geq 0$. Then $(\text{the function arctan})(r) = (\sum_{\alpha=0}^{\kappa} (\text{the Leibniz series of } r)(\alpha))_{\kappa \in \mathbb{N}}(n) + \int_A ((-1)^{n+1} \cdot (\frac{\square^{2 \cdot (n+1)}}{\square^0 + \square^2}))(x) dx$.

PROOF: Set $Z_0 = \square^0$. Set $Z_2 = \square^2$. Set $r_1 = \text{the Leibniz series of } r$. Define $\mathcal{P}[\text{natural number}] \equiv (\text{the function arctan})(r) = (\sum_{\alpha=0}^{\kappa} r_1(\alpha))_{\kappa \in \mathbb{N}}(\$1) + \int_A ((-1)^{\$1+1} \cdot (\frac{\square^{2 \cdot (\$1+1)}}{Z_0 + Z_2}))(x) dx$. $\mathcal{P}[0]$ by (5), [14, (43)], [13, (4)], [9, (21)].

If $\mathcal{P}[i]$, then $\mathcal{P}[i + 1]$ by [13, (11)], [2, (11)], (6). $\mathcal{P}[i]$ from [2, Sch. 2]. \square

(13) If $0 \leq r \leq 1$, then $(\text{the function arctan})(r) = \sum (\text{the Leibniz series of } r)$.

PROOF: Set $r_1 =$ the Leibniz series of r . Set $P = (\sum_{\alpha=0}^{\kappa} r_1(\alpha))_{\kappa \in \mathbb{N}}$. Set $A = (\text{the function arctan})(r)$. Define $\mathcal{I}(\text{natural number}) = \frac{\square^{2 \cdot s_1}}{\square^0 + \square^2}$. P is convergent. For every s such that $0 < s$ there exists n such that for every m such that $n \leq m$ holds $|P(m) - A| < s$ by [12, (3)], (4), [7, (11), (10)]. \square

(14) LEIBNIZ SERIES FOR π :

$$\frac{\pi}{4} = \sum(\text{the Leibniz series}).$$

(15) $(\sum_{\alpha=0}^{\kappa}(\text{the Leibniz series})(\alpha))_{\kappa \in \mathbb{N}}(2 \cdot n + 1) \leq \sum(\text{the Leibniz series}) \leq (\sum_{\alpha=0}^{\kappa}(\text{the Leibniz series})(\alpha))_{\kappa \in \mathbb{N}}(2 \cdot n)$. The theorem is a consequence of (9), (10), and (8).

(16) (i) $(\sum_{\alpha=0}^{\kappa}(\text{the Leibniz series})(\alpha))_{\kappa \in \mathbb{N}}(1) = \frac{2}{3}$, and

(ii) if n is odd, then $(\sum_{\alpha=0}^{\kappa}(\text{the Leibniz series})(\alpha))_{\kappa \in \mathbb{N}}(n+2) = (\sum_{\alpha=0}^{\kappa}(\text{the Leibniz series})(\alpha))_{\kappa \in \mathbb{N}}(n) + \frac{2}{4 \cdot n^2 + 16 \cdot n + 15}$.

(17) π APPROXIMATION:

$$\frac{313}{100} < \pi < \frac{315}{100}. \text{ The theorem is a consequence of (16), (14), and (15).}$$

REFERENCES

- [1] George E. Andrews, Richard Askey, and Ranjan Roy. *Special Functions*. Cambridge University Press, 1999.
- [2] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [3] Czesław Byliński. The complex numbers. *Formalized Mathematics*, 1(3):507–513, 1990.
- [4] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [5] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [6] Lokenath Debnath. *The Legacy of Leonhard Euler: A Tricentennial Tribute*. World Scientific, 2010.
- [7] Noboru Endou, Katsumi Wasaki, and Yasunari Shidama. Definition of integrability for partial functions from \mathbb{R} to \mathbb{R} and integrability for continuous functions. *Formalized Mathematics*, 9(2):281–284, 2001.
- [8] Konrad Knopp. *Infinite Sequences and Series*. Dover Publications, 1956. ISBN 978-0-486-60153-3.
- [9] Jarosław Kotowicz. Partial functions from a domain to the set of real numbers. *Formalized Mathematics*, 1(4):703–709, 1990.
- [10] Jarosław Kotowicz. Monotone real sequences. Subsequences. *Formalized Mathematics*, 1(3):471–475, 1990.
- [11] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(2):269–272, 1990.
- [12] Jarosław Kotowicz. Convergent real sequences. Upper and lower bound of sets of real numbers. *Formalized Mathematics*, 1(3):477–481, 1990.
- [13] Rafał Kwiatek. Factorial and Newton coefficients. *Formalized Mathematics*, 1(5):887–890, 1990.
- [14] Xiquan Liang and Bing Xie. Inverse trigonometric functions arctan and arccot. *Formalized Mathematics*, 16(2):147–158, 2008. doi:10.2478/v10037-008-0021-3.
- [15] Akira Nishino and Yasunari Shidama. The Maclaurin expansions. *Formalized Mathematics*, 13(3):421–425, 2005.

- [16] Chanapat Pacharapokin, Kanchun, and Hiroshi Yamazaki. Formulas and identities of trigonometric functions. *Formalized Mathematics*, 12(**2**):139–141, 2004.
- [17] Konrad Raczkowski. Integer and rational exponents. *Formalized Mathematics*, 2(**1**):125–130, 1991.
- [18] Konrad Raczkowski and Andrzej Nędzusiak. Real exponents and logarithms. *Formalized Mathematics*, 2(**2**):213–216, 1991.
- [19] Piotr Rudnicki and Andrzej Trybulec. Abian’s fixed point theorem. *Formalized Mathematics*, 6(**3**):335–338, 1997.
- [20] Yuguang Yang and Yasunari Shidama. Trigonometric functions and existence of circle ratio. *Formalized Mathematics*, 7(**2**):255–263, 1998.

Received October 18, 2016
