HELMINTHOLOGIA, 55, 1: 60 - 69, 2018

Diplodiscus mehrai Pande, 1937 and *D. japonicus* (Yamaguti, 1936): morphology of developmental stages and molecular data

V. V. BESPROZVANNYKH¹, K. V. ROZHKOVAN^{1*}, A. V. ERMOLENKO¹, A. V. IZRAILSKAYA²

¹Federal Scientific Center of the East Asia Terrestrial Biodiversity FEB RAS, Vladivostok, 690022, Russia, *E-mail: 27.tomcat@gmail.com; ²Department of Biodiversity and marine bioresoures, Far Eastern Federal University, 690051, Vladivostok, Russia

Article info	Summary
Received March 29, 2017 Accepted August 23, 2017	 Specimens of the snails <i>Anisus centrifugops</i> were infected with two types of cercariae relating to the genus <i>Diplodiscus</i>, differing from each other by body and organ sizes were found in a pond in the territory of Vladivostok (Primorsky region, Russia). Further study of their morphology and life cycles allowed us to establish that these flukes belong to <i>Diplodiscus japonicus</i> (Yamaguti, 1936) and <i>D. mehrai</i> Pande, 1937. Morphological and molecular data confirmed the validity of <i>D. japonicus</i> which had previously been synonymized with <i>D. amphichrus</i> Tubangui, 1933 and <i>D. mehrai</i> Pande, 1937. Keywords: <i>Diplodiscus mehrai</i>; <i>Diplodiscus japonicus</i>; life cycle; cercariae; metacercariae; adult worm

Introduction

The family Diplodiscidae Cohn, 1904 is a small group of paramphistomoids, found predominantly in amphibians but with representatives in reptiles and in fish (Jones, 2005). The concept of *Diplodiscus* Diesing, 1836, as well as the taxonomy of individual species of this genus, is based on the morphology of adult worms (Sey, 1991). After revision of Amphistomida, and particularly, Diplodiscus, most of the species and subspecies from this genus described before 1991 were synonymized to different species (Sey, 1991). At present, there is a small amount of data on developmental stages of Diplodiscus. For D. subclavatus (Goeze, 1882), D. amphichrus Tubangui, 1933, D. brevicoeca Richard, Chabaud et Brygoo, 1968 (Sey, 1991) and D. japonicus (Yamaguti, 1936), only cercariae, metacercariae, and parthenitae have been described (Sey, 1991; Skrjabin, 1949). However, the morphometric data for these stages are incomplete. Besides the life cycles of Diplodiscus, data on the circulation features of D. subclavatus were obtained. The life span of cercariae after it emergence from a snail (usually in the morning) is approximately 28 h (Grabda-Kazubska, 1980). Cercariae encyst at the end of the free-living phase. Grabda-Kazubska also showed that cercariae encyst in water, but not on amphibian skin as proposed by Lang (1892), and that the life span of metacercariae is 5 days.

The only species of *Diplodiscus* known from Russia is *D. subclavatus*. The literature also reports *Diplodiscus sphincterostoma* at the territory of the Primorsky Region, discovered by E.V. Belous. However, this fluke is not described; thus, this species was considered a "nomen nudum" by Ryzhikov *et al.* (1980).

In a pond in the territory of Vladivostok, we found the snail *Anisus centrifugops* (Prosorova et Starobogatov) with two types of cercariae of *Diplodiscus*. These cercariae differed in body and organs sizes. Further studies of the life cycles, morphology of different life stages of adult worms allowed us to establish that these flukes belonged to *D. japonicus* and *D. mehrai* Pande, 1937.

^{* -} corresponding author

Material and Methods

Morphology and life cycle

A total of 250 specimens of Anisus centrifugops naturally infected with parthenitae of species of Diplodiscus were taken from a small pond in the territory of Vladivostok, Primorsky Region, Russia. Adult flukes were also found in naturally infected Rana dybowskii caught in the same place in the spring and autumn. In order to identify these trematodes and their circulation, four infected snails were separately placed in dishes with 1 I of tap water. The snails were divided according to the size of cercariae: two reservoirs contained mollusks with smaller cercariae, and two others contained mollusks with larger ones. Fifteen tadpoles of Rana dybowskii were then placed in each reservoir. Tadpoles were collected in natural ponds free of snails; 30 tadpoles were checked to verify absence of infection before the experiment. During the experiment, tadpoles were periodically dissected to check the level of infection and time of development to the adult worm stage. The process of trematode invasion into the definitive host was studied by placing infected snails into deep Petri dishes. After the peak of cercariae emission, the snail in each dish was replaced with three tadpoles. The process of invasion was observed under a stereo microscope. To check the infection in tadpoles, they were dissected on the first and the subsequent two days from the beginning of the experiment.

To determine the life span of cercariae, infected snails were placed in Petri dishes with tap water for 20 – 30 min. Cercariae emerging within this time were counted as even-aged. Maximal cercariae life span was defined by the time of emergence from the mollusk until the death of all specimens. Phototaxy was tested by placing the cercariae into long, narrow glass vials (20 × 2 × 3 cm), 1/3 of which were covered by black paper from one side, and a bright spotlight placed on the other side. The diurnal rhythm of cercariae emission was observed by placing a single snail in a Petri dish with 50 ml of tap water. Dishes were shaken every 2 h, and 10 aliquots of 1 ml were taken from each dish and placed in smaller vessels. Dishes were then refilled with water to 50 ml. Cercariae were immobilized by iodine and their number in each vessel was counted. The average number of larvae was then multiplied by 5. Experimental studies were conducted at water temperatures ranging between 18 and 22 °C.

All measurements, in millimeters (mm), of metacercariae and

parthenitae were made on living specimens, and before measurements, cercariae were fixed in hot 4 % formalin. Adult worms were fixed in 70 % ethanol and then placed in 96 % ethanol. For whole mounts, worms were stained in alcoholic hydrochloric carmine, dehydrated in a graded ethanol series, cleared in clove oil, and mounted in Canada balsam.

Extraction of total genomic DNA, PCR, and sequencing

Two and four adult specimens of D. mehrai and D. japonicus, respectively, were washed several times in a physiological solution immediately after dissection and stored in 96 % ethanol. Genomic DNA was extracted from individual worms using hot sodium hydroxide and Tris (HotSHOT) technique (Truett et al., 2000). The complete sequences of 18S rDNA, ITS1-5.8S-ITS2 rDNA region and partial sequences of 28S rDNA were amplified using the following primer sets: 18S-E and 18S-F (Littlewood & Olson, 2001) for 18S rDNA, BD1 and BD2 (Morgan & Blair, 1995) for ITS1-5.8S-ITS2 rDNA, and digl2 and 1500R (Tkach et al., 2003) for 28S rDNA. PCR reaction was conducted in a 25 µl reaction mixture containing 10 pM of each primer, 12.5 µl of PCR Master Mix (2X) (Thermo Fisher Scientific Baltics), and 5 µl of extracted DNA. Sequencing was performed on an ABI 3130 automated capillary sequencer (Life Technologies, Grand Island, New York, USA) using PCR primers for 18S rDNA and the following internal primers: 373C, 892C, 1262, 1262C, 1/F (Krieger et al., 2006), Pace-A, 18S-A27, 18S-4 (Littlewood & Olson, 2001). ITS1-5.8S-ITS2 rDNA region was sequenced using PCR primers and the internal primer 3S (Littlewood & Olson, 2001). 28S rDNA was sequenced with ECD2, 900F and 1200R internal primers (Waeschenbach et al., 2007).

Sequence alignment and phylogenetic analyses

The ribosomal DNA sequences were assembled and aligned manually using the MEGA 5 (Tamura *et al.*, 2011). Sequences have been submitted to GenBank under the following accession numbers: KX506856-KX506857 (*D. mehrai*) and KX506852-KX506855 (*D. japonicus*).

For the purpose of phylogenetic analysis we used the combined set of 18S and 28S rDNA sequences. Prior to phylogenetic analyses sequences were edited, the ends of each sequence were trimmed to match the shortest sequence in the alignment. The final

		•	•
Species	n	Reference	Accession number
Diplodiscus mehrai	2	Current study	KX506856-KX506857
Diplodiscus japonicus	4	Current study	KX506852-KX506855
Diplodiscus subclavatus	1	Olson <i>et al.</i> , 2003	AY222212
Fischoederius cobboldi	3	Ghatani <i>et al.</i> , 2014	JX518961-JX518963
Fischoederius elongatus	3	Ghatani <i>et al.</i> , 2014	JX518964-JX518966
Fasciola hepatica	1	Olson et al., 2003	AY222244

Table 1. List of taxa, incorporated into the molecular analysis

alignment yielded was 2874 nucleotide long and consisted of 13 sequences (Table 1). Sequences of *Fasciola hepatica* (Linnaeus, 1758) were used as outgroup. Distance matrices were constructed with the absolute pairwise character difference. Pairwise comparisons of absolute sequence divergence for all taxa were calculated with gaps treated as missing data. Maximum Likelihood (ML) using PhyML 3.1 (Guindon & Gascuel, 2003), and Bayesian Inference (BI) using Mr. Bayes 3.1.2 (Ronquist & Huelsenbeck, 2003) analyses of the 18S-28S rDNA data were conducted to explore relationships between the newly generated data and nucleotide sequences available in NCBI Genbank. ML and BI phylogenetic analyses were performed using TIM3+G substitution model with gamma

correction value 0.1190. This model showed the best fit to our data using Bayesian information criterion (BIC) in jModeltest 2.1.5 software (Darriba *et al.*, 2012). Bayesian inference was used with the following nucleotide substitution parameters: Iset nucmodel = 4by4, nst = 6, and rates = gamma. Markov chain Monte Carlo (MCMC) chains were run for 1,000,000 generations, log-likelihood scores were plotted and only the final 75 % of trees were used to produce the consensus tree. This number of generations was considered sufficient because the standard deviation of spitted sequences value dropped below 0.01 at the end of the run. The significance of the phylogenetic relationship was estimated using a bootstrap analysis (Felsenstein, 1985) with 100 replications for ML



Fig. 1. Adult Diplodiscus: a - D. mehrai Pande, 1937, b - D. japonicus (Yamaguti, 1936).

			Table 2. Measure	ments (mm) of adult Diplodiscus.				
Feature	D. japonicus (present study)	<i>D. japonicus</i> (from: Yamaguti 1936)	<i>D. japonicus</i> (according to Bravo, 1941; from: Skrjabin 1949)	D. mehrai (present study)	<i>D. mehrai</i> (according to Pande, 1937; from Skrjabin 1949)	<i>D. mehrai</i> (from: Sey, 1991)	<i>D. amphichrus</i> (according to Tubangi, 1933; from: Skrjabin 1949)	D. amphichrus (from: Sey, 1991)
Body length Body width	1.25 – 2.00 (0.80 – 0.89) 0.57 – 0.99 (0.41 – 0.49)	1.5 - 2.5	1.8 – 2.75 0 74 – 0 975	2.37 – 2.77 (0.98 – 1.03)* 0 91 – 1 23 (0 36 – 0 48)	2.1-3.2 10-13	2.1 – 3.2 1 0 – 1 3	1.60 – 3.45 0.60 – 0.86	1.6 – 3.4 0 2 – 0 8
Pharvnx length	0.20 - 0.31 (0.112 - 0.168)		0.156 - 0.292	0.25 - 0.28 (0.123 - 0.168)	0.36	0.36	0.110 – 0.292	0.29 - 0.32
Pharynx width	0.18 - 0.31 (0.145 - 0.168)	0.15 – 0.23	0.195 – 0.214	0.22 - 0.31 (0.134 - 0.145)	0.2 – 0.32	I	0.100 - 0.330	I
Pharyngeal sacs length	0.18 – 0.20 (0.10 – 0.11)	I	0.117	0.17 – 0.24 (0.14 – 0.15)	0.2	0.28	0.136	0.13
Pharyngeal sacs width	0.13 – 0.22 (0.09 – 0.10)	I	0.078 – 0.097	0.19 – 0.22 (0.089 – 0.112)	0.18	I	0.136	I
Pharynx together with sacs length	0.26 – 0.43	0.25 - 0.37		0.32 – 0.43				
Esophagus length	0.18 – 0.39	0.33	0.273 – 0.33	0.28 - 0.50	0.38 – 0.50	0.38 – 0.50	0.270 – 0.440	0.35
Esophageal bulb length	0.120 – 0.154 (0.041 – 0.056)	0.12 – 0.15	0.102 – 0.175	0.123 – 0.169 (0.084)	0.11 – 0.14	0.11 – 0.150	0.156	I
Esophageal bulb width	0.108 – 0.123 (0.041 – 0.056)	0.10 – 0.13	0.085 – 0.117	0.108 – 0.139 (0.056 – 0.072)	0.07 – 0.10	I		I
Acetabulum length	0.45 – 0.65 (0.26 – 0.35)	0.47 – 0.80	0.2	0.62 – 0.95 (0.32 – 0.40)	0.56 – 0.74	0.56 – 0.74	0.50 – 1.08	0.54 – 1.16
Acetabulum width	0.54 – 0.89 (0.39 – 0.59)	0.47 – 0.80	0.7 - 0.975	0.82 – 1.19 (0.40 – 0.54)	0.90 – 1.11	0.90 – 1.14	0.50 – 1.08	0.54 – 1.16
Sucker-like protuberance length	0.17 – 0.31	I	I	0.34 – 0.40	I	I		I
Sucker-like protuberance width	0.20 – 0.35	I	I	0.38 – 0.47	I	I		I
Ovary length	0.11 – 0.13 (0.05 – 0.069)	0.13 – 0.25	0.102 – 0.175	0.15 – 0.17 (0.061 – 0.072)	0.16 – 0.18	0.16 – 0.18	0.12 – 0.24	I
Ovary width	0.12 – 0.17 (0.05 – 0.069)	0.16 – 0.20	0.062 – 0.128	0.15 – 0.17 (0.056 – 0.072)	0.16 - 0.18	0.16 – 0.18	0.12 – 0.24	I
Testis length	0.32 – 0.37 (0.084 – 0.089)	0.18 – 0.37	0.198 – 0.292	0.36 – 0.45 (0.112 – 0.123)	0.23 – 0.40	0.23 – 0.40	0.140 – 0.1487	0.14
Testis width	0.30 – 0.50 (0.112 – 0.123)	0.20 – 0.33	0.253 – 0.312	0.38 – 0.42 (0.100 – 0.170)	0.34 – 0.45	0.34 – 0.45	0.140 – 0.1487	0.14
Cirrus-sac length	0.127 – 0.189 (0.089 – 0.108)	0.16	I	0.21 – 0.23 (0.092 – 0.139)	0.16	0.18	0.12 – 0.22	0.13-0.15
Cirrus-sac width	0.060 – 0.135 (0.054 – 0.069)	0.10	I	0.14 – 0.20 (0.058 – 0.081)	0.18	0.16	0.10 – 0.14	0.09 – 0.12
Number of vitelline follicles	17 – 21	30 – 40	21	17 – 21	35 – 40	17 – 20	45	I
Eggs length	0.095 – 0.104	0.102 – 0.126	0.105 - 0.120	0.112 – 0.126	0.09 – 0.14	0.146	0.104 – 0.112	0.113 – 0.119
Eggs width	0.061 – 0.073	0.066 – 0.078	0.058 - 0.097	0.067 – 0.078		0.093	0.062 – 0.070	0.068 – 0.079
Body length/length of pharynx with sacs	4.5 – 4.9	I	I	6.2 - 7.4	I	I	I	I
Acetabulum/pharynx	1.93 – 2.4	I	I	2.48 – 3.39	I	I	I	I
								Postbifurcal
Genital pore	Bifurcal or postbifurcal	Bifurcal	Postbifurcal	Bifurcal or postbifurcal	Bifurcal	Bifurcal	Postbifurcal	(on some distance – Sey 1991, Fig. 314)

Table 2. Measurements (mm) of adult Diplodiscus.

* - Sizes at the time of puberty

and by the amount of a posterior probability for BI. Trees were visualized using FigTree 1.4 software (http://tree.bio.ed.ac.uk/software/figtree/).

Results

Description

Diplodiscus mehrai Pande, 1937

Definitive host: tadpoles and adult Rana dybowskii Günther (Ranidae, Rafinesque).

Infection rate of tadpoles and frogs in natural conditions: 100 %.

Infection intensity of tadpoles in natural conditions: 4 – 23 specimens.

Infection intensity of frogs in natural conditions: 1 – 15 specimens. Infection rate of tadpoles in experiment: 20 %.

Infection intensity of tadpoles in experiment: 1 – 7 specimens. Site: rectum.

Intermediate host: freshwater snail Anisus centrifugops (Prosorova et Starobogatov) (Planorbidae, Rafinesque).

Infection rate: 4.2 %.

Locality: Vladivostok, Primorsky Region, Russia; 43°12' N, 131°55'E.

Adult worm (based on 10 specimens; Table 2, Figs. 1a, 3a): Body conical, non-spined with narrow anterior end and wide posterior

end. Pharynx terminal with two pharyngeal sacs. Specimens from tadpoles with diffuse pigment spots on both sides of these sacs. Specimens from frogs with pigment spots or not. Oesophagus ventral to pharyngeal sacs. Oesophageal bulb with wall thickness 0.027 - 0.031. Oesophageal bifurcation on border of anterior and middle third of body. Caeca reach anterior margin of acetabulum. Acetabulum terminal, on posterior end of body, with central sucker-like protuberance. Testis single, spherical, smooth to irregular, between caeca on median line of body before acetabulum. Cirrus-sac in area of oesophageal bifurcation, with small internal seminal vesicle. External seminal vesicle present, size depends on its occupancy with sexual products. Genital pore bifurcal to postbifurcal, right or left from bifurcation. Ovary dextral from medial line of body on level of posterior margin of testis. Mehlis' gland left from ovary, 0.16 × 0.15. Uterus between caeca, genital pore and testis. Each of two vitelline fields form two groups of follicles, one between level of middle of oesophagus and middle of testis, second between posterior margin of testis and acetabulum. Most of second group of follicles in both vitelline fields across body, and follicles reach median line of body. Each vitelline field comprises 17 - 21 follicles. Eggs not numerous, operculated. Excretory bladder T-shaped.

Redia (based on 10 specimens; Fig. 2a.) Body sac-shaped, $0.10 - 0.13 \times 0.28 - 0.3$. Pharinx $0.09 - 0.12 \times 0.11 - 0.12$. Caecum short. *Cercaria* (based on 10 specimens; Table 3, Figs. 2b, 3b) Body leaf-shaped, slightly transparent, grey by pigment. Pigment eyes



Fig. 2. Diplodiscus mehrai Pande, 1937: a – redia, b – cercaria, c and d – metacercaria, dorso-ventrally and laterally, accordingly; D. japonicus (Yamaguti, 1936): e – cercaria.



Fig. 3. Diplodiscus mehrai Pande, 1937: a – adult worm, b – cercaria; D. japonicus (Yamaguti, 1936): c – adult worm, d – cercaria; e – metacercaria found in feces of snail.

on both sides of body. Anterior end of body with long sensillae. Pharynx terminal with two sacs. Oesophagus short, with bulb. Oesophageal bifurcation on border of anterior and middle third of body. Caeca terminate at level of middle of acetabulum. Acetabulum terminal, on posterior end of body, with central sucker-like protuberance. Primordium of testis cross-oval, immediately before acetabulum. Primordium of ovary left from median line of body, on level of testis. Cystogenous glands from pharyngeal sacs to posterior end of body. Excretory bladder T-shaped. Two main collective channels, each with 20 - 21 granules, left and right from median line of body, turn in area of pigment eyes to posterior end of body. On midline of body each channel divided on two secondary channels; one of them reaches pharynx and second reaches acetabulum, and further divided into three branches, each includes three ducts terminated by small expansion. Tail 1.7 - 2.1 times longer than body.

 $\label{eq:metacercaria} \begin{array}{l} \textit{Metacercaria} \mbox{ (based on 10 specimens; Fig. 2c,d). Cyst 0.28 \times 0.28 \\ - 0.29, \mbox{ spherical, concaved on one side.} \end{array}$

Molecular sequence data 18S rDNA (complete) – 1991 bp, ITS1 (complete) – 490 bp, 5.8S rDNA (complete) – 159 bp, ITS2 (complete) – 281 bp, 28S rDNA (partial) – 1391 bp.

Diplodiscus japonicus (Yamaguti, 1936)

Definitive host: tadpoles and adult Rana dybowskii Günther (Ranidae, Rafinesque). Infection rate of tadpoles and frogs in natural conditions: 100 %. Infection intensity of tadpoles in natural conditions: 6 – 32 specimens.

Infection intensity of frogs in natural conditions: 1 – 21 specimens. Infection rate of tadpoles in experiment: 43 %.

Infection intensity of tadpoles in experiment: 1 – 95 specimens Site: rectum.

Intermediate host: freshwater snail Anisus centrifugops (Prosorova et Starobogatov) (Planorbidae, Rafinesque).

Rate of infection: 6.4 %.

Locality: Vladivostok, Primorsky Region, Russia; 43° 12'N, 131° 55'E.

Adult worm (based on 10 specimens; Table 2, Figs. 1a, 3c). Body conical or trapezoidal, non spined with narrow anterior end and wide posterior end. Pharynx terminal with two pharyngeal sacs. Specimens from tadpoles with diffuse pigment spots on both sides of these sacs. Specimens from frogs with pigment spots or not. Oesophagus ventral to pharyngeal sacs. Oesophageal bulb oval, with wall thickness 0.039 – 0.042. Oesophageal bifurcation on border of anterior and middle third of body. Caeca reach anterior margin of acetabulum. Acetabulum terminal, on posterior end of body, with central sucker-like protuberance, can partly cover ovary. Testis spherical, smooth to irregular, between caeca on median line of body before acetabulum. Cirrus-sac in area of oesophageal

bifurcation, with small internal seminal vesicle. External seminal vesicle present. Genital pore bifurcal to postbifurcal. Ovary between testis and acetabulum, on median line of body, on some distance from testis. Mehlis' gland left from ovary, 0.11×0.12 . Uterus between caeca, genital pore and ovary. Each of two vitelline fields rorm two groups of follicles. One between level of middle of oesophagus and middle of testis, second between posterior margin of testis and acetabulum Most of second group of follicles in both vitelline fields across body, and follicles reach median line of body. Each vitelline field comprises 17 - 21 follicles. Eggs not numerous, operculated. Excretory bladder T-shaped.

Redia (based on 10 specimens). Body sac-shaped, $0.10 - 0.13 \times 0.28 - 0.3$. Pharinx $0.09 - 0.12 \times 0.11 - 0.12$. Caecum short.

Cercaria (based on 10 specimens; Table 3, Figs. 2e, 3d). Body leaf-shaped, transparent. Pigment eyes on both sides of body. Anterior end of body with poorly visible, short sensillae. Pharynx terminal with two sacs. Oesophagus short, with bulb. Oesophageal bifurcation on border of anterior and middle third of body. Caeca terminate at level of anterior margin of acetabulum. Acetabulum terminal, on posterior end of body, with central sucker-like protuberance. Primordium of testis cross-oval, immediately before acetabulum. Primordium of ovary right from median line of body, on level of testis. Cystogenous glands from pharyngeal sacs to posterior end of body. Excretory bladder T-shaped. Two main collective channels, each with 21 - 23 granules, left and right from median line of body, turn to posterior end of body. On midline of body each channel divided on two secondary channels; one of them reaches pharynx and second reaches acetabulum, and further divided into three branches, each includes three ducts terminated by small expansions. Tail 2.3 – 2.4 times longer than body.

Metacercaria (based on 10 specimens). Cyst $0.23 - 0.24 \times 0.25 - 0.26$, spherical, concaved on one side.

Molecular sequence data 18S rDNA (complete) – 1991 bp, ITS1 (complete) – 469 bp, 5.8S rDNA (complete) – 159 bp, ITS2 (complete) – 281 bp, 28S rDNA (partial) – 1391 bp.

Life cycles

For both studied species of Diplodiscus, as for other representatives of this genus, the two-host life cycle is usual. Cercariae (1 - 50 specimens) emerged from snails between 11 a.m. and 2 p.m. The most intensive emergences, up to 50 specimens per day, of cercariae occurred during spring. In August - September, this value decreased to 1 - 10 cercariae per day. Larvae are active in water and have positive phototaxis. This allows them to concentrate in warm and sunny pond regions. Such places contain many tadpoles. These factors presumably improve the possibility of definitive host infection. The life span of the free-living form in water temperature 18 - 22 °C is 8 - 12 h. At the end of this time, some cercariae are encysted. The rest, up to 80 %, do not encyst and die. Metacercariae did not attach to tadpole skin nor to the bottom of the vessel nor to plants and sand in the vessel, and their life span under laboratory conditions was no longer than 3 - 4 days. Larvae can frequently be swallowed by snails but are not digested and stay alive. Such metacercariae were found in snail feces (Fig. 3e). The results of experiments reported here showed that tadpoles infection occurred mainly by swallowing the cercariae. Tadpoles, after being placed with cercariae, swallowed all of them within 20 - 30 min. At this moment, cercariae were not yet encysted and

metacercariae were absent. Tadpoles were dissected over 3 days

Features	D. mechrai (present study)	<i>D. japonicus</i> (present study)	<i>D. japonicus</i> (from: Yamaguti 1940)	D. amphichrus (according to Tubangi & Masilungan; from: Sey 1991)	D. amphichrus (according to Odening; from: Sey 1991)		
		Cerca	ria				
Body length	0.51 – 0.67	0.30 - 0.36	0.36 - 0.75	0.21 – 0.24	0.41 – 0.59		
Body width	0.30 - 0.37	0.19 - 0.26	0.16 – 0.35	0.20 - 0.22	0.27 – 0.29		
Pharynx with sacs length	0.12 - 0.14	0.084 - 0.127	0.080 - 0.120	-	0.12		
Pharynx width	0.089 - 0.095	0.061 - 0.072	0.054 - 0.084	-	0.09 - 0.15		
Pharyngeal bulb length	0.038 - 0.045	0.033 - 0.035	0.027 - 0.045	-	-		
Pharyngeal bulb width	0.028 - 0.034	0.014 - 0.019	0.020 - 0.033	-	-		
Acetabulum length	0.12 – 0.15	0.112 - 0.140	0.096 - 0.165	-	0.15 – 0.18		
Acetabulum width	0.22 - 0.30	0.150 – 0.178	0.130 - 0.204	-	0.15 – 0.18		
Primordium of ovary length	0.035 - 0.036	0.028 - 0.034	-	-	-		
Primordium of ovary width	0.056 - 0.058	0.045 - 0.056	-	-	-		
Primordium of testis length	0.059 - 0.062	0.055 - 0.058	0.030	-	-		
Primordium of testis width	0.089 - 0.091	0.061 - 0.063	0.060	-	-		
Tail length	1.12 – 1.16	0.72 – 0.83	0.60 - 0.90	0.76 – 1.00	0.56 - 0.69		
Tail widht	0.11 – 0.13	0.078 - 0.089	0.075 - 0.130	-	-		
Metacercaria							
Cyst	0.28 – 0.29 in diameter	0.23 – 0.24 × 0.25 – 0.26	0.27 × 0.25	0.24 - 0.25 × 0.20 - 0.23	-		

able 3	Sizes (mm) of	the cerc:	ariae and	metacercaria	e of I	Dinlodiscus
able J.	JIZES (11111) 01		anae anu	melacercana		JIDIOUISCUS



0.02

Fig. 4. Bayesian tree inferred from combined 18S and 28S rDNA sequence data. Results of statistical support are shown as follows: BI/ML. Sequences obtained in this study are in bold.

post-exposure, and 7 - 20 flukes were found in their intestines. In other cases, cercariae and metacercariae were found in the tadpoles' oesophagus. Thus, tadpole infection can occur via swallowing either cercariae or metacercariae. However, because only some cercariae are encysted, and the life span of metacercariae is up to 4 days, the primary way of infection occurs through the swallowing of cercariae. Metacercariae can only increase the probability of infection for a short period of time. The first adult *D. mehrai* and *D. japonicus* were both recovered on 30^{th} day of the experiment. *Diplodiscus mehrai* infected 6 tadpoles with an intensity of 1 to 7 flukes, and *D. japonicus* infected 13 tadpoles with an intensity of 1 to 95 flukes.

According to the natural conditions of the Primorsky Region and biological features of *Rana dybowskii*, the maximal number of tadpoles in ponds is observed in May – June, and the minimal one is in August – September (Kuzmin, 2012). These periods correspond to maximal and minimal emission of *Diplodiscus* cercariae, respectively. The tadpoles collected in May – June were infected by both adult flukes and larvae, whereas adult frogs studied in September were infected by adult worms only. This allows us to propose that frog infection in nature occurs at the tadpole stage, and not at adult *Rana*.

Molecular data

The 18S-ITS1-5.8S-ITS2-28S rDNA sequence length for *D. mehrai* was 2412 bp, and 2391 bp for *D. japonicus*. No intraindividual variation was found in any rDNA region examined. For genetic distances calculation and phylogenetic reconstructions we chose the 18S and 28S rDNA genes as the most prevalent in Genbank for the selected group. The main differences between *D. mehrai* and *D. japonicus* were in one G \leftrightarrow A transition for 18S rDNA and 8 transitions

(5 G \leftrightarrow A and 3 C \leftrightarrow T) and one G \leftrightarrow T transversion for 28S rDNA. The genetic divergence between *D. mehrai* and *D. japonicus* was 0.35 %, which is equal to that between *Fischoederius cobboldi* and *F. elongatus*. The differences relative to other representative of *Diplodiscus*, *D. subclavatus*, were 0.87 and 0.8 % for *D. mehrai* and *D. japonicus*, respectively. Phylogenetic analysis (Fig. 4) showed that *D. mehrai* and *D. japonicus* are separate species and they are more closely related to each other than to *D. subclavatus*.

Discussion

Sey (1991) revised all species belonging to superfamily Paramphistomoidea Fischoeder, 1901 and included 11 species in Diplodiscus. As a result, specimens that had previously been attributed to a single species were synonymized to different species within Diplodiscus (Sey, 1991). In part, this was related to D. mehrai and D. amphichrus japonicus Yamaguti, 1936. From these two species, only D. mehrai, described by Pande in 1937 was included in the 11 valid species within *Diplodiscus* (Sey, 1991). Specimens found later and identified as D. mehrai were synonymized to D. magnus (Srivastava, 1934) (Sey, 1991). Li (1937) D. amphichrus japonicus was elevated to species rank and its name was changed to D. japonicus. Later, specimens described as D. amphichrus japonicus and D. japonicus were synonymized either to D. amphichrus or D. mehrai (Sey, 1991). All flukes listed above were discovered in East-Asia and India. D. mehrai was also found in Georgia (Petriashvili, 1964).

The results of our research demonstrate the presence of two species of *Diplodiscus* in the territory of the Primorsky Region. Adult worms of one of these species by metric and morphological characteristics are identical to *D. mehrai* described by Pande (1937) (Table 2). The second species has much in common morphologically with D. japonicus described by Yamaguti (1936, 1941) (Tables 2, 3). The only significant difference between D. japonicus from this study and that of Yamaguti is maximal cercariae body size (Yamaguti, 1940). According to Yamaguti the maximal body size is twice the minimal one. Such differences in size of cercariae are not typical for trematodes from one species. All other morphological traits of cercariae, metacercariae, and adult worms and the participation of snails from Anusis in the circulation of flukes, points to that the second trematode discovered in Primorsky Region, in our opinion, is D. japonicus. This species differs from both adult worms and cercariae of D. mehrai and D. amphichrus by morphometric features. Diplodiscus japonicus differs from adult D. mehrai by smaller maximal body size, length of oesophagus, width of ventral sucker and cirrus sac, and maximal egg length. Adult specimens of D. japonicus differ from D. amphichrus by maximal body size and ventral sucker, testis size, length of pharyngeal sac and position of genital pore (Table 2). Cercariae of D. japonicus also have sufficient metric differences from those from D. amphichrus and D. ehrai (Table 3).

Thus, in most morphometric characters of the developmental stages, *D. japonicus* differs from both *D. mehrai* and *D. amphichrus* (Tables 2, 3), which allows us to count *D. japonicus* as a separate species within *Diplodiscus*. The validity of *D. mehrai* and *D. japonicus* is also confirmed by molecular data. Genetic *p*-distances and phylogenetic analysis (Fig. 4) showed that *D. mehrai* and *D. japonicus* are separate species and that they are more closely related to each other than to *D. subclavatus*.

References

DARRIBA, D., TABOADA, G.L., DOALLO, R., POSADA, D. (2012): jModel-Test 2: more models, new heuristics and parallel computing. *Nat. Methods*, 9, 772. DOI:10.1038/nmeth.2109

FELSENSTEIN, J. (1985): Confidence limits on phylogenies: an approach using bootstrap. *Evolution*, 39, 783 – 791. DOI: 10.2307/2408678

GHATANI, S., SHYLLA, J.A., ROY, B., TANDON, V. (2014): Multilocus sequence evaluation for differentiating species of the trematode Family Gastrothylacidae, with a note on the utility of mitochondrial COI motifs in species identification. *Gene*, 548, 277 – 284. DOI: 10.1016/j.gene.2014.07.046

GRABDA-KAZUBSKA, B. (1980): Observation on life-cycle of *Diplodiscus subclavatus* (Pallas, 1760) (Trematoda, Diplodiscidae). *Acta Parasitol.*, 27, 261 – 271

GUINDON, S., GASCUEL, O. (2003): A simple, fast and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst. Biol.*, 52, 696 – 704. DOI: 10.1080/10635150390235520

JONES, A. (2005): Family Diplodiscidae Cohn, 1904. In: JONES, A., BRAY, R.A., GIBSON, D.I. (Eds), *Keys to the Trematoda*. Volume 2. London, UK: CABI Publishing and The Natural History Museum, pp. 319 – 324

KRIEGER, J., HETT, A.K., FUERST, P.A., BIRSTEIN, V.J., LUDWIG, A.

(2006): Unusual intraindividual variation of the nuclear 18S rRNA gene is widespread within the Acipenseridae. *J. Hered.*, 97, 218 – 225. DOI:10.1093/jhered/esj035

KUZMIN, S.L. (2012): [Amphibians of former USSR] Moscow, Fellowship of scientific books. 370 p. (In Russian)

LANG, A. (1892): Über die Cercariae von Amphistomum subclavatum, Ber. Naturforsch. Ges. Freiburg im Breisgau. 9 p.

Lı, L.Y. (1937): Some Trematode parasites of frogs with a description of *Diplodiscus sinicus* sp. nov. *Lingnan Sci. Journ. Canton*, 19, 61 – 70

LITTLEWOOD, D.T.J., OLSON, P.D. (2001): Small subunit rDNA and the Platyhelminthes: signal, noise, conflict and compromise. In: LITTLE-WOOD, D.T.J. & BRAY, R.A. (Eds), *Interrelationships of the Platyhelminthes*. London, Taylor & Francis. pp. 262 – 278

MORGAN, J.A.T., BLAIR, D. (1995): Nuclear rDNA ITS sequence variation in the trematode genus *Echinostoma*: an aid to establishing relationships within the 37-collar-spine group. *Parasitology*, 111, 609 – 615. DOI: 10.1017/S003118200007709X

OLSON, P.D., CRIBB, T.H., TKACH, V.V., BRAY, R.A., LITTLEWOOD, D.T.J. (2003): Phylogeny and classification of the Digenea (Platyhelminthes: Trematoda). *Int. J. Parasitol.*, 33, 733 – 755. DOI: 10.1016/S0020-7519(03)00049-3

PETRIASHVILI, L.G. (1964) [Helminth fauna of the marsh-frog (Rana ridibunda Pall.) in Lake Basaletsky] Soobshch. Akad. Nauk. Gruz. GSSR Vol.36. pp. 457 – 462 (In Russian)

RONQUIST, F., HUELSENBECK, J.P. (2003): MrBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics*, 19, 1572 – 1574. DOI: 10.1093/bioinformatics/btg180

RYZHIKOV, K.M., SHARPILO, V.P., SHEVTSCHENKO, N.N. (1980): [Helminthes of Amphibia of fauna USSR] Moscow, Nauka. pp. 1 – 278 (In Russian)

SEY, O. (1991): CRC Handbook of the zoology of Amphistomes. CRC Press, Inc., Boca Raton, Florida. 480 p.

SKRJABIN, K.I. (1949): Podotriad Paramphistomatata (Szidat, 1936) Skrjabin et Schulz, 1937 [Suborder Paramphistomatata (Szidat, 1936) Skrjabin et Schulz, 1937]. In: SKRJABIN K. I. *Trematody zhivotnykh i cheloveka. Osnovy trematodologii* [*Trematodes of animals and humans*]. Moscow, Leningrad, AN SSSR Publ. Volume 3, pp. 1 – 623 (In Russian)

TAMURA, K., PETERSON, D., PETERSON, N., STECHER, G., NEI, M., KU-MAR, S. (2011): MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.*, 28, 2731 – 2739. DOI: 10.1093/molbev/msr121

TKACH, V.V., LITTLEWOOD, D.T.J., OLSON, P.D., KINSELLA, J.M., SWIDERSKI, Z. (2003): Molecular phylogenetic analysis of the Microphalloidea Ward, 1901 (Trematoda: Digenea). *Syst. Parasitol.*, 56, 1 – 15. DOI: 10.1023/A:1025546001611

TRUETT, G.E., HEEGER, P., MYNATT, R.L., TRUETT, A.A., WALKER, J.A., WARMAN, M.L. (2000): Preparation of PCR-Quality Mouse Genomic DNA with Hot Sodium Hydroxide and Tris (HotSHOT). *BioTechniques*, 29, 52 – 54

WAESCHENBACH, A., WEBSTER, B.L., BRAY, R.A., LITTLEWOOD, D.T.J. (2007): Added resolution among ordinal level relationships of tapeworms (Platyhelminthes: Cestoda) with complete small and large subunit nuclear ribosomal RNA genes. *Mol Phyl Evol* 45, 311 – 325. DOI: 10.1016/j.ympev.2007.03.019

YAMAGUTI, S. (1936): Studies on the helminth fauna of Japan. Part 14. Amphibian Trematodes. *Jap. J. Zool.*, 6, 559 – 561

YAMAGUTI, S. (1941): Zur Entwicklungsgeschichte von Diplodiscus

amphichrus japonicus Yamaguti, 1936 [History of the development of *Diplodiscus amphichrus japonicus* Yamaguti, 1936]. *Z. Parasitenkd.*, 11, 652 – 656. DOI: 10.1007/BF02120746 (In German) ZHAO, J.H., WANG, S.S., TU, G.J., ZHOU, Y.K., WU, X.B. (2016): Morphological and molecular characterization of *Ortleppascaris sinensis* sp. nov. (Nematoda: Ascaridoidea) from the Chinese alligator *Alligator sinensis. J. Helminthol.*, 90,303 – 311. DOI: 10.1017/ S0022149X15000255