

## ORIGINAL ARTICLE

# Mitigation of diazinon-induced cardiovascular and renal dysfunction by gallic acid

Temitayo Olabisi AJIBADE<sup>1</sup>, Ademola Adetokunbo OYAGBEMI<sup>1</sup>, Temidayo Olutayo OMOBOWALE<sup>2</sup>, Ebinoluwa Racheal ASENUGA<sup>3</sup>, Jeremiah Moyinoluwa AFOLABI<sup>2</sup>, Adeolu Alex ADEDAPO<sup>1</sup>

<sup>1</sup> Department of Veterinary Physiology, Biochemistry and Pharmacology, Faculty of Veterinary Medicine, University of Ibadan, Nigeria

<sup>2</sup> Department of Veterinary Medicine, Faculty of Veterinary Medicine, University of Ibadan, Nigeria

<sup>3</sup> Department of Veterinary Physiology and Biochemistry, University of Benin, Benin City, Nigeria

ITX090216A03 • Received: 20 July 2015 • Revised: 31 March 2016 • Accepted: 04 April 2016

## ABSTRACT

Studies of the link between environmental pollutants and cardiovascular dysfunction, neglected for decades, have recently provided new insights into the pathology and consequences of these killers. In this study, rats were divided into four groups, each containing 10 rats. The rats in group one served as controls and were administered normal saline, whereas the rats in group two were orally gavaged with 3 mg/kg of diazinon (DZN) alone for twenty one consecutive days. The rats in groups 3 and 4 were administered respective 60 mg/kg and 120 mg/kg gallic acid (GA) in addition to DZN for twenty one consecutive days. Exposure of rats to diazinon significantly ( $p < 0.05$ ) reduced the activities of superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx), glutathione-S-transferase (GST) and reduced glutathione (GSH) content. Malondialdehyde, hydrogen peroxide ( $H_2O_2$ ) and nitric oxide (NO) contents were also significantly ( $p < 0.05$ ) elevated following DZN exposure. DZN further caused a significant ( $p < 0.05$ ) decrease of heart rate and QT interval prolongation. Hematologic analysis revealed significant reduction ( $p < 0.05$ ) in packed cell volume (PCV), hemoglobin concentration (Hb), red blood cell (RBC) count, and total white blood cell count of rats administered only DZN. Observations in this study suggest a modulatory role of gallic acid in diazinon-induced anemia and associated cardiovascular dysfunction in rats. Treatment with gallic acid reversed the oxidative stress markers studied, increased the antioxidant defence system and reduced deleterious effects on hematological parameters in rats. Pathologic findings of the heart and kidney were also found to be lessened.

**KEY WORDS:** diazinon; gallic acid; heart; kidney; oxidative stress; cardiovascular dysfunction

## Introduction

The use of and exposure to pesticides in developing countries is becoming increasingly uncontrollable, making the environment more dangerous to the living system (Mostafalou & Abdollahi, 2013). The increasing incidence of hypertension, coronary heart disease and renal failure has become distressing over the last two decades (Sexton & Salinas, 2014). There is a probability of positive correlation between cardiorenal dysfunction and environmental pollutants such as organophosphate (Hutcheson & Rocic, 2012). Exposure to pesticides can be environmental,

occupational, intentional or suicidal in nature (Paudyal, 2008). Since exposure of humans and animals to these agents is inevitable, it constitutes a major global health risk (Kumar *et al.*, 2008).

Diazinon (DZN) is an organophosphate insecticide used extensively in agriculture for control of pests (Jafari *et al.*, 2012). It is used also in veterinary medicine for prophylactic and therapeutic management of infestation with ticks, mites and fleas. Unfortunately, DZN is highly toxic to man and animals. The primary mechanism of toxicity for DZN to animals involves inhibition of acetylcholinesterase (Kousba *et al.*, 2007). Induction of free radical production and oxidative stress was reported in vital organs of rats, including the brain, heart and spleen (Jafari *et al.*, 2012). DZN causes deleterious effects on hematological and serum biochemical parameters in rodents (Kalender *et al.*, 2005). Decreased red blood cell count, hemoglobin concentration and packed cell volume were reported in DZN treated mice (El-Shenawy *et al.*, 2009).

Correspondence address:

**Dr. Ademola Adetokunbo Oyagbemi**

Department of Veterinary Physiology, Biochemistry and Pharmacology,  
Faculty of Veterinary Medicine, University of Ibadan, Nigeria  
TEL.: +234833639776

E-MAIL: ademola.oyagbemi778@gmail.com; aa.oyagbemi@ui.edu.ng

Acute organophosphate poisoning (OP) was reported to increase creatinine kinase and lactate dehydrogenase levels accompanied by myocardial necrosis (Repplinger *et al.*, 2014; Juliana *et al.*, 2009). Toxicities associated with OP may manifest as sinus tachycardia, sinus bradycardia, hypertension, hypotension, impaired heart rate and force of contraction (Juliana *et al.*, 2009). ECG changes associated with OP include prolonged QTc interval, ST segment elevation, low amplitude T waves, and prolonged PR interval (Yusuf *et al.*, 2009). Suicidal use of OP was reported by Hulse *et al.* (2014) and OP compounds have been linked with the highest incidence of suicidal poisoning (Tsai *et al.*, 2014). The typical toxidrome in OP poisoning comprises salivation, lacrimation, urination, defecation, gastric cramps, emesis, collectively known as 'SLUDGE' symptoms (Peter *et al.*, 2014). Furthermore, high frequency of neurologic manifestations involving parasympathetic hyperactivity due to the accumulated acetylcholine (ACh) resulting from acetyl cholinesterase (AChE) inhibition has been documented (Rastogi *et al.*, 2010).

Different treatment regimens for OP have been adopted for decades to circumvent, delay or prevent toxicities associated with OP, with timely intervention yielding positive results (Moon *et al.*, 2015; Iyer *et al.*, 2015; Sun *et al.*, 2015; Kumar *et al.*, 2014; Buckley *et al.*, 2005; Sungur & Güven, 2001). Buckley *et al.* (2011) reported standard treatment to involve administration of intravenous atropine and oxime to reactivate inhibited acetylcholinesterase.

The use of medicinal plants to ameliorate various disease conditions has gained global acceptance in the last two decades. Humans and animals are constantly exposed to reactive oxygen species (ROS) and reactive nitrogen species (RNS) that are generated daily by endogenous and exogenous sources (Bak and Weerapana, 2015). These dangerous species are generated via different mechanisms/reactions, including mitochondrial electron transport chain, NADPH-oxidases, xanthine oxidase, metal dependent and independent pathways. Furthermore, ROS and RNS can damage the DNA, RNA and proteins and therefore serve as mediators of many disease conditions (Phaniendra *et al.*, 2015; Kielland *et al.*, 2009). In contrast, antioxidants of natural or synthetic origin can quench ROS, RNS and their products, thereby maintaining balanced redox signalling. The imbalance in the production of ROS and the antioxidant defence system is known as oxidative stress (Valko *et al.*, 2006).

Gallic acid (GA) is a natural phenolic agent endowed with antioxidant, anti-inflammatory, anticancer, and free radical scavenging activities (Subramanian *et al.*, 2014; Cedó *et al.*, 2014; Liang *et al.*, 2014; Nair & Nair, 2013; Bak *et al.*, 2013; Verma *et al.*, 2013). Furthermore, orally administered antioxidant agents are currently under intensive scientific investigation for their roles in the amelioration and/or prevention of wide varieties of cardiovascular diseases (Faure *et al.*, 2007). However, few reports exist on the amelioration of DZN-induced hemotoxicity and cardiovascular dysfunction with orally administered antioxidants. This study was designed to evaluate the presumed protective effect of GA on DZN-induced

cardiorenal dysfunction in an experimental animal model and to assess the mechanism of its action.

## Materials and methods

### Chemicals

Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), hydrochloric acid, sulphuric acid, xylene orange, sodium hydroxide, potassium iodide, reduced glutathione (GSH), potassium dichromate, O-dianisidine, sodium potassium tartrate, copper sulphate, glacial acetic acid, ethanol, sodium azide, 2-dichloro-4-nitrobenzene (CDNB), thiobarbituric acid (TBA), trichloroacetic acid, Ellman's reagent (DTNB), ammonium ferrous sulphate, and sorbitol were purchased from Sigma (St Louis, MO, USA). All other chemicals used were of analytical grade and obtained from British Drug Houses (Poole, Dorset, UK).

### Care of animals

All the experimental animals received humane care according to the criteria outlined in the Guide for the Care and the Use of Laboratory Animals prepared by the National Academy of Science and published by the National Institute of Health. The ethics regulations were followed in accordance with national and institutional guidelines for the protection of the animals' welfare during experiments according to Public Health Service (PHS, 1996).

### Experimental design

In this study, forty male Wistar rats with the weight range of 130–180 g were used. The rats were divided into four groups each containing 10 rats. Rats in group one served as controls and were administered normal saline, whereas rats in group two were orally gavaged with 3 mg/kg of diazinon (DZN) alone for twenty one consecutive days. The rats in groups 3 and 4 were co-administered 60 mg/kg and 120 mg/kg gallic acid (GA) with DZN for twenty one consecutive days. Rats were sacrificed 24 hours after the last administration.

### Blood collection, serum preparation and hematological analysis

From each rat, fresh whole blood (5 ml) was collected through the retro-orbital venous plexus into heparinized bottles and 2 ml were used for hematological analysis. Hematological analysis was done according to the instruction manual of an automated hematology analyzer (MINDRAY 3000) for the determination of packed cell volume (PCV), hemoglobin concentration (Hb), red blood cell count (RBC), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC), and white blood cell counts. The remaining three milliliters (3 ml) of blood were also collected into sterile tubes and left for about 30 minutes to clot. The clotted blood was thereafter centrifuged at 4000 rpm for 10 minutes. Serum was harvested into sample bottles and stored at –20°C till the time of analysis. The animals were not anesthetized.

### Cardiac and renal tissue preparation

The rats were sacrificed 24 hours after the last administration. The hearts and the kidney (left and right) were removed, rinsed in 1.15% KCl and homogenized in potassium phosphate buffer (0.1 M, pH 7.4), centrifuged at 12 000 g for 15 minutes to obtain the post mitochondrial fraction (PMF). The obtained PMF of the heart and kidneys was subsequently stored at  $-20^{\circ}\text{C}$  until the time of use.

### Biochemical assays

#### *Determination of cardiac and renal catalase (CAT) activity*

Catalase (CAT) activity was determined according to the method of Sinha (1972). Briefly, 1 ml portion from the reaction mixture – 2 ml of  $\text{H}_2\text{O}_2$  solution, 2.5 ml of 0.01 M potassium phosphate buffer (pH 7.0) – and 1 ml of properly diluted PMF was blown into 1 ml dichromate/acetic acid solution by a gentle swirl at room temperature at 60-second intervals for 3 times. The mixture was incubated in the water bath at  $100^{\circ}\text{C}$  for 10 minutes. The absorbance was read at 570 nm using distilled water as blank. One unit of CAT activity represents the amount of enzyme required to decompose  $1\ \mu\text{mol}$  of  $\text{H}_2\text{O}_2$ /minute.

#### *Determination of cardiac and renal superoxide dismutase (SOD) activity*

Superoxide dismutase (SOD) was determined by the method of Misra and Fridovich (1972) with modification from our laboratory (Oyagbemi *et al.*, 2015; Omobowale *et al.*, 2014), giving credence to the validation of the method. Briefly, 100 mg of epinephrine was dissolved in 100 ml distilled water and acidified with 0.5 ml concentrated hydrochloric acid. This preparation prevents oxidation of epinephrine and is stable for 4 weeks. 0.01 ml of cardiac or renal PMF was added to 2.5 ml 0.05 M carbonate buffer (pH 10.2) followed by the addition of 0.3 ml of 0.3 mM adrenaline. The increase in absorbance at 480 nm was monitored every 30 seconds for 150 seconds. One unit of SOD activity was given as the amount of SOD necessary to cause 50% inhibition of the auto-oxidation of adrenaline to adrenochrome during 1 minute.

#### *Determination of cardiac and renal reduced glutathione (GSH)*

The cardiac and renal reduced glutathione (GSH) was estimated by the method of Jollow *et al.* (1974). Briefly, 0.5 ml of 4% sulfosalicylic acid (precipitating agent) was added to 0.5 ml of PMF and centrifuged at 4000 rpm for 5 minutes. To 0.5 ml of the resulting supernatant, 4.5 ml of Ellman's reagent (0.04 g of DTNB in 100 ml of 0.1 M phosphate buffer, pH 7.4) was added. The absorbance was read at 412 nm against distilled water as blank.

#### *Determination of cardiac and renal glutathione peroxidase (GPx) activity*

The cardiac and renal glutathione peroxidase (GPx) activity was also measured according to Buetler *et al.* (1963). The reaction mixtures contained 0.5 ml of potassium phosphate buffer (pH 7.4), 0.1 ml of sodium azide, 0.2 ml of GSH solution, 0.1 ml of  $\text{H}_2\text{O}_2$ , 0.5 ml of PMF and 0.6 ml of distilled water. The mixture was incubated in the water bath at  $37^{\circ}\text{C}$  for 5 minutes and 0.5 ml of

trichloroacetic acid (TCA) was added and centrifuged at 4000 rpm for 5 mins. One ml of the supernatant was taken and 2 ml of  $\text{K}_2\text{PHO}_4$  and 1 ml of Ellman's reagent were added. The absorbance was read at 412 nm using distilled water as blank.

#### *Determination of cardiac and renal thiobarbituric acid reactive substances (TBARS)*

Thiobarbituric acid reactive substance (TBARS) was quantified as malondialdehyde (MDA) in the cardiac and renal PMF. The MDA was determined according to the method of Varshney and Kale (1990). To 1.6 ml of Tris-KCl, 0.5 ml of 30% TCA, 0.4 ml of samples and 0.5 ml of 0.75% thiobarbituric acid (TBA) prepared in 0.2 M HCl were added. The reaction mixture was incubated in the water bath at  $80^{\circ}\text{C}$  for 45 minutes, cooled on ice and centrifuged at 4000 rpm for 15 minutes. The absorbance was measured against a blank of distilled water at 532 nm. Lipid peroxidation in units/mg protein was calculated with a molar extinction coefficient of  $1.56 \times 10^5\ \text{M/Cm}$ .

#### *Measurement of cardiac and renal nitric oxide (NO) contents*

Cardiac and renal nitric oxide (NO) was measured as described by Olaleye *et al.* (2007) by indirectly measuring the nitrite concentration. After incubation at room temperature for 20 minutes, the absorbance at 540 nm was measured by spectrophotometer. The concentration of nitrite in the sample was determined from a sodium nitrite ( $\text{NaNO}_2$ ) standard curve and was expressed as  $\mu\text{mol}$  nitrite/mg protein.

#### *Measurement of cardiac and renal hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) generation*

Hydrogen peroxide generation was determined according to Woff (1994). To 2.5 ml of 0.1 M potassium phosphate buffer (pH 7.4), 0.250 ml of ammonium ferrous sulphate (AFS), 0.1 ml of sorbitol, 0.1 ml of xylenol orange (XO), 0.025 ml of  $\text{H}_2\text{SO}_4$  and 0.050 ml of cardiac or renal PMF was added. The mixture was mixed thoroughly by vortexing till it foamed and a light pink color of the reaction mixture was observed. The reaction mixture was subsequently incubated at room temperature for 30 minutes. The absorbance was assessed at 560 nm, using distilled water as blank. The hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) generated was extrapolated from the hydrogen peroxide standard curve.

#### *Determination of serum aminotransferases (ALT and AST) and alkaline phosphatase (ALP)*

The activities of aspartate aminotransferase (AST), alanine aminotransferase (ALT) and alkaline phosphatase (ALP) were determined according to Reitma and Frankel (1957), using the Randox Kit (Randox Laboratories Limited, UK).

#### *Protein determination*

Protein concentrations were determined as described by Gornal *et al.* (1949). Briefly, 1 ml of diluted serum was added to 3 ml of the biuret reagent. The reaction mixture was incubated at room temperature for 30 minutes. The mixture was thereafter read with a spectrophotometer

at 540 nm using distilled water as blank. The final value for total protein was extrapolated from the total protein standard curve.

### Electrocardiogram

Standard lead II electrocardiogram was recorded in conscious rats using a 7-lead ECG machine (EDAN VE-1010, Shanghai, China). From the electrocardiogram, parameters such as heart rate, P-wave duration, PR-interval, QRS duration, R-amplitude, QT segment and Bazett's correction of the QT interval were determined.

### Histopathology

Small pieces of heart and kidney tissues were collected in 10% formal saline buffer for proper fixation. These tissues were processed and embedded in paraffin wax. Sections of 5–6  $\mu\text{m}$  in thickness were made and stained with hematoxylin and eosin for histopathological examination (Drury *et al.*, 1976).

### Statistical analysis

All values are expressed as mean $\pm$ S.D. One-way ANOVA with Dunnett's post-test was also performed using GraphPad Prism version 4.00. The level of statistical significance was considered as  $p < 0.05$ .

## Results

In this study, administration of DZN in the dose of 3mg/kg led to a significant ( $p < 0.05$ ) increase in cardiac malondialdehyde (MDA) content, hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) generated and nitric oxide (NO) level, whereas co-administration of DZN and GA reversed the given

markers of oxidative and nitrosative stress (Table 1). Similarly, rats exposed to a toxic dose of DZN showed a significant ( $p < 0.05$ ) increase in renal and cardiac MDA content and hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) generation compared to controls (Tables 1 and 2). Nitric oxide (NO) level, however, increased significantly in renal tissues but it significantly decreased in cardiac tissues (Tables 1 and 2). Furthermore, renal markers of oxidative stress were significantly decreased following GA (60 & 120 mg/kg) co-administered with DZN.

Our data showed that both cardiac and renal glutathione (GSH) reduced significantly following DZN treatment (Tables 3 and 4). The enzymatic antioxidants glutathione peroxidase (GPx) and glutathione-S-transferase (GST) decreased significantly in both cardiac and renal tissues in rats treated with DZN alone (Tables 3 and 4). The administration of GA dose-dependently improved the antioxidant defence system and attenuated the elevated markers of oxidative stress. The activities of cardiac superoxide dismutase (SOD) and catalase (CAT) activity decreased significantly following administration of diazinon compared to the control group (Figures 1 and 2). In contrast, rats co-administered GA (60 and 120 mg/kg) showed significantly improved activity of SOD as compared to DZN treatment only (Figures 1 and 2). Similar results were obtained in renal tissues with a significant reduction in both SOD and CAT activity (Figures 3 and 4). The cardiac SOD responded less than did the renal SOD following treatment with GA (Figures 1 and 3). DZN treatment also significantly reduced the antioxidant activity of catalase (CAT) compared to the control, whereas co-administration of DZN with GA (60 and 120 mg/kg) led to a significant increase in cardiac and renal CAT activity (Figures 2 and 4).

**Table 1.** Effect of Gallic acid on Diazinon-induced cardiac oxidative stress (Mean $\pm$ S.D.).

Experimental Groups (n=10)	Control	Diazinon (3 mg/kg)	Diazinon + Gallic acid (60 mg/kg)	Diazinon + Gallic acid (120 mg/kg)
<sup>a</sup> LPO	9.65 $\pm$ 0.32	16.73 $\pm$ 0.38*	7.60 $\pm$ 0.43*,#	7.28 $\pm$ 0.37*,#
<sup>b</sup> $\text{H}_2\text{O}_2$ generated	15.33 $\pm$ 0.26	16.00 $\pm$ 0.35*	15.41 $\pm$ 0.14#	15.38 $\pm$ 0.18#
<sup>c</sup> Nitric oxide (NO)	0.25 $\pm$ 0.03	0.19 $\pm$ 0.04*	0.19 $\pm$ 0.02*,#	0.17 $\pm$ 0.03*,#

Superscripts (\*, #) indicate significant difference at  $p < 0.05$ . Superscript (\*) indicates significant difference when Diazinon (3 mg/kg), Diazinon (3 mg/kg) + Gallic acid (60 mg/kg) and Diazinon (3 mg/kg) + Gallic acid (120 mg/kg) are compared to the control. Superscript (#) indicates significant difference when Diazinon (3 mg/kg) + Gallic acid (60 mg/kg) and Diazinon (3 mg/kg) + Gallic acid (120 mg/kg) are compared to the Diazinon (3 mg/kg).

<sup>a</sup>MDA/LPO (Malondialdehyde;  $\mu\text{mole}$  of MDA formed/mg protein), <sup>b</sup> $\text{H}_2\text{O}_2$  (Hydrogen peroxide generation;  $\mu\text{mole}$ /mg protein), <sup>c</sup>NO (Nitric oxide;  $\mu\text{mole}$ /L)

**Table 2.** Effect of Gallic acid on Diazinon-induced renal oxidative stress (Mean $\pm$ S.D.).

Experimental Groups (n=10)	Control	Diazinon (3 mg/kg)	Diazinon + Gallic acid (60 mg/kg)	Diazinon + Gallic acid (120 mg/kg)
<sup>a</sup> LPO	6.42 $\pm$ 0.65	9.47 $\pm$ 0.51*	8.04 $\pm$ 0.91*,#	8.82 $\pm$ 0.23*
<sup>b</sup> $\text{H}_2\text{O}_2$ generated	15.44 $\pm$ 0.13	17.13 $\pm$ 0.18*	15.94 $\pm$ 0.24*	15.42 $\pm$ 0.14#
<sup>c</sup> Nitric oxide (NO)	0.40 $\pm$ 0.03	1.31 $\pm$ 0.25*	0.88 $\pm$ 0.05*,#	0.71 $\pm$ 0.03*,#

Superscripts (\*, #) indicate significant difference at  $p < 0.05$ . Superscript (\*) indicates significant difference when Diazinon (3 mg/kg), Diazinon (3 mg/kg) + Gallic acid (60 mg/kg) and Diazinon (3 mg/kg) + Gallic acid (120 mg/kg) are compared to the control. Superscript (#) indicates significant difference when Diazinon (3 mg/kg) + Gallic acid (60 mg/kg) and Diazinon (3 mg/kg) + Gallic acid (120 mg/kg) are compared to the Diazinon (3 mg/kg).

<sup>a</sup>MDA/LPO (Malondialdehyde;  $\mu\text{mole}$  of MDA formed/mg protein), <sup>b</sup> $\text{H}_2\text{O}_2$  (Hydrogen peroxide generation;  $\mu\text{mole}$ /mg protein), <sup>c</sup>NO (Nitric oxide;  $\mu\text{mole}$ /mg protein)



**Table 3.** Effect of Gallic acid on Diazinon on cardiac non-enzymic and enzymic antioxidant defence system (Mean±S.D.).

Experimental Groups (n=10)	Control	Diazinon (3 mg/kg)	Diazinon + Gallic acid (60 mg/kg)	Diazinon + Gallic acid (120 mg/kg)
<sup>a</sup> GSH	60.67±0.14	60.00±0.25*	61.17±0.29*,#	61.19±0.13*,#
<sup>b</sup> GPx	43.75±0.18	37.77±2.63*	38.43±1.48*	39.37±1.79*
<sup>c</sup> GST	0.16±0.008	0.13±0.006*	0.17±0.002*,#	0.14±0.011*,#

Superscripts (\*, #) indicate significant difference at  $p<0.05$ . Superscript (\*) indicates significant difference when Diazinon (3 mg/kg), Diazinon (3 mg/kg) + Gallic acid (60 mg/kg) and Diazinon (3 mg/kg) + Gallic acid (120 mg/kg) are compared to the control. Superscript (#) indicates significant difference when Diazinon (3 mg/kg) + Gallic acid (60 mg/kg) and Diazinon (3 mg/kg) + Gallic acid (120 mg/kg) are compared to the Diazinon (3 mg/kg).


<sup>a</sup>GSH (Reduced glutathione;  $\mu$ mole/mg protein), <sup>b</sup>GPx (Glutathione peroxidase; units/mg protein), <sup>c</sup>GST (Glutathione S-transferase; mmole1-chloro-2, 4-dinitrobenzene-GSH complex formed/min/mg protein).

**Table 4.** Effect of Gallic acid on Diazinon on renal non-enzymic and enzymic antioxidant defence system (Mean±S.D.).


Experimental Groups (n=10)	Control	Diazinon (3 mg/kg)	Diazinon + Gallic acid (60 mg/kg)	Diazinon + Gallic acid (120 mg/kg)
<sup>a</sup> GSH	64.87±0.18	64.25±0.25*	64.33±0.29*	64.63±0.18#
<sup>b</sup> GPx	37.59±0.89	30.53±0.67*	33.18±1.01*,#	33.02±0.74*,#
<sup>c</sup> GST	0.209±0.031	0.131±0.012*	0.191±0.015#	0.194±0.008#

Superscripts (\*, #) indicate significant difference at  $p<0.05$ . Superscript (\*) indicates significant difference when Diazinon (3 mg/kg), Diazinon (3 mg/kg) + Gallic acid (60 mg/kg) and Diazinon (3 mg/kg) + Gallic acid (120 mg/kg) are compared to the control. Superscript (#) indicates significant difference when Diazinon (3 mg/kg) + Gallic acid (60 mg/kg) and Diazinon (3 mg/kg) + Gallic acid (120 mg/kg) are compared to the Diazinon (3 mg/kg).

<sup>a</sup>GSH (Reduced glutathione;  $\mu$ mole/mg protein), <sup>b</sup>GPx (Glutathione peroxidase; units/mg protein), <sup>c</sup>GST (Glutathione S-transferase; mmole1-chloro-2, 4-dinitrobenzene-GSH complex formed/min/mg protein).



**Figure 1.** Effect of diazinon toxicity on cardiac superoxide dismutase (SOD) activity (Mean±S.D; n=10). Superscripts (\*, #) indicate significant difference at  $p<0.05$ . Superscript (\*) indicates significant difference when diazinon (3 mg/kg), diazinon (3 mg/kg) + gallic acid (60 mg/kg), and diazinon (3 mg/kg) + gallic acid (120 mg/kg) are compared to the control. Superscript (#) indicates significant difference when diazinon (3 mg/kg) + gallic acid (60 mg/kg) and diazinon (3 mg/kg) + gallic acid (120 mg/kg) are compared to diazinon (3 mg/kg).



**Figure 2.** Effect of diazinon toxicity on cardiac catalase (CAT) activity (Mean±S.D; n=10). Superscripts (\*, #) indicate significant difference at  $p<0.05$ . Superscript (\*) indicates significant difference when diazinon (3 mg/kg), diazinon (3 mg/kg) + gallic acid (60 mg/kg) and diazinon (3 mg/kg) + gallic acid (120 mg/kg) are compared to the control. Superscript (#) indicates significant difference when diazinon (3 mg/kg) + gallic acid (60 mg/kg) and diazinon (3 mg/kg) + gallic acid (120 mg/kg) are compared to diazinon (3 mg/kg).


#### Aminotransferases and alkaline phosphatase

In addition, the administration of DZN led to a significant increase in serum alanine aminotransferase (ALT), aspartate aminotransferase (AST) and alkaline phosphatase (ALP), as shown in table 5. Treatment with GA (60 and 120mg/kg) however significantly lowered these enzymes (ALT, AST and ALP) in rats compared to those on DZN alone and exhibited values close to normal compared to the control (Table 5). The toxicity of DZN was also demonstrated on cardiac and renal tissues. The toxic dose of


DZN significantly increased the heart and kidney weight compared to the control and other treatment groups (Table 6).

#### Hematological parameters

In this study, values observed for the packed cell volume (PCV), hemoglobin concentration (Hb), and red blood cell (RBC) count, were significantly ( $p<0.05$ ) lowered in rats administered DZN alone and in those that received DZN and 60mg/kg GA combination, compared to the control



**Figure 3.** The effect of diazinon toxicity on renal superoxide dismutase (SOD) activity (Mean±S.D; n=10). Superscripts (\*, #) indicate significant difference at  $p<0.05$ . Superscript (\*) indicates significant difference when diazinon (3 mg/kg), diazinon (3 mg/kg) + gallic acid (60 mg/kg) and diazinon (3 mg/kg) + gallic acid (120 mg/kg) are compared to the control. Superscript (#) indicates significant difference when diazinon (3 mg/kg) + gallic acid (60 mg/kg) and diazinon (3 mg/kg) + gallic acid (120 mg/kg) are compared to diazinon (3 mg/kg).



**Figure 4.** Effect of diazinon toxicity on renal catalase (CAT) activity (Mean±S.D; n=10). Superscripts (\*, #) indicate significant difference at  $p<0.05$ . Superscript (\*) indicates significant difference when diazinon (3 mg/kg), diazinon (3 mg/kg) + gallic acid (60 mg/kg) and diazinon (3 mg/kg) + gallic acid (120 mg/kg) are compared to the control. Superscript (#) indicates significant difference when diazinon (3 mg/kg) + gallic acid (60 mg/kg) and diazinon (3 mg/kg) + gallic acid (120 mg/kg) are compared to diazinon (3 mg/kg).

**Table 5.** Effect of Diazinon on serum aminotransferases and phosphatase (Mean±S.D.).

Experimental Groups (n=10)	Control	Diazinon (3 mg/kg)	Diazinon + Gallic acid (60 mg/kg)	Diazinon + Gallic acid (120 mg/kg)
ALT (U/L)	0.12±0.02	1.41±0.04*	0.58±0.07*, #	0.55±0.04*, #
AST (U/L)	161.77±1.99	175.28±2.60*	157.21±4.47#	121.42±7.31*, #
ALP (U/L)	289.80±19.52	358.80±39.03*	294.40±15.93#	266.80±31.87#

Superscripts (\*, #) indicate significant difference at  $p<0.05$ . Superscript (\*) indicates significant difference when Diazinon (3 mg/kg), Diazinon (3 mg/kg) + Gallic acid (60 mg/kg) and Diazinon (3 mg/kg) + Gallic acid (120 mg/kg) are compared to the control. Superscript (#) indicates significant difference when Diazinon (3 mg/kg) + Gallic acid (60 mg/kg) and Diazinon (3 mg/kg) + Gallic acid (120 mg/kg) are compared to the Diazinon (3 mg/kg).

**Table 6.** Effect of Diazinon on heart and kidney weight (Mean±S.D.).

Experimental Groups (n=10)	Control	Diazinon (3 mg/kg)	Diazinon + Gallic acid (60 mg/kg)	Diazinon + Gallic acid (120 mg/kg)
Heart weight (gram)	0.29±0.02	0.41±0.02*	0.35±0.03*, #	0.41±0.03#
Kidney weight (gram)	0.56±0.02	0.74±0.04*	0.77±0.03*	0.81±0.04#

Superscripts (\*, #) indicate significant difference at  $p<0.05$ . Superscript (\*) indicates significant difference when Diazinon (3 mg/kg), Diazinon (3 mg/kg) + Gallic acid (60 mg/kg) and Diazinon (3 mg/kg) + Gallic acid (120 mg/kg) are compared to the control. Superscript (#) indicates significant difference when Diazinon (3 mg/kg) + Gallic acid (60 mg/kg) and Diazinon (3 mg/kg) + Gallic acid (120 mg/kg) are compared to the Diazinon (3 mg/kg).

group. In contrast, reductions in the values of these hematological parameters were not significant in rats treated with 120 mg/kg GA and DZN. Furthermore, the PCV and hemoglobin concentration of these rats significantly increased relative to that of rats administered only DZN. Concerning hematometric indices, the following observations were also recorded: significant increase in the mean corpuscular volume (MCV) in rats administered only DZN and in rats with DZN and 60 mg/kg GA combination compared to the control group; significant decrease in MCV of rats administered DZN and 120 mg/kg, relative to that of rats administered DZN alone, and non-significant changes in the mean corpuscular hemoglobin

(MCH), and mean corpuscular hemoglobin concentration (MCHC), as shown in Table 7. The total white blood cell count (TWBC), lymphocyte and neutrophil counts of rats administered DZN alone was significantly ( $p<0.05$ ) decreased relative to controls, but these changes were not significant in groups of rats administered 60 mg/kg and 120 mg/kg DZN. Further, lymphocyte and neutrophil count of these rats increased significantly compared to those of rats administered DZN alone.

#### ECG results

The ECG results showed that DZN treatment led to a significant reduction in heart rate (HR) compared to the

**Table 7.** Effect of Diazinon on haematological parameters of rats.

Parameter	Control	Diazinon	Diazinon+60 mg/kg GA	Diazinon+120 mg/kg
PCV %	38.42±5.39	31.59±1.22 <sup>a</sup>	33.24±1.37 <sup>a</sup>	34.07±2.22 <sup>b*</sup>
Hb g/dl	12.62±2.30	8.11±0.26 <sup>a</sup>	8.11±1.00 <sup>a</sup>	10.37±1.38 <sup>b</sup>
RBC × 10 <sup>12</sup> /l	7.90±1.01	5.37±1.44 <sup>a</sup>	5.01±1.13 <sup>a</sup>	6.84±1.00
MCV fl	48.63±3.12	60.40±2.03 <sup>a</sup>	66.35±7.09 <sup>a</sup>	49.81±1.08 <sup>b</sup>
MCH pg	15.97±1.02	15.51±0.95	16.19±1.44	15.16±0.27
MCHC g/dl	32.85±3.21	25.67±1.06	24.39±2.41	30.43±1.01
TWBC × 10 <sup>9</sup> /l	11.50±1.03	8.20±0.04 <sup>a</sup>	10.19±2.34	10.53±1.62
Lymphocytes × 10 <sup>9</sup> /l	6.92±1.15	4.26±1.08 <sup>a</sup>	6.03±0.21	6.11±0.23
Monocytes × 10 <sup>9</sup> /l	0.20±0.10	0.10±0.10	0.10±0.10	0.10±0.10
Neutrophils × 10 <sup>9</sup> /l	4.33±1.82	1.93±0.15 <sup>a</sup>	4.22±0.13 <sup>b</sup>	3.92±1.08 <sup>b</sup>
Eosinophils × 10 <sup>9</sup> /l	0.10±0.01	0.10±0.01	0.10±0.01	0.10±0.01
Basophils × 10 <sup>9</sup> /l	0.01±0.00	0.01±0.00	0.01±0.00	0.01±0.00

<sup>a</sup>Significant as compared groups 2, 3 and 4 with group 1 ( $p<0.05$ )

<sup>b</sup>Significant as compared groups 3 and 4 with group 2 ( $p<0.05$ )

**Table 8.** Effect of Gallic acid on Diazinon-induced alteration of electrocardiograph (ECG) (Mean±S.D.).

Experimental Groups	Heart rate (rates/min)	P-Duration (ms)	PR (ms)	QRS (ms)	QT (ms)	QTc (ms)	R-Amplitude (mv)
Control	286±28.66	26.83±5.78	45.17±6.49	20.00±2.68	77.33±16.60	146.00±12.88	0.20±0.08
Diazinon (3 mg/kg)	231±18.88 <sup>*</sup>	29.33±4.27	49.00±9.36	26.40±4.51 <sup>*</sup>	83.33±11.81	173.25±15.97 <sup>*</sup>	0.33±0.04 <sup>*</sup>
Diazinon + Gallic acid (60 mg/kg)	245.75±5.12 <sup>*, #</sup>	26.25±5.44	47.75±3.59	18.00±3.81 <sup>#</sup>	79.00±9.76	150.33±9.61 <sup>#</sup>	0.22±0.02 <sup>#</sup>
Diazinon + Gallic acid (120 mg/kg)	241.25±7.59 <sup>*</sup>	23.00±2.65	40.25±4.88 <sup>#</sup>	19.75±1.71 <sup>#</sup>	57.25±6.608, <sup>#</sup>	120.67±11.02 <sup>*, #</sup>	0.24±0.08 <sup>#</sup>

Superscripts (\*, #) indicate significant difference at  $p<0.05$ . Superscript (\*) indicates significant difference when Diazinon (3 mg/kg), Diazinon (3 mg/kg) + Gallic acid (60 mg/kg) and Diazinon (3 mg/kg) + Gallic acid (120 mg/kg) are compared to the control. Superscript (#) indicates significant difference when Diazinon (3 mg/kg) + Gallic acid (60 mg/kg) and Diazinon (3 mg/kg) + Gallic acid (120 mg/kg) are compared to the Diazinon (3 mg/kg).

values obtained from control animals and rats administered GA (60 and 120 mg/kg) body weight (Table 8). Similarly, other ECG changes observed included prolongation of QRS, QT and QTc intervals as indicated by significant ( $p<0.05$ ) increase in the values of the these parameters compared to GA treated rats and controls (Table 8).

### Histopathology

The histology of the cardiac tissue showed mild infiltration of inflammatory cells in cardiomyocytes of the DZN treated group (Figure 5). However, administration of GA (60 and 120 mg/kg) reduced the level of cellular infiltration accompanied with no visible lesion on cardiomyocytes (Figure 5). This cellular infiltration might also indicate inflammation. A similar result was obtained from the histology of the kidney of rats exposed to DZN with renal tissue showing infiltration of the perivascular and interstitial spaces with infiltration of inflammatory cells (Figure 6). However, the renal tissues could not recover from the toxicity of DZN, as evidenced by the presence of infiltration of inflammatory cells with tubular degenerative changes (Figure 6). The glomeruli exhibited a normal appearance, with normal urinary


space in rats that obtained DZN co-administered with GA (Figure 6).

### Discussion

The increase in the incidence of organophosphate (OP) poisoning in developing countries can be attributed to lack or inadequate control, to ignorance, poverty and illiteracy. However, abuse and misuse of these killer agents cannot be underestimated. This work sought to understand the relationship between cardiovascular dysfunction and OP and the use of phytonutrients present in medicinal plants. Induction of oxidative stress has been implicated in the pathogenesis of OP (El-Demerdash *et al.*, 2014, Oksay *et al.*, 2013, Messarah *et al.*, 2013). Accumulating evidence is emerging on the benefits of antioxidants that reside in medicinal plants to quench free radicals and oxidative stress associated with OP poisoning (Kristofco *et al.*, 2015, Oğuzhanoglu *et al.*, 2014, Razavi *et al.*, 2013, Hariri *et al.*, 2010).

NO has been extensively documented as a signalling molecule. It has been implicated both in physiology and





**Figure 5.** Group A (control) cardiac tissue shows mild infiltration of inflammatory cell (black arrow) into the myocardium. Group B (diazinon 3 mg/kg) indicates cardiac cell swelling (black arrow) of the cardiac tissue. Group C (diazinon 3 mg/kg + 60 mg/kg gallic acid) shows photomicrographs of cardiac tissue with no significant lesion. Similarly, rats in Group D (diazinon 3 mg/kg + 120 mg/kg gallic acid), show cardiac tissue with no significant lesion. Histologic slides were stained with hematoxylin & eosin (magnification X400).


pathology. Its low level was reported to elicit hypertension, whereas excessive production could also precipitate hypotension. Administration of DZN might cause overproduction of NO in renal tissues and this may be attributed to cardiovascular dysfunction. Surprisingly, administration of GA reversed the excessive production of NO and reduced the tissue level to near normal values.

In parallel, our findings showed that administration of DZN significantly reduced the activity of SOD, CAT, GPx, GST activities and GSH content. SOD is the first line of defence against free radical generation. SOD dismutates superoxide anion radical ( $O_2^{\bullet-}$ ) to hydrogen peroxide ( $H_2O_2$ ). The reduction in the activity of SOD by DZN will allow  $O_2^{\bullet-}$  to accumulate and then to react with NO to generate peroxynitrite ( $ONOO^-$ ), which is more reactive and damaging.  $ONOO^-$  can damage DNA, RNA, lipids and proteins.

From this study, administration of DZN to rats led to significant elevation of MDA and  $H_2O_2$  in both cardiac and renal tissues and of NO only in renal tissues. These

are markers of oxidative and nitrosative stress. On balance, this is an indication of peroxidation of membrane phospholipid via oxidation. The increase in these markers in renal tissues has been associated with oxidative and nitrosative stress, which is an imbalance between generation of reactive oxygen or nitrogen species and the antioxidant defence system. This imbalance favors the production of ROS and RNS rather than the antioxidant defence system. Furthermore, we speculate that DZN enhanced the production of ROS and RNS, reducing thereby the *in vivo* antioxidant capacity in the heart and kidney, vital organs of the body. In parallel, the ROS and RNS generated by DZN could reduce the ability of the kidney to excrete sodium, thus facilitating sodium retention accompanied by hypertension. The functional capacity of the nephron could also be disrupted via reduction in the glomerular filtration rate (GFR) and enhancement of sodium retention. However, co-administration of GA was able to significantly reduce these markers of ROS and RNS. The ability of GA to quench these radicals is





**Figure 6.** Group A (control) shows no visible lesion. Group B (diazinon 3 mg/kg) indicates renal tissue infiltration of the perivascular and interstitial spaces with inflammatory cells (black arrow). Group C (diazinon 3 mg/kg + 60 mg/kg gallic acid) shows the architecture of renal tissue with slight periglomerular inflammation and desquamation of the tubules, whereas in rats in Group D (diazinon 3 mg/kg + 120 mg/kg gallic acid), the renal tissue shows severe infiltration of inflammatory cells (black arrow) with tubular degenerative changes (black arrow). However, the glomeruli look normal with normal urinary space. Histologic slides were stained with hematoxylin & eosin (magnification X400).

an indication of its cardioprotective and renoprotective property. DZN also reduced myocardial contractility and precipitated left ventricular hypertrophy with resultant heart failure. These results thus corroborate the report of Cakici and Akat (2013), Boroushaki *et al.* (2013), Shah and Iqbal (2010), Gokel (2010), Wu *et al.* (2008), Stonys *et al.* (2007), and Agostini and Bianchin (2003) who reported renal failure and hepatic damage following DZN toxicity.

The  $\text{ONOO}^-$  that is formed via reaction between  $\text{O}_2^{\bullet-}$  and NO will reduce the bioavailability of NO and ultimately precipitate hypertension. We have also suggested that DZN could stimulate NADPH oxidase-mediated  $\text{O}_2^{\bullet-}$  production. This pathway was found to be upregulated in hypertension, atherosclerosis and diabetes (Griendling *et al.*, 2000). Although, blood pressure was not recorded in this study, we hypothesized that reduction

in the bioavailability of NO induced hypertension. NO is known to mediate vascular tone, prevent platelet activation, limit leukocyte adhesion to the endothelium, and regulate myocardial contractility. Furthermore, it plays a critical role in the pathogenesis of common cardiovascular disorders, including hypotension, hypertension, and atherosclerosis. Yet co-administration of GA with DZN improved the antioxidant capacity of cardiorenal SOD. Similarly, the activities of CAT and GPx were significantly inhibited following DZN treatment. CAT and GPx are enzymes involved in the second line of defence against ROS and RNS. CAT and GPx catalyze the conversion of  $\text{H}_2\text{O}_2$  to  $\text{H}_2\text{O}$  and  $\text{O}_2$ . However, reduction in the activity of these enzymes will cause accumulation of  $\text{H}_2\text{O}_2$ . The accumulated  $\text{H}_2\text{O}_2$  will disrupt the functional capacity of the heart and kidney with resultant vascular remodeling

and endothelial dysfunction. Thus our study exhibits important mechanisms that underlie the exposure to environmental pollutants such as DZN with a negative impact on the cardiovascular system. Cardiotoxicity associated with DZN toxicity was reported (Pizzurro *et al.*, 2014, Razavi *et al.*, 2014, Watson *et al.*, 2014, Razavi *et al.*, 2013). We further proposed a positive correlation between exposure to OP such as DZN and the incidence of hypertension/hypotension, heart failure, arteriosclerosis and renal failure. Karasu-Minareci *et al.* (2012) reported that OP poisoning could result in myocardial infarction. Interestingly, our data showed that GA administration reversed and increased the activity of CAT and GPx that appeared so far decline by the toxicity of DZN. GST is a phase 2 detoxification and antioxidant enzyme. The activity of GST and its co-factor GSH was reduced by DZN. In contrast, GA by its strong antioxidant capacity increased the activity of GST and GSH content of the heart and the kidney. GSH belongs to the intracellular antioxidant defence system. It is important to note that the ability of GA to increase GSH content might occur through increase in the novel synthesis of GSH. Other observable cardiotoxic effects of DZN in this study included reduction in heart rate accompanied with both QT QTc interval prolongations. Accumulating evidence has shown that QTc interval prolongation is a predictive indicator of the severity of poisoning and is also associated with cardiovascular collapse (Lin *et al.*, 2014; Liu *et al.*, 2012; Anand *et al.*, 2009; Yurumez *et al.*, 2009). However, administration of GA corrected the QT interval prolongation by reducing the QTc interval caused by DZN administration.

We also found that DZN caused increase in the heart and kidney weight. Co-administration of GA was able to reduce the size of the heart yet not of the kidney, as compared to the control, suggesting that the anti-inflammatory property of GA exerts a more profound effect on the heart than on the kidney. We may also speculate that the toxicity of DZN affects more the kidney and that the heart has the capacity to withstand the toxicity of DZN better than the kidney. On balance, DZN toxicity could also reduce cardiac output, thereby affecting normal cardiovascular function and renal perfusion.

An increase in alkaline phosphatase and in the aminotransferases ALT and particularly AST has been associated with cardiotoxicity. The elevation of this enzyme in blood or tissues is known to occur due to membrane breakdown and subsequent leakage into the systemic circulation. An elevated AST level in particular is a pointer of cardiac damage, suggesting that DZN disrupted the membranes covering the heart (pericardium) allowing AST to accumulate in blood. The above results suggest that GA provided a cardioprotective effect through improvement of the cardiorenal antioxidant defence system and its capacity to quench and scavenge ROS and RNS.

Baydin *et al.* (2007) reported a negative correlation between blood cholinesterase and QTc interval in patients with acute organophosphate poisoning.

In this study, the establishment of anemia in the group of rats administered DZN alone can be inferred from the

significant reduction observed in the values of PCV, Hb, and RBC counts of this group compared to the control group. The significant increase in MCV and normal values of MCH and MCHC suggest macrocytic normochromic anemia. This anemia was effectively reversed in rats following co-administration of GA, especially at the 120 mg/kg dose. Observations in this study are similar to the reports of El-Shenawy *et al.* (2009) and Kalender *et al.* (2006) who reported amelioration of DZN-induced hemotoxicity following administration of vitamin E, an orally active antioxidant, in mice and rats. The ameliorative activity of GA on DZN-induced hemotoxicity observed in this study might be due to a direct inhibition of oxidative stress and free radical production in the erythrocyte membrane. GA has been reported to suppress lipid peroxidation in erythrocytes (Nabavi *et al.*, 2013) and to increase the resistance of the erythrocyte membrane to hydrogen peroxide-induced peroxidation (Ramkumar *et al.*, 2014). Inhibition of lipid peroxidation may expectedly lead to amelioration of anemia since lipid peroxidation is an important step before cell membrane blebbing, hemolysis, and consequent shortening of the lifespan of erythrocytes (Betrosian *et al.*, 1995; Bowman *et al.*, 2005).

Another important component of the ameliorative potential of GA on DZN-induced anemia observed in this study was the restoration of reduced hemoglobin concentration to near normal values in rats in which DZN was co-administered to rats with 120 mg/kg body weight GA. Concomitant reduction in hemoglobin and red blood cell counts is sometimes attributable to the depressant effects of toxicants on hemopoietic organs, especially bone marrow, and interference with specific steps in the heme synthetic pathway (Ray, 1992). Thus, results from this study suggest the ability of GA to counteract or ameliorate the toxic effects of DZN on hemopoietic organs and processes. Moreover, the observed decrease in total white blood cell count, lymphopenia and neutropenia in rats administered DZN alone relative to that of the control group, suggests a depression of bone marrow. A potential anti-immunosuppressive effect of GA is reflected in this study by the significantly increased lymphocyte and neutrophil count of rats in which GA was co-administered with DZN, in contrast to those that received DZN only.

The study concludes that GA holds great promise in the amelioration of DZN-induced hemotoxicity and associated cardiovascular dysfunction. On balance then, inclusion of GA-rich feed supplements in diets of humans and animals potentially at risk of DZN exposure is advocated.

## Acknowledgements

The authors acknowledge the technical assistance of our Research Assistant Dr. E. R. Asenuga in this study.

**Conflict of interest.** There is no conflict of interest.



## REFERENCES

- Agostini M, Bianchin A. (2003). Acute renal failure from organophosphate poisoning: a case of success with haemofiltration. *Hum Exp Toxicol* **22**: 165–167.
- Anand S, Singh S, Nahar Saikia U, Bhalla A, Paul Sharma Y, Singh D. (2009). Cardiac abnormalities in acute organophosphate poisoning. *Clin Toxicol (Phila)* **47**: 230–235.
- Bak DW, Weerapana E. (2015). Cysteine-mediated redox signalling in the mitochondria. *Mol Biosyst* **11**: 678–697.
- Bak EJ, Kim J, Jang S, Woo GH, Yoon HG, Yoo YJ, et al. (2013). Gallic acid improves glucose tolerance and triglyceride concentration in diet-induced obesity mice. *Scand J Clin Lab Invest* **73**: 607–614.
- Baydin A, Aygun D, Yazici M, Karatas A, Deniz T, Yardan T. (2007). Is there a relationship between the blood cholinesterase and QTc interval in the patients with acute organophosphate poisoning? *Int J Clin Pract* **61**: 927–930.
- Betrosian A, Balla M, Kafiri G, Kofinas G, Makri R, Kakouri A. (1995). Multiple system organ failure from organophosphate poisoning. *J Clin Toxicol* **33**: 257–260.
- Boroushaki MT, Arshadi D, Jalili-Rasti H, Asadpour E, Hosseini A. (2013). Protective effect of pomegranate seed oil against acute toxicity of diazinon in rat kidney. *Iran J Pharm Res* **12**: 821–817.
- Bowman ZB, Jason DM, David JJ, David CM. (2005). Primaquine-induced haemolytic anaemia: Role of membrane lipid peroxidation and cytoskeleton protein alterations in the haemotoxicity of 5-hydroxylprimaquine. *J Pharmacol Exp Ther* **314**: 838–845.
- Buckley NA, Eddleston M, Li Y, Bevan M, Robertson J. (2011). Oximes for acute organophosphate pesticide poisoning. *Cochrane Database Syst Rev* **16**(2): CD005085. doi: 10.1002/14651858.CD005085.pub2.
- Buckley NA, Eddleston M, Szinicz L. (2005). Oximes for acute organophosphate pesticide poisoning. *Cochrane Database Syst Rev* **25**(1): CD005085.
- Buetler E, Duron O, Kelly BM. (1963). Improved method for the determination of blood glutathione. *J Lab Clin Med* **61**: 882–888.
- Cakici O, Akat E. (2013). Effects of oral exposure to diazinon on mice liver and kidney tissues: biometric analyses of histopathologic changes. *Anal Quant Cytopathol Histopathol* **35**: 7–16.
- Cedó L, Castell-Auví A, Palleràs V, Macià A, Blay M, Ardévol A, et al. (2014). Gallic acid is an active component for the anticarcinogenic action of grape seed procyanidins in pancreatic cancer cells. *Nutr Cancer* **66**: 88–96.
- Drury RA, Wallington EA, Cancerson R. (1976). Carlton's Histopathological Techniques, fourth ed. Oxford University Press, Oxford, London, New York.
- El-Demerdash FM, Nasr HM. (2014). Antioxidant effect of selenium on lipid peroxidation, hyperlipidemia and biochemical parameters in rats exposed to diazinon. *J Trace Elem Med Biol* **28**: 89–93.
- El-Shenawy NS, Al-Eisa RA, El-Salmy F, Salah O. (2009). Prophylactic effect of vitamin E against hepatotoxicity, nephrotoxicity, haematological indices and histopathology induced by diazinon insecticide in mice. *Curr Zool* **55**: 219–226.
- Farombi EO, Tahnteng JG, Agboola AO, Nwankwo JO, Emerole GO. (2000). Chemoprevention of 2-acetylaminofluorene-induced hepatotoxicity and lipid peroxidation in rats by kolaviron--a Garcinia kola seed extract. *Food Chem Toxicol* **38**: 535–541.
- Gokel Y. (2002). Subarachnoid hemorrhage and rhabdomyolysis induced acute renal failure complicating organophosphate intoxication. *Ren Fail* **24**: 867–871.
- Gornal AG, Bardawill JC, David MM. (1949). Determination of serum proteins by means of Biuret reaction. *Journal of Biological Chemistry* **177**: 751–766.
- Griendling KK, Sorescu D, Ushio-Fukai M. (2000). NADPH oxidase. Role in cardiovascular biology and disease. *Circ Res* **86**: 494–501.
- Habig WH, Pabst MJ, Jacoby WB. (1974). Glutathione-S-transferase Activity: The Enzymic step in Mercapturic Acid formation. *J Biol Chem* **249**: 130–139.
- Hariri AT, Moallem SA, Mahmoudi M, Memar B, Hosseinzadeh H. (2010). Subacute effects of diazinon on biochemical indices and specific biomarkers in rats: protective effects of crocin and safranal. *Food Chem Toxicol* **48**: 2803–2808.
- Hulse EJ, Davies JO, Simpson AJ, Sciuto AM, Eddleston M. (2014). Respiratory complications of organophosphorus nerve agent and insecticide poisoning. Implications for respiratory and critical care. *Am J Respir Crit Care Med* **190**: 1342–1354.
- Hulse EJ, Clutton RE, Drummond G, Eddleston M. (2014). Translational toxicological research: investigating and preventing acute lung injury in organophosphorus insecticide poisoning. *J R Army Med Corps* **160**: 191–192.
- Hutcheson R and Rocic P. (2012). The metabolic syndrome, oxidative, environment, and cardiovascular disease. The Great Exploration. *Exp Diabetes Res* **2012**: 271028. .
- Ilyer R, Iken B, Leon A. (2015). Developments in alternative treatments for organophosphate poisoning. *Toxicol Lett* **233**: 200–206.
- Jafari M, Salehi M, Ahmadi S, Asgari A, Abasnezhad M, Hajigholamali M.. (2012). The role of oxidative stress in diazinon-induced tissues toxicity in Wistar and Norway rats. *Toxicol Mech Methods* **22**: 638–647
- Jollow DJ, Mitchell JR, Zampaglione N, Gillette JR. (1974). Bromobenzene-induced liver necrosis. Protective role of glutathione and evidence for 3,4-bromobenzene oxide as the hepatotoxic metabolite. *Pharmacology* **11**(3): 151–69.
- Kalender SA, Ogutcu M, Uzunhisarcikli F, Acikgoz D, Durak Y, Ulusoy Y, et al. (2012). What may be happen after an organophosphate exposure: acute myocardial infarction? *J Forensic Leg Med* **19**: 94–96.
- Kielland A, Blom T, Nandakumar KS, Holmdahl R, Blomhoff R, Carlsen H. (2009). In vivo imaging of reactive oxygen and nitrogen species in inflammation using the luminescent probe L-012. *Free Radic Biol Med* **47**: 760–766.
- Kousba AA, Poet TS, Timchalk C. (2007). Age-related brain cholinesterase inhibition kinetics following in vitro incubation with chlorpyrifos-oxon and diazinon-oxon. *Toxicol Sci* **95**: 147–155.
- Kristofco LA, Du B, Chambliss CK, Berninger JP, Brooks BW. (2015). Comparative pharmacology and toxicology of pharmaceuticals in the environment: diphenhydramine protection of diazinon toxicity in Danio rerio but not Daphnia magna. *AAPS J* **17**: 175–183.
- Kumar LP and Bnnserselvam. (2008). Toxic effects of pesticides. A review on cytogenetic biomonitoring studies. *Facta Universitatis series: Med Biol* **15**: 46–50.
- Kumar MR, Kumar GP, Babu PR, Kumar SS, Subrahmanyam BV, Veeraprasad M, et al. (2014). A retrospective analysis of acute organophosphorus poisoning cases admitted to the tertiary care teaching hospital in South India. *Ann Afr Med* **13**: 71–75.
- Liang W, Li X, Li Y, Li C, Gao B, Gan H, et al. (2014). Gallic acid induces apoptosis and inhibits cell migration by upregulating miR-518b in SW1353 human chondrosarcoma cells. *Int J Oncol* **44**: 91–98.
- Messarah M, Amamra W, Boumendjel A, Barkat L, Bouasla I, Abdenmour C, et al. (2013). Ameliorating effects of curcumin and vitamin E on diazinon-induced oxidative damage in rat liver and erythrocytes. *Toxicol Ind Health* **29**: 77–88.
- Misra HP, Fridovich I. (1972). The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase. *J Biol Chem* **217**: 3170–3175.
- Moon J, Chun B, Song K. (2015). An exploratory study; the therapeutic effects of premixed activated charcoal-sorbitol administration in patients poisoned with organophosphate pesticide. *Clin Toxicol (Phila)* **53**: 119–126.
- Mostafalou S, Abdollahi M. (2013). Pesticides and human chronic diseases: evidences, mechanisms, and perspectives. *Toxicol Appl Pharmacol* **268**: 157–177.
- Nair GG, Nair CK. (2013). Radioprotective effects of gallic acid in mice. *Biomed Res Int* **2013**: 953079.
- Nabavi SF, Habtemariam S, Sureda A, Hajizadeh Moghaddam A, Daglia M, Nabavi SM. (2013). In vivo protective effects of Gallic acid isolated from *Peltiphyllum peltatum* against sodium fluoride-induced oxidative stress in rat erythrocytes. *Arh Hig Rada Toksikol* **64**: 553–539.
- Oğuzhanoglu E, Andaç AC, Tüfek A, Yavuz L, Vural H, Gökalp O. (2014). Protective role of caffeic acid phenethyl ester on serum cholinesterase inhibition by acute exposure to diazinon in rats. *Turk J Med Sci* **44**: 115–120.
- Oksay T, Naziroğlu M, Ergün O, Doğan S, Özatik O, Armağan A, et al. (2013). N-acetyl cysteine attenuates diazinon exposure-induced oxidative stress in rat testis. *Andrologia* **45**: 171–177.
- Olaleye SB, Adaramoye OA, Erigbali PP, Adeniyi OS. (2007). Lead exposure increases oxidative stress in the gastric mucosa of HCl/ethanol-exposed rats. *World J Gastroenterol* **13**: 5121–5126.
- Omobowale TO, Oyagbemi AA, Akinrinde AS, Saba AB, Daramola OT, Ogunpolu BS, et al. (2014). Failure of recovery from lead induced hepatotoxicity and disruption of erythrocyte antioxidant defence system in Wistar rats. *Environ Toxicol Pharmacol* **37**: 1202–1211.
- Oyagbemi AA, Omobowale TO, Akinrinde AS, Saba AB, Ogunpolu BS, Daramola O. (2015). Lack of reversal of oxidative damage in renal tissues of lead acetate-treated rats. *Environ Toxicol* **30**(11): 1235–43.



- Paudyal BP. (2008). Organophosphorus poisoning. *JNMA J Nepal Med Assoc* **47**: 251–258.
- Phaniendra A, Jestadi DB, Periyasamy L. (2015). Free radicals: properties, sources, targets, and their implication in various diseases. *Indian J Clin Biochem* **30**: 11–26.
- PHS (PUBLIC HEALTH SERVICE). (1996). Public health service policy on humane care and the use of laboratory animals. US Department of Health and Humane services, Washington, DC, pp. 99–158.
- Pizzurro DM, Dao K, Costa LG. (2015). Astrocytes protect against diazinon- and diazoxon-induced inhibition of neurite outgrowth by regulating neuronal glutathione. Organophosphate pesticides induce morphological abnormalities and decrease locomotor activity and heart rate in *Danio rerio* and *Xenopus laevis*. *Toxicol* **318**: 59–68.
- Ramkumar KM, Vijayakumar RS, Vanitha P, Suganya N, Manjula C, Rajaguru P. (2014). Protective effect of gallic acid on alloxan-induced oxidative stress and osmotic fragility in rats. *Hum Exp Toxicol* **33**: 638–649.
- Rastogi SK, Tripathi S, Ravishanker D. (2010). A study of neurologic symptoms on exposure to organophosphate pesticides in the children of agricultural workers. *Indian J Occup Environ Med* **14**: 54–57.
- Ray G. (1992). Pollution and health. Wiley Eastern Ltd. New-Delhi p. 45.
- Razavi BM, Hosseinzadeh H, Abnous K, Imenshahidi M. (2014). Protective effect of crocin on diazinon induced vascular toxicity in subchronic exposure in rat aorta ex-vivo. *Drug Chem Toxicol* **37**: 378–383.
- Razavi BM, Hosseinzadeh H, Movassaghi AR, Imenshahidi M, Abnous K. (2013). Protective effect of crocin on diazinon induced cardiotoxicity in rats in subchronic exposure. *Chem Biol Interact* **203**: 547–555.
- Razavi M, Hosseinzadeh H, Abnous K, Motamedshariaty VS, Imenshahidi M. (2013). Crocin restores hypotensive effect of subchronic administration of diazinon in rats. *Iran J Basic Med Sci* **16**: 64–72.
- Reitmann S, Frankel S. (1957). Colorimetric method for the determination of serum transaminase activity. *Am J Clin Pathol* **28**: 56–68.
- Repplinger D, Su MK, McKinnon K. (2014). Troponin elevations and organophosphate poisoning: direct cardiac injury or demand ischemia? *Clin Toxicol (Phila)* **52**: 1298.
- Sexton K, Salinas JJ. (2014). Concurrent fetal exposure to multiple environmental chemicals along the U.S.-Mexico border: an exploratory study in Brownsville, Texas. *Int J Environ Res Public Health* **11**: 10165–10181.
- Shah MD, Iqbal M. (2010). Diazinon-induced oxidative stress and renal dysfunction in rats. *Food Chem Toxicol* **48**: 3345–3353.
- Shinha KA. (1972). Colorimetric assay of Catalase. *Anal Biochem* **47**: 389–394.
- Stonys A, Kuzminskis V, Seputyte A, Astauskaite S, Jeseviciute. (2007). [Acute renal failure in patients with alcoholic surrogate intoxication]. *Medicina (Kaunas) Suppl* **1**: 36–39.
- Subramanian V, Venkatesan B, Tumala A, Vellaichamy E. (2014). Topical application of Gallic acid suppresses the 7, 12-DMBA/Croton oil induced two-step skin carcinogenesis by modulating anti-oxidants and MMP-2/MMP-9 in Swiss albino mice. *Food Chem Toxicol* **66**: 44–55.
- Sun L, Li GQ, Yan PB, Liu Y, Li GF, Wei LQ. (2015). Clinical management of organophosphate poisoning in pregnancy. *Am J Emerg Med* **33**: 305.e1–3.
- Sungur M, Güven M. (2001). Intensive care management of organophosphate insecticide poisoning. *Crit Care* **5**: 211–215.
- Thomaz JM1, Martins ND, Monteiro DA, Rantin FT, Kalinin AL. (2009). Cardio-respiratory function and oxidative stress biomarkers in Nile tilapia exposed to the organophosphate insecticide trichlorfon (NEGUVON). *Ecotoxicol Environ Saf* **72**: 1413–1424.
- Tsai CY, Chang AY, Chan JY, Chan SH. (2014). Activation of PI3K/Akt signaling in rostral ventrolateral medulla impairs brain stem cardiovascular regulation that underpins circulatory depression during mevinphos intoxication. *Biochem Pharmacol* **88**: 75–85.
- Peter JV, Sudarsan TI, Moran JL. (2014). Clinical features of organophosphate poisoning: A review of different classification systems and approaches. *Indian J Crit Care Med* **18**: 735–745.
- Valko M, Rhodes CJ, Moncol J, Izakovic M, Mazur M. (2006). Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chem Biol Interact* **160**: 1–40.
- Verma S, Singh A, Mishra A. (2013). Gallic acid: molecular rival of cancer. *Environ Toxicol Pharmacol* **35**: 473–485.
- Watson FL, Schmidt H, Turman ZK, Hole N, Garcia H, Gregg J, et al. (2014). Organophosphate pesticides induce morphological abnormalities and decrease locomotor activity and heart rate in *Danio rerio* and *Xenopus laevis*. *Environ Toxicol Chem* **33**: 1337–1345.
- Woff SP. (1994). Ferrous ion oxidation in the presence of ferric ion indicator xylenol orange for measurement of hydroperoxides. *Methods Enzymol* **233**: 182–189.
- Wu IW, Wu MS, Lin JL. (2008). Acute renal failure induced by bentazone: 2 case reports and a comprehensive review. *J Nephrol* **21**: 256–260.