



CARDIOLOGY // IMAGING

# Left Ventricle Noncompaction in a Young Adult

András Mester<sup>1</sup>, Laura Jáni<sup>1</sup>, Monica Chițu<sup>2</sup>

<sup>1</sup> Center of Advanced Research in Multimodality Cardiac Imaging, Cardio Med Medical Center, Tîrgu Mureş, Romania
<sup>2</sup> Clinic of Cardiology, University of Medicine and Pharmacy, Tîrgu Mureş, Romania

### CORRESPONDENCE

#### Laura Jáni

Str. 22 Decembrie 1989 nr. 76 540124 Tîrgu Mureş, Romania Tel +40 265 217 333 E-mail: jlaura15@yahoo.com

#### **ARTICLE HISTORY**

Received: 27 August, 2016 Accepted: 8 September, 2016

András Mester • Str. 22 Decembrie 1989 nr. 76, 540124 Tirgu Mureş, Romania. Tel +40 265 217 333 E-mail: andras.mester@yahoo.com

Monica Chiţu • Str. Gheorghe Marinescu nr. 38, 540139 Tîrgu Mureş, Romania. Tel +40 265 215 551. E-mail: iuliachitu@yahoo.com We report the case of a 23-year-old young male adult, without any previous personal or family medical history, who was admitted to our department with excessive fatigue, shortness of breath and a presyncope. The physical examination revealed peripheral edemas and bilateral fine basal crackles, cyanosis and hypotension. The resting ECG showed sinus rhythm, 130 bpm, with negative T waves in II, aVF, V4–V6.

Transthoracic echocardiography revealed significant left ventricle (LV) enlargement (75/64 mm), with moderate left atrium (LA, 53 mm) and right ventricle (RV, 44 mm) enlargement and significant systolic dysfunction (LV ejection fraction 30%). LV noncompaction (LVNC) was also revealed by echocardiography (Panel A) with prominent trabeculation of the LV (green arrow) and deep recesses (red arrow), with a relatively mobile 13/15mm echo-dense mass at the apex of the LV, highly specific for thrombus.

Transesophageal echocardiography (Panel B) identified an apical thrombus (red arrow).

As LVNC is frequently associated with other congenital abnormalities, computed tomography (CT) was performed (Panel C), which revealed a ratio of 2.41 of the noncompacted to compacted layer, with no other structural or coronary artery origin abnormality (Panel D).<sup>1,2</sup> The apical thrombus (red arrow) and the mid- and apical segment noncompaction (blue arrows) were also noted.

LVNC is a rare genetic cardiomyopathy, with possible late onset of the symptoms and it has to be considered at any age.<sup>3</sup> Proper imaging techniques must be used for correct diagnosis, management and prevention of complications.

## **CONFLICT OF INTEREST**

Nothing to declare.

## REFERENCES

1. Bennett CE, Freudenberger R. The Current Approach to Diagnosis and Management of Left Ventricular Noncompaction Cardiomyopathy: Review of the Literature. *Cardiol Res Pract.* 2016;2016:5172308.



- Arbustini E, Weidemann F, Hall JL. Left Ventricular Noncompaction: A Distinct Cardiomyopathy or a Trait Shared by Different Cardiac Diseases? J Am Coll Cardiol. 2014;64(17):1840-1850.
- Stöllberger C, Blazek G, Dobias C, Hanafin A, Wegner C, Finsterer J. Frequency of stroke and embolism in left ventricular hypertrabeculation/ noncompaction. Am J Cardiol. 2011;108(7):1021-1023.