

ON THE DIOPHANTINE EQUATION

$$11 + 2^{x+2} + (7)3^y = z^2$$

IDIR SADANI

ABSTRACT. In this note, we investigate solutions of the Diophantine equation

$$11 + 2^{x+2} + (7)3^y = z^2 \quad (x, y, z) \in \mathbb{N}^3.$$

1. Introduction

In 1844, Catalan in [2] conjectured that $(a, b, x, y) = (3, 2, 2, 3)$ is a unique solution for the exponential Diophantine equation

$$a^x - b^y = 1 \quad (a, b, x, y) \in \mathbb{N}^4 \quad \text{and} \quad \min(a, b, x, y) > 1.$$

The proof of this conjecture is due to Mihăilescu [3, 2004]. In 2011, Suvarnamani [1] found all non-negative integer solutions of diophantine equation

$$2^x + p^y = z^2.$$

In 2012, Chotchaisthit [4], using this theorem, found all solutions for the Diophantine equation $4^x + p^y = z^2$. In 2014, Yahui Yu and Xiaoxue Li [5] investigated solutions of equations of type

$$2^x + b^y = c^z, \quad x, y, z \in \mathbb{N},$$

where b and c are fixed coprime odd positive integers with $\min(b, c) > 1$.

In this note, we give a proof of the following result.

2. Main theorem

THEOREM 2.1. *The Diophantine equation $11 + 2^{x+2} + (7)3^y = z^2$ has a unique solution given by $(x, y, z) = (3, 1, 8)$.*

Proof. Let x, y be non-negative integers and z be an even non-negative integer such that

$$11 + 2^{x+2} + (7)3^y = z^2.$$

We need to consider several cases:

Case 1. If $y = 0$. Then, $2^{x+2} + 18 = z^2$. If we take $z = 2k$, we obtain $2^{x+1} = 2k^2 - 9$. But $2^{x+1} = 0 \pmod{2}$. This is a contradiction.

Case 2. If $y = 1$. Then, the equation becomes $2^{x+2} + 2^5 = z^2$. The solution of this equation is $(x, z) = (3, 8)$ (see [1]). Thus, the triplet $(3, 1, 8)$ is a solution of our equation.

Case 3. If $y > 1$. We have $11 + 2^{x+2} + (7)3^y = 4(1 + 2^x) + 7(1 + 3^y) = z^2$. Since z is even, then, y is odd because $7(1 + 3^{2l+1}) = 0 \pmod{4}$. Next, we can write $4(1 + 2^x) + 7(1 + 3^y)$ as $2^j c + 2^{j'} c'$. Then, we deduce that $j = 2$ and $1 + 3^y = 2^{j'}$. The unique solution of the last equation is $(y, j') = (1, 2)$. Hence, $(3, 1, 8)$ is the unique solution.

COROLLARY 2.1. $(3, 1, 2)$ is a unique non-negative integer solution (x, y, t) for the Diophantine equation $11 + 2^{x+2} + (7)3^y = t^6$, where x, y and t are non-negative integers.

Proof. Let x, y and t be non-negative integers such that $11 + 2^{x+2} + (7)3^y = t^6$. Let $z = t^3$. Then $11 + 2^{x+2} + (7)3^y = z^2$. By Theorem 2.1, we obtain that $(x, y, z) = (1, 3, 8)$. Then $t^3 = z = 8$. Hence, $t = 2$. \square

3. Open problem

Let a, b and c be positive odd prime numbers such that $(a, b, c) = 1$. We may ask for the set of all solutions (x, y, z) for the Diophantine equation

$$a + b2^x + c3^y = z^2,$$

where x, y and z are non-negative integers. \square

Acknowledgement. We would like to thank the editor and the referee for the opportunity to revise our manuscript.

ON THE DIOPHANTINE EQUATION $11 + 2^{x+2} + (7)3^y = z^2$

REFERENCES

- [1] SUVARNAMANI, A.: *Solutions of the Diophantine equation $2^x + p^y = z^2$* , Int. J. Math. Sci. Appl. **1** (2011), 1415–1419.
- [2] CATALAN, E.: *Note extraite d'une lettre adressée a l'éditeur*, J. Reine Angew. Math. **27** (1844), p. 192.
- [3] MIHAILESCU, P.: *Primary cyclotomic units and a proof of Catalan's conjecture*, J. Reine Angew. Math. **27** (2004), 167–195.
- [4] CHOTCHAISTHIT, S.: *On the Diophantine equation $4^x + p^y = z^2$ where p is a prime number*, Amer. J. Math. Sci. **1** (2012), 191–193.
- [5] YU, Y.—LI, X.: *The exponential Diophantine equation $2^x + b^y = z^c$* , Sci. World J. **2014** (2014), 1–3. <http://dx.doi.org/10.1155/2014/401816>

Received February 20, 2017

Department of Mathematics
University of Mouloud Mammeri
15000 Tizi-Ouzou
ALGERIA
E-mail: sadani.idir@yahoo.fr