

SIMPLIFICATION OF COEFFICIENTS IN DIFFERENTIAL EQUATIONS ASSOCIATED WITH HIGHER ORDER FROBENIUS-EULER NUMBERS

FENG QI^{1,2,3} — DA-WEI NIU⁴ — BAI-NI GUO³

¹School of Mathematical Sciences, Tianjin Polytechnic University, Tianjin, CHINA

²College of Mathematics, Inner Mongolia University for Nationalities, Tongliao, CHINA

³School of Mathematics and Informatics, Henan Polytechnic University, Jiaozuo, Henan, CHINA

⁴Department of Mathematics, East China Normal University, Shanghai, CHINA

ABSTRACT. In the paper, the authors apply Faà di Bruno formula, some properties of the Bell polynomials of the second kind, the inversion formulas of binomial numbers and the Stirling numbers of the first and the second kind, to significantly simplify coefficients in two families of ordinary differential equations associated with the higher order Frobenius–Euler numbers.

1. Motivations

In [4, Theorem 2.2], it was inductively and recursively established that the family of differential equations

$$F^{(n)}(t) = \left[\sum_{i=0}^n a_i(n) \left(\frac{u}{e^t - u} \right)^i \right] F(t) \quad (1)$$

for $n \geq 0$, $r \in \mathbb{N}$, and $u \in \mathbb{C} \setminus \{1\}$ has a solution

$$F(t) = F(t; r, u) = \left(\frac{1}{e^t - u} \right)^r, \quad (2)$$

where $a_0(n) = (-r)^n$,

© 2018 Mathematical Institute, Slovak Academy of Sciences.

2010 Mathematics Subject Classification: Primary: 34A05; Secondary: 05A16, 11A25, 11B37, 11B68, 11B73, 11B83, 33B10, 34A34.

Keywords: simplification; coefficient; ordinary differential equation; Faà di Bruno formula; higher order Frobenius–Euler number; Bell polynomial of the second kind; inversion formula.

$$a_i(n) = (-1)^n (r+i-1)_i \sum_{k_i=0}^{n-i} \sum_{k_{i-1}=0}^{n-i-k_i} \cdots \sum_{k_1=0}^{n-i-k_i-\cdots-k_2} r^{n-i-\sum_{\ell=1}^i k_\ell} \prod_{\ell=1}^i (r+\ell)^{k_\ell} \quad (3)$$

for $1 \leq i \leq n$, and

$$(x)_n = \prod_{\ell=0}^{n-1} (x+\ell) = \begin{cases} x(x+1)(x+2) \cdots (x+n-1), & n \geq 1 \\ 1, & n = 0 \end{cases}$$

is the rising factorial. Hereafter, the following results were deduced.

(1) For $k, n \geq 0$, we have

$$H_{k+n}^{(r)}(u) = \sum_{i=0}^n a_i(n) \left(\frac{u}{1-u} \right)^i H_k^{(r+i)}(u),$$

where $H_k^{(r)}$, which can be generated by

$$\left(\frac{1-u}{e^t-u} \right)^r = \sum_{k=0}^{\infty} H_k^{(r)}(u) \frac{t^k}{k!},$$

stand for the Frobenius-Euler numbers of order r . See [4, Theorem 2.3].

(2) For $k, n \geq 0$, we have

$$E_{k+n}^{(r)} = \sum_{i=0}^n \left(-\frac{1}{2} \right)^i a_i(n) E_k^{(r+i)},$$

where $E_k^{(r)}$, which can be generated by

$$\left(\frac{2}{e^t+1} \right)^r = \sum_{k=0}^{\infty} E_k^{(r)} \frac{t^k}{k!},$$

stand for the Euler numbers of order r . See [4, Corollary 2.5].

(3) When $0 \leq k \leq r-1$ and $k \geq r+n$,

$$B_k^{(r)} = \frac{1}{(k-r)_n} \sum_{i=\max\{n-k, 0\}}^n a_i(n) B_{k+i-n}^{(r+i)}(k)_{n-i};$$

when $r \leq k \leq r-1+n$,

$$\sum_{i=\max\{n-k, 0\}}^n a_i(n) B_{k+i-n}^{(r+i)}(k)_{n-i} = 0;$$

where $B_k^{(r)}$, which can be generated by

$$\left(\frac{t}{e^t-1} \right)^r = \sum_{k=0}^{\infty} B_k^{(r)} \frac{t^k}{k!},$$

stand for the Bernoulli numbers of order r . See [4, Theorem 2.7].

COEFFICIENTS IN DIFFERENTIAL EQUATIONS

In [5, Theorem 2.1], it was inductively and recursively proved that the family of differential equations

$$(-1)^{n-1}(r)_n \left(\frac{u}{e^t - u} \right)^n F(t) = \sum_{i=0}^n b_i(n) F^{(i)}(t) \quad (4)$$

for $u \in \mathbb{C}$ and $r \in \mathbb{N}$ has a solution $F(t)$ defined in (2), where $b_0(n) = -\langle r+n-1 \rangle_n$,

$$b_i(n) = - \sum_{k_i=0}^{n-i} \sum_{k_{i-1}=0}^{n-i-k_i} \cdots \sum_{k_1=0}^{n-i-k_i-\cdots-k_2} \prod_{\ell=1}^i \left\langle r+n-i-1 - \sum_{j=\ell+1}^i k_j + \ell \right\rangle_{k_\ell} \quad (5)$$

$$\times \left\langle r+n-i-1 - \sum_{j=1}^i k_j \right\rangle_{n-i-\sum_{j=1}^i k_j}, \quad 1 \leq i \leq n,$$

and

$$\langle x \rangle_n = \prod_{\ell=0}^{n-1} (x - \ell) = \begin{cases} x(x-1)(x-2)\dots(x-n+1), & n \geq 1 \\ 1, & n = 0 \end{cases}$$

is the falling factorial of $x \in \mathbb{R}$ for $n \in \{0\} \cup \mathbb{N}$. Hereafter, the following conclusions were derived:

(1) For $k, n \geq 0$, we have

$$(-1)^{n-1}(r)_n \left(\frac{u}{1-u} \right)^n H_k^{(r+n)}(u) = \sum_{i=0}^n b_i(n) H_{k+i}^{(r)}(u). \quad (6)$$

See [5, Theorem 3.1]. In particular, taking $u = -1$ in (6) leads to

$$(-1)^{n-1}(r)_n \left(-\frac{1}{2} \right)^n E_k^{(r+n)} = \sum_{i=0}^n b_i(n) E_{k+i}^{(r)}.$$

See [5, Corollary 3.3].

(2) When $0 \leq k \leq n+r-1$, we have

$$B_k^{(r+n)} = (-1)^{n-1} \frac{1}{(r)_n} \sum_{i=\max\{n-k, 0\}}^{\min\{r+n-1-k, n\}} b_i(n) B_{k+i-n}^{(r)} \frac{\langle k+i-n-r \rangle_i k!}{(k+i-n)!};$$

when $k \geq n+r$, we have

$$B_k^{(r+n)} = (-1)^{n-1} \frac{1}{(r)_n} \sum_{i=0}^n b_i(n) B_{k+i-n}^{(r)} \frac{\langle k+i-n-r \rangle_i k!}{(k+i-n)!}.$$

See [5, Theorem 3.4].

(3) The matrices $(a_i(j))_{0 \leq i, j \leq n}$ and $\left(\frac{b_i(j)}{(-1)^{j-1}(r)_j}\right)_{0 \leq i, j \leq n}$ are inverse to each other for all n . See [5, Remark 3.2].

It is easy to see that expressions (3) and (5) of the quantities $a_i(n)$ and $b_i(n)$ are too complicated to be computed by hand and computer software. Therefore, can one find simple, meaningful, and significant expressions for the quantities $a_i(n)$ and $b_i(n)$ in (3) and (5)?

2. Lemmas

For answering the above question and proving our main results, we need the following lemmas.

LEMMA 1 ([1, p. 134, Theorem A] and [1, p. 139, Theorem C]). *For $n \geq k \geq 0$, the Bell polynomials of the second kind, or say, partial Bell polynomials, denoted by $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$, are defined by*

$$B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) = \sum_{\substack{1 \leq i \leq n-k+1 \\ \ell_i \in \{0\} \cup \mathbb{N} \\ \sum_{i=1}^{n-k+1} i \ell_i = n \\ \sum_{i=1}^{n-k+1} \ell_i = k}} \frac{n!}{\prod_{i=1}^{n-k+1} \ell_i!} \prod_{i=1}^{n-k+1} \left(\frac{x_i}{i!}\right)^{\ell_i}.$$

The Faà di Bruno formula can be described in terms of the Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ by

$$\frac{d^n}{dt^n} f \circ h(t) = \sum_{k=0}^n f^{(k)}(h(t)) B_{n,k} \left(h'(t), h''(t), \dots, h^{(n-k+1)}(t) \right). \quad (7)$$

LEMMA 2 ([1, p. 135]). *For $n \geq k \geq 0$, we have*

$$B_{n,k}(abx_1, ab^2x_2, \dots, ab^{n-k+1}x_{n-k+1}) = a^k b^n B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) \quad (8)$$

and

$$B_{n,k}(1, 1, \dots, 1) = S(n, k), \quad (9)$$

where a and b are any complex numbers and $S(n, k)$ for $n \geq k \geq 0$, which can be generated by

$$\frac{(e^x - 1)^k}{k!} = \sum_{n=k}^{\infty} S(n, k) \frac{x^n}{n!},$$

stand for the Stirling numbers of the second kind.

LEMMA 3 ([27, p. 171, Theorem 12.1]). *If b_α and a_k are a collection of constants independent of n , then*

$$a_n = \sum_{\alpha=0}^n S(n, \alpha) b_\alpha \quad \text{if and only if} \quad b_n = \sum_{k=0}^n s(n, k) a_k,$$

where $s(n, k)$ for $n \geq k \geq 0$, which can be generated by

$$\frac{[\ln(1+x)]^k}{k!} = \sum_{n=k}^{\infty} s(n, k) \frac{x^n}{n!}, \quad |x| < 1,$$

stand for the Stirling numbers of the first kind.

LEMMA 4 ([27, p. 83, Eq. (7.12)]). *If a_k and b_k for $k \geq 0$ are a collection of constants independent of n , then*

$$a(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} b(k) \quad \text{if and only if} \quad b(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} a(k).$$

3. Main results and their proofs

Now we are able to answer the above question and to state and prove our main results.

THEOREM 1. *For $n \geq 0$, $r \in \mathbb{R}$, and $u \in \mathbb{C}$, the function $F(t)$ defined by (2) satisfies*

$$F^{(n)}(t) = \sum_{\ell=0}^n \left[\sum_{k=\ell}^n \binom{k}{\ell} S(n, k) \langle -r \rangle_k \right] \left(\frac{u}{e^t - u} \right)^\ell F(t) \quad (10)$$

and

$$\sum_{k=0}^n \left[\sum_{\ell=k}^n (-1)^\ell \binom{n}{\ell} \frac{s(\ell, k)}{\langle -r \rangle_\ell} \right] F^{(k)}(t) = (-1)^n \left(\frac{u}{e^t - u} \right)^n F(t). \quad (11)$$

Proof. Let

$$F(t) = \frac{1}{w^r} \quad \text{and} \quad w = w(t) = w(t; u) = e^t - u.$$

Then, by the Faà di Bruno formula (7) and the identities (8) and (9) in sequence,

$$\begin{aligned} F^{(n)}(t) &= \sum_{k=0}^n \left(\frac{1}{w^r} \right)^{(k)} B_{n,k}(e^t, e^t, \dots, e^t) \\ &= \sum_{k=0}^n \frac{\langle -r \rangle_k}{w^{r+k}} e^{kt} B_{n,k}(1, 1, \dots, 1) \end{aligned}$$

$$\begin{aligned}
&= \sum_{k=0}^n \frac{\langle -r \rangle_k}{(e^t - u)^{r+k}} e^{kt} S(n, k) \\
&= \frac{1}{(e^t - u)^r} \sum_{k=0}^n \frac{\langle -r \rangle_k}{(e^t - u)^k} e^{kt} S(n, k) \\
&= F(t) \sum_{k=0}^n \langle -r \rangle_k S(n, k) \left(\frac{e^t}{e^t - u} \right)^k \\
&= F(t) \sum_{k=0}^n \langle -r \rangle_k S(n, k) \left(1 + \frac{u}{e^t - u} \right)^k \\
&= F(t) \sum_{k=0}^n \langle -r \rangle_k S(n, k) \sum_{\ell=0}^k \binom{k}{\ell} \left(\frac{u}{e^t - u} \right)^\ell \\
&= F(t) \sum_{\ell=0}^n \left[\sum_{k=\ell}^n \langle -r \rangle_k S(n, k) \binom{k}{\ell} \right] \left(\frac{u}{e^t - u} \right)^\ell.
\end{aligned}$$

Therefore, the identity (10) follows immediately.

From the above proof of the identity (10), it can be deduced that

$$F^{(n)}(t) = F(t) \sum_{k=0}^n S(n, k) \langle -r \rangle_k \sum_{\ell=0}^k \binom{k}{\ell} \left(\frac{u}{e^t - u} \right)^\ell, \quad n \geq 0.$$

Utilizing Lemma 3 we obtain

$$F(t) \langle -r \rangle_n \sum_{\ell=0}^n \binom{n}{\ell} \left(\frac{u}{e^t - u} \right)^\ell = \sum_{k=0}^n s(n, k) F^{(k)}(t), \quad n \geq 0$$

which can be rearranged as

$$\sum_{\ell=0}^n (-1)^\ell \binom{n}{\ell} \left(-\frac{u}{e^t - u} \right)^\ell = \frac{1}{F(t) \langle -r \rangle_n} \sum_{k=0}^n s(n, k) F^{(k)}(t), \quad n \geq 0.$$

Further use of Lemma 4 derives

$$\sum_{\ell=0}^n (-1)^\ell \binom{n}{\ell} \frac{1}{F(t) \langle -r \rangle_\ell} \sum_{k=0}^\ell s(\ell, k) F^{(k)}(t) = \left(-\frac{u}{e^t - u} \right)^n, \quad n \geq 0$$

which can be rewritten as (11). The required proof is complete. \square

4. Remarks

In this section, we give several remarks and some explanation about our main results.

Remark 1. Theorem 1 extends the range of r from \mathbb{N} to \mathbb{R} .

Remark 2. Comparing (1) with (10) one reveals that

$$a_i(n) = \sum_{k=i}^n \binom{k}{i} S(n, k) \langle -r \rangle_k, \quad 0 \leq i \leq n. \quad (12)$$

This implies that the identity (10) is more meaningful, more significant, more computable than the one (1).

Remark 3. It is not difficult to see that

$$a_0(n) = \sum_{k=0}^n S(n, k) \langle -r \rangle_k = (-r)^n, \quad n \geq 0.$$

Then it is natural to ask a question: Is the finite sum

$$a_i(n) = \sum_{k=i}^n \binom{k}{i} S(n, k) \langle -r \rangle_k, \quad 1 \leq i \leq n$$

summarizable?

Remark 4. Comparing (4) with (11) one obtains

$$b_i(n) = (r)_n \sum_{\ell=i}^n (-1)^{\ell+1} \binom{n}{\ell} \frac{s(\ell, i)}{\langle -r \rangle_\ell}, \quad n \geq i \geq 0. \quad (13)$$

This means that the identity (11) is more meaningful, more significant, more computable than (4).

Remark 5. By virtue of the expressions (12) and (13), all the above mentioned results in the papers [4, 5] can be reformulated simpler, more meaningfully, and more significantly. For the sake of saving the space and shortening the length of this paper, we do not rewrite them in detail here.

Remark 6. Till now we can see that the method used in this paper is simpler, shorter, nicer, more meaningful, and more significant than the inductive and recursive method used in [4, 5] and closely related references therein.

Remark 7. In the papers and preprints [2, 3, 6–20, 22–26, 28], there are similar ideas, methods, techniques, and purposes to this paper.

Remark 8. This paper is a slightly revised version of the preprint [21].

Acknowledgement. The authors thank Professor T a e k y u n K i m at Kwang-woon University in South Korea for his supplying electronic versions of the papers [4, 5] through e-mail on 31 July 2017.

REFERENCES

- [1] COMTET, L.: *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, Revised and Enlarged Edition, D. Reidel Publishing Co., Dordrecht and Boston 1974.
<https://doi.org/10.1007/978-94-010-2196-8>.
- [2] GUO, B.-N.—QI, F.: *Explicit formulae for computing Euler polynomials in terms of Stirling numbers of the second kind*, J. Comput. Appl. Math. **272** (2014), 251–257;
<http://dx.doi.org/10.1016/j.cam.2014.05.018>.
- [3] ——— *Some identities and an explicit formula for Bernoulli and Stirling numbers*, J. Comput. Appl. Math. **255** (2014), 568–579;
<http://dx.doi.org/10.1016/j.cam.2013.06.020>.
- [4] KIM, D. S.—KIM, T.—PARK, J.-W.—SEO, J.-J.: *Differential equations associated with higher-order Frobenius-Euler numbers*, Glob. J. Pure Appl. Math. **12** (2016), no. 3, 2537–2547.
- [5] KIM, T.—KIM, D. S.—JANG, L.-C.—KWON, H. I.: *Differential equations associated with higher-order Frobenius-Euler numbers revisited*, Differ. Equ. Dyn. Syst. (2019) (to appear); <https://doi.org/10.1007/s12591-017-0380-8>.
- [6] QI, F.: *A simple form for coefficients in a family of nonlinear ordinary differential equations*, Adv. Appl. Math. Sci. **17** (2018), no. 8, 555–561.
- [7] ——— *A simple form for coefficients in a family of ordinary differential equations related to the generating function of the Legendre polynomials*, Adv. Appl. Math. Sci. **17** (11) (2018), no. 11, 693–700.
- [8] ——— *Notes on several families of differential equations related to the generating function for the Bernoulli numbers of the second kind*, Turkish J. Anal. Number Theory **6** (2018), no. 2, 40–42; <https://doi.org/10.12691/tjant-6-2-1>.
- [9] ——— *Simple forms for coefficients in two families of ordinary differential equations*, Glob. J. Math. Anal. **6** (2018), no. 1, 7–9. <https://doi.org/10.14419/gjma.v6i1.9778>.
- [10] ——— *Simplification of coefficients in two families of nonlinear ordinary differential equations*, Turkish J. Anal. Number Theory **6** (2018), no. 6, 116–119;
<https://doi.org/10.12691/tjant-6-4-2>.
- [11] ——— *Simplifying coefficients in a family of nonlinear ordinary differential equations*, Acta Comment. Univ. Tartu Math. **22** (2018), no. 2, 293–297;
<https://doi.org/10.12697/ACUTM.2018.22.24>.
- [12] ——— *Simplifying coefficients in a family of ordinary differential equations related to the generating function of the Laguerre polynomials*, Appl. Appl. Math. **13** (2018), no. 2, 750–755.

- [13] QI, F.: *Simplifying coefficients in differential equations related to generating functions of reverse Bessel and partially degenerate Bell polynomials*, Bol. Soc. Paran. Mat. **39**(2021), no. 4, (to appear); <http://dx.doi.org/10.5269/bspm.41758>.
- [14] QI, F.—GUO, B.-N.: *A diagonal recurrence relation for the Stirling numbers of the first kind*, Appl. Anal. Discrete Math. **12** (2018), no. 1, 153–165; <https://doi.org/10.2298/AADM170405004Q>.
- [15] ——— *Explicit formulas and recurrence relations for higher order Eulerian polynomials*, Indag. Math. **28** (2017), no. 4, 884–891, <https://doi.org/10.1016/j.indag.2017.06.010>.
- [16] ——— *Some properties of the Hermite polynomials*. In: *Advances in Special Functions and Analysis of Differential Equations* (Praveen Agarwal, Ravi P. Agarwal, and Michael Ruzhansky, eds.) CRC Press, Taylor & Francis Group, 2019 (to appear).
- [17] ——— *Viewing some ordinary differential equations from the angle of derivative polynomials*, Iran. J. Math. Sci. Inform. **14** (2019), no. 2 (to appear).
- [18] QI, F.—LIM, D.—GUO, B.-N.: *Explicit formulas and identities for the Bell polynomials and a sequence of polynomials applied to differential equations*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM **113** (2019), no. 1, 1–9; <http://dx.doi.org/10.1007/s13398-017-0427-2>.
- [19] ——— *Some identities related to Eulerian polynomials and involving the Stirling numbers*, Appl. Anal. Discrete Math. **12** (2018), no. 2., 467–480; <https://doi.org/10.2298/AADM171008014Q>.
- [20] QI, F.—LIM, D.—LIU, A.-Q.: *Explicit expressions related to degenerate Cauchy numbers and their generating function*, In: *Proceeding on Mathematical Modelling, Applied Analysis and Computation* In: Springer Proceedings in Mathematics and Statistics 2019 (to appear); <https://www.springer.com/series/10533>.
- [21] QI, F.—NIU, D.-W.—GUO, B.-N.: *Simplification of coefficients in differential equations associated with higher order Frobenius-Euler numbers*, Preprints **2017**, 2017080017, 7 pp; <http://dx.doi.org/10.20944/preprints201708.0017.v1>.
- [22] ——— *Some identities for a sequence of unnamed polynomials connected with the Bell polynomials*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Math. RACSAM **112** (2018) (to appear); <https://doi.org/10.1007/s13398-018-0494-z>.
- [23] QI, F.—WANG, J.-L.—GUO, B.-N.: *Notes on a family of inhomogeneous linear ordinary differential equations*, Adv. Appl. Math. Sci. **17** (2018), no. 4, 361–368.
- [24] ——— *Simplifying and finding ordinary differential equations in terms of the Stirling numbers*, Korean J. Math. **26** (2018), no. 4, 675–681; <https://doi.org/10.11568/kjm.2018.26.4.675>.
- [25] ——— *Simplifying differential equations concerning degenerate Bernoulli and Euler numbers*, Trans. A. Razmadze Math. Inst. **172** (2018), no. 1, 90–94; <http://dx.doi.org/10.1016/j.trmi.2017.08.001>.
- [26] QI F.—ZHAO, J.-L.: *Some properties of the Bernoulli numbers of the second kind and their generating function*, Bull. Korean Math. Soc. **55** (2018), no. 6, 1909–1920; <https://doi.org/10.4134/BKMS.b180039>.

- [27] QUAINANCE J.—GOULD, H. W.: : *Combinatorial Identities for Stirling Numbers*. The unpublished notes of H. W. Gould. With a foreword by George E. Andrews. World Scientific Publishing Co. Pte. Ltd., Singapore 2016.
- [28] ZHAO, J.-L.—WANG, J.-L.—QI, F.: : *Derivative polynomials of a function related to the Apostol–Euler and Frobenius–Euler numbers*, J. Nonlinear Sci. Appl. **10** (2017), no. 4; 1345–1349, <http://dx.doi.org/10.22436/jnsa.010.04.06>.

Received July 9, 2018

Feng Qi
School of Mathematical Sciences
Tianjin Polytechnic University
Tianjin 300387

College of Mathematics
Inner Mongolia University for Nationalities
Tongliao 028043, Inner Mongolia

School of Mathematics and Informatics
Henan Polytechnic University
Jiaozuo 454010, Henan
CHINA
E-mail: qifeng618@gmail.com

Da-Wei Niu
Department of Mathematics
East China Normal University
Shanghai 200241
CHINA
E-mail: nnddww@gmail.com

Bai-Ni Guo
School of Mathematics and Informatics
Henan Polytechnic University
Jiaozuo 454010, Henan
CHINA
E-mail: bai.ni.guo@hotmail.com