

Scientific paper

DOI: 10.2478/10004-1254-64-2013-2319

MUTAGENICITY AND DNA DAMAGE OF BISPHENOL A AND ITS STRUCTURAL ANALOGUES IN HEPG2 CELLS

Anja FIC¹, Bojana ŽEGURA², Marija SOLLNER DOLENC¹, Metka FILIPIČ², and Lucija PETERLIN MAŠIČ¹

Faculty of Pharmacy, University of Ljubljana¹, National Institute of Biology, Department for Genetic Toxicology and Cancer Biology², Ljubljana, Slovenia

Received in October 2012

CrossChecked in November 2012

Accepted in April 2013

Environmental oestrogen bisphenol A (BPA) and its analogues are widespread in our living environment. Because their production and use are increasing, exposure of humans to bisphenols is becoming a significant issue. We evaluated the mutagenic and genotoxic potential of eight BPA structural analogues (BPF, BPAF, BPZ, BPS, DMBPA, DMBPS, BP-1, and BP-2) using the Ames and comet assay, respectively. None of the tested bisphenols showed a mutagenic effect in *Salmonella typhimurium* strains TA98 and TA100 in either the presence or absence of external S9-mediated metabolic activation (Aroclor 1254-induced male rat liver). Potential genotoxicity of bisphenols was determined in the human hepatoma cell line (HepG2) at non-cytotoxic concentrations (0.1 $\mu\text{mol L}^{-1}$ to 10 $\mu\text{mol L}^{-1}$) after 4-hour and 24-hour exposure. In the comet assay, BPA and its analogue BPS induced significant DNA damage only after the 24-hour exposure, while analogues DMBPS, BP-1, and BP-2 induced a transient increase in DNA strand breaks observed only after the 4-hour exposure. BPF, BPAF, BPZ, and DMBPA did not induce DNA damage.

KEY WORDS: *Ames test, bisphenols, comet assay, genotoxicity*

Bisphenol A (BPA; 4,4'-propane-2,2-diylidiphenol, CAS no. 80-05-7) is used in the production of food contact materials, such as baby bottles, food containers, and protective coatings for canned food and beverages, and for metal lids on glass jars and bottles. At higher temperatures, longer contact with, and higher pH of the contact medium, BPA monomer can hydrolyse and leach into food and beverages. Average exposure concentrations range from 10 $\mu\text{g kg}^{-1}$ to 70 $\mu\text{g kg}^{-1}$ in solid canned food and from 1 $\mu\text{g L}^{-1}$ to 23 $\mu\text{g L}^{-1}$ in liquid canned food (1). BPA has been under scrutiny due to concerns over potential adverse health effects related to its endocrine-disrupting activity (2-3). The first regulatory risk assessment report on BPA published by the Canadian government (4) has resulted

in the ban of BPA in baby bottles in Canada. In January 2010, the US Food and Drug Administration expressed concern about potential adverse health effects of BPA in infants and children. The use of BPA in food contact materials has been banned in Japan, Canada, and many US states, and the US Congress is considering federal ban on BPA in all food and beverage containers (5). In 2011, the European Union prohibited the manufacture, marketing, and import of baby bottles containing BPA (6).

Current efforts are focused on replacing BPA with safer food contact materials. All of these alternative materials need to be assessed for appropriate functionality and safety using state-of-the-art methodology and scientific knowledge (1).

Bisphenols (Table 1) are a class of chemicals known as diphenylmethanes, which contain two benzene rings separated by one central carbon atom, usually with a 4-OH substituent on both benzene rings (e.g. BPA, BPF, BPAF, BPZ, and DMBPA). In some bisphenols, the central carbon atom is replaced by a sulphone group (e.g. BPS, DMBPS, or BP-1) or sulphide moiety (e.g. BP-2). Some BPA analogues seem to be safer alternatives to BPA in industrial applications (7). For example, the production of bisphenol S (BPS), which is stable at high temperatures and resistant to sunlight, is increasing from year to year (7-8). The largest US manufacturer of thermal paper has been using BPS as a replacement for BPA since 2006 (9). However, insufficient data are available to tell whether these BPS-containing papers are safer than BPA-containing papers. While BPA is moderately susceptible to environmental breakdown, BPS may be more persistent (10-11). Another example is BP-1, whose use as polymer bottle component was first reported more than 30 years ago (12). Lotti et al. (13) reported that poly(butylene terephthalate) modified with BP-1 showed improved glass transition temperature and thermal stability. Bisphenol F (BPF) and bisphenol AF (BPAF) are also used for polycarbonate resin production (14). From the viewpoint of biodegradability in the aquatic environment, BPF is more biodegradable under aerobic and anaerobic conditions than BPA, and may replace BPA to lower environmental risks (11). BPF also occurs as a monomer of phenol-formaldehyde resin. BPAF is a component of certain plasters and is used as a rubber bridging material, while DMBPA is a monomer of polycarbonate, epoxy, and polyester resins (15). Due insufficient toxicity data and structural similarity to BPA, BPAF has been nominated for a comprehensive toxicological characterisation by the US National Institute of Environmental Health Sciences (16).

While the toxicity of BPA has received a lot of attention, BPA analogues have not, despite the fact that they are threatening to become dominant environmental pollutants in the near future and that their impact on the environment and human health requires urgent attention (7). The main mechanism underlying BPA-induced adverse effects is endocrine disruption that may lead to developmental and/or reproductive disorders (2-3). Moreover, endocrine disruptors may induce carcinogenic effects due to epigenetic events or due to genotoxic effects. One potent endocrine disruptor and evidenced carcinogen, diethylstilbestrol (DES), is structurally related to BPA

(17-18). Both DES and BPA have a hydrophobic core with two OH groups at each end of the backbone whose distance is similar (9.0 Å to 9.2 Å for BPA and 9 Å to 11.5 Å for DES at different conformations) (19). Studies of BPA genotoxicity have yielded conflicting results. BPA is considered non-genotoxic because it was negative to a set of basic genotoxicity tests. It was not mutagenic in the *Salmonella*/microsome assay (20-21), did not induce gene mutations (21-22) or chromosomal aberrations (23) in mammalian cells *in vitro*, and failed to induce chromosomal aberration and micronucleus formation *in vivo* in mice bone marrow (24). In contrast, BPA induced numerical chromosomal aberrations and morphological changes in cultured SHE cells (22) and in mice it induced achromatic lesions and c-mitotic effects in bone marrow cells (24). In addition, BPA metabolite(s) were shown to bind to DNA in a cellular system (25-26), in cultured SHE cells (22), and in rodent liver *in vivo* (27-28). Moreover, in oestrogen receptor (ER)-positive MCF-7 cells, BPA caused DNA strand breaks that were ER-dependent (29). In turn, BPF has been reported to induce DNA strand breaks, but not micronuclei, in human hepatoma HepG2 cells (30). Audebert et al. (31) have recently found that BPF genotoxicity depended on the metabolic capabilities of cells. In human HepG2 cells it induced histone H2AX phosphorylation, an indicator of DNA double-strand breaks. BPAF induced metaphase arrest and micronucleus formation in V79 cells (32). In SHE cells, BPAF did not induce gene mutation or chromosomal aberrations, but induced aneuploidy and morphological changes (15, 33).

The aim of this *in vitro* study was to investigate the mutagenic and genotoxic potential of a series of BPA analogues selected from three structural groups based on the bridging moiety between two phenolic rings and substitution pattern on phenolic rings. The first group has a central carbon atom (BPA, BPF, BPAF, BPZ, and DMBPA), the second group contains sulphones (BPS, DMBPS and BP-1), and the third group a sulphide analogue (BP-2). BP-1 and BP-2 were denoted by numbers to avoid confusion between abbreviations found in literature.

MATERIALS AND METHODS

Chemicals

Bisphenol A (BPA, >99 % pure; CAS # 80-05-7), bisphenol F (BPF, 98 % pure; CAS # 620-92-8), bisphenol AF (BPAF, 97 % pure; CAS # 1478-61-1),

bisphenol Z (BPZ, 98 % pure; CAS # 843-55-0), 2,2-bis(4-hydroxy-3-methylphenyl)propane (DMBPA, 97 % pure; CAS # 79-97-0), bisphenol S (BPS, 98 % pure; CAS # 80-09-1), 4,4'-sulfonylbis(2-methylphenyl) (DMBPS, 97 % pure; CAS # 16346-97-7), [sulphonylbis(benzene-4,1-diyloxy)]diethanol (BP-1, 95 % pure; CAS # 27205-03-4), 4,4'-sulphanedioldiphenol (BP-2, 99 % pure; CAS # 2664-63-3), William's medium E, ampicillin, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT), glucose-6-phosphate (disodium salt), benzo[a]pyrene (B[a]P; CAS # 50-32-8), ethidium bromide solution (CAS # 1239-45-8), dimethyl sulphoxide (DMSO; CAS # 67-68-5), EDTA (CAS # 6381-92-6), and Triton X-100 were obtained from Sigma (St. Louis, MO, USA). Normal melting point (NMP) agarose and low melting point agarose (LMP) were from Invitrogen (Carlsbad, CA, USA). Lyophilised Aroclor 1254-induced male rat liver post-mitochondrial fraction (S9) was obtained from Moltax (Boone, NC, USA). Penicillin/streptomycin, foetal bovine serum (FBS), L-glutamine, and phosphate buffered saline (PBS) were sourced from PAA Laboratories (Dartmouth, MA, USA) and trypsin from BD-Difco (Le Pont-De-Claix Cedex, France). All other reagents were of the purest grade available and all solutions were made using distilled water.

Bacterial strains

Salmonella typhimurium strains TA98 (frame shift mutations) and TA100 (base pair substitutions) were obtained from Professor B. N. Ames (University of California, Berkeley, USA) and were regularly checked for their phenotypic characteristics [histidine/biotin dependence, rfa marker (crystal violet), uvrB deletion (UV sensitivity), and the presence of plasmid pKM101 (ampicillin resistance)]. The working bacterial cultures were prepared from frozen permanents by overnight incubation (37 °C) in the Oxoid nutrient broth no. 2 (Oxoid, Basingstoke, UK) in the presence of 25 µg mL⁻¹ ampicillin.

Determination of mutagenicity with the Ames test

Bisphenol mutagenicity was tested with the *Salmonella*/microsomal reverse mutation assay (34-35). Prior to the testing, BPs were dissolved and diluted in 100 % DMSO to give final concentrations of (0.004, 0.02, 0.1, and 0.5) mg per plate. Overnight cultures of *S. typhimurium* strains TA98 and TA100 (100 µL) as well as corresponding BP dilutions

(100 µL) were added to 2 mL of molten top agar containing a limited amount of histidine/biotin (42 °C), gently mixed, and poured onto minimal agar plates. For the assay with metabolic activation, 500 µL of S9 mix (containing 4 % S9 – Aroclor-induced rat liver microsomal fraction) was also added to 2 mL of molten soft agar. Benzo[a]pyrene (B[a]P; final concentration 10 µg per plate) and 4-nitroquinoline-*N*-oxide (4-NQNO; final concentration 0.5 µg per plate) were used as positive controls for testing in the presence and absence of S9 mix, respectively. 100 % DMSO was used as a solvent control. The number of His⁺ revertants was counted after 48 h (TA100) and 72 h (TA98) of incubation at 37 °C. Three plates were used per treatment point. The mutagenic potential of the samples was expressed as an induction factor (IF), where IF = (number of revertants in the presence of the sample)/(number of revertants in solvent control).

For the purposes of this study, a non-statistical procedure was used to evaluate the results of the Ames test. A compound was considered a mutagen if it produced a reproducible, dose-related increase in the number of revertant colonies in one or more strains and induced at least a twofold increase in the number of revertants in respect to solvent control (IF ≥ 2) (34).

Human HepG2 cells

These cells were selected as they express a range of xenobiotic-metabolising enzymes (36-40) as well as ERα and ERβ receptors (41). The HepG2 cells were provided by Professor Firouz Darroudi (Leiden University Medical Centre, Department of Toxicogenetics, Leiden, The Netherlands). They were grown in William's medium E containing 15 % FBS, 2 mmol L⁻¹ L-glutamine, and 100 U mL⁻¹ penicillin/streptomycin in 5 % CO₂ at 37 °C. The cells were used at passages between 3 and 10. For sub-cultivation, the cells were trypsinised, washed with phosphate-buffered saline (PBS, pH 7.4), centrifuged at 100 g for 5 min, and separated by pressing the suspensions through a syringe (needle 0.9x40 mm, Becton Dickinson, S.A., Fraga, Spain).

Cell viability

Bisphenol cytotoxicity was determined with the MTT assay in accordance with the procedure used by Mosmann (42) with minor modifications (43). HepG2 cells were seeded into 96-well microtitre plates at a density of 8,000 cells per well. After a 24-hour

Table 1 BPA and its structural analogues

Compound	Structure	Chemical name
BPA		4,4'-(propane-2,2-diyl)diphenol
BPF		4,4'-methylenediphenol
BPAF		4,4'-(perfluoropropane-2,2-diyl)diphenol
BPZ		4,4'-(cyclohexane-1,1-diyl)diphenol
DMBPA		2,2-bis(4-hydroxy-3-methylphenyl)propane
BPS		4,4'-sulfonyldiphenol
DMBPS		4,4'-sulfonylbis(2-methylphenol)
BP-1		((sulfonylbis(4,1-phenylene))bis(oxy))dimethanol
BP-2		4,4'-thiodiphenol

incubation at 37 °C, the growth medium was replaced with fresh medium containing from 12.5 $\mu\text{mol L}^{-1}$ to 100 $\mu\text{mol L}^{-1}$ of BPs, and the cells were incubated for additional 24 h. The final concentration of DMSO in solvent control and dilutions was 0.1 %. MTT was then added to a final concentration 0.5 mg mL^{-1} , and the cells further incubated at 37 °C for 3 h. The medium was removed and formazan crystals dissolved in DMSO. The optical density (OD) of the solution in each well was measured against a blank (a well with

DMSO) at 570 nm (the formazan absorption peak) and at 690 nm (measurement of the medium turbidity caused by cell debris) with a GENios™ microplate spectrofluorometer (Tecan, Trappes, France). The viability of cells was determined by comparing relative formazan concentrations (OD570-OD690) of the treated cells with those of untreated solvent control cells. Five individual wells were measured per treatment point. The experiment was repeated twice. Statistical significance between the treated groups and

control was determined using a two-tailed Student's *t*-test, where $P < 0.01$ was considered significant.

Determination of genotoxicity with the comet assay

Stock solutions of BPs (100 mmol L^{-1}) were prepared in DMSO, and dilutions were prepared in the culture medium. The final concentration of DMSO in dilutions did not exceed 0.1 %. Solvent control (cell growth medium containing 0.1 % DMSO) and positive control ($30 \text{ } \mu\text{mol L}^{-1}$ B[a]P) were included in each experiment. HepG2 cells were seeded at a density of 40,000 cells/well into 12-well microtitre plates (Corning Costar Corporation, Corning, NY, USA) and left overnight at 37°C in 5 % CO_2 to attach to the plates. The growth medium was then replaced with fresh medium containing BPs in the following concentrations: $0.1 \text{ } \mu\text{mol L}^{-1}$, $1 \text{ } \mu\text{mol L}^{-1}$, and $10 \text{ } \mu\text{mol L}^{-1}$. The cells were incubated for 4 h and 24 h. At the end of exposure, the cells were washed, trypsinised, and resuspended in fresh medium for the comet assay.

The comet assay was performed as described by Singh et al. (44) with minor modifications (45). Briefly, $30 \text{ } \mu\text{L}$ of cell suspension was mixed with $70 \text{ } \mu\text{L}$ of 1 % LMP agarose and added to fully frosted slides coated with $80 \text{ } \mu\text{L}$ of 1 % NMP agarose. The cells were then incubated in a lysis solution (2.5 mol L^{-1} NaCl, 100 mmol L^{-1} EDTA, 10 mmol L^{-1} Tris, 1 % Triton X-100, pH 10) at 4°C for at least 1 h, at which point the slides were placed into an alkaline solution (300 mmol L^{-1} NaOH, 1 mmol L^{-1} EDTA, pH 13) at 4°C for 20 min so as to allow DNA unwinding, and electrophoresed at 25 V (300 mA) for 20 min. Finally, the slides were neutralised in a 400 mmol L^{-1} Tris buffer (pH 7.5) for 15 min and stained with EtBr ($5 \text{ } \mu\text{g mL}^{-1}$). Images of 50 randomly selected nuclei per experimental point were captured using a fluorescence microscope (Eclipse 800, Nikon, Tokyo, Japan) and analysed with image analysis software (Comet Assay IV, Perceptive Instruments, Suffolk, UK).

Statistical analysis

The statistical analyses were performed using GraphPad Prism 5 software (GraphPad Software, Inc., San Diego, CA, USA). A one-way analysis of variance (one-way ANOVA) was used to analyse the differences in tail intensity between treatments within each experiment. Dunnett's multiple comparison test was used to compare sample groups with control. Fifty

cells were analysed per experimental point in each of at least two independent experimental cultures. Data are presented as quantile box plots. The edges of the box represent the 25th and 75th percentiles, a line in the box presents the median, and the bars represent 95 % confidence intervals. $P < 0.05$ was considered to be statistically significant. In all experiments, the results of BP-treated cells are compared with those of solvent control cells.

RESULTS AND DISCUSSION

Ames test

Table 2 shows that neither BPA nor any of its analogues were mutagenic. Four of the bisphenols tested in our study, namely BPA, BPF, BPZ, and BPS, have already tested non-mutagenic in *S. typhimurium* TA98 and TA100 either with or without S9 metabolic activation, and our results have confirmed these findings (20-21, 30, 46-48). As the remaining five bisphenols (BPAF, DMBPA, DMBPS, BP-1 and BP-2) have not been tested for mutagenicity using the Ames test, our findings are the first to show that BPAF, DMBPA, DMBPS, BP-1, and BP-2 are not mutagenic in *S. typhimurium* TA98 and TA100 with and without S9 metabolic activation at the tested concentrations.

However, judging by the density of the background lawn, analogues DMBPA and BP-2 at 0.5 mg per plate, and BPZ and BPAF at 0.1 mg and 0.5 mg per plate were toxic to both *S. typhimurium* strains in the presence and in the absence of metabolic activation.

In order to detect cross-linking and/or oxidative properties of BPA and its analogues, other bacterial strains like *Salmonella typhimurium* TA102 or *Escherichia coli* WP2 or WP2 (*pKMI01*) should be used.

Bisphenol cytotoxicity

At the concentrations of up to $100 \text{ } \mu\text{mol L}^{-1}$, BPA, BPF, BPZ, BPS, DMBPS, BP-1, and BP-2 did not affect cell viability after 24 h. DMBPA reduced cell viability by 35 % at the highest tested concentration, whereas BPAF reduced cell viability by 50 % at $50 \text{ } \mu\text{mol L}^{-1}$ and by 70 % at $100 \text{ } \mu\text{mol L}^{-1}$ (Table 3).

In their recent study, Audebert et al. (31) reported that 24-hour exposure to BPA ($50 \text{ } \mu\text{mol L}^{-1}$ and $100 \text{ } \mu\text{mol L}^{-1}$) and BPF ($100 \text{ } \mu\text{mol L}^{-1}$) reduced the viability of several cell lines, including HepG2 cells.

Table 2 Mutagenic effects of nine bisphenols, determined by the Ames test with *S. typhimurium* strains TA98 and TA100 in the presence and absence of an S9 mix. Revertants are presented as means of triplicate plates \pm standard deviation.

Sample	Added / μg per plate	TA98 + S9 mix		TA98 - S9 mix		TA100 + S9 mix		TA100 - S9 mix	
		Revertants	IF ^a	Revertants	IF	Revertants	IF	Revertants	IF
C ^b (DMSO)		32.3 \pm 1.5	1.0	26.7 \pm 5.1	1.0	97.3 \pm 9.6	1.0	103.0 \pm 6.6	1.0
PC ^c (B[a]P)	10	228 \pm 42.3	7.1			784.0 \pm 215.3	8.1		
PC ^d (4-NQNO)	0.5			408.0 \pm 69.4	15.3			2725.3 \pm 193.5	26.5
BPA	4	42.3 \pm 9.5	1.3	27.3 \pm 2.1	1.0	92.3 \pm 17.2	1.0	97.3 \pm 7.2	0.9
	20	42.7 \pm 7.0	1.3	24.3 \pm 2.5	0.9	79.7 \pm 7.2	0.8	105.3 \pm 22.0	1.0
	100	48.3 \pm 4.2	1.5	26.0 \pm 4.4	1.0	105.3 \pm 18.4	1.1	97.7 \pm 1.5	1.0
	500	32.3 \pm 3.2	1.0	22.0 \pm 2.8	0.8	91.7 \pm 5.5	1.0	T ^e	
BPF	4	41.3 \pm 7.8	1.3	27.0 \pm 7.0	1.0	108.0 \pm 16.7	1.1	102.7 \pm 9.5	1.0
	20	40.0 \pm 6.1	1.2	31.0 \pm 8.0	1.1	100.3 \pm 19.4	1.0	102.3 \pm 19.9	1.0
	100	38.3 \pm 2.1	1.2	29.5 \pm 6.4	1.1	109.3 \pm 5.9	1.1	100.7 \pm 9.9	1.0
	500	33.7 \pm 0.6	1.0	23.3 \pm 2.1	0.9	103.7 \pm 11.7	1.1	112.7 \pm 16.7	1.1
BPAF	4	34.3 \pm 8.1	1.1	33.0 \pm 2.6	1.2	94.7 \pm 7.6	1.0	92.0 \pm 10.4	0.9
	20	41.3 \pm 5.9	1.3	33.7 \pm 7.6	1.3	114.3 \pm 5.7	1.2	107.7 \pm 11.6	1.1
	100	T		T		T		T	
	500	T		T		T		T	
BPZ	4	37.0 \pm 7.9	1.1	31.3 \pm 6.4	1.2	98.3 \pm 8.6	1.0	98.0 \pm 4.6	1.0
	20	50.3 \pm 4.5	1.6	32.7 \pm 3.2	1.2	89.0 \pm 15.1	0.9	99.7 \pm 14.3	1.0
	100	43.7 \pm 5.1	1.4	T		T		107.7 \pm 10.5	1.1
	500	T + P ^f		T + P		T + P		T + P	
DMBPA	4	32.3 \pm 10.6	1.0	29.0 \pm 3.5	1.1	95.7 \pm 7.8	1.0	114.3 \pm 13.6	1.1
	20	44.3 \pm 2.1	1.4	21.7 \pm 0.6	0.8	107.7 \pm 6.4	1.1	115.3 \pm 5.7	1.1
	100	43.0 \pm 4.6	1.3	24.0 \pm 7.5	0.9	113.3 \pm 15.3	1.1	114.7 \pm 15.0	1.1
	500	T		T		T		T	
BPS	4	38.7 \pm 4.0	1.2	23.0 \pm 3.6	0.9	83.7 \pm 10.0	0.9	98.7 \pm 7.0	1.0
	20	37.3 \pm 5.1	1.1	25.7 \pm 1.2	1.0	107.0 \pm 11.3	1.1	93.0 \pm 13.0	0.9
	100	33.7 \pm 1.5	1.0	27.3 \pm 3.8	1.0	101.3 \pm 13.9	1.0	90.0 \pm 6.6	0.9
	500	36.3 \pm 5.0	1.1	21.7 \pm 4.6	0.8	97.3 \pm 7.2	1.0	78.0 \pm 6.2	0.8
DMBPS	4	35.0 \pm 7.2	1.1	26.3 \pm 4.6	1.0	108.0 \pm 20.0	1.1	105.7 \pm 4.7	1.0
	20	42.7 \pm 3.1	1.3	33.7 \pm 4.0	1.3	114.3 \pm 6.5	1.2	110.3 \pm 6.7	1.1
	100	38.7 \pm 8.3	1.2	28.0 \pm 3.0	1.1	105.0 \pm 20.0	1.1	115.7 \pm 2.1	1.1
	500	35.7 \pm 0.6	1.1	30.5 \pm 0.7	1.1	111.0 \pm 4.0	1.1	112.7 \pm 9.1	1.1
BP-1	4	34.0 \pm 6.0	1.1	27.0 \pm 4.6	1.0	112.3 \pm 9.0	1.2	125.3 \pm 14.5	1.2
	20	32.7 \pm 5.5	1.0	22.7 \pm 5.0	0.9	105.3 \pm 19.9	1.1	110.7 \pm 13.1	1.1
	100	30.0 \pm 1.7	0.9	26.0 \pm 2.0	1.0	92.7 \pm 17.2	1.0	111.0 \pm 18.3	1.1
	500	39.0 \pm 7.0	1.2	28.5 \pm 0.7	1.1	100.0 \pm 1.0	1.0	108.0 \pm 13.5	1.1
BP-2	4	40.3 \pm 1.5	1.3	31.0 \pm 4.2	1.1	101.0 \pm 4.0	1.0	104.0 \pm 10.8	1.0
	20	39.7 \pm 4.5	1.2	27.7 \pm 3.5	1.0	103.0 \pm 9.5	1.1	114.0 \pm 5.2	1.1
	100	42.7 \pm 3.1	1.3	24.3 \pm 1.2	0.9	97.7 \pm 8.5	1.0	113.0 \pm 6.9	1.1
	500	T		T		T		T	

^a IF: induction factor;^b solvent control (100 μL per plate);^c positive control benzo[a]pyrene (+S9);^d positive control 4-nitroquinoline (-S9);^e T: toxic effect (reduced background lawn);^f P: precipitation;

However, they determined cell viability by measuring DNA content with in-cell western (ICW) assays, while we used an MTT assay that is based on the measurement

of the metabolic activity of living cells. This may explain the differences in the observed cytotoxicity of BPA and BPF between the two studies.

Bisphenol-induced DNA strand breaks in HepG2 cells

HepG2 cells showed a significant increase in DNA strand breaks after four hours of exposure to DMBPS, BP-1, and BP-2. This increase was observed even at the lowest tested concentration of $0.1 \mu\text{mol L}^{-1}$, but no dose-response relationship was observed. In contrast, BPA, BPF, BPAF, BPZ, DMBPA, and BPS did not induce a significant increase in DNA strand breaks (Figure 1).

After 24 h of exposure, a significant increase in DNA strand breaks was observed only in the cells exposed to BPA in all concentrations and to BPS at $0.1 \mu\text{mol L}^{-1}$ and $10 \mu\text{mol L}^{-1}$. Again, no dose-response relationship was observed (Figure 1).

The selected concentrations of BPs are below cytotoxic but are still relevant for or even higher than human exposure (1). The issues surrounding low-dose effects of endocrine disrupting chemicals have been discussed at the National Program (NTP) Workshop on Low Dose Effects of Endocrine Disrupting Chemicals (49-50). Types of DNA damage detected with the alkaline comet assay include single- and double-strand breaks, alkali-labile sites, DNA-DNA/DNA-protein cross-links, and single-strand breaks associated with incomplete excision repair (51). However, in most cases the detected DNA strand breaks and alkali-labile sites are the intermediates formed during the repair of different types of DNA lesions by base or nucleotide excision repair (52). As DNA lesions detected with the comet assay may be transient due to the effective DNA repair, we observed short-term (4-hour) and long-term (24-hour) exposure to BPs.

Our BPA findings are in agreement with several recent studies. Iso et al. (53) reported that BPA induced DNA strand breaks in ER-positive MCF-7 cells (at $1 \mu\text{mol L}^{-1}$ and $100 \mu\text{mol L}^{-1}$) and that its genotoxicity was ER-dependent, as evidenced by much lower effect in ER-negative MDA-MB-231 cells. The increase in DNA strand breaks was significant as soon as after three hours of exposure and increased even further up to hour 24, which is in agreement with our study. When applied *in vivo* in rats, BPA induced micronucleus formation and structural chromosome aberrations in bone marrow as well as DNA damage in lymphocytes (46). Tiwari et al. (46) also observed increased plasma levels of 8-hydroxydeoxyguanosine, an increase in lipid peroxidation, and a decrease in glutathione activity in the liver, suggesting that oxidative stress

could be one of the mechanisms of BPA genotoxicity. In a study by Audebert et al. (31) on the other hand, BPA failed to induce H2AX histone phosphorylation in HepG2 cells, but induced it in human renal adenocarcinoma (ACHN) cells. The authors explained this difference between the two cell lines with differences in the biotransformation of BPA.

In our study, BPF at concentrations up to $10 \mu\text{mol L}^{-1}$ did not induce DNA damage. This result is in line with the study by Cabaton et al. (30), who did not detect DNA strand breaks in HepG2 cells exposed to BPF at concentrations lower than $50 \mu\text{mol L}^{-1}$. At higher concentrations, however, they observed a significant increase in DNA strand breaks. BPF also induced a significant increase in H2AX histone phosphorylation in HepG2, ACHN, and human epithelial colorectal adenocarcinoma (LS174T) cells, a phenomenon that was observed only at the two highest concentrations tested ($50 \mu\text{mol L}^{-1}$ and $100 \mu\text{mol L}^{-1}$) (31).

In HepG2 cells exposed to low concentrations of DMBPS, BP-1, and BP-2, the increase in DNA strand breaks was significant after four, but not after 24 h of exposure, whereas in cells exposed to BPA and BPS, DNA strand breaks were observed only after the 24-hour exposure. However, these changes were small and not dose-related. Even so, our results do not undermine the genotoxic potential of BPA analogues and call for further research. Scientific evidence supports the hypothesis that natural oestrogens, synthetic oestrogen diethylstilbestrol, as well as BPA generate reactive oxygen species (ROS) during biotransformation and that certain reactive species, predominantly quinones, can react with DNA and cause DNA damage (54). One pathway of BPA metabolism is the hydroxylation of one of its symmetric phenyl rings to form its catechol, o-OH BPA, which can oxidise to o-quinone BPA (55) and, in turn, react with DNA. o-Quinone BPA forms predominantly depurinating adducts o-OH-BPA-6-N3Ade and o-OH-BPA-6-N7Gua (56-58). Sakuma et al. (59) found that o-quinone BPA could increase ROS formation and oxidise the guanine moiety of deoxyguanosine in the DNA of primary rat hepatocyte cultures. Adducts such as these, formed during BPA metabolism as well as oxidative DNA damage are readily detected by the alkaline comet assay.

BPA analogues included in our study differ in the bridging atom between the two phenyl rings [sulfone (BPS) and sulfide (BP-2) moieties instead of a carbon atom], in the functional groups on the bridging carbon




Figure 1 DNA damage (comet assay) induced by bisphenols in HepG2 cells after 4 h and 24 h of exposure to concentrations of 0.1 µmol L⁻¹, 1 µmol L⁻¹, and 10 µmol L⁻¹.

B[a]P (30 µmol L⁻¹) was used as the positive control. The level of DNA strand breaks is expressed as tail intensity. Fifty cells were analysed per experimental point in each of at least two independent experimental cultures. Data are presented as quantile box plots. The edges of the box represent the 25th and 75th percentile, a line in the box represents the median value, and the bars represent 95 % confidence intervals.

* significant difference (one-way ANOVA; Dunnett's multiple comparison test) between BPs-treated cells and solvent control (0),

** (P<0.01) and *** (P<0.001)

Table 3 Viability of HepG2 cells exposed to bisphenols, as determined by the MTT assay.^a

Concentration μmol L ⁻¹	Viability / %								
	BPA	BPF	BPAF	BPZ	DMBPA	BPS	DMBPS	BP-1	BP-2
12.5	110±6	114±5	107±16	99±7	121±8	99 ±2	101±7	108±8	101±9
25	114±19	112±6	100±21	105±12	123±10	100±7	108±8	106±5	103±9
50	116±9	118±12	50±4*	113±9	100±13	98±8	118±10	109±3	113±5
100	90±15	123±13	31±4*	125±14	67±12*	94±4	129±12	113±13	117±10

^a Independent experiments were performed in 5 replicates and repeated two times. Data show one out of two independent experiments, which showed the same trend. The survival was normalized to the solvent control (100 % viability). Values are expressed as mean ± SD.

* Significant difference between the solvent control and BP-treated cells (Student's t-test, P<0.001).

atom [without methyl groups (BPF), with two trifluoromethyl groups (BPAF), and with a cyclohexyl ring (BPZ)], and in substitutions on both phenol rings [methyl groups on positions 3 of phenyl rings (DMBPA and DMBPS) and substituted both phenol groups (BP-1)] (Table 1). These structural differences may explain the differences in their metabolism in HepG2 cells and the extent of formation of reactive quinone intermediates and ROS. However, structural differences alone cannot clearly explain the structure-activity relationship in this series of BPA analogues or why BPF, BPAF, BPZ, and DMBPA did not induce DNA damage, and why BPA and BPS did induce significant DNA damage only after 24 h of exposure. It has recently been demonstrated that BPA and BPF are metabolised in HepG2 cells predominantly to conjugated sulphate metabolites (31, 60). However, we still do not know to what extent BPA and its analogues are biotransformed into reactive quinone intermediates in HepG2 cells, and how stable these intermediates are. Less stable intermediates are probably less harmful, due to their rapid reaction with water molecules, while more stable intermediates can also react with biological molecules and be more harmful. It has been hypothesised that those natural products that undergo oxidation to quinones and are then rapidly hydrated are unlikely to damage important biological macromolecules (61).

In conclusion, neither BPA nor its analogues induced bacterial mutations, and the minor and transient DNA damage induced by BPA, DMBPS, BP-1, and BP-2 in HepG2 cells was observed at concentrations that are higher than human exposure. Our data suggest that the toxic potential of BPA lies in the formation of reactive quinone metabolites and oxidative stress and warrant further investigation of

genotoxic and mutagenic effects of other BPA analogues.

Acknowledgements

This work was supported by the Slovenian Research Agency (Grant No. Program P1-0208 and Program P1-0245). We wish to thank Professor Roger Pain for helpful comments and critical reading of the manuscript.

REFERENCES

1. World Health Organization (WHO). Joint FAO/WHO Expert Meeting to Review Toxicological and Health Aspects of Bisphenol A [displayed 30 May 2011]. Available at http://www.who.int/entity/foodsafety/chem/chemicals/BPA_Summary2010.pdf
2. Maffini MV, Rubin BS, Sonnenschein C, Soto AM. Endocrine disruptors and reproductive health: The case of bisphenol-A. *Mol Cell Endocrinol* 2006;254:179-86.
3. Rasier G, Toppari J, Parent AS, Bourguignon JP. Female sexual maturation and reproduction after prepubertal exposure to estrogens and endocrine disrupting chemicals: A review of rodent and human data. *Mol Cell Endocrinol* 2006;254:187-201.
4. Screening Assessment for the Challenge Phenol, 4,4' -(1-methylethylidene)bis- (Bisphenol A). Chemical Abstracts Service Registry Number 80-05-7. Environment Canada 2008 [displayed 11 Sept 2012]. Available at http://www.ec.gc.ca/ese-ees/3C756383-BEB3-45D5-B8D3-E8C800F35243/batch2_80-05-7_en.pdf
5. Appleton. Nation's Largest Maker of Thermal Receipt Paper Does Not Use BPA [displayed at 23 May 2013]. Available at <http://www.appletonideas.com/pdf/Appleton%20BPA%20free%20news%20release.7.27.2010.pdf>
6. EU directive 2011/8/EU. Commission directive 2011/8/EU of 28 January 2011. [displayed 3 June 2011]. Available at <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2011:026:0011:0014:EN:PDF>

7. Chen MY, Ike M, Fujita M. Acute toxicity, mutagenicity, and estrogenicity of bisphenol-A and other bisphenols. *Environ Toxicol* 2002;17:80-6. doi: 10.1002/tox.10035
8. Kuruto-Niwa R, Nozawa R, Miyakoshi T, Shiozawa T, Terao Y. Estrogenic activity of alkylphenols, bisphenol S, and their chlorinated derivatives using a GFP expression system. *Environ Toxicol Pharmacol* 2005;19:121-30. doi: 10.1016/j.etap
9. Raloff J. Receipts a large - and largely ignored - source of BPA. *ScienceNews.org*. [displayed 30 August 2010]. Available at http://www.sciencenews.org/view/generic/id/61764/title/Receipts_a_large_%E2%80%94_and_largely_ignored_%E2%80%94_source_of_BPA
10. Danzl E, Sei K, Soda S, Ike M, Fujita M. Biodegradation of bisphenol A, bisphenol F and bisphenol S in Seawater. *Int J Environ Res Public Health* 2009;6:1472-84. doi: 10.3390/ijerph6041472
11. Ike M, Chen MY, Danzl E, Sei K, Fujita M. Biodegradation of a variety of bisphenols under aerobic and anaerobic conditions. *Water Sci Technol* 2006;53:153-9. doi: 10.3390/ijerph6041472
12. US Patent. 1978. 4,098,769 [displayed 8 December 2011]. Available at <http://www.patents.com/us-4098769.html>
13. Lotti N, Colonna M, Fiorini M, Finelli L, Berti C. Poly(butylene terephthalate) modified with ethoxylated bisphenol S with increased glass transition temperature and improved thermal stability. *Polymer* 2011;52:904-11. doi: 10.1016/j.polymer.2011.01.018
14. Kitamura S, Suzuki T, Sanoh S, Kohta R, Jinno N, Sugihara K, Yoshihara S, Fujimoto N, Watanabe H, Ohta S. Comparative Study of the Endocrine-Disrupting Activity of Bisphenol A and 19 Related Compounds. *Toxicol Sci* 2005;84:249-59. doi: 10.1093/toxsci/kfi074
15. Tsutsui T, Tamura Y, Suzuki A, Hirose Y, Kobayashi M, Nishimura H, Metzler M, Barrett JC. Mammalian cell transformation and aneuploidy induced by five bisphenols. *Int J Cancer* 2000;86:151-4. doi: 10.1002/(SICI)1097-0215(20000415)86:2<151::AID-IJC1>3.0.CO;2-0
16. National Toxicology Program (NTP). Chemical Information Profile for Bisphenol AF [CAS No. 1478-61-1], Supporting Nomination for Toxicological Evaluation by the National Toxicology Program [displayed 30 March 2011]. Available at http://ntp.niehs.nih.gov/ntp/htdocs/Chem_Background/ExSumPdf/BisphenolAF_093008_508.pdf
17. Doherty LF, Bromer JG, Zhou Y, Aldad TS, Taylor HS. In utero exposure to diethylstilbestrol (DES) or bisphenol-A (BPA) increases EZH2 expression in the mammary gland: an epigenetic mechanism linking endocrine disruptors to breast cancer. *Horm Cancer* 2010;1:146-55. doi: 10.1007/s12672-010-0015-9
18. International Agency for Research on Cancer (IARC). International Agency for Research on Cancer, Sex Hormones (II). IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Vol. 21. Lyon: IARC; 1979.
19. ChemBio3D Ultra 13.0 [Computer software] in ChemBioOffice® Ultra 13.0, PerkinElmer, www.cambridgesoft.com/, Release Date: August, 2012.
20. Ashby J, Tennant RW. Chemical-structure, Salmonella mutagenicity and extent of carcinogenicity as indicators of genotoxic carcinogenesis among 222 chemicals tested in rodents by the United-States NCI/NTP. *Mutat Res* 1988;204:17-115.
21. Schweikl H, Schmalz G, Rackebrandt K. The mutagenic activity of unpolymerized resin monomers in Salmonella typhimurium and V79 cells. *Mutat Res* 1998;415:119-30.
22. Tsutsui T, Tamura Y, Yagi E, Hasegawa K, Takahashi M, Maizumi N, Yamaguchi F, Barrett JC. Bisphenol-A induces cellular transformation, aneuploidy and DNA adduct formation in cultured Syrian hamster embryo cells. *Int J Cancer* 1998;75:290-4. doi: 10.1002/(SICI)1097-0215(19980119)75:2<290::AID-IJC19>3.0.CO;2-H
23. Ivett JL, Brown BM, Rodgers C, Anderson BE, Resnick MA, Zeiger E. Chromosomal aberrations and sister chromatid exchange tests in Chinese hamster ovary cells *in vitro*. IV. Results with 15 chemicals. *Environ Mol Mutagen* 1989;14:165-87. doi: 10.1002/em.2850140306
24. Naik P, Vijayaaxmi KK. Cytogenetic evaluation for genotoxicity of Bisphenol-A in bone marrow cells of Swiss albino mice. *Mutat Res* 2009;676:106-12. doi: 10.1016/j.mrgentox.2009.04.010
25. Atkinson A, Roy D. *In vitro* conversion of environmental estrogenic chemical bisphenol A to DNA binding metabolite(s). *Biochem Biophys Res Commun* 1995;210:424-33.
26. Edmonds JS, Nomachi M, Terasaki M, Morita M, Skelton BW, White AH. The reaction of bisphenol A 3,4-quinone with DNA. *Biochem Biophys Res Commun* 2004;319:556-61. doi: 10.1016/j.bbrc.2004.05.024
27. Atkinson A, Roy D. *In vivo* DNA adduct formation by bisphenol A. *Environ Mol Mutagen* 1995;26:60-6. doi: 10.1002/em.2850260109
28. Izzotti A, Kanitz S, D'Agostini F, Camoirano A, De Flora S. Formation of adducts by bisphenol A, an endocrine disruptor, in DNA *in vitro* and in liver and mammary tissue of mice. *Mutat Res* 2009;679:28-32. doi: 10.1016/j.mrgentox.2009.07.011
29. Iso T, Watanabe T, Iwamoto T, Shimamoto A, Furuichi Y. DNA damage caused by bisphenol A and estradiol through estrogenic activity. *Biol Pharm Bull* 2006;29:206-10. doi:10.1248/bpb.29.206
30. Cabaton N, Dumont C, Severin I, Perdu E, Zalko D, Cherkaoui-Malki M, Chagnon MC. Genotoxic and endocrine activities of bis(hydroxyphenyl)methane (bisphenol F) and its derivatives in the HepG2 cell line. *Toxicology* 2009;255:15-24. doi: 10.1016/j.tox.2008.09.024
31. Audebert M, Dolo L, Perdu E, Cravedi JP, Zalko D. Use of the gamma H2AX assay for assessing the genotoxicity of bisphenol A and bisphenol F in human cell lines. *Arch Toxicol* 2011;85:1463-73. doi: 10.1007/s00204-011-0721-2
32. Pfeiffer E, Rosenberg B, Deuschel S, Metzler M. Interference with microtubules and induction of micronuclei *in vitro* by various bisphenols. *Mutat Res* 1997;390:21-31.
33. Kanai H, Barrett JC, Metzler M, Tsutsui T. Cell-transforming activity and estrogenicity of bisphenol-A and 4 of its analogs in mammalian cells. *Int J Cancer* 2001;93:20-5. doi: 10.1002/ijc.1303
34. Maron DM, Ames BN. Revised methods for the Salmonella mutagenicity test. *Mutat Res* 1983;113:173-215.
35. Kenyon MO, Cheung JR, Dobo KL, Ku WW. An evaluation of the sensitivity of the Ames assay to discern low-level mutagenic impurities. *Regul Toxicol Pharmacol* 2007;48:75-86. doi: 10.1016/j.yrtph.2007.01.006
36. Westerink WM, Schoonen WG. Phase II enzyme levels in HepG2 cells and cryopreserved primary human hepatocytes

- and their induction in HepG2 cells. *Toxicol In Vitro* 2007;21:1592-602. doi: 10.1016/j.tiv.2007.06.017
37. Westerink WM, Schoonen WG. Cytochrome P450 enzyme levels in HepG2 cells and cryopreserved primary human hepatocytes and their induction in HepG2 cells. *Toxicol In Vitro* 2007;21:1581-91. doi: 10.1016/j.tiv.2007.05.014
 38. Wilkening S, Stahl F, Bader A. Comparison of primary human hepatocytes and hepatoma cell line HepG2 with regard to their biotransformation properties. *Drug Metab Dispos* 2003;31:1035-42. doi: 10.1124/dmd.31.8.1035
 39. Hreljac I, Filipič M. Organophosphorus pesticides enhance the genotoxicity of benzo(a)pyrene by modulating its metabolism. *Mutat Res* 2009;671:84-92. doi: 10.1016/j.mrfmmm.2009.09.011
 40. Knasmüller S, Parzefall W, Sanyal R, Ecker S, Schwab C, Uhl M, Mersch-Sundermann V, Williamson G, Hietsch G, Langer T, Darroudi F, Natarajan AT. Use of metabolically competent human hepatoma cells for the detection of mutagens and antimutagens. *Mutat Res* 1998;402:185-202.
 41. Solakidi S, Sarra AMG, Sekeris CE. Differential subcellular distribution of estrogen receptor isoforms: Localization of ER α in the nucleoli and ER β in the mitochondria of human osteosarcoma SaOS-2 and hepatocarcinoma HepG2 cell lines. *Biochim Biophys Acta - Mol Cell Res* 2005;1745:382-92. doi: 10.1016/j.bbamcr.2005.05.010
 42. Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Meth* 1983;65:55-63. doi: 10.1016/0022-1759(83)90303-4
 43. Žegura B, Zaje I, Lah TT, Filipič M. Patterns of microcystin-LR induced alteration of the expression of genes involved in response to DNA damage and apoptosis. *Toxicol* 2008;51:615-23. doi: 10.1016/j.toxicol.2007.11.009
 44. Singh NP, McCoy MT, Tice RR, Schneider EL. A simple technique for quantitation of low levels of DNA damage in individual cells. *Exp Cell Res* 1988;175:184-91.
 45. Žegura B, Filipič M. Application of *in vitro* comet assay for genotoxicity testing. In: Yan Z, Caldwell G, editors. *Optimization in drug discovery: In vitro methods*. Totowa (NJ): Humana Press; 2004. p. 301-13.
 46. Tiwari D, Kamble J, Chilgunde S, Patil P, Maru G, Kawle D, Bhartiya U, Joseph L, Vanage G. Clastogenic and mutagenic effects of bisphenol A: an endocrine disruptor. *Mutat Res* 2012;743:83-90. doi: 10.1016/j.mrgentox.2011.12.023
 47. Yamaguchi T, Yamauchi A, Yamazaki H, Kakiuchi Y. Mutagenicity of rubber additives in tire. *Eisei Kagaku* 1991;37:6-13.
 48. Japan Chemical Industry Ecology-Toxicology & Information Center (JETOC). *Mutagenicity Test Data of Existing Chemical Substances Based on the Toxicity Investigation System of the Industrial Safety and Health Law (I)*. Tokyo: JETOC; 1996.
 49. National Institute of Environmental Health Sciences (NIH). National Toxicology Program. 2001. National Toxicology Program's Report of the Endocrine Disruptors Low-Dose Peer Review [displayed 20 Dec 2011]. Available at www.epa.gov/endo/pubs/edmvslowdosepeerfinalrpt.pdf
 50. Richter CA, Birnbaum LS, Farabollini F, Newbold RR, Rubin BS, Talsness CE, Vandenberg JG, Walser-Kuntz DR, vom Saal FS. *In vivo* effects of bisphenol A in laboratory rodent studies. *Reprod Toxicol* 2007;24:199-224.
 51. Tice RR, Agurell E, Anderson D, Burlinson B, Hartmann A, Kobayashi H, Miyamae Y, Rojas E, Ryu JC, Sasaki YF. Single cell gel/comet assay: guidelines for *in vitro* and *in vivo* genetic toxicology testing. *Environ Mol Mutagen* 2000;35:206-21.
 52. Collins AR, Dobson VL, Dušinska M, Kennedy G, Štetiina R. The comet assay: what can it really tell us? *Mutat Res* 1997;375:183-93.
 53. Iso T, Futami K, Iwamoto T, Furuichi Y. Modulation of the expression of Bloom helicase by estrogenic agents. *Biol Pharm Bull* 2007;30:266-71. doi: 10.1248/bpb.30.266
 54. Cavalieri EL, Rogan EG. Is Bisphenol A a weak carcinogen like the natural estrogens and diethylstilbestrol? *IUBMB Life* 2010;62:746-51. doi: 10.1002/iub.376
 55. Schmidt J, Kotnik P, Trontelj J, Knez Z, Mašič LP. Bioactivation of bisphenol A and its analogs (BPF, BPAF, BPZ and DMBPA) in human liver microsomes. *Toxicol In Vitro* 2013;27:1267-76. doi: 10.1016/j.tiv.2013.02.016
 56. Jaeg JP, Perdu E, Dolo L, Debrauwer L, Cravedi JP, Zalko D. Characterization of new bisphenol A metabolites produced by CD1 mice liver microsomes and S9 fractions. *J Agric Food Chem* 2004;52:4935-42. doi: 10.1021/jf049762u
 57. Nakagawa Y, Suzuki T. Metabolism of bisphenol A in isolated rat hepatocytes and oestrogenic activity of a hydroxylated metabolite in MCF-7 human breast cancer cells. *Xenobiotica* 2001;31:113-23.
 58. Kolšek K, Mavri J, Sollner Dolenc M. Reactivity of bisphenol A-3,4-quinone with DNA. A quantum chemical study. *Toxicol in Vitro* 2012;26:102-6. doi: 10.1016/j.tiv.2011.11.003
 59. Sakuma S, Nakanishi M, Morinaga K, Fujitake M, Wada S, Fujimoto Y. Bisphenol A 3,4-quinone induces the conversion of xanthine dehydrogenase into oxidase *in vitro*. *Food Chem Toxicol* 2010;48:2217-22. doi: 10.1016/j.fct.2010.05.051
 60. Bursztyka J, Perdu E, Pettersson K, Pongratz I, Fernandez-Cabrera M, Olea N, Debrauwer L, Zalko D, Cravedi JP. Biotransformation of genistein and bisphenol A in cell lines used for screening endocrine disruptors. *Toxicol in Vitro* 2008;22:1595-604. doi: 10.1016/j.tiv.2008.06.013
 61. Tu T, Gliblin D, Gross mL. Structural determinant of chemical reactivity and potential health effects of quinones from natural products. *Chem Res Toxicol* 2011;24:1527-39. doi: 10.1021/tx200140s

Povzetek**MUTAGENOST IN POŠKODBE DNA POVZROČENE Z BISFENOLOM A IN NJEGOVIMI STRUKTURNIMI ANALOGI V CELIČNI LINIJI HEPG2**

Okoljski estrogen, bisfenol A (BPA), in njegovi strukturni analogi so v veliki meri prisotni v našem okolju. Ker njihova proizvodnja in uporaba naraščata, je vse pomembneje ovrednotiti njihovo toksičnost zaradi izpostavljenosti ljudem. Z Amesovim in kometnim testom smo ovrednotili mutagenost in genotoksičnost osmih strukturnih analogov BPA (BPF, BPAF, BPZ, BPS, DMBPA, DMBPS, BP-1 in BP-2). Nobeden od testiranih bisfenolov ni izkazoval mutagenega delovanja na sevih TA98 in TA100 *Salmonella typhimurium* v prisotnosti in odsotnosti metabolne aktivacije (z Aroklorom 1254 inducirani encimi podganjih jeter). Potencialno genotoksičnost pa smo določali s kometnim testom na celični liniji humanega hepatoma (HepG2) pri necitotoksičnih koncentracijah ($0.1 \mu\text{mol L}^{-1}$ do $10 \mu\text{mol L}^{-1}$) po 4-urni in 24-urni izpostavljenosti. BPA in njegov analog BPS sta pri kometnem testu povzročila poškodbe DNA samo po 24-urni izpostavljenosti, medtem ko so analogi DMBPS, BP-1 in BP-2 povzročili prehodne poškodbe DNA (samo po 4-urni izpostavljenosti). BPF, BPAF, BPZ in DMBPA niso povzročili poškodb DNA.

KLJUČNE BESEDE: *Amesov test, bisfenoli, kometni test*

CORRESPONDING AUTHOR:

Lucija Peterlin Mašič
Faculty of Pharmacy, University of Ljubljana
Aškerčeva 7, 1000 Ljubljana, Slovenia
E-mail address: lucija.peterlin@ffa.uni-lj.si