COMPARISON OF TWO METHODS FOR THE ESTIMATION OF STABILITY OF COPPER(II) BIS-COMPLEXES WITH AROMATIC LIGANDS RELEVANT TO ALZHEIMER'S DISEASE

Ante MILIČEVIĆ and Nenad RAOS

Institute for Medical Research and Occupational Health, Zagreb, Croatia

Received in July 2013
CrossChecked in July 2013
Accepted in August 2013

In order to compare the density functional theory (DFT) and the method with the molecular valence connectivity index of the 3rd order (${}^3\chi^v$) in the estimation of stability constants $\log \beta_2$, we used copper(II) complexes with thioflavin-based intercalation compounds designed for application in Alzheimer's disease. Correlation of ten $\log \beta_2$ values calculated by DFT with ${}^3\chi^v$ yielded $r=0.988$ and $S.E.=0.85$. The correlations of indices I_{NG} , HOMA and ${}^3\chi^v$ with $\log \beta_2$, and the intercorrelations between the indices revealed the crucial influence of ${}^3\chi^v$ on the prediction of the stability of these complexes.

KEY WORDS: *aromaticity, density-functional theory (DFT), regression models, stability constants, topological indices*

Alzheimer's disease (AD), the leading mental disorder of the elderly population in developed countries, is closely related to copper metabolism (1). Next to the liver, the brain contains the second highest cellular concentration of copper, which is more frequent in grey matter ($60 \mu\text{mol L}^{-1}$ to $110 \mu\text{mol L}^{-1}$) than in white matter ($25 \mu\text{mol L}^{-1}$ to $79 \mu\text{mol L}^{-1}$) (2). Patients with AD have elevated copper(II) concentrations in whole blood (3). Even more importantly, copper(II) concentrations in the senile plaques of AD patients reach $400 \mu\text{mol L}^{-1}$, which is nearly four times as much as usual grey matter content (2). These findings led George J. Brewer, Emeritus Professor at the Michigan Medical School, to the hypothesis that AD could be caused by copper(II) poisoning (4, 5). This hypothesis is further supported by the finding that this disease is still virtually unknown in developing countries, as was the case in developed countries 50 to 60 years ago. Professor

Brewer found the principal reason for the epidemic of AD in elevated copper concentrations in drinking water caused by copper plumbing (6) and extensive red meat (especially beef) (7, 8) and multi-vitamin mineral pill consumption (9).

This hypothesis is also supported by research on the molecular level. The main mechanism in the development of AD is the formation of extracellular amyloid plaques in the brain. These plaques are composed of β -amyloid peptides ($A\beta$) (10), a normal cleavage product of the larger membrane amyloid precursor protein (APP) (11). Human $A\beta$ binds copper, in contrast to rodent peptides (due to mutations $\text{Arg5} \rightarrow \text{Gly}$, $\text{Tyr10} \rightarrow \text{Phe}$, $\text{His13} \rightarrow \text{Arg}$). This serves as a very reasonable explanation for the virtual absence of $A\beta$ deposits in normal rodent brains (12).

NMR (13) and AFM (14) studies have shown that copper(II) influences the aggregation behaviour of the $A\beta$ peptide. The second, more vital, influence of

copper in the development of AD is the reduction of Cu^{2+} to Cu^+ via the A β peptide. This process directly produces hydrogen peroxide (15) and, subsequently, reactive oxygen species (ROS) (16, 17). Copper(II) chelators, especially ligands capable of binding to the A β peptide, are therefore viewed as potential drugs and diagnostic tools for the treatment of AD (18, 19). Among these compounds is a class of ligands with a common core of the thioflavin T (ThT) and clioquinol (CQ) molecule (Figure 1). They were designed with the notion that they possess the lipophilicity adequate for crossing the blood-brain barrier as well as potential antioxidant properties. They also have a structure suitable for iodine substitution to the ligating group at the para position, which enables their use in diagnostics (20, 21).

After testing the $^3\chi^v$ index on various copper(II) (23), bivalent transition metal (24), and lanthanide complexes (25), but not yet on aromatic compound complexes, we found these copper(II) chelators suitable for testing and developing our method for the prediction of stability constants of coordination compounds based on the connectivity index $^3\chi^v$ (22). Unfortunately, among the ten complexes we studied, only two overall stability constants (β_2) were measured (20). Thus, we correlated the $^3\chi^v$ index to the stability constants ($\log \beta_2$) calculated by the density functional theory (DFT) (21). Also, we studied the referred correlations (21) of aromatic indices I_{NG} and HOMA (used to investigate the influence of Möbius metalloaromaticity (26, 27) on stability) vs. $\log \beta_2$, to compare their influence with that of the $^3\chi^v$ index.

METHODS

Calculation of topological indices

The molecular valence connectivity index of the 3rd order $^3\chi^v$ was calculated using the E-Dragon program system, developed by R. Todeschini and coworkers, which is capable of yielding 119 topological indices in a single run, along with many other molecular descriptors (28, 29). The connectivity matrices were constructed with the aid of the *Online SMILES Translator and Structure File Generator* (30).

The $^3\chi^v$ connectivity index (31-36) was defined as:

$$^3\chi^v = \sum_{\text{path}} [\delta(i) \delta(j) \delta(k) \delta(l)]^{-0.5} \quad (1)$$

Figure 1 Constitutional formulas of aromatic ligands; *X* is NH, O or S.

Figure 2 Graph representation of $[\text{Cu}(\text{N}_\text{O}-\text{O}_2)_2]$. Heteroatoms are marked with $\circ$ (Cu), $\bullet$ (N), $\odot$ (O), and $\odot$ (I).

where $\delta(i)$, $\delta(j)$, $\delta(k)$, and $\delta(l)$ are weights (valence values) of vertices (atoms) i, j, k , and l making up the path of length 3 (three consecutive chemical bonds) in a vertex-weighted molecular graph. The valence value, $d(i)$, of the vertex i was defined by:

$$\delta(i) = [Z^v(i) - H(i)]/[Z(i) - Z^v(i) - 1] \quad (2)$$

where $Z^v(i)$ is the number of valence electrons belonging to the atom corresponding to vertex i , $Z(i)$ its atomic number, and $H(i)$ the number of hydrogen atoms attached to it. It should be stressed that $^3\chi^v$ is only a member of the family of valence connectivity indices $^n\chi^v$, which differ among each other by path length, i.e. the number of δ 's in the summation term (Equation 1).

The $^3\chi^v$ index for all bis-complexes was calculated from their graph representations assuming that the Cu(II) ion is tetracoordinated (Figure 2) (24).

Regression calculations

Regression calculations, including the leave-one-out procedure (LOO) of cross validation were done using the CROMRsel program (37). The standard error of the cross validation estimate was defined as:

$$S.E._{cv} = \sqrt{\sum_i \frac{\Delta X_i^2}{N}} \quad (3)$$

where ΔX and N denote cv residuals and the number of reference points, respectively.

RESULTS AND DISCUSSION

In order to compare the efficiency of the two methods for the estimation of stability constants, DFT and the $^3\chi^v$ index, we referred to $\log \beta_2$ DFT values (21) (Table 1). Although the DFT values {15.1 and 11.0 for $[\text{Cu}(\text{N}_\text{O}-\text{O1})_2]$ and $[\text{Cu}(\text{N}_\text{S}-\text{O1})_2]$, respectively} were far from their experimental values (23.1 and 20.3), as obtained by UV-VIS spectrometry (20), the

authors had, by mistake, stated that the agreement was remarkable (21). However, if we direct our attention to the relative values of $\log \beta_2$ s obtained by either DFT or spectrophotometry, it is evident that $\log \beta_2 [\text{Cu}(\text{N}_\text{O}-\text{O1})_2] > \log \beta_2 [\text{Cu}(\text{N}_\text{S}-\text{O1})_2]$ for both methods, being 4.1 and 2.8 for DFT and spectrophotometry, respectively.

Bearing all this in mind, we reproduced $\log \beta_2$ (DFT) values through our method based on the $^3\chi^v$ index. Simple regression (Figure 3) yielded a correlation coefficient of $r=0.988$ and a standard error of cross-validation of $S.E._{cv}=0.97$ (Model 1, Table 2). Due to such a good correlation, we can deem our method equally successful as the DFT method, but much less demanding.

Table 1 Logarithm of stability constants of copper(II) bis-complexes with 10 aromatic ligands, and values of their metalloaromaticity (I_{NG} and HOMA) and connectivity $^3\chi^v$ indices.

No.	complex	$\log \beta_2^{[a]}$	$\log \beta_2^{[b]}$	$I_{\text{NG}}^{[c]}$	HOMA ^[c]	$^3\chi^v$
1	$[\text{Cu}(\text{N}_{\text{NH}}-\text{O1})_2]$	15.1	15.7	0.0174	0.658	11.753
2	$[\text{Cu}(\text{N}_{\text{NH}}-\text{O2})_2]$	19.1	19.0	0.0167	0.545	9.701
3	$[\text{Cu}(\text{N}_{\text{NH}}-\text{O3})_2]$	24.1	24.6	0.0183	0.848	6.412
4	$[\text{Cu}(\text{N}_\text{O}-\text{O1})_2]$	15.1	16.4	0.0174	0.659	11.404
5	$[\text{Cu}(\text{N}_\text{O}-\text{O2})_2]$	18.8	19.6	0.0172	0.641	9.406
6	$[\text{Cu}(\text{N}_\text{O}-\text{O3})_2]$	24.7	25.0	0.0188	0.915	6.130
7	$[\text{Cu}(\text{N}_\text{S}-\text{O1})_2]$	11.0	11.0	0.0170	0.555	14.504
8	$[\text{Cu}(\text{N}_\text{S}-\text{O2})_2]$	16.1	14.9	0.0172	0.572	12.036
9	$[\text{Cu}(\text{N}_\text{S}-\text{O3})_2]$	22.8	20.6	0.0188	0.894	8.637
10	$[\text{Cu}(\text{N}-\text{O4})_2]$	30.8	30.9	0.0202	0.939	2.613

[a] DFT values from Ref. 21

[b] Cross-validated values (Model 1, Table 2)

[c] Ref. 21

Table 2 Regression models for the estimation of the $\log \beta_2$ of Cu(II) bis-complexes.

model	descriptor	complexes	regression coefficients		r	S.E.	S.E. _{cv}
			slope(S.E.)	intercept(S.E.)			
1	$^3\chi^v$	all	-1.667(91)	35.20(89)	0.988	0.85	0.97
2		1, 2, 3, 10	-1.693(46)	35.17(39)	0.999	0.23	0.40
3		4, 5, 6, 10	-1.785(17)	35.54(14)	0.999	0.08	0.21
4		7, 8, 9, 10	-1.65(13)	35.8(14)	0.994	0.84	2.44
5	I_{NG}	all	4711(897)	-65(16)	0.880	2.64	3.24
6		1, 2, 3, 10	4025(1372)	-51(25)	0.901	2.55	4.50
7		4, 5, 6, 10	4736(974)	-65(18)	0.960	1.67	3.06
8		7, 8, 9, 10	5608(768)	-82(14)	0.982	1.41	2.63
9	HOMA	all	31.9(66)	-3.3(49)	0.863	2.81	3.66
10		1, 2, 3, 10	32(14)	-1(11)	0.854	3.05	7.01
11		4, 5, 6, 10	39(12)	-8.6(98)	0.915	2.40	4.84
12		7, 8, 9, 10	39(11)	-8.7(83)	0.931	2.72	5.45

Figure 3 Dependence of stability constant $\log \beta_2$ (DFT) on the ${}^3\chi^v$ index for Cu(II) bis-complexes with ten aromatic ligands (Table 1, Model 1 in Table 2).

We also analysed reported correlations (24) of indices defining metalloaromaticity, I_{NG} , and HOMA vs. $\log \beta_2$ (Models 6-8, 10-12, Table 2), on three subsets, imida-complexes (1-3 and 10), oxa-complexes (4-6 and 10), and thia-complexes (7-10). Although the stability was positively correlated with I_{NG} and HOMA for all three subsets, the differences in $\log \beta_2$ between complexes 1 and 2 and 4 and 5 showed an opposite trend. From the comparison of these correlations (Models 6-8, 10-12, Table 2) with the correlations obtained with the ${}^3\chi^v$ index (Models 2-4, Table 2), it is clear that ${}^3\chi^v$ is a substantially better index for modelling the stability of the studied complexes than I_{NG} and HOMA. Specifically, standard errors for ${}^3\chi^v$ regressions are an order of magnitude smaller than the standard errors of the regressions with other two indices, with the exception of the thia-complexes subset [7-10, Table 1); S.E. (${}^3\chi^v$):S.E. (I_{NG})=1:1.68 and S.E. (${}^3\chi^v$):S.E. (I_{NG})=1:1.08 (for HOMA the ratios are 1:3.24 and 1:2.23)]. Accordingly, the regression of ${}^3\chi^v$ to $\log \beta_2$ on all complexes (Model 1, Table 2, Figure 3), yielded much better results than the regression with metalloaromaticity indices I_{NG} and HOMA (Models 5 and 9, Table 2, Figure 4).

Bivariate linear regressions of I_{NG} and ${}^3\chi^v$ vs. $\log \beta_2$ ($r=0.990$, S.E.=0.77, S.E._{cv}=1.11) and HOMA and ${}^3\chi^v$ vs. $\log \beta_2$ ($r=0.990$, S.E.=0.79, S.E._{cv}=1.19), did not yield improvement over univariate regression with ${}^3\chi^v$. To the contrary, S.E._{cv} rose from 0.97 to 1.11 and 1.19, respectively. In addition, the intercorrelations between ${}^3\chi^v$ and indices I_{NG} and HOMA (Figure 5) show that the better the correlation of I_{NG} and HOMA to ${}^3\chi^v$, the better the correlation of I_{NG} and HOMA to $\log \beta_2$ (Models 6-8, 10-12 in Table 2). All of the

a)

b)

Figure 4 Dependence of stability constant $\log \beta_2$ (DFT), of Cu(II) bis-complexes with ten aromatic ligands, on a) I_{NG} index (Model 5, Table 2) and b) HOMA index (Model 9, Table 2).

Figure 5 The correlation coefficient of I_{NG} and HOMA to ${}^3\chi^v$ vs. the correlation coefficient of I_{NG} and HOMA to $\log \beta_2$ (Models 6-8, 10-12 in Table 2).

mentioned strongly suggests that the information needed for stability modelling of the studied complexes stems almost exclusively from ${}^3\chi''$.

CONCLUSION

We correlated the ${}^3\chi''$ index to the stability constant $\log \beta_2$ calculated by the density functional theory (DFT). Assuming that DFT $\log \beta_2$'s were fairly well correlated with experimental stability constants, we concluded that the method with ${}^3\chi''$ was equally good but far simpler for the estimation of stability constants in comparison to the DFT method. As DFT also proved unable to estimate the absolute $\log \beta_2$ values (21), it could not be ascertained, at least in our case, that DFT had any advantage over our method.

We also showed that ${}^3\chi''$ correlates to the $\log \beta_2$ of the studied complexes much better than metalloaromaticity indices I_{NG} and HOMA. Moreover, adding indices I_{NG} or HOMA into regression along with ${}^3\chi''$ worsened the estimates. This indicates that ${}^3\chi''$ encompasses structural information relevant to stability, including metalloaromaticity.

Acknowledgements

This work was supported by the Croatian Ministry of Science, Education and Sports (project 022-1770495-2901).

REFERENCES

- Kowalik-Jankowska T, Ruta-Dolejsz M, Wisniewska K, Lankiewicz L, Kozłowski H. Possible involvement of copper(II) in Alzheimer disease. *Environ Health Perspect* 2002;110(Suppl 5):869-70. PMID: 12426149
- Lovell MA, Robertson JD, Teesdale WJ, Campbell JL, Markesbery WR. Copper, iron and zinc in Alzheimer's disease senile plaques. *J Neurol Sci* 1998;158:47-52. doi: 10.1016/s0022-510x(98)00092-6
- Bocca B, Forte G, Petrucci F, Pino A, Marchione F, Bomboi G, Senofonte O, Giubilei F, Alimonti A. Monitoring of chemical elements and oxidative damage in patients affected by Alzheimer's disease. *Ann Ist Super Sanità* 2005;41:197-203. PMID: 16244393
- Brewer GJ. The risks of copper toxicity contributing to cognitive decline in the aging population and to Alzheimer's disease. *J Am Coll Nutr* 2009;28:238-42. PMID: 20150596
- Brewer GJ, Newsome DA. Copper Proof: How Chronic Copper Toxicity is Causing the Epidemic of Alzheimer's Disease and Dementia. Ann Arbor: George J. Brewer Inc; 2009.
- Sparks D, Friedland R, Retanceska S, Schreurs BG, Shi J, Perry G, Smith MA, Sharma A, Derosa S, Ziolkowski C, Stankovic G. Trace copper levels in the drinking water, but not zinc or aluminium, influence CNS Alzheimer's like pathology. *J Nutr Health Aging* 2006;10:247-54. PMID: 16886094
- Waldman M, Lamb M. Dying for a Hamburger: Modern Meat Processing and the Epidemic of Alzheimer's Disease. New York (NY): Thomas Dunne Books, St. Martin's Press; 2004.
- Sinha R, Cross AJ, Graubard BI, Leitzmann MF, Schatzkin A. Meat intake and mortality: A perspective study of over half a million people. *Arch Intern Med* 2009;169:562-71. doi: 10.1001/archinternmed.2009.6
- Morris MC, Evans DA, Tangney CC, Benias JL, Schneider JA, Wilson RS, Scherr PA. Dietary copper and high saturated and trans fat intakes associated with cognitive decline. *Arch Neurol* 2006;63:1085-8. doi: 10.1001/archneur.63.8.1085
- Masters CL, Simms G, Weinman A, Multhaup G, McDonald BL, Beyreuther K. Amyloid plaque core protein in Alzheimer disease and Down syndrome. *Proc Natl Acad Sci USA* 1985;82:4245-9. PMID: 3159021
- Esch F, Keim PS, Beattie EC, Blacher RW, Culwell AR, Oltersdorf T, McClure D, Ward PJ. Cleavage of amyloid beta peptide during constitutive processing of its precursor. *Science* 1990;244:1122-4. doi: 10.1126/science.2111583
- Otvos LJ, Szendrei GI, Lee VM, Mantsch HH. Human and rodent Alzheimer β -amyloid peptides acquire distinct conformations in membrane-mimicking solvents. *Eur J Biochem* 1993;211:249-57. doi: 10.1111/j.1432-1033.1993.tb19893.x
- Hou L, Zagorski MG. NMR reveals anomalous copper(II) binding to amyloid A β peptide of Alzheimer's disease. *J Am Chem Soc* 2006;128:9260-1. doi: 10.1021/ja046032u
- Innocenti M, Salvietti E, Guidotti M, Casini A, Bellandi S, Foresti ML, Gabbiani C, Pozzi A, Zatta P, Messori L. Trace copper(II) and zinc(II) ions drastically modify the aggregation behavior of amyloid- β_{1-42} : An AFM study. *J Alzheimer's Dis* 2010;19:1323-9. doi: 10.3233/JAD-2010-1338
- Huang X, Atwood CS, Hartshorn MA, Multhaup G, Goldstein LE, Scarpa RC, Cuajungco MP, Gray DN, Lim J, Moir RD, Tanzi RE, Bush AI. The A β peptide of Alzheimer's disease directly produces hydrogen peroxide through metal ion reduction. *Biochemistry* 1999;38:7609-16. doi: 10.1021/bi990438f
- Opazo C, Ruiz FH, Inestrosa NC. Amyloid- β -peptide reduces copper(II) to copper(I) independent of its aggregation state. *Biol Res* 2000;33:125-31. PMID: 15693279
- Da Silva GFZ, Ming LJ. Alzheimer's disease related copper(II)- β -amyloid peptide exhibits phenol monooxygenase and catechol oxidase activities. *Angew Chem Int Ed* 2005;44:5501-4. doi: 10.1002/anie.200501013
- Ferrada E, Arancibia V, Loeb B, Norambuena E, Olea-Azar C, Huidobro-Toro JP. Stoichiometry and conditional stability constants of Cu(II) or Zn(II) clioquinol complexes; implications for Alzheimer's and Huntington's disease therapy. *Neurotoxicology* 2007;28:445-9. doi: 10.1016/j.neuro.2007.02.004
- Lim S, Paterson BM, Fodero-Tavoletti MT, O'Keefe GJ, Cappai R, Barnham KJ, Villemagne VL, Donnelly PS. A copper radiopharmaceutical for diagnostic imaging of Alzheimer's disease: a bis(thiosemicarbazonato)copper(II)

- complex that binds to amyloid-beta plaques. *Chem Commun (Camb)* 2010;46:5437-9. doi: 10.1039/c0cc01175d
20. Rodríguez-Rodríguez C, de Groot NS, Rimola A, Álvarez-Larena Á, Lloveras V, Vidal-Gancedo J, Ventura S, Vendrell J, Sodupe M, González-Duarte P. Design, selection, and characterization of thioflavin-based intercalation compounds with metal chelating properties for application in Alzheimer's disease. *J Am Chem Soc* 2009;131:1436-51. doi: 10.1021/ja806062g.
21. Rimola A, Ali-Torres J, Rodríguez-Rodríguez C, Poater J, Matito E, Solà M, Sodupe M. Ab initio design of chelating ligands relevant to Alzheimer's disease: Influence of metalloaromaticity. *J Phys Chem A* 2011;115:12659-66. doi: 10.1021/jp203465h
22. Raos N, Miličević A. Estimation of stability constants of coordination compounds using models based on topological indices. *Arh Hig Rada Toksikol* 2009;60:123-8. doi: 10.2478/10004-1254-60-2009-1923
23. Miličević A, Raos N. Estimation of stability constants by graph-theoretical models. Binding of amino acids to copper(II) and nickel(II) complexes of iminodiacetates and pyridyl derivatives of aspartic acid. *Int J Chem Model* 2008;1:395-403.
24. Miličević A, Branica G, Raos N. Irving-Williams order in the framework of connectivity index $^3\chi^v$ enables simultaneous prediction of stability constants of bivalent transition metal complexes. *Molecules* 2011;16:1103-12. doi: 10.3390/molecules16021103
25. Miličević A, Raos N. Modeling of stability constants of *mono*-complexes of La^{3+} , Ce^{3+} , Pr^{3+} , and Nd^{3+} with carboxylic acids in water-dioxane solutions by using connectivity index $^3\chi^v$. *Chem Phys Lett* 2012;528:63-7. doi: 10.1016/j.cplett.2012.01.023
26. Herges R. Topology in chemistry: designing Möbius molecules. *Chem Rev* 2006;106:4820-42. doi: 10.1021/cr0505425
27. Mauksch M, Tsogoeva SB. Demonstration of "Möbius" aromaticity in planar metallocycles. *Chem Eur J* 2010;16:7843-51. doi: 10.1002/chem.201000396
28. Tetko IV, Gasteiger J, Todeschini R, Mauri A, Livingstone D, Ertl P, Palyulin VA, Radchenko EV, Zefirov NS, Makarenko AS, Tanchuk VY, Prokopenko VV. Virtual computational chemistry laboratory-design and description. *J Comput Aid Mol Des* 2005;19:453-63. doi: 10.1007/s10822-005-8694-y
29. VCCLAB, Virtual Computational Chemistry Laboratory [displayed 30 August 2013]. Available at <http://www.vcclab.org>
30. NCI/CADD Group. Online SMILES Translator and Structure File Generator [displayed 30 August 2013]. Available at <http://cactus.nci.nih.gov/services/translate/>
31. Randić M. On history of the Randić index and emerging hostility toward chemical graph theory. *MATCH Commun Math Comput Chem* 2008;59:5-124.
32. Kier LB, Hall LH. Molecular connectivity VII: specific treatment to heteroatoms. *J Pharm Sci* 1976;65:1806-9. PMID: 1032667
33. Kier LB, Hall LH. *Molecular Connectivity in Structure-Activity Analysis*. New York (NY): John Wiley & Sons; 1986.
34. Kier LB, Hall LH. *Molecular Connectivity in Chemistry and Drug Research*. New York (NY): Academic Press; 1976.
35. Randić M. On characterization of molecular branching. *J Am Chem Soc* 1975;97:6609-15. doi: 10.1021/ja00856a001
36. Lučić B, Trinajstić N. Comparison between the sum-connectivity index and product-connectivity index for benzenoid hydrocarbons. *Chem Phys Lett* 2009;475:146-8. doi: 10.1016/j.cplett.2009.05.022
37. Lučić B, Trinajstić N. Multivariate regression outperforms several robust architectures of neural networks in QSAR modeling. *J Chem Inf Comput Sci* 1999;39:121-32. doi: 10.1021/ci980090f

Sažetak

USPOREDBA DVIJU METODA ZA PROCJENU STABILNOSTI BIS-KOMPLEKSA BAKRA(II) S AROMATSKIM LIGANDIMA RELEVANTNIMA ZA ALZHEIMEROVU BOLEST

Izabrali smo komplekse bakra(II) s interkalacijskim spojevima temeljenima na tioflavinu, koji su namijenjeni za liječenje Alzheimerove bolesti, kako bismo usporedili teoriju funkcionala gustoće (DFT) s metodom molekularnog valencijskog indeksa 3. reda (${}^3\chi^v$) za procjenu vrijednosti konstanti stabilnosti $\log \beta_2$. Korelacija deset vrijednosti $\log \beta_2$ izračunatih metodom DFT s ${}^3\chi^v$ indeksom dala je $r=0,988$ i S.E.=0,85. Korelacije indeksa I_{NG} , HOMA i ${}^3\chi^v$ s vrijednostima $\log \beta_2$ te međukorelacije indeksa pokazuju bitan utjecaj ${}^3\chi^v$ na predviđanje stabilnosti istraživanih kompleksa.

KLJUČNE RIJEČI: *aromatičnost, konstante stabilnosti, regresijski modeli, teorija DFT, topološki indeksi*

CORRESPONDING AUTHOR:

Ante Miličević, PhD
Institute for Medical Research and Occupational Health
Ksaverska cesta 2, 10000 Zagreb, Croatia
E-mail: *antem@imi.hr*