Letter to the Editor

DOI: 10.1515/aiht-2015-66-2647

Hydroxyethyl starch should not be used to manage severe aluminium phosphide poisoning

Zvonko Rumboldt

Split University School of Medicine, Split, Croatia

In the *Archives* January issue (1), I have read with interest and surprise a letter by Marashi and Nasrabadi concerning the use of hydroxyethyl starch (HES, hetastarch; best known under the registered name Plasmasteril) to treat severe aluminium phosphide (ALP) poisoning.

At the current level of scientific evidence such a claim seems unacceptable.

Aluminium phosphide poisoning (the compound is commercially available as a fumigant for stored cereals) is believed to be caused by phosphine (PH_3), as it induces mitochondrial damage and severe multi-organ failure (2). The poisoning is mostly reported from the Indian subcontinent, and is largely unknown in Europe. The management is exclusively supportive (gastric lavage with KMnO_4 solution, treatment of shock, lactic acidosis, etc.), since there is no specific antidote (2, 3). Some traditional (e.g. coconut oil) or “modern” interventions (e.g. haemodialysis) have not been adequately assessed, and rest instead on pathophysiological assumptions, anecdotal reports, and wishful thinking (2, 3).

Lactic acidosis type A, a form of high anion gap metabolic acidosis, is just one of the many aspects of ALP poisoning with fatal prognosis. Treatment with NaHCO_3 in this situation is highly controversial, as the authors (1) appropriately underscore. However, the suggested administration of colloid solutions is potentially even more dangerous. The authors corroborate their proposal by quoting just one hypothetical paper of their own on the role of HES in acute ALP poisoning (4) with no further experimental evidence. It is not clear whether they are advocating starch solution administration at arterial $\text{pH} > 7$ (as stated in the letter) or < 7 , or what the dosage of 500–1000 mL of HES really means (i.e. isotonic, 6 g L^{-1} , or hypertonic, 9 g L^{-1} solution)?

Contrary to their claim (1), colloidal solutions such as dextrans, pentastarch, polygeline, and HES in particular may worsen metabolic acidosis and shock, possibly due to

the substance’s deposition in the tissue and its interaction with fibrinogen, leading to even more severe coagulopathy and organ failure (5). Crystalloid solutions are better tolerated and equally effective plasma expanders (6).

Beware of ill-founded therapeutic enthusiasm!

REFERENCES

1. Marashi SM, Nasri-Nasrabadi Z. Can sodium bicarbonate really help in treating metabolic acidosis caused by aluminium phosphide poisoning? *Arh Hig Rada Toksikol* 2015;66:83–4. doi: 10.1515/aiht-2015-66-2637
2. Agrawal VK, Bansal A, Singh RK, Kumawat BL, Mahajan P. Aluminium phosphide poisoning: possible role of supportive measures in the absence of specific antidote. *Indian J Crit Care Med* 2015;19:109–12. doi: 10.4103/0972-5229.151019
3. Gurjar M, Baronia AK, Azim A, Sharma K. Managing aluminium phosphide poisoning. *J Emerg Trauma Shock* 2011;4:378–84. doi: 10.4103/0974-2700.83868
4. Marashi SM, Arefi M, Behnouch B, Nasrabadi MG, Nasri Nasrabadi Z. Could hydroxyethyl starch be a therapeutic option in the management of acute aluminium phosphide toxicity. *Med Hypotheses* 2011;76:596–8. doi: 10.1016/j.mehy.2011.01.009
5. Hartog CS, Natanson C, Sun J, Klein HG, Reinhart K. Concerns over use of hydroxyethyl starch solutions. *BMJ* 2014;349:g5891. doi: 10.1136/bmj.g5891
6. The European Medicines Agency’s Pharmacovigilance Risk Assessment Committee. Hydroxyethyl-starch solutions (HES) should no longer be used in patients with sepsis or burn injuries or in critically ill patients [displayed 14 July 2015]. Available at: http://www.ema.europa.eu/ema/index.jsp?curl=pages/medicines/human/referrals/Hydroxyethyl_starch-containing_solutions/human_referral_prac_000012.jsp&mid=WC0b01ac05805c516f