

The influence of royal jelly and human interferon-alpha (HuIFN- α N3) on proliferation, glutathione level and lipid peroxidation in human colorectal adenocarcinoma cells *in vitro*

Bratko Filipič¹, Lidija Gradišnik⁴, Klemen Rihar², Eugen Šooš⁵, Adriana Pereyra³, and Jana Potokar³

Institute for Microbiology and Immunology, Medical Faculty, University of Ljubljana¹, Chengdujska 4², Medex d.o.o.³, Ljubljana, Medical Faculty, University of Maribor, Maribor⁴, Slovenia, Trg Sv. Ivana 5, Kloštar Ivanič, Croatia⁵

[Received in March 2015; CrossChecked in March 2015; Accepted in November 2015]

Among royal jelly's (RJ) various biological activities, its possible antitumour activity deserves particular attention. The purpose of this study was to investigate the influence of RJ, its bioactive component 10-hydroxy-2-decenoic acid (10-HDA), and human interferon-alpha (HuIFN- α N3) on the proliferation of human colorectal adenocarcinoma cells (CaCo-2), and ascertain their effect on intracellular glutathione (GSH) level and lipid peroxidation. We studied the antiproliferative (AP) activity of RJ [(0.1 g/10 mL phosphate buffer saline (PBS)], HuIFN- α N3 (1000 I.U. mL⁻¹), 10-HDA at 100.0 μ mol L⁻¹, and their different combinations, in the ratio 1:1, 1:2, and 2:1 on CaCo-2 cells. The GSH level was measured by glutathione assay. The lipid peroxidation was measured by malondialdehyde (MDA) assay. Single RJ had a low AP activity: 2.0 (0.5 mg mL⁻¹). HuIFN- α N3 had an AP activity of 2.5 (208.33 I.U. mL⁻¹), while 10-HDA had an AP activity of 1.5 (37.5 μ mol mL⁻¹). The highest AP activity of 3.8 was obtained when RJ and HuIFN- α N3 were applied at the ratio 2:1. In that combination the level of GSH was 24.9 \pm 2.4 nmol g⁻³ of proteins (vs. 70.2 \pm 3.2 nmol g⁻³ in the control) and the level of MDA was 72.3 \pm 3.1 nmol g⁻³ (vs. 23.6 \pm 9.1 nmol g⁻³ in the control). It is generally assumed that 10-HDA, an important constituent of RJ, together with HuIFN- α N3, is responsible for the inhibition of CaCo-2 cells proliferation *in vitro*. In our study, however, RJ and HuIFN- α N3 applied at 2:1 decreased the level of GSH the most and significantly increased lipid peroxidation via MDA in CaCo-2 cells. Future studies should show whether these GSH- and MDA-related activities of RJ, HuIFN- α N3, 10-HDA, and their combinations may decrease the tumorigenicity index and tumorigenic potential of various tumour cells *in vitro*.

KEY WORDS: *antiproliferative activity; antitumour activity; CaCo-2 cells; 10-hydroxy-2-decenoic acid; malondialdehyde*

Royal jelly (RJ) is a milky material secreted by the hypopharyngeal and mandibular glands of young worker bees between the sixth and twelfth day of their life. It is food exclusively for the queen honey bee (*Apis mellifera*) larva, which leads to the development of a sexually mature queen bee (1, 2). Chemically, RJ comprises water (50-60%), different proteins (18.0%), carbohydrates (15.0%), lipids (3-6%), mineral salts (1.5%), and vitamins, together with a large number of bioactive substances, such as 10-hydroxy-2-decenoic (10-HDA) acid (3). RJ exhibits various immunomodulatory and antiproliferative properties, as well as possible antitumour activity. It also contains different antibacterial proteins like 350-kDa protein stimulating the proliferation of human monocytes (4-6). In addition, the RJ Protein₃₀ fraction exhibits a clear cytotoxic effect on HeLa cells by decreasing the initial cell population by 50% at the end of treatment (7).

HuIFN- α N3, a type I interferon, is a multisubtype protein (8, 9) with antiviral, antiproliferative, and antitumour activity (10). It has been used clinically in the treatment of various cancers for over 30 years (11). The molecular mechanism behind its cytoreductive action is still not clear. The antiproliferative effect of HuIFN- α N3 plays a central role in its chemotherapeutic effect. Recent research (12) has also indicated its action in apoptosis pathways as a possible anti-tumour mechanism. It was also shown that HuIFN- α N3 can exert a direct cytotoxic effect on different malignant cells and tumour cell lines *in vitro* (13).

Glutathione (GSH), a low molecular weight thiol, has a central role in the control of the cellular thiol/disulphide redox state, bearing significance for normal redox signalling in cells, among them also in the CaCo-2 cells used in this study (14). Intracellular GSH concentration, cell proliferation, and apoptosis are mutually connected. A high level of GSH enhances cell proliferation, while GSH

Correspondence to: Dr Bratko Filipič, Institute for Microbiology and Immunology, Medical Faculty, University of Ljubljana, Zaloška 4, 1105 Ljubljana, Slovenia, E-mail: Bratko.Filipic@gmail.com

depletion results in the inhibition of CaCo-2 cell growth and proliferation, due to the increase of apoptosis.

Lipid peroxidation, or oxidative degradation of lipids, is a process by which free radicals “steal” electrons from the lipids in cellular membranes inducing cell damage. It proceeds by a free radical chain reaction mechanism (15, 16). The end-products of lipid peroxidation are reactive aldehydes with carcinogenic potential. Among them, the most important is malondialdehyde (MDA), a major bioactive marker of lipid peroxidation that exerts numerous biological activities resembling activities of reactive oxygen species.

The aim of this study was to investigate the effect of RJ and 10-HDA on HuIFN- α 3-induced inhibition of CaCo-2 cells proliferation *in vitro* and ascertain their effect on the intracellular level of GSH and lipid peroxidation via MDA activity. We assumed that such an approach might serve to establish possible antiproliferative/antitumor mechanisms in human colorectal adenocarcinoma cells, which could be of value for developing future anticancer treatments based on the use of these bioactive compounds.

METHODS

Material

The following materials were used: Human Interferon - α 3, (HuIFN- α 3) (Institute of Immunology, Zagreb, Croatia) applied at 1000 I.U. mL⁻¹, which was a standard concentration previously used in experiments (17). Royal jelly-fresh (Mižigoj, Ljubljana, Slovenia) (RJ-F(M)) (MEDEX d.o.o., Ljubljana, Slovenia), was applied at 0.1 g/10 mL (18). 10-hydroxy-2-decenoic acid (10-HDA) (Sigma-Aldrich, Missouri, USA) was applied at 100 μ mol L⁻¹ (19). All of the reagents were dissolved in the phosphate buffer saline (PBS), pH=7.2 and then filtered through a 0.2 μ m syringe filter (Millipore, USA).

Cell culture

CaCo-2 cells (Institute for Microbiology and Immunology, Ljubljana, Slovenia) were cultivated in Eagle's medium with L-Glutamine (2.0 mmol L⁻¹) (Sigma-Aldrich, Missouri, USA) and antibiotics: Penicillin (100 units mL⁻¹), Streptomycin (100 μ g mL⁻¹), and Gentamycin (50 μ g mL⁻¹) (Sigma-Aldrich, Missouri, USA) and supplemented with 10 % of foetal calf serum (FCS) (Sigma-Aldrich, Missouri, USA). Before the experiment, the cells were multiplied and their viability and capability for proliferation was assayed by the MTT Cell Proliferation Assay Kit (K299-100) (BIOVISION, Milpitas, California, USA). The cell cultivation was performed in 96-well flat microtiter plates in 5 % CO₂ at 37 °C or in 25 cm² flasks (Sterilin, Sigma-Aldrich, Missouri, USA) in 5 % CO₂ at 37 °C.

Design of the study

The experiments were designed as follows: single substances: RJ-F(M), HuIFN- α 3 and 10-HDA in the previously stated concentrations, were added alone in a volume of 100 μ L/well. When different combinations between them were used, for the ratio 1:1 100+100 μ L/well was added. For the ratio 1:2, 66.8+133.2 μ L/well and for the ratio 2:1, 133.2+66.8 μ L/well was added.

Antiproliferative (AP) activity

Single substances or their combinations, as described in the design of the study, were added in a volume of 200 μ L per well in the first well in row of 96-well flat microtiter plates, and the samples were serially transferred per 100 μ L from 1:2 to 1:4096 in Eagle's medium with L-Glutamine and antibiotics. After the substances, the cells (CaCo-2) were added (10⁴ cells/well/100 μ L) in Eagle's medium with L-Glutamine and antibiotics and 10 % FCS. The cells without substances were added separately (the negative control). As positive controls, single tested substances (RJ, HuIFN- α 3, 10-HAD) were used. The microtiter plates were incubated for 72 hours at 37 °C in a 5 % CO₂ atmosphere. Afterward, the supernatants were discharged and the cells were fixed with the addition of 100 μ L/well of 10 % formalin (Sigma-Aldrich, Missouri, USA) in PBS. After two hours, the fixative was removed and the cells were washed twice with the PBS. After that, 2 % Rhodamine B (Sigma-Aldrich, Missouri, USA) (100 μ L/well) was added for 15 minutes. This was then removed, and the cells were washed twice with PBS and air-dried. On the dried plates, the optical density (OD) at 550 nm was measured (Synergy HTX Multi Mode Reader with Gen 5 software, Biotek, Winooski, USA). The AP activity was determined with the well in rows where 50 % cell growth inhibition was found. The AP₅₀ was calculated for each separate substance (RJ, HuIFN α 3, and 10-HDA) (20).

Glutathione determination

CaCo-2 cells were cultivated in 25 cm² flasks (Sterilin, Sigma-Aldrich, Missouri, USA) in Eagles' medium with L-Glutamine and antibiotics and 10 % FCS. When monolayers were developed, the cells in flasks were treated with substances alone or their combinations in a volume of 1.0 mL when single substances were added, and a total of 2.0 mL when the combinations were added. The cells were treated for 24 hours at 37 °C and 5% CO₂. The medium was removed, and the cells were detached with trypsin (Sigma-Aldrich, Missouri, USA) and treated with 1 mL of 10 mmol mL⁻¹ Tris-HCl solution (pH=6.0) containing 0.5 mol mL⁻¹ diethylene triamine pentacetic acid (DTPA) (Sigma-Aldrich, Missouri, USA), and syringed several times with an insulin syringe for their lysis. The cell protein amount was determined by Bio-Rad protein assay (Bio-Rad Laboratories, Hercules, California, USA) and Bovine serum albumin (BSA) (Sigma-Aldrich, Missouri, USA) as a

standard. For total GSH determination, 100 μ L of DL-Dithiothreitol (DTT), 25 μ mol L⁻¹, and 150 μ L of 0.1 mol mL⁻¹ Tris-HCl (pH 8.5) were added to 50 μ L of the cell lysate. After 30 minutes on ice, the proteins were precipitated by adding 750 μ L of 2.5 % (wt./vol) 5-sulfosalicylic acid and centrifuged at 13000 g (Centric, Tehnica d.o.o., Železniki, Slovenia) for 4 minutes at 4 °C. The cellular supernatants were used in Glutathione Assay Kit (Sigma-Aldrich, Missouri, USA) to measure the GSH level at OD at 412 nm (Synergy HTX Multi Mode Reader with Gen 5 software, Biotek, Winooski, USA) and expressed as nmol of GSH g⁻³ of proteins.

Measurements of lipid peroxidation

CaCo-2 cells were cultivated in 25 cm² flasks (Sterilin, Sigma-Aldrich, Missouri, USA) in Eagles' medium with L-Glutamine and antibiotics and 10 % FCS. When the monolayer was formed, the cells were treated with substances alone or their combinations as described in the design of the study, in a quantity of 1.0 mL when single substances were added, and a total of 2.0 mL when the combinations were added. The treated/non-treated cells were incubated for 24 hours at 37 °C and 5 % CO₂. The medium was removed and cells detached with trypsin washed and resuspended in 5 mL of PBS. Cell protein concentration was determined using the Bio-Rad protein assay (Bio-Rad Laboratories, Hercules, California, USA) and BSA as a standard. A measure of 1 mL of thiobarbituric acid (TBA) reagent (0.38 % 2-TBA, 15.0 % TBA, 0.3 mol mL⁻¹ HCl) was added to the cell suspension. The samples were heated at 95 °C for 20 minutes, chilled to room temperature and centrifuged at 1500 g for 10 minutes. The

TBA reactive substances (RS) developed by lipid peroxidation were measured in the supernatant at OD at 535 nm (Synergy HTX Multi Mode Reader with Gen 5 software, Biotek, Winooski, USA), according to the TBA method (21, 22). The results were expressed as MDA nmol g⁻³ of protein.

Data analysis

T-test was used for significance determination (**p*<0.1, ***p*<0.05); data are shown as mean value±standard deviation. Each of the tests was performed in triplicate and each experiment was repeated three to four times.

RESULTS AND DISCUSSION

Antiproliferative activity

The following results of AP activity and concentrations at AP₅₀ for single substances were obtained: RJ: 2.0 (0.5 mg mL⁻¹), HuIFN- α 3: 2.5 (208.33 I.U. mL⁻¹) and 10-HDA: 1.5 (37.5 μ mol L⁻¹). The observed AP activity was relatively low. The AP activity of their combinations (1:1, 1:2, 2:1) is shown on Figure 1.

The highest AP activity was obtained when the combination of RJ-F (M) and HuIFN- α 3 was used in a 2:1 ratio. In this case the AP activity was 3.8. When the combination of the RJ-F (M) component 10-HDA and HuIFN- α 3 2:1 was tested, the AP activity was 2.4-2.6, which was much lower than with RJ-F (M) and HuIFN- α 3. Such results suggest that RJ-F (M) could contain some




Figure 1 The AP effect of RJ-F (M), HuIFN- α 3, 10-HDA and their combinations (1:1, 1:2 and 2:1) on the CaCo-2 cell line. Inhibition of proliferation of CaCo-2 cell line treated with RJ-F (M) (0.1 g /10 mL PBS), HuIFN- α 3 (1000 I.U. mL⁻¹), 10-HDA (100 μ mol mL⁻¹) and different combinations between them (1:1, 1:2 and 2:1) was analysed. The AP activity was determined by Rhodamine B staining method after 72 h of incubation at 37 °C and 5 % CO₂. The t-test was used for significance determination (**p*<0.1, ***p*<0.05); data are shown as mean value ± standard deviation

other components responsible for the relatively strong AP activity of the combination RJ-F (M) and HuIFN- α N3 in the ratio 2:1. In this respect, the possible role of the RJ Protein₃₀ water soluble fraction that previously exhibited the clear cytotoxic effect on HeLa cells by decreasing the initial cell population by 50 % at the end of treatment (7) should be also anticipated.

It is known that the AP activity of the RJ-F (M), HuIFN- α N3, 10-HDA on the CaCo-2 cells is connected with the induction of apoptosis and cytotoxicity (14). Also their influence on the glutathione level and lipid peroxidation was found (23).

GSH determination and measurement of lipid peroxidation

The results obtained in our study show that RJ-F (M), HuIFN- α N3, 10-HDA and their combinations decreased the level of glutathione and increased the lipid peroxidation via the MDA. Detailed data are shown in Table 1.

Glutathione (GSH) plays an important role in many cellular processes, like cell differentiation, proliferation, and apoptosis and cancer. While GSH deficiency, or a decrease in the GSH/glutathione disulphide (GSSG) ratio, leads to an increased susceptibility to oxidative stress in the progression of cancer, elevated GSH levels increase the antioxidant capacity and resistance to oxidative stress found in cancer cells. It is important to stress that RJ-F (M), HuIFN- α N3, 10-HDA, and their combinations decreased the level of glutathione and significantly increased the lipid peroxidation via the MDA.

Some of the mechanisms of the possible antitumour mechanisms of royal jelly are connected with the modulation of the oxidative stress and induction of apoptosis. (24). Practically the same effects were found

after the treatment of the Pancreatic Cancer Cell tumour cells PaCa-44 with 10-HDA, where the induction of the apoptosis was found (25). It is interesting to note that the antitumour activity of HuIFN- α N3 was also connected with the induction of apoptosis and the modulation of the oxidative stress in rats with breast cancer (26).

The most important finding of our study is that royal jelly, especially its constituent 10-HDA and HuIFN- α N3 had similar active points in the antitumor activity, which in proper ratio can be enhanced. It can be concluded, that the most active was combination of RJ-F(M) and HuIFN- α N3 2:1, where the level of the GSH was $24.9 \pm 2.4 \text{ nmol g}^{-3}$ of proteins (vs. $70.2 \pm 3.2 \text{ nmol g}^{-3}$ in the control) and the level of MDA $72.3 \pm 3.1 \text{ nmol g}^{-3}$ (vs. $23.6 \pm 9.1 \text{ nmol g}^{-3}$ in the control).

Future experiments will show whether these GSH- and MDA-related activities of RJ-F (M), HuIFN- α N3, 10-HDA and their combinations may cause the decrease of the tumorigenicity index of different tumour cells *in vitro*, as previously reported in the literature (27, 28), also through their tumorigenic potential. This is important for the practical use of royal jelly (10-HDA) and HuIFN- α N3 in a combination that could be of value for future development of tumour therapy based on the use of bioactive compounds.

Conflict of Interests

All the authors declare no conflict of interest.

Acknowledgement

The research was supported by a grant from the Slovenian Research Agency: ARRS-NRU/P3-0083-0381-2014. Authors are indebted to Tomaž Velnar for English suggestions.

Table 1 Glutathione (GSH) determination and measurement of lipid peroxidation (MDA) after the CaCo-2 cells treatment with RJ, HuIFN- α N3, 10-HDA and their combinations: 1:1, 1:2 and 2:1^a

Sample	Glutathion (GSH) ^b	Malondialdehyde (MDA) ^c
	(Mean \pm SE)	(Mean \pm SE)
Cell control	70.2 \pm 3.2	23.6 \pm 9.1
RJ-F(M) ^d	43.8 \pm 2.8	30.2 \pm 4.3
HuIFN- α N3	28.7 \pm 6.4	38.6 \pm 4.2
10-HDA ^e	33.6 \pm 5.8	50.7 \pm 4.6
RJ-F(M)+HuIFN- α N3 1:1	45.2 \pm 4.7	43.6 \pm 4.1
RJ-F(M)+HuIFN- α N3 1:2	40.8 \pm 3.1	58.3 \pm 5.2
RJ-F(M)+HuIFN- α N3 2:1	24.9 \pm 2.4**	72.3 \pm 3.1**
RJ-F(M)+10-HDA 1:1	40.6 \pm 4.5	43.1 \pm 2.6
RJ-F(M)+10-HDA 1:2	37.2 \pm 2.1	50.6 \pm 4.5
RJ-F(M)+10-HDA 2:1	30.3 \pm 3.7*	61.6 \pm 5.2*
10-HDA+HuIFN- α N3 1:1	29.5 \pm 1.7	49.6 \pm 4.2
10-HDA+HuIFN- α N3 1:2	42.6 \pm 5.3	57.2 \pm 2.6
10-HDA+HuIFN- α N3 2:1	25.6 \pm 3.1	55.6 \pm 6.2

^at-test was used for significance determination; * p<0.1, **p<0.05

^bMeasured as nmol g⁻³ of proteins; ^cMeasured as nmol g⁻³ of proteins; ^dRJ-F(M)=royal jelly – Fresh(Mižigoj); ^e10-HDA=10-hydroxy-2-decenoic acid

REFERENCES

1. Ramadan FM, Al-Ghamdi. Bioactive compounds and health-promoting properties of royal jelly: a review. *J Funct Foods* 2012;4:39-52. doi: 10.1016/j.jff.2011.11.12.007
2. Pavel IC, Margharitas AL, Bobis O, Dezmirean DS, Sapcaliu A, Radoi I, Madas NM. Biological activities of royal jelly - review. *Anim Sci Biotechnol* 2011;44:108-18.
3. Tahashi K, Hiroshi M. Royal jelly acid 10-hydroxy-trans-2-decenoic acid as a modulator of the innate immune responses. *Endocr Metab Immune Disord Drug Targets* 2012;12:368-76. doi: 10.2174/187153012803832530
4. Townsend GF, Morgan JF, Tolnai S, Hazlett B, Morton HJ, Shuel RW. Studies on the *in vitro* antitumor activity of atty acids. I. 10-Hdroxy-2-decenoic acid from royal jelly. *Cancer Res* 1960;20:503-10. PMID: 13839101
5. Tamura T, Fuji A, Kuboyama N. [Antitumor effects of royal jelly (RJ), in Japanese]. *Folia Pharmacol Japon* 1987;89:73-80. doi: 10.1254/fpj.89.73
6. Watanabe K, Shinmoto H, Kobori M, Tsushida T, Shinohara K, Kanaeda J, Yonekura M. Stimulation of cell growth in the U937 human myeloid cell line by honey royal jelly protein. *Cytotechnology* 1998;26:23-7. doi: 10.1023/A:1007928498128
7. Salazar-Olivio LA, Paz-González V. Screening of biological activities present in honeybee (*Apis mellifera*) royal jelly. *Toxicology in Vitro* 2005;19:645-51. doi: 10.1016/j.tiv.2005.03.001
8. Mocellin S, Lens MB, Pasquali S, Pilati P, Chiarion Sileni V. Interferon alpha for the adjuvant treatment of cutaneous melanoma. *Cochrane Database Syst Rev* 2013;6:CD008955. doi: 10.1002/14651858.CD008955
9. Kalliolias DG, Ivashkiv BL. Overview of the biology of type I interferons. *Arthritis Res Ther* 2010;12(Suppl 1):S1-9. doi: 10.1186/ar2881
10. Liu X, Lu J, He ML, Li Z, Zhang B, Zhou LH, Li Q, Li G, Wang L, Tian WD, Peng Y, Li XP. Antitumor effect of interferon alpha on cell growth and metastasis in human nasopharyngeal carcinoma. *Curr Cancer Durg Targets* 2013;12:561-70. doi: 10.2174/156800912800673293
11. Jonasch E, Hluska GF. Interferon in oncological practice: review of interferon biology, clinical application and toxicities. *Oncologist* 2001;6:34-55. doi: 10.1634/theoncologist.6-1-34
12. Jedema I, Barge RMY, Willemze R, Falkenburg JH. High susceptibility of human leukemic cells to Fas-induced apoptosis is restricted to G1 phase of the cell cycle and can be increased by interferon treatment. *Leukemia* 2003;17:576-84. doi: 10.1038/sj.leu.2402844
13. Peek SF, Bonds MD, Gangemi DG, Thomas CB, Schultz RD. Evaluation of cytotoxicity and antiviral activity of recombinant human interferon alfa-2a and recombinant human interferon alfa-B/D hybrid against bovine viral diarrhoea virus, infectious bovine rhinotracheitis virus, and vesicular stomatitis virus *in vitro*. *Am J Vet Res* 2004;65:871-4. doi: 10.2460/ajvr.2004.65.871
14. Ziegler TR, Nkabyo SY, Gu LH, Watson WH, Jones DP. Glutathione and the redoxin during the differentiation in human colon epithelial (CaCo-2) cells. *Am J Physiol Gastrointest Liver Physiol* 2002;283:G1352-59. doi: 10.1152/ajpgi.00183.2002
15. Gago-Dominguez M, Jiang X, Castela EJ. Lipid peroxidation, oxidative stress genes and dietary factors in breast cancer protection: a hypothesis. *Breast Cancer Res* 2007;9:201-11. doi: 10.1186/bcr1628
16. Niki E, Yosida Y, Saito Y, Noguchi N. Lipid peroxidation: mechanism, inhibition and biological effects. *Biochem Biophys Res Commun* 2005;338:668-76. doi: 10.1016/j.bbrc.2005.08.072
17. Fidler IJ, Heicappel R, Saiki I, Grutter GM, Horisberger AM, Nuesch J. Direct antiproliferative effects of recombinant human interferon- α B/D hybrids on human tumor cell lines. *Cancer Res* 1987;47:2020-7. PMID: 3828990
18. Yanagita M, Kojima Y, Mori K, Yamada S, Murakami S. Osteoinductive effect of royal jelly on periodontal ligament cells. *Biomed Res* 2011;32:285-91. doi: 10.220/biomedres.32.285
19. Mihajlović D, Vučević D, Chinou J, Čolić M. Royal jelly fatty acid modulate proliferation and cytokine production by human peripheral blood mononuclear cells. *Eur Food Res Technol* 2014;238:881-7. doi: 10.1007/s00217-014-2154-7
20. Borden EC, Hogan TF, Voekel JG. Comparative antiproliferative activity *in vitro* of natural interferons α and β for diploid and transformed human cells. *Cancer Res* 1982;42:4948-53. PMID: 7139598
21. Buege JA, August SD. Microsomal lipid peroxidation. *Methods Enzymol* 1978;52:302-10. doi: 10.1016/S0076-6879(78)52032-6
22. Devasagayam TPA, Bloor KK, Ramasarma T. Methods for estimating the lipid peroxidation: an analysis of merits and demerits. *Indian J Biochem Biophys* 2003;40:300-8. PMID: 22900323
23. Traverso N, Ricciarelli R, Nitti M, Marengo B, Furfaro AL, Pronzato A, Marinari UM, Domenicotti C. Role of glutathione in cancer progression and chemoresistance. *Oxid Med Cell Longev* Volume 2013; Article ID 972913: doi: 10.1155/2013/972913
24. Karadeniz A, Simsek N, Karakus E, Yildirim S, Kara A, Can I, Kisa F, Emre H, Turkeli M. Royal jelly modulates oxidative stress and apoptosis in liver and kidneys of rats treated with cisplatin. *Oxid Med Cell Longev* 2011; Article ID 981793, doi: 10.1155/2011/981793
25. Merendino N, Loppi B, D'Aquino M, Molinari R, Pessino G, Romano C, Velotti F. Docohexaenoic acid induces apoptosis in the human PaCa-44 pancreatic cancer cell line by active reduced glutathione extrusion and lipid peroxidation. *Nutr Cancer* 2005;52:225-33. doi: 10.1207/s15327914nc5202_12
26. Eghdami A, Sohi HMS. Investigation of α -IFN-SWNT and α -IFN-PLGA effects on breast cancer in rats induced by DMBA by using CAIS-3 tumor marker. *Adv Biores* 2014;5:9-13.
27. Swierenga HHS, Whitfield FJ, Karasakti S. Loss of proliferative calcium dependence: simple *in vitro* indicator of tumorigenicity. *Proc Natl Acad Sci USA* 1978;75:6069-72. doi: 10.1073/pnas.75.12.6069
28. Thepot A, Desanlis A, Venet E, Thivillier L, Justin V, Morel AP, DeFraipont, Till M, Krutovskikh V, Tommasino M, Damour O, Hainaut P. Assessment of transformed properties *in vitro* and of tumorigenicity *in vivo* in primary keratinocytes cultured for epidermal sheet transplantation. *J Skin Cancer* Volume 2011, Article ID 936546, doi: 10.1155/2011/936546

Vpliv matičnega mlečka in humanega interferona-alfa (HuIFN- α N3) na proliferacijo, nivo glutationa in na preoksidacijo lipidov v humanih kolorektalnih adenokarcinomskih celicah *in vitro*

Kot del biološke aktivnosti MM (Matičnega mlečka) so avtorji preučevali njegovo protitumorsko delovanje kot tudi možno interakcijo s humanim interferonom alfa (HuIFN- α N3). Cilj opravljenih poskusov je bil preučiti vpliv kombinacije med MM in HuIFN- α N3 na proliferacijo celic Humanega kolorektalnega adenokarcinoma (CaCo-2) in njun vpliv na znotrajcelični nivo glutationa (GSH) in peroksidacijo lipidov. Avtorji so preučevali AP (Antiproliferativno) delovanje MM (0.1 g/10 mL fosfatnega pufru) (PBS), HuIFN- α N3, (1000 I.U. mL⁻¹), 10-hidroxy-2-decenoične kisline (10-HDA) (100.0 μ mol L⁻¹) in različne kombinacije med njimi (1:1, 1:2 in 2:1) na celice CaCo-2 *in vitro*. Njihov vpliv na znotrajcelični nivo GSH so merili s pomočjo komercialnega kita. Peroksidacijo lipidov so merili s pomočjo meritve vrednosti malondialdehida (MDA). MM sam kaže AP aktivnost 2.0 (0.5 mg mL⁻¹). HuIFN- α N3 ima AP aktivnost 2.5 (208.33 IU mL⁻¹) medtem ko ima 10-HDA AP aktivnost 1.5 (37.5 μ mol mL⁻¹). AP aktivnost kombinacije MM:HuIFN- α N3 (2:1) je bila 3.8. Pri tej kombinaciji je bil viden vpliv na nivo GSH: 24.9 \pm 2.4 nmol g⁻³ proteinov (70.2 \pm 3.2 nmol g⁻³ pri kontroli). Nivo MDA je bil 72.3 \pm 3.1 nmol g⁻³ pri kontroli). 10-HDA je glavna sestavina MM, ki v kombinaciji s HuIFN- α N3 deluje antiproliferativno na CaCo-2 celice. MM in HuIFN- α N3 v kombinaciji 2:1 pospešujeta peroksidacijo lipidov (MDA) in zmanjšujeta nivo glutationa (GSH). Nadaljni poskusi bodo pokazali ali z GSH- in MDA- povezane aktivnosti MM, HuIFN- α N3, 10-HDA in kombinacij med njimi, zmanjšujejo indeks tumorigenosti in s tem tumorigeni potencial različnih tumorskih celic *in vitro*.

KLJUČNE BESEDE: *antiproliferativno delovanje; CaCo-2 celice; protitumorsko delovanje; 10-hidroksi-2-decenoična kislina; malondialdehidi*