

## Imidacloprid as reproductive toxicant and endocrine disruptor: investigations in laboratory animals

Anja Mikolić and Irena Brčić Karačonji

*Institute for Medical Research and Occupational Health, Zagreb, Croatia*

[Received in May 2018; Similarity Check in May 2018; Accepted in May 2018]

Imidacloprid, a neonicotinoid insecticide, has been used worldwide due to its selective toxicity for insects. Its residues may enter the food chain, which is why it is important to investigate the potential adverse effects of imidacloprid exposure. This review summarises current knowledge of the reproductive toxicity and disruptive endocrine effects of imidacloprid in laboratory animals. Investigations, conducted mostly on laboratory rats, have shown adverse effects of imidacloprid on the reproductive ability in both parental and offspring generation as well as on the development of the offspring. Like many pesticides, imidacloprid may also act as endocrine disrupting chemical (EDC). It may disrupt the metabolic homeostasis, contribute to obesity, and disrupt steroidogenesis by inhibiting cytochrome P450 (CYP) enzyme activities. All these adverse effects of imidacloprid may pose a serious risk for reproduction and development with long-term consequences in adulthood.

**KEY WORDS:** *endocrine disruption; insecticide; laboratory rats; neonicotinoids; reproductive toxicity*

Neonicotinoid insecticides, the most important class of synthetic neuroactive insecticides, are widely used to treat crops and livestock against a broad range of pests (1, 2). Imidacloprid (IMI) [1-(6-chloro-3-pyridylmethyl)-N-nitroimidazolidin-2-ylideneamine] is a chlorinated analogue of nicotine and the first neonicotinoid registered for use as pesticide by the United States Environmental Protection Agency (US EPA). Since its introduction to the insecticide market in 1992, imidacloprid's use has been growing every year due to its selective toxicity and apparent safety for humans. Today, it accounts for 41.5 % of neonicotinoid use (2, 3). Imidacloprid is classified as a "Group E" carcinogen, which means that there is no evidence of carcinogenicity in humans (4). Neonicotinoid insecticides act as agonists of insect nicotinic acetylcholine receptors (nAChRs), which play an important role in synaptic transmission in the central nervous system (5). Due to structural differences between mammal and insect nAChRs and the higher binding affinity for insect nAChR, imidacloprid is considered a selective toxicant (6-8). Its oral LD<sub>50</sub> in rats and mice is 450 and 131 mg kg<sup>-1</sup> of body weight (bw), respectively (9). Even though imidacloprid is mildly toxic to mammals, it may still affect their heart, kidney, and other organs and cause gastrointestinal irritation and neurological symptoms or even death. There are many reports indicating adverse effects of imidacloprid in mammals, such as teratogenic (10, 11), mutagenic (12), neurotoxic (13-15), and immunotoxic effects (16, 17). The toxicity of imidacloprid

is dose-dependent, as reported by a study that evaluated effects of 30 days oral administration of 1/85 LD<sub>50</sub> and 1/120 LD<sub>50</sub> of imidacloprid on thyroid toxicity in female rats (18) and a study of haematological and biochemical parameters in male rats after oral administration of 0.125 LD<sub>50</sub> and 0.1 LD<sub>50</sub> of imidacloprid, which correspond to the levels found in and on treated cucumbers one hour and one day after the application of imidacloprid, respectively (19).

In this mini-review, however, we focus on the current knowledge of adverse effects of imidacloprid on the reproductive and endocrine function as evidenced by recent animal studies.

### REPRODUCTIVE TOXICITY

Studies conducted in male and female laboratory animals, mostly rats, have established that oral administration of imidacloprid could affect reproductive function, as summarised in Table 1 (20-28). Subchronic and chronic exposure to different doses of imidacloprid can cause histological damage of the testicular tissue, changes in sperm morphology, increased sperm mortality and abnormality, decreased sperm count and motility, inhibited spermatogenesis, and reduced testosterone levels in male rats, all of which could lead to infertility (20-24).

More specifically, studies in male rats by Bal et al. (20, 21) showed that a three-month exposure to imidacloprid doses lower than the no-observable-effect-level (NOEL), which is 5-10 mg kg<sup>-1</sup> for rats, adversely affected their testicular function during early postnatal development and in adulthood. The elevated fatty acids and lipid peroxidation

**Correspondence to:** Anja Mikolić, PhD, Institute for Medical Research and Occupational Health, Ksaverska cesta 2, HR-10001 Zagreb, Croatia; E-mail: akatic@imi.hr

**Table 1** Summary of studies of imidacloprid (IMI) effect on reproduction in laboratory rats

Reference, year of publication	Sex/age	Exposure conditions (duration and route of administration)	Dose(s)	Effects
Bal et al. (20) 2012	male developing	90 days oral (gavage)	0.5, 2 and 8 mg kg <sup>-1</sup>	<p>↓ weights of epididymis, right cauda epididymis and seminal vesicle</p> <p>↓ epididymal sperm concentration at IMI – 8 mg kg<sup>-1</sup></p> <p>↑ abnormal sperm rates (head, tale and total values) at IMI – 8 mg kg<sup>-1</sup></p> <p>↑ apoptosis and seminal DNA fragmentation at IMI – 2 and 8 mg kg<sup>-1</sup></p> <p>↓ serum T</p> <p>↑ serum total cholesterol at IMI – 2 and 8 mg kg<sup>-1</sup></p> <p>↑ testicular fatty acids</p> <p>↑ MDA;</p> <p>↓ GSH in testis at IMI – 8 mg kg<sup>-1</sup></p>
Bal et al. (21) 2012	male adult	90 days oral (gavage)	0.5, 2 and 8 mg kg <sup>-1</sup>	<p>↓ weight of epididymis, right cauda epididymis and seminal vesicle</p> <p>↓ epididymal sperm concentration at IMI – 2 and 8 mg kg<sup>-1</sup></p> <p>↓ sperm motility at IMI – 8 mg kg<sup>-1</sup></p> <p>↑ abnormal sperm rates (head, tale and total values) at IMI – 8 mg kg<sup>-1</sup></p> <p>↑ apoptosis and seminal DNA fragmentation at IMI – 8 mg kg<sup>-1</sup></p> <p>↓ serum T at IMI – 8 mg kg<sup>-1</sup></p> <p>↓ GSH in testis at IMI – 8 mg kg<sup>-1</sup></p> <p>↑ testicular fatty acids change in disturbance of fatty acid composition</p>
Najafi et al. (22) 2010	male adult	60 days oral (gavage)	112 and 225 mg kg <sup>-1</sup>	<p>adverse histological changes on testicular tissue</p> <p>sperm mortality</p> <p>↑ abnormal sperm velocity</p> <p>↓ sperm viability</p> <p>↓ motile sperm velocity</p> <p>↓ T serum levels</p>
Hafez et al. (23) 2016	male adult	15 days oral (gavage)	45, 90 and 450 mg kg <sup>-1</sup>	<p>↓ LH, FSH, T, E2 and prolactin in testis</p> <p>↓ sperm count at 90 mg kg<sup>-1</sup></p> <p>↓ motility and vitality</p> <p>affected sperm morphology at 45 and 90 mg kg<sup>-1</sup></p> <p>↓ spermatogenesis</p>
Lonare et al. (24) 2015	male adult	28 days oral (gavage)	45 and 90 mg kg <sup>-1</sup>	<p>↓ total epididymal sperm count</p> <p>↓ sperm motility</p> <p>↓ live sperm count</p> <p>↑ sperm abnormalities (head and tail)</p> <p>↑ γ-GT, LDH-x, SDH in testis</p> <p>↓ T in plasma and testis</p> <p>↓ 3β-HSD, 17β-HSD in testis</p> <p>↑ LPO; ↓ GSH in testis</p> <p>↓ CAT, SOD at 90 mg kg<sup>-1</sup> in testis</p> <p>↓ GPx and GST in testis</p> <p>histopathological alterations in testis and epididymis</p>

Reference, year of publication	Sex/age	Exposure conditions (duration and route of administration)	Dose(s)	Effects
Kapoor et al. (25) 2011	female adult	90 days oral	5, 10 and 20 mg kg <sup>-1</sup> day <sup>-1</sup>	↓ ovarian weight at 20 mg kg <sup>-1</sup> significant pathomorphological changes in follicles, antral follicles and atretic follicles at 20 mg kg <sup>-1</sup> ↑ FSH; ↓ LH and P4 in serum at 20 mg kg <sup>-1</sup> ↑ LPO in ovary at 20 mg kg <sup>-1</sup> ↓ GSH, SOD, CAT and GPX in ovary at 20 mg kg <sup>-1</sup>
Nabiuni et al. (26) 2015	pregnant female	from gestation day 7 to 21, via tail vein	10 mg kg <sup>-1</sup> bw	↓ E2 and P in serum of 55-day-old offspring of IMI-treated mothers ↓ ovarian weight, ovary diameter, number and diameter of follicles and corpus luteum in the offspring of IMI-treated mothers ↓ rate of successful mating and number of fetus in the offspring of IMI-treated mothers down regulation of DAX1 gene in ovary of the offspring of IMI-treated mothers
Suter et al. (27) 1990	female adult	P1/P2: 84 days/105 days before mating, through mating, gestation and lactation oral	100, 250 and 700 ppm	↔ no reproductive effects ↓ body weight gain in P1 at 700 ppm ↓ food consumption in P1 at 700 ppm ↓ body weight gain in P1 – offspring until weaning at 700 ppm ↓ pre-mating body weights in P1-F1 males at 700 ppm
Vohra et Khera (28) 2016	female adult	F0 females: 10 weeks or more F1 parents: 8 weeks or more oral	10 and 20 mg kg <sup>-1</sup> bw day <sup>-1</sup>	↓ ovarian weight in F0 and F1 generation at 20 mg kg <sup>-1</sup> bw day <sup>-1</sup> ↑ body weight of F0 pups (day 0) ↓ body weight of F1 pups (day 21) histopathology of ovary revealed different stages of follicles ↑ ALT in F1 and F2 females ↑ body weight of F1 pups on day 0 at 20 mg kg <sup>-1</sup> bw day <sup>-1</sup> ↓ body weight in F2 pups on day 21 at 20 mg kg <sup>-1</sup> bw day <sup>-1</sup> ↓ food consumption in IMI treated females of F0 and F1 generation

Abbreviations: ↑=increase; ↓= decrease; ↔ = no effect; ALT=alanine aminotransferase; bw=body weight; CAT=catalase; E2=oestradiol; F1/2=first/second generation of offspring; FSH=follicular stimulating hormone; GPx=glutathione peroxidase; γ-GSH=reduced glutathione; GST= glutathione-S-transferase; GT= gamma-glutamyl transpeptidase; 3β-HSD= 3β-hydroxysteroid dehydrogenase; 17β-HSD= 17β-hydroxysteroid dehydrogenase; IMI = imidacloprid; LDH-x= lactate dehydrogenase-x; LH=luteinizing hormone; LPO=lipid peroxidation; P4=progesterone; P/F0=parental generation; SDH=sorbitol dehydrogenase; SOD=superoxide dismutase; T=testosterone

(LPO) accompanied by lower glutathione (GSH) levels in the testes of the exposed rats resulted in impaired fertility, as evidenced by increased germ cell apoptosis and fragmentation of seminal DNA, lower serum testosterone level and higher rate of sperm abnormalities. The authors associated these adverse effects with the induction of oxidative stress in the testes, which are particularly sensitive to oxidative damage due to the presence of polyunsaturated fatty acids.

In female rats, Kapoor et al. (25) reported that exposure to imidacloprid (20 mg kg<sup>-1</sup> day<sup>-1</sup>) for 90 days caused

changes in the ovarian morphology, reduced its weight, and adversely affected serum follicle stimulating hormone (FSH), luteinizing hormone (LH), and progesterone levels. The authors also reported induced oxidative stress, evidenced by lower activity of superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx), and GSH and increased LPO level.

These studies were the first to suggest the mechanisms of adverse imidacloprid effects on male and female reproduction. Later research reported higher LPO and lower GSH levels, as well as lower CAT, SOD, GPx, and

glutathione *S*-transferase (GST) activities in male rats exposed to higher doses of imidacloprid (45 and 90 mg kg<sup>-1</sup> bw, corresponding to 1/10 and 1/5 of LD<sub>50</sub>) for 28 days and confirmed that oxidative stress is the mechanism that leads to reproductive toxicity of imidacloprid (24).

Reproductive toxicity of imidacloprid was also studied following *in utero* exposure of the female offspring of pregnant rats receiving 10 mg kg<sup>-1</sup> bw of imidacloprid through the tail vein from gestation day 7 to 21 (26). The offspring showed disrupted follicular development, lower oestradiol and progesterone levels, lower ovarian weight and diameter as well as decrease in the rate of successful mating and reduction in foetal number due to down-regulation of ovarian *DAX1* gene compared to controls.

In a two-generation, two-litter study in rats Suter et al. (27) reported no adverse effects on the reproductive function or birth defects of imidacloprid exposure before and during mating, gestation, and lactation. However, the authors found decreased body weight in mother rats and male and female offspring that was associated with reduced food consumption at dietary imidacloprid dose of 700 ppm (corresponding to 47 mg kg<sup>-1</sup> bw day<sup>-1</sup>). In a three-generation reproductive study (28) oral administration of 20 mg kg<sup>-1</sup> bw day<sup>-1</sup> of imidacloprid affected body weight and food consumption and caused only minimal effects on the reproductive performance such as reduction in fertility index in the second and third generation.

Neurotoxicity as the consequence of imidacloprid exposure *in utero* has been investigated in rat offspring by Abou-Donia et al. (29). Gestational exposure to a single large dose (0.75 of the LD<sub>50</sub>) of imidacloprid caused developmental neurobehavioral deficits and increased the expression of glial fibrillary acidic protein (GFAP) in the brain regions of 30-day-old offspring with possible long-term adverse health effects. An *in vitro* investigation by Gu et al. (30) has shown adverse effects of direct exposure to imidacloprid on the sperm function and embryonic development that may also present a risk for human reproductive health.

Investigations in other laboratory animals showed no adverse effects of imidacloprid exposure on the early development of the zebrafish (*Danio rerio*) (31), but in male rabbits (*Oryctolagus cuniculus*) it was associated with testicular damage that can lead to infertility and reproductive disorders (32).

## ENDOCRINE DISRUPTION

Many pesticides, including imidacloprid, can affect the normal functioning of animal endocrine systems by interfering with natural hormones. Due to their strong affinity for oestrogen and androgen receptors, they can activate these receptors and mimic hormonal action or block the receptors and inhibit their action. In addition, endocrine disrupting pesticides can interfere with the synthesis,

secretion, transport, metabolism, binding, or elimination of natural hormones (32).

Some investigations imply that exposure to environmental endocrine disruptors could be the generative cause of obesity (34, 35). This may also be true for endocrine disrupting pesticides. Imidacloprid's role in obesity was investigated in a study in which mice were exposed to low doses (0.5 % of the LD<sub>50</sub>) of dithiocarbamate mancozeb and imidacloprid from postnatal day 1 to weaning (postnatal day 28) through lactating mother. The exposure to individual pesticides did not disrupt hormonal/metabolic homeostasis or affect body weight gain, but exposure to their combination did (36). An *in silico* study by the same authors showed that imidacloprid and mancozeb disrupted steroid hormone biosynthesis by impairing the activities of the cytochrome P450 (CYP) enzymes that have a key role in the synthesis and breakdown of various steroid hormones (37). Several studies have shown that imidacloprid potentiates adipogenesis in 3T3-L1 adipocytes and induces insulin resistance in C2C12 myotubes (38, 39). The same group of authors reported that oral exposure to low doses (0.6 and 6 mg kg<sup>-1</sup> bw day<sup>-1</sup>) of imidacloprid for 12 weeks in male and female C57BL/6J mice enhanced high fat-diet adiposity by mechanisms that still need to be clarified. Insulin resistance observed in male mice was not confirmed in females, which was explained by potential sex differences (40, 41).

## CONCLUSION

Animal study evidence is pretty clear about the risks involved in imidacloprid exposure for male and female reproduction. It may alter essential reproductive functions by affecting hormone levels and cause reproductive organ disruption and suppression of their function, which may lead to infertility. *In utero* exposure to imidacloprid can significantly affect offspring development with lasting adverse consequences in adulthood. The European Commission has restricted the use of imidacloprid and two other neonicotinoids since 2013 due to their potential role in the collapse of the bee population (42). The updated risk assessment report regarding neonicotinoids was issued by the European Food Safety Authority in February 2018 (43). Although it points to risks for honeybees, the decision regarding a possible EU-wide ban of neonicotinoids was not made and remains the responsibility of the European Commission and Member state authorities. Nevertheless, since imidacloprid is still widely used, more measures should be taken to minimise its adverse effects.

### Acknowledgements

This work was financially supported by Project Organic Pollutants in Environment – Markers and Biomarkers of Toxicity (OPENTOX), funded by the Croatian Science Foundation (grant number 8366).

### Conflicts of interest

The authors declare no conflict of interest.

### REFERENCES

1. Matsuda K, Buckingham SD, Kleier D, Rauh JJ, Grauso M, Sattelle DB. Neonicotinoids: insecticides acting on insect nicotinic acetylcholine receptors. *Trends Pharmacol Sci* 2001;22:573-80. doi: 10.1016/S0165-6147(00)01820-4
2. Tomizawa M, Casida JE. Neonicotinoid insecticide toxicology: mechanisms of selective action. *Annu Rev Pharmacol* 2005;45:247-68. doi: 10.1146/annurev.pharmtox.45.120403.095930
3. California Environmental Protection Agency (CEPA). Imidacloprid risk characterization document: Dietary and drinking water exposure. Sacramento (CA): Department of Pesticide Regulation; 2006.
4. US Environmental Protection Agency (US EPA). Imidacloprid; Pesticide tolerances. 22245-52. Washington (DC): US EPA; 2003.
5. Di Muccio A, Fidente P, Barbini DA, Dommarco R, Seccia S, Morrica P. Application of solid-phase extraction and liquid chromatography-mass spectrometry to the determination of neonicotinoid pesticide residues in fruit and vegetables. *J Chromatogr A* 2006;1108:1-6. doi: 10.1016/j.chroma.2005.12.111
6. Liu MY, Casida JE. High affinity binding of [<sup>3</sup>H]-imidacloprid in the insect acetylcholine receptor. *Pestic Biochem Phys* 1993;46:40-6. doi: 10.1006/pest.1993.1034
7. Chao S-L, Casida JE. Interaction of imidacloprid metabolites and analogs with nicotinic acetylcholine receptor of mouse brain in relation to toxicity. *Pestic Biochem Phys* 1997;58:77-88. doi: 10.1006/pest.1997.2284
8. Zhang A, Kaiser H, Maienfisch P, Casida JE. Insect nicotinic acetylcholine receptor: Conserved neonicotinoid specificity of [<sup>3</sup>H]-imidacloprid binding site. *J Neurochem* 2000;75:1294-303. doi: 10.1046/j.1471-4159.2000.751294.x
9. Tomlin CDS, editor. *The Pesticide Manual*. 11<sup>th</sup> ed. Surrey: British Crop Protection Council; 1997. p. 706-8.
10. Becker H, Vogel W, Terrier C. Embryotoxicity study (including teratogenicity) with NTN 33893 technical in the rat. Unpublished Report no. R 4582, 1988, submitted to WHO by Bayer AG, Mannheim, Germany. INCHEM Toxicological Evaluations: Imidacloprid; International Programme on Chemical Safety. Geneva: WHO; 1988.
11. Becker H, Vogel W, Terrier C. Embryotoxicity study (including teratogenicity) with NTN 33893 technical in the rabbit. Unpublished Report no. R 4583, 1988, submitted to WHO by Bayer AG, Mannheim, Germany. INCHEM Toxicological Evaluations: Imidacloprid; International Programme on Chemical Safety. Geneva: WHO; 1988.
12. Bagri P, Kumar V, Sikka AK. Assessment of imidacloprid-induced mutagenic effects in somatic cells of Swiss albino male mice. *Drug Chem Toxicol* 2016;39:412-7. doi: 10.3109/01480545.2015.1137301
13. Duzguner V, Erdogan S. Acute oxidant and inflammatory effects of imidacloprid on the mammalian central nervous system and liver in rats. *Pestic Biochem Phys* 2010;97:13-8. doi: 10.1016/j.pestbp.2009.11.008
14. Duzguner V, Erdogan S. Chronic exposure to imidacloprid induces inflammation and oxidative stress in the liver & central nervous system of rats. *Pestic Biochem Phys* 2010;104:58-64. doi: 10.1016/j.pestbp.2012.06.011
15. Lonare M, Kumar M, Raut S, Badgujar P, Doltade S, Telang A. Evaluation of imidacloprid-induced neurotoxicity in male rats: a protective effect of curcumin. *Neurochem Int* 2014;78:122-9. doi: 10.1016/j.neuint.2014.09.004
16. Gawade L, Dadarkar SS, Husain R, Gatne M. A detailed study of developmental immunotoxicity of imidacloprid in Wistar rats. *Food Chem Toxicol* 2013;51:61-70. doi: 10.1016/j.fct.2012.09.009
17. Badgujar PC, Jain SK, Singh A, Punia JS, Gupta RP, Chandratre GA. Immunotoxic effects of imidacloprid following 28 days of oral exposure in BALB/c mice. *Environ Toxicol Phar* 2013;35:408-18. doi: 10.1016/j.etap.2013.01.012
18. Saadi L, Mahboubi Y, Matallah R. Inflammatory effects of imidacloprid on thyroid activity in rats. In: *International Conference on Civil, Biological and Environmental Engineering (CBEE-2014)*; 27-28 May 2014; Istanbul, Turkey [displayed 23 May 2018]. Available at <http://iicbe.org/upload/9427C514572.pdf>
19. Abbassy MA, Marzouk MA, Nasr HM, Mansy ASM. Effect of imidacloprid and tetraconazole on various haematological and biochemical parameters in male albino rats (*Rattus Norvegicus*). *J Pol Sci Pub Aff* 2014;2:122. doi:10.4172/2332-0761.1000122
20. Bal R, Naziroğlu M, Türk G, Yimalz Ö, Kuloğlu T, Etem E, Baydas G. Insecticide imidacloprid induces morphological and DNA damage through oxidative toxicity on the reproductive organs of developing male rats. *Cell Biochem Funct* 2012;30:492-9. doi: 10.1002/cbf.2826
21. Bal R, Türk G, Tuzcu M, Yimalz Ö, Kuloğlu T, Gundogdu R, Gür S, Agca A, Ulas M, Çambay Z, Tuzcu Z, Gencoglu H, Guvenc M, Ozsahin AD, Kocaman N, Aslan A, Etem E. Assessment of imidacloprid toxicity on reproductive organ system of adult male rats. *J Environ Sci Health B* 2012;47:434-44. doi: 10.1080/03601234.2012.663311
22. Najafi G, Razi M, Hoshyar A, Shahmohammadloo S, Feyzi S. The effect of chronic exposure with imidacloprid insecticide on fertility in mature male rats. *Int J Fertil Steril* 2010;4:9-16.
23. Hafez EM, Issa SY, Al-Mazroua MK, Ibrahim KT, Rahman SMA. The neonicotinoid insecticide imidacloprid: a male reproductive system toxicity inducer-human and experimental study. *Toxicol Open Access* 2016;2:1000109. doi: 10.4172/tyoa.1000109
24. Lonare M, Kumar M, Raut S, More A, Doltade S, Badgujar P, Telang A. Evaluation of ameliorative effect of curcumin on imidacloprid-induced male reproductive toxicity in Wistar rats. *Environ Toxicol* 2016;10:1250-63. doi: 10.1002/tox.22132
25. Kapoor U, Srivastava MK, Srivastava LP. Toxicological impact of technical imidacloprid on ovarian morphology, hormones and antioxidant enzymes in female rats. *Food Chem Toxicol* 2011;49:3086-9. doi: 10.1016/j.fct.2011.09.009
26. Nabuini M, Parivar K, Noorinejad R, Falahati Z, Khalili F, Krimzadeh L. The reproductive side effects of imidacloprid in pregnant Wistar rat. *Int J Cell Mol Biotechnol* 2015;1:10-8. doi: 10.5899/2015/ijcmb-00017
27. Suter P, Biedermann K, Luetkemeier H, Wilson J, Terrier C. NTN 33893 technical. Multiple generation study in rats. RCC, Research and Consulting Company AG. Itingen,

- Switzerland. Study No. 100647. DPR Vol.51950-0019# 119496, 1990.
28. Vohra P, Khera KS. Effect of imidacloprid on reproduction on female albino rats in three generation study. *J Vet Sci Technol* 2016;7:340. doi:10.4172/2157-7579.1000340
  29. Abou-Donia MB, Goldstein LB, Bullman S, Tu T, Khan WA, Dechkovskaia AM, Abdel-Rahman AA. Imidacloprid induces neurobehavioral deficits and increases expression of glial fibrillary acidic protein in the motor cortex and hippocampus in offspring rats following *in utero* exposure. *J Toxicol Environ Health A* 2008;71:119-30. doi: 10.1080/15287390701613140
  30. Gu YH, Li Y, Huang XF, Zheng JF, Yang J, Diao H, Yuan Y, Xu Y, Liu M, Shi HJ, Xu WP. Reproductive effects of two neonicotinoid insecticides on mouse sperm function and early embryonic development *in vitro*. *PloS One* 2013;8:e70112. doi: 10.1371/journal.pone.0070112
  31. Scheil V, Köhler HR. Influence of nickel chloride, chlorpyrifos, and imidacloprid in combination with different temperatures on the embryogenesis of the zebrafish *Danio rerio*. *Arch Environ Contam Toxicol* 2009;56:238-43. doi: 10.1007/s00244-008-9192-8
  32. Memon SA, Memon N, Mal B, Shaikh SA, Shah MA. Histopathological changes in the gonads of Male rabbits (*Oryctolagus cuniculus*) on exposure to imidacloprid insecticide. *J Entomol Zool Stud* 2014;2:159-63.
  33. Mnif W, Hassine AIH, Bouaziz A, Bartegi A, Thomas O, Roig B. Effect of endocrine disruptor pesticides: a review. *Int J Environ Res Public Health* 2011;8:2265-303. doi: 10.3390/ijerph8062265
  34. Janesick A, Blumberg B. Endocrine disrupting chemicals and the developmental programming of adipogenesis and obesity. *Birth Defects Res C Embryo Today* 2011;93:34-50. doi: 10.1002/bdrc.20197
  35. Legler J, Hamers T, van Eck van der Sluijs-van de Bor M, Schoeters G, van der Ven L, Eggesbo M, Koppe J, Feinberg M, Trnovec T. The OBELIX project: early life exposure to endocrine disruptors and obesity. *Am J Clin Nutr* 2011;94:1933S-8S. doi: 10.3945/ajcn.110.001669
  36. Bhaskar R, Mohanty B. Pesticides in mixture disrupt metabolic regulation: *In silico* and *in vivo* analysis of cumulative toxicity of mancozeb and imidacloprid on body weight of mice. *Gen Comp Endocrinol* 2014;205:226-34. doi: 10.1016/j.ygcen.2014.02.007
  37. Bhaskar R, Mishra AK, Mohanty B. Effects of mancozeb and imidacloprid pesticides on activities of steroid biosynthetic enzymes cytochromes P450. *J Kalash Sci* 2014;2:1-6.
  38. Park Y, Kim Y, Kim J, Yoon KS, Clark J, Lee J, Park Y. Imidacloprid, a neonicotinoid insecticide, potentiates adipogenesis in 3T3-L1 adipocytes. *J Agric Food Chem* 2013;61:255-9. doi: 10.1021/jf3039814
  39. Kim J, Park Y, Yoon KS, Clark JM, Park Y. Imidacloprid, a neonicotinoid insecticide, induces insulin resistance. *J Toxicol Sci* 2013;38:655-60. doi: 10.2131/jts.38.655
  40. Sun Q, Xiao X, Kim Y, Kim D, Yoon KS, Clark JM, Park Y. Imidacloprid promotes high fat diet-induced adiposity and insulin resistance in male C57BL/6J mice. *J Agric Food Chem* 2016;64:9293-306. doi: 10.1021/acs.jafc.6b04322
  41. Sun Q, Qi W, Xiao X, Yang SH, Kim D, Yoon KS, Clark JM, Park Y. Imidacloprid promotes high fat diet-induced adiposity in female C57BL/6J mice and enhances adipogenesis in 3T3-L1 adipocytes via the AMPK $\alpha$ -mediated pathway. *J Agric Food Chem* 2017;65:6572-81. doi: 10.1021/acs.jafc.7b02584
  42. The European Commission. Commission Implementing Regulation (EU) No 485/2013 of 24 May 2013 amending Implementing Regulation (EU) No 540/2011, as regards the conditions of approval of the active substances clothianidin, thiamethoxam and imidacloprid, and prohibiting the use and sale of seeds treated with plant protection products containing those active substances. Text with EEA relevance [displayed 20 March 2018]. Available at <http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32013R0485&from=EN>
  43. European Food Safety Authority. Evaluation of the data on clothianidin, imidacloprid and thiamethoxam for the updated risk assessment to bees for seed treatments and granules in the EU [displayed 6 June 2018]. Available at <https://efsa.onlinelibrary.wiley.com/doi/epdf/10.2903/sp.efsa.2018.EN-1378>

### Imidacloprid kao reprodukcijski toksikant i endokrini disruptor: istraživanja na laboratorijskim životinjama

Imidacloprid, neonicotinoidni insekticid, koristi se diljem svijeta zbog svoje selektivne toksičnosti za insekte. S obzirom na to da njegovi ostaci mogu ući u hranidbeni lanac, važno je istražiti potencijalno štetne učinke izloženosti imidaclopridu. U ovom preglednom radu prikazani su podaci o reprodukcijskoj toksičnosti imidacloprida i njegovim učincima kao hormonskog otrova u laboratorijskih životinja. Istraživanja koja su provedena uglavnom na laboratorijskim štakorima pokazala su štetne učinke imidacloprida na reprodukciju roditelja i potomstva, kao i na razvoj potomstva. Poput mnogih pesticida, imidacloprid može djelovati i kao hormonski otrov te poremetiti ne samo metaboličku ravnotežu i pridonijeti povećanju pretilosti nego i steroidogenezu inhibicijom aktivnosti sustava enzima citokrom P-450 (CYP). Svi ti štetni učinci imidacloprida mogu predstavljati ozbiljan rizik za reprodukciju i razvoj s dugoročnim štetnim posljedicama u odrasloj dobi.

KLJUČNE RIJEČI: endokrini disruptor; insekticid; laboratorijski štakori; neonicotinoidi; reproduktivna toksičnost