

## Effect of pretreatments on mycotoxin profiles and levels in dried figs

Jasenka Petrić<sup>1</sup>, Bojan Šarkanj<sup>2</sup>, Ibrahim Mujić<sup>3</sup>, Aida Mujić<sup>4</sup>, Michael Sulyok<sup>5</sup>, Rudolf Krška<sup>5</sup>,  
Drago Šubarić<sup>6</sup>, and Stela Jokić<sup>6</sup>

Croatian Food Agency, Osijek, Croatia<sup>1</sup>, University North, Center Koprivnica, Department of Food Technology, Koprivnica, Croatia<sup>2</sup>, Collegium Fluminense Polytechnic of Rijeka, Rijeka, Croatia<sup>3</sup>, Clinical Hospital Centre Rijeka, Clinical Department of Transfusion Medicine, Rijeka, Croatia<sup>4</sup>, Center for Analytical Chemistry, Department for Agrobiotechnology, IFA, Tulln, University of Natural Resources and Life Sciences, Vienna, Austria<sup>5</sup>, Josip Juraj Strossmayer University of Osijek, Faculty of Food Technology, Osijek, Croatia<sup>6</sup>

[Received in May 2018; Similarity Check in May 2018; Accepted in December 2018]

The aim of this explorative study was to investigate how effective drying preservation methods are in reducing mycotoxin content in figs. Dried autochthonous varieties of white and dark figs (Petrovača Bijela and Šaraguja, respectively) were analysed for mycotoxins using an LC-MS/MS “dilute and shoot” method capable of determining 295 fungal and bacterial secondary metabolites. Before drying in a cabinet dryer the figs were preserved with 0.5 % citric acid solution or 0.5 % ascorbic acid solution or 0.3 % L-cysteine solution or 0.2 % chestnut extract solution or 0.15 % Echinacea extract solution by immersion. We found nine metabolites: aflatoxin B1 (AFB<sub>1</sub>), ochratoxin A, ochratoxin alpha, kojic acid, emodin, altenuene, alternariol methyl ether, brevianamide F, and tryptophol. The most efficient preserver was L-cysteine (15 % reduction), while ascorbic acid favoured mycotoxin production (158 % increase). However, all pretreatment solutions reduced AFB<sub>1</sub>, which is a major fig contaminant.

KEY WORDS: antimycotoxigenic effect; drying; LC-MS/MS; multi-mycotoxin analysis

Mycotoxins are naturally occurring secondary metabolites of various moulds, some of which are the natural microbiota of figs and are unavoidable contaminants. However, through the implementation of good agricultural and manufacturing practices it is possible to minimise the contamination. According to Heperkan et al. (1), the dominant fungal flora in the Mediterranean dried figs are the *Aspergillus*, *Fusarium*, and *Penicillium* species and the related *Aspergillus* (2–5), *Fusarium* (6, 7), *Penicillium* (8, 9), and *Alternaria* mycotoxins (10). The last, *Alternaria alternata* has been reported in figs from Saudi Arabia and Montenegro. The Saudi study reported the presence of altenuene and alternariol (11), but the Montenegrin mycotoxin analysis found no mycotoxins (12). Other authors (13) reported high amounts of kojic acid (KA) in dried figs, as well as methoxysterigmatocystin, roquefortine A, fumagillin, fumigaclavine B, malformins (peptides), aspergillilic acid, nigragillin, terrein, terrestric acid, and penicillilic acid.

Mycotoxin production in figs begins on the tree and depends on a number of extrinsic and intrinsic factors, including stress and physical damage (1). Mycotoxin levels may increase during transportation, storage, and processing,

if the conditions are favourable. However, contamination can be controlled with effective measures, starting with the orchards through transportation, storage, and processing like drying in a solar drier (1, 3). Drying is one of the oldest preservation methods that can also reduce and/or control mycotoxin contamination. Traditional sun drying is undesirable because figs are prone to contamination with fungal spores from the soil (14). An alternative to traditional drying are mechanical air-drying systems which provide controlled environment (15).

Drying is often combined with pretreatment with water and chemical solutions (14, 16, 17) to counter undesired effects and improve the quality of dried products. L-cysteine has proved its worth in preventing the oxidation (and related browning) of apples and potatoes (18), and other natural antioxidants, such as ascorbic acid, Echinacea (19), citric acid (16), and phenolic compounds extracted from chestnuts (20, 21) have also been successful antioxidants. Since oxidative stress is one of the prerequisites for mycotoxin biosynthesis (22), antioxidants such as these also exert antimycotoxigenic effects (23). However, knowledge about these antimycotoxigenic effects is still scarce (24). The aim of this study was therefore to determine the efficiency of pretreatment with the most common antioxidants in reducing mycotoxin content in dried autochthonous Croatian fig varieties.

**Correspondence to:** Dr Bojan Šarkanj, Sveučilište Sjever, Sveučilišni centar Koprivnica, Trg dr. Žarka Dolinara 1, 48000 Koprivnica, Croatia  
E-mail: bsarkanj@unin.hr

## MATERIALS AND METHODS

### Chemicals and reagents

HPLC-grade methanol and acetonitrile as well as MS-grade ammonium acetate and glacial acetic acid (p.a.) were purchased from Sigma-Aldrich (Vienna, Austria). Ultrapure water was obtained by purification of reverse osmosis water with a Purelab Ultra system (ELGA LabWater, Celle, Germany). Ascorbic acid, citric acid, and L-cysteine were purchased from Sigma-Aldrich Chemie GmbH (Taufkirchen, Germany). The chestnut extract was produced from whole local chestnut variety Lovranski marun, and the Echinacea extract was produced from the flower of *Echinacea purpurea*. Both were obtained by adding 250 mL of 50 % ethanol to 50 g of the sample (sample-to-solvent ratio was 1:5) and applying ultrasonication at 50 Hz and 125 W for 30 min. The liquid extract was obtained by filtration through Whatman® Grade 4 filter paper (Whatman, Maidstone, United Kingdom) and removal of solvent by evaporation under vacuum at 40 °C.

### Sampling

Figs were obtained from a pilot orchard in Kazela, Istria, Croatia, harvested in August 2015. We used two most common autochthonous varieties of fresh white and dark figs, namely Petrovača Bijela and Šaraguja (Figure 1, Table 1). Two random batches (5 kg) per treatment were used and mixed before mycotoxin measurements. Control samples were pretreated with distilled water.

### Drying

Untreated (control) and pretreated whole figs were dried on 17 trays with 2.5 kg of figs on each in a cabinet dryer

(2x4 kW heater power, 150 kg drying capacity, Rasadnik Skink Rovinj, Croatia). The dryer was equipped with temperature and airflow velocity controllers. Drying temperature was 60-70 °C.

Before drying, the figs intended for pretreatment were immersed in 0.5 % citric acid solution or 0.5 % ascorbic acid solution or 0.3 % L-cysteine solution or 0.2 % chestnut extract solution or 0.15 % Echinacea extract solution for 5 min.


The measurements started as soon as the fig samples were placed onto the trays in the cabinet dryer. Sample weight loss was measured with a digital scale every 10 min, and drying stopped when the moisture content dropped to 25 % of the fresh fruit content. The samples were then stored in paper bags at room temperature for 30 days before mycotoxin analysis.

### Mycotoxin analysis

Standards for mycotoxin analysis were prepared as described elsewhere (24). Briefly, 34 stock solutions of analyte standards were mixed into one working solution, which was used for spiking and calibration.

We used an LC-MS/MS “dilute-and-shoot” method capable of determining 295 fungal and bacterial secondary metabolites in one run as described elsewhere (24).

Frozen, dried, and cut figs were transported to Austria, ground there with an Osterizer blender (Sunbeam Oster Household Products, Fort Lauderdale, FL, USA), and 5 g samples weighed and put in 50 mL tubes. Then 20 mL of extraction solvent (acetonitrile/water/acetic acid 79:20:1, v/v/v) was added and the content thoroughly mixed by hand. Followed a 90-minute extraction at room temperature with a GFL 3017 rotary shaker set at 180 rpm (GFL, Burgwedel, Germany). After extraction, the samples were allowed to precipitate, and 500 µL of the extract was diluted with 500 µL of dilution solvent (acetonitrile/water/acetic acid 20:79:1, v/v/v) in 1.5 mL vials. The vials were mixed by vortexing for 10 s, and 5 µL of the diluted sample extract was directly injected into a liquid chromatography – mass spectrometry (LC-MS/MS) system. The LC part consisted of an Agilent 1290 binary UHPLC pump (Agilent Technologies, Waldbronn, Germany) coupled with a Sciex QTRAP® 5500 triple quadrupole MS/MS detector (Sciex, Foster City, CA, USA). The chromatographic conditions were strictly followed as described elsewhere (24). The sample with the lowest concentration of mycotoxins was used for the recovery experiment by spiking 250 mg of the sample with 100 µL of the working standard solution. Mycotoxin levels used for recovery were as follows: aflatoxin B1 (AFB<sub>1</sub>) – 4.57 µg kg<sup>-1</sup>; OTA – 26.1 µg kg<sup>-1</sup>; ochratoxin α (OTα) – 3.30 µg kg<sup>-1</sup>; alternariolmethylether (AME) – 14.4 µg kg<sup>-1</sup>; emodin (EMO) – 7.21 µg kg<sup>-1</sup>; altenuen (ATN) – 7.21 µg kg<sup>-1</sup>; KA – 206 µg kg<sup>-1</sup>; tryptophol (TryOH) – 26.1 µg kg<sup>-1</sup>; and brevianamide F (BREF) – 5.23 µg kg<sup>-1</sup>. The exact mycotoxin levels in the mastermix



**Figure 1** Autochthonous white (left) and dark (right) fig varieties in Istria used for testing the efficiency of pretreatment in reducing mycotoxin levels

**Table 1** Characteristics of the investigated fig varieties

	Petrovača Bijela	Šaraguja
Fruit volume (cm <sup>3</sup> )	94	45
Fruit weight (g)	82	51
Fruit width (mm)	55	46
Fruit length (mm)	57	45
Fruit skin thickness (mm)	11	4
Fruit skin firmness	Medium	Firm
Fruit skin cracks	Absent	Minute crack
Resistance to cracks	Resistant	Intermediate
Fruit skin ground colour	Green	Green
Overcolour – regular bands	Yellow, green	Purple
Pulp internal colour	Pink, amber	Red
Dry weight (%)	25.5	26.5

All values are the means of three measurements

have been described elsewhere (24). The sample was left at room temperature in the dark overnight to allow for the solvent to evaporate and to ensure complete sample-spike equilibration of the mycotoxins in the matrix.

All results were corrected for the obtained recovery. Measurements were performed as described elsewhere (24) and quantified with the Analyst<sup>®</sup> 1.6.2 and Multiquant<sup>®</sup> 3.0.2. software (Sciex). The limits of detection (LOD) and quantification (LOQ) were determined as the signal-to-noise ratio of 3 and 10 in the tested matrices, respectively, while the linearity of the calibration curve was calculated automatically with the Multiquant<sup>®</sup> 3.0.2. software.

The accuracy of the method had been verified through regular participation in proficiency testing schemes organised by the Bureau Interprofessionnel d'Etudes Analytiques - BIPEA (Gennevilliers, France). All six results submitted for fig samples exhibited a z-score between -2 and 2.

#### Statistical analysis

The data were checked for normality of distribution with the Shapiro-Wilk test and compared with nonparametric methods (Mann-Whitney U test for the comparison between two fig varieties and Kruskal-Wallis ANOVA for the comparison of pretreatments). For the analysis we used Statistica<sup>®</sup> 12.0 (Dell Inc., Round Rock, TX, USA) and Microsoft Excel<sup>®</sup> 2016 (Microsoft, Redmond, WA, USA). Differences <0.05 were considered significant.

## RESULTS AND DISCUSSION

Of the 295 bacterial and fungal metabolites detectable by the method, we found only nine mycotoxins above LOD (Table 2). Of these, TryOH, BREF, EMO, AME, ATN, and OTα were evidenced for the first time in figs.

Table 3 shows the effects of pretreatment compared to control samples. L-cysteine pretreatment achieved the best

overall results, while all pretreatments proved to be effective against the most problematic mycotoxin in figs, AFB<sub>1</sub>, with reductions ranging from 25 to 93 %.

AFB<sub>1</sub> levels were below the LOD in half of the samples. In the rest of them they were much lower than the EU limit of 6 µg kg<sup>-1</sup> (14). The dark Šaraguja variety had higher median and maximum values than the white variety, but the difference was not statistically significant (Table 2). The best reduction (to below LOD) was achieved with ascorbic acid (both levels were below LOD).

OTA was detected in only one sample (at level below the LOQ), unlike its main major metabolite OTα, whose levels were higher than those of AFB<sub>1</sub> in all but the white fig sample pretreated with citric acid. This finding indicates that OTA had been present in the figs at some point, but the phenylalanine part was cleaved off by other microflora in the fig (25). The dark Šaraguja variety had significantly higher OTα levels than the white Petrovača variety (p=0.005).

AFB<sub>1</sub> and OTA have often been reported in dried figs (1, 3, 23), and particularly high contamination was reported in dried figs from Turkey and Egypt (1). In contrast, all of our samples from Croatia were below MLs (26). This may be related to harsher climate that does not favour *Aspergillus* spp. growth. However, if the global temperature will continue to increase, aflatoxigenic *Aspergilli* will inevitably take a stronger hold in southeast Europe, including Croatia (27). OTA, on the other hand, is also produced by the *Penicillium* spp. that grow in colder climates, which calls for regular OTA monitoring in figs, regardless of their region of origin.

Even though this is the first evidence of AME in dried figs, its concentrations were very low or below LOD. The acute toxicity of AME is low, and little is known about its mutagenic and genotoxic properties (10). AME, ATN, alternariol (AOH), tentoxin (TTX), and tenuazonic acid (TeA) are produced by the *Alternaria* spp., whose occurrence is rare in dried figs (10). Again, the dark variety

**Table 2** Mass fractions ( $\mu\text{g kg}^{-1}$ ) of mycotoxins above LOD

Sample	Sort	AFB1	OTA	OT $\alpha$	AME	ATN	EMO	KA	TryOH	BREF
Control	white	0.41	<LOD	2.00*	0.30	12.0	0.64	449	21.2*	8.02
Control	dark	0.65	<LOD	3.81	0.10*	16.7	<LOD	5.575	<LOD	4.34
L-cysteine	white	0.12*	<LOD	0.92*	<LOD	9.20	<LOD	1.191	<LOD	5.27
L-cysteine	dark	<LOD	<LOD	5.71	<LOD	10.9	<LOD	4.636	17.4*	5.65
Ascorbic acid	white	<LOD	0.5*	<LOD	0.36	16.5	0.24*	1.388	20.8*	7.71
Ascorbic acid	dark	<LOD	<LOD	4.53	0.34	21.9	1.88	5.257	27.0*	10.9
Citric acid	white	0.32	<LOD	<LOD	<LOD	30.0	0.26*	1.148	30.7*	8.38
Citric acid	dark	<LOD	<LOD	6.81	<LOD	30.8	<LOD	4.853	<LOD	5.09
Chestnut extract	white	<LOD	<LOD	2.54*	0.11*	12.1	0.71	1.414	19.0*	7.99
Chestnut extract	dark	0.33	<LOD	6.58	0.23*	24.8	<LOD	8.594	99.1	7.21
Echinacea extract	white	<LOD	<LOD	2.87	<LOD	6.80	<LOD	850	20.2*	8.56
Echinacea extract	dark	0.44	<LOD	3.59	0.23*	14.7	<LOD	5.416	<LOD	5.71
Recovery		91 %	96 %	71 %	89 %	111 %	95 %	89 %	79 %	76 %
LOD		0.08	0.40	0.80	0.08	0.80	0.16	5.00	16.0	0.40
LOQ		0.30	1.30	2.60	0.30	2.60	0.50	16.5	52.8	1.30
r <sup>2</sup>		0.998	0.999	0.998	0.999	0.994	0.999	0.997	0.999	0.999

All values are the means of three measurements. AFB1=afatoxin B1; OTA=ochratoxin A; OT $\alpha$ =ochratoxin alpha; AME=alternariol methyl ether; ATN=altenuene; EMO=emodin; KA=kojic acid; TryOH=tryptophol; BREF=brevianamide F; LOD=limit of detection; LOQ=limit of quantification. \*values below LOQ

had higher concentrations of AME and ATN, but the difference was not statistically significant. L-cysteine and citric acid were the most effective against AME, and ATN concentrations were reduced by L-cysteine and Echinacea extract.

Kojic acid in our study was found in all samples above LOD, and its levels in pretreated figs were the highest compared to other mycotoxins. However, many studies claim unlikely or no adverse effects of KA, as it acts as an antioxidant and is widely used in food and cosmetic industry (5).

EMO was detected in several samples above LOD, most notably in the control white fig variety and in the dark variety pretreated with ascorbic acid. EMO was first confirmed in chestnuts (28, 29) and afterwards in many other matrices (30, 31) but never in dried figs. This anthraquinone polyphenol can be produced by various species (plant and fungal), but in most food it occurs as the *Aspergillus* metabolite (30). From the toxicological point of view, EMO has been documented in a wide range of actions, from cytotoxic to laxative and antitumour (32). The Echinacea extract lowered its levels below LOD in both varieties, but there was no significant difference between the treatments.

To the best of our knowledge, this is the first study to report TryOH in dried figs, but concerning the producing microbiota, this finding is not a surprise. TryOH is not strictly classified as a mycotoxin, as it has important functions in filamentous fungi and is one of the most prevalent fungal metabolites (30, 31, 33). In one study (34) it was also determined as one of the fermentation products in bread.

BREF was determined in all samples with concentrations above LOQ. This *Aspergillus* metabolite is quite common in a number of matrices (33), but it has not been reported in dried figs until now. Pretreatment was more effective in the white variety, whereas in the dark variety it only increased BREF levels. However, little is known about its toxicity.

## CONCLUSIONS

Our results suggest that L-cysteine is a potent antimycotoxigenic agent for fig treatment before drying, but the winner against aflatoxins is citric acid. L-cysteine potently reduced the *Alternaria*, EMO, and KA in the dark variety, while the Echinacea extract was the best against EMO. The dark fig variety Šaraguja was more prone to mycotoxin contamination, most likely because of thinner skin with many more cracks in it, which facilitates fungal infection.

Before these pretreatments are applied on the industrial scale, their efficiency should be tested on intentionally infected figs (by different mycotoxin-producing fungi) and on a larger scale to obtain relevant practical findings. Our study has just given a glimpse of potential issues with dry fig mycotoxin contamination and treatment possibilities.

**Table 3** Increase or reduction of mycotoxin levels by fig variety and pretreatment agent

Sample	Fig variety	AFB <sub>1</sub>	OTA	OT $\alpha$	AME	ATN	EMO	KA	TryOH	BREF
L-cysteine	white	25	100	45	13	77	13	265	38	65
L-cysteine	dark	7	100	150	40	65	100	83	218	130
Ascorbic acid	white	10	250	20	100	138	33	309	98	96
Ascorbic acid	dark	7	100	118	300	131	2250	94	338	253
Citric acid	white	75	100	20	13	250	33	256	145	104
Citric acid	dark	7	100	179	40	184	100	87	100	116
Chestnut extract	white	10	100	125	33	101	117	315	90	99
Chestnut extract	dark	50	100	171	200	149	100	154	1239	167
Echinacea extract	white	10	100	140	13	57	13	189	95	106
Echinacea extract	dark	67	100	92	200	88	100	97	100	133

The figures show the percentage of control levels, which are 100 %. Figures below 100 % show reduction and above 100 % increase. AFB<sub>1</sub>=aflatoxin B<sub>1</sub>; OTA=ochratoxin A; OT $\alpha$ =ochratoxin alpha; AME=alternariol methyl ether; ATN=altenuene; EMO=emodin; KA=kojic acid; TryOH=tryptophol; BREF=brevianamide F; all results are mean values of three measurements

#### REFERENCES

- Heperkan D, Moretti A, Dikmen CD, Logrieco AF. Toxigenic fungi and mycotoxin associated with figs in the Mediterranean area. *Phytopathol Mediterr* 2012;51:119-30. doi: 10.14601/Phytopathol\_Mediterr-9467
- Karbancioğlu-Güler F, Heperkan D. Natural occurrence of ochratoxin A in dried figs. *Anal Chim Acta* 2008;617:32-6. doi: 10.1016/j.aca.2008.01.009
- Trucksess MW, Scott PM. Mycotoxins in botanicals and dried fruits: A review. *Food Addit Contam A* 2008;25:181-92. doi: 10.1080/02652030701567459
- Oktay Basegmez HI, Heperkan D. Aflatoxin, cyclopiazonic acid and  $\beta$ -nitropropionic acid production by *Aspergillus* section *Flavi* from dried figs grown in Turkey. *Qual Assur Saf Crop* 2015;7:477-85. doi: 10.3920/QAS2013.0320
- Steiner W, Brunschweiler K, Lembacher E, Schneider R. Aflatoxine B<sub>1</sub> und G<sub>1</sub>, cyclopiazonsäure, kojisaure und ochratoxin A in trockenfeigen mit BGY-fluoreszenz [Aflatoxins B<sub>1</sub> and G<sub>1</sub>, cyclopiazonic acid, kojic acid and ochratoxin A in dry figs with BGY fluorescence, in German]. *Mitt Gebiete Lebens Hyg* 1993;84:523-36.
- Karbancioğlu-Güler F, Heperkan, D. Natural occurrence of fumonisin B<sub>1</sub> in dried figs as an unexpected hazard. *Food Chem Toxicol* 2009;47:289-92. doi: 10.1016/j.fct.2008.11.003
- Azaiez I, Font G, Mañes J, Fernández-Franzón M. Survey of mycotoxins in dates and dried fruits from Tunisian and Spanish markets. *Food Control* 2015;51:340-6. doi: 10.1016/j.foodcont.2014.11.033
- Karaca H, Nas S. Aflatoxins, patulin and ergosterol contents of dried figs in Turkey. *Food Addit Contam* 2006;23:502-8. doi: 10.1080/02652030600550739
- Aziz NH, Moussa LAA. Influence of gamma-radiation on mycotoxin producing moulds and mycotoxins in fruits. *Food Control* 2002;13:281-8. doi: 10.1016/S0956-7135(02)00028-2
- López P, Venema D, de Rijk T, de Kok A, Scholten JM, Mol HGJ, de Nijs M. Occurrence of *Alternaria* toxins in food products in The Netherlands. *Food Control* 2016;60:196-204. doi: 10.1016/j.foodcont.2015.07.032
- Alomari A-H, Bahkali AH, Mongy A-AA, Al-Sum BA, Elgorban AM. Quantitative analysis of mycotoxins produced by three fungi in dry fruits in Saudi Arabia using high-performance liquid chromatography. *J Pure Appl Microbio* 2014;8:577-81.
- Latinović N, Rasišek S, Latinović J. First report of *Alternaria alternata* causing fruit rot on fig (*Ficus carica*) in Montenegro. *Plant Dis* 2014;98:424. doi: 10.1094/PDIS-07-13-0770-PDN
- Şenyuva HZ, Gilbert J, Öztürkoğlu Ş. Rapid analysis of fungal cultures and dried figs for secondary metabolites by LC/TOF-MS. *Anal Chim Acta* 2008;617:97-106. doi: 10.1016/j.aca.2008.01.019
- Molnar-Perl I, Friedman M. Inhibition of browning by sulfur amino acids. 3. Apples and potatoes. *J Agr Food Chem* 1990;38:1652-6. doi: 10.1021/jf00098a006
- Mujić I, Bavcon Kralj M, Jokić S, Jarni K, Tjaša J, Prgomet Ž. Changes in aromatic profile of fresh and dried fig - the role of pre-treatments in drying process. *Int J Food Sci Tech* 2012;47:2282-8. doi: 10.1111/j.1365-2621.2012.03099.x
- Hiranvarachat B, Devahastin S, Chiewchan N. Effects of acid pretreatments on some physicochemical properties of carrot undergoing hot air drying. *Food Bioprod Process* 2011;89:116-27. doi: 10.1016/j.fbp.2010.03.010
- Villalobos MC, Serradilla MJ, Martín A, Pereira C. Lopez-Corrales M, Cordoba MG. Evaluation of different drying systems as an alternative to sun drying for figs (*Ficus carica* L.). *Innov Food Sci Emerg* 2016;36:156-65. doi: 10.1016/j.ifset.2016.06.006
- Barreira JCM, Ferreira ICFR, Oliveira MBPP, Pereira JA. Antioxidant activities of the extracts from chestnut flower, leaf, skins and fruit. *Food Chem* 2008;107:1106-13. doi: 10.1016/j.foodchem.2007.09.030
- Lewicki PP. Design of hot air drying for better foods. *Trends Food Sci Tech* 2006;17:153-63. doi: 10.1016/j.tifs.2005.10.012
- Sapkota K, Park SE, Kim JE, Kim S, Choi HS, Chun HS, Kim SJ. Antioxidant and antimelanogenic properties of chestnut flower extract. *Biosci Biotech Bioch* 2010;74:1527-33. doi: 10.1271/bbb.100058
- Reverberi M, Zjalic S, Ricelli A, Punelli F, Camera E, Fabbri C, Picardo M, Fanelli C, Fabbri AA. Modulation of antioxidant defense in *Aspergillus parasiticus* is involved in aflatoxin biosynthesis: a role for the *ApyapA* gene. *Eukaryot Cell* 2008;7:988-1000. doi: 10.1128/EC.00228-07

22. Özay G, Aran N, Pala M. Influence of harvesting and drying techniques on microflora and mycotoxin contamination of figs. *Die Nahrung* 1995;39:156-65. doi: 10.1002/food.19950390209
23. Karaca H, Nas S. Combined effect of pH and heat treatment on degradation of aflatoxins in dried figs. *J Food Process Pres* 2009;33(Suppl 1):329-39. doi: 10.1111/j.1745-4549.2008.00344.x
24. Malachova A, Sulyok M, Beltrán E, Berthiller F, Krska R. Optimization and validation of a quantitative liquid chromatography-tandem mass spectrometric method covering 295 bacterial and fungal metabolites including all regulated mycotoxins in four model food matrices. *J Chromatogr A* 2014;1362:145-56. doi: 10.1016/j.chroma.2014.08.037
25. Klapac T, Šarkanj B, Banjari I, Strelec I. Urinary ochratoxin A and ochratoxin alpha in pregnant women. *Food Chem Toxicol* 2012;50:4487-92. doi: 10.1016/j.fct.2012.09.030
26. Commission Regulation (EC) No 1881/2006 of 19 December 2006 setting maximum levels for certain contaminants in foodstuffs [displayed 24 October 2018]. Available at <https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32006R1881&from=EN>
27. Battilani P, Toscano P, Van der Fels-Klerx HJ, Moretti A, Camardo Leggieri M, Brera C, Rortais A, Goumperis T, Robinson T. Aflatoxin B<sub>1</sub> contamination in maize in Europe increases due to climate change. *Sci Rep-UK* 2016;6:24328. doi: 10.1038/srep24328
28. Ezekiel CN, Sulyok M, Warth B, Odebo AC, Krska R. Natural occurrence of mycotoxins in peanut cake from Nigeria. *Food Control* 2012;27:338-42. doi: 10.1016/j.foodcont.2012.04.010
29. Overy DP, Seifert KA, Savard ME, Frisvad JC. Spoilage fungi and their mycotoxins in commercially marketed chestnuts. *Int J Food Microbiol* 2003;88:69-77. doi: 10.1016/S0168-1605(03)00086-2
30. Abdallah MF, Krska R, Sulyok M. Occurrence of ochratoxins, fumonisin B<sub>2</sub>, aflatoxins (B<sub>1</sub> and B<sub>2</sub>), and other secondary fungal metabolites in dried date palm fruits from Egypt: A mini-survey. *J Food Sci* 2018;83:559-64. doi: 10.1111/1750-3841.14046
31. Yunus AW, Sulyok M, Böhm J. Mycotoxin cocktail in the samples of oilseed cake from early maturing cotton varieties associated with cattle feeding problems. *Toxins* 2015;7:2188-97. doi: 10.3390/toxins7062188
32. Liu X, Liu Y, Qu Y, Cheng M, Xiao H. Metabolomic profiling of emodin-induced cytotoxicity in human liver cells and mechanistic study. *Toxicol Res-UK* 2015;4:948-55. doi: 10.1039/C4TX00246F
33. Kosalec I, Šafranić A, Pepeljnjak S, Bačun-Družina V, Ramić S, Kopjar N. Genotoxicity of tryptophol in a battery of short-term assays on human white blood cells *in vitro*. *Basic Clin Pharmacol* 2008;102:443-52. doi: 10.1111/j.1742-7843.2007.00204.x
34. Jiang D, Peterson DG. Identification of bitter compounds in whole wheat bread. *Food Chem* 2013;141:1345-53. doi: 10.1016/j.foodchem.2013.03.021
35. Narasaiah KV, Sashidhar RB, Subramanyam C. Biochemical analysis of oxidative stress in the production of aflatoxin and its precursor intermediates. *Mycopathologia* 2006;162:179-89. doi: 10.1007/s11046-006-0052-7

### Utjecaj predtretmana sušenja na pojavnost i količinu mikotoksina u suhim smokvama

Za određivanje pojavnosti mikotoksina u osušenim autohtonim sortama bijelih i tamnih smokava petrovača bijela i šaraguja korištena je metoda LC-MS/MS "razrijedi i mjeri", kojom je moguće analizirati 295 gljivičnih i bakterijskih sekundarnih metabolita. Smokve su sušene u pilot-pogonu uz korištenje različitih predtretmana za održavanje sušenog voća: uranjanje u 0,5 %-tnu otopinu limunske kiseline, 0,5 %-tnu otopinu askorbinske kiseline, 0,3 %-tnu otopinu L-cisteina, 0,2 %-tnu otopinu ekstrakta kestena i 0,15 %-tnu otopinu ekstrakta Echinaceae. Ovim se istraživanjem otkrilo devet različitih metabolita u koncentracijama iznad njihove granice detekcije: aflatoksin B<sub>1</sub>, okratoksin A, okratoksin α, kojična kiselina, emodin, altenuen, alternariolmetileter, brevianamid F i triptofol. Sveukupno, najbolji antimikotoksigeni učinak postignut je predtretmanom s L-cisteinom (prosječno smanjenje od 15 %), a tretman s askorbinskom kiselinom pokazao je najveći učinak na indukciju mikotoksina (povećanje od 158 %). Svi predtretmani smanjili su koncentracije aflatoksina, glavnog kontaminanta smokava.

KLJUČNE RIJEČI: antimikotoksigeni učinak; LC-MS/MS; multimikotoksinska analiza; sušenje