



# Applied Mathematics and Nonlinear Sciences

<https://www.sciendo.com>

## New Complex Hyperbolic Structures to the Lonngren-Wave Equation by Using Sine-Gordon Expansion Method

Haci Mehmet Baskonus<sup>1†</sup>, Hasan Bulut<sup>2</sup>, Tukur Abdulkadir Sulaiman<sup>3</sup>

<sup>1</sup>Department of Mathematics and Science Education, Harran University, Sanliurfa, Turkey

<sup>2,3</sup>Department of Mathematics, Firat University, Elazig, Turkey

<sup>3</sup>Department Mathematics, Federal University Dutse, Jigawa, Nigeria

### Submission Info

Communicated by Juan Luis García Guirao

Received February 16th 2019

Accepted April 29th 2019

Available online June 25th 2019

### Abstract

In this paper, a powerful sine-Gordon expansion method (SGEM) with aid of a computational program is used in constructing a new hyperbolic function solutions to one of the popular nonlinear evolution equations that arises in the field of mathematical physics, namely; longren-wave equation. We also give the 3D and 2D graphics of all the obtained solutions which are explaining new properties of model considered in this paper. Finally, we submit a comprehensive conclusion at the end of this paper.

**Keywords:** The sine-Gordon expansion method; the longren-wave equation; hyperbolic function solutions.

**PACS codes:** 02.60.Lj; 02.70.Wz; 02.90.+p; 03.65.-w.

## 1 Introduction

Over some decades, the field of nonlinear evolution equations (NEEs) has attracted the attention of many researchers. NEEs are broadly used to describe problems in science, engineering and mathematical physics such as fluid dynamics, plasma physics, hydro magnetic waves, optic fibers, solid state physics and many others. NEEs can also be used to describe the propagation of a nonlinear dispersive waves in inhomogeneous media [1,2]. It has become an important bottom-line to find the analytical solutions to these types of equations. Several methods for finding the solutions of various NEEs have been proposed and/or improved by many scholars [3-71].

The aim of this paper was to apply the sine-Gordon expansion method (SGEM) to find a new solutions to the Lonngren-wave equation [14].

$$(u_{xx} - \alpha u + \beta u^2)_{tt} + u_{xx} = 0, \quad (1)$$

<sup>†</sup>Corresponding author E-mail: hmbaskonus@gmail.com

where  $\alpha$  and  $\beta$  are real constants. The equation describes the electric signals in telegraph lines on the basis of the tunnel diode [15,16]. The Lonngren-wave equation was used as an example by Akcagil and Aydemir [14] to show the existence of strong connection between the  $(\frac{G'}{G})$ -expansion method and the modified extended tanh method.

SGEM is a method for solving different nonlinear partial differential equations that is developed based on wave transformation and the sine-Gordon expansion method [17]. A new hyperbolic function solutions to the Davey-Stewartson equation with power-law nonlinearity was obtained in [18] by using SGEM. With the aid of symbolic computation, a new transformation was developed using the general sine-Gordon travelling wave reduction equation and a generalized transformation to obtain the solutions of various types of nonlinear differential equations [19]. A considerable investigation has been implemented by Yan [20] to sine-Gordon-type equations where the equations are systematically solved by using the Jacobi elliptic function expansion method.

The remaining parts of this paper are organized as follows: In Section 2, we discuss the general facts of the SGEM. In Section 3, we apply the SGEM to the Lonngren-wave equation given in Eq. (1). Section 4 is about results, discussion and some remarks. We finally, give the conclusion of this paper in Section 5.

## 2 General Facts of the SGEM

In this section we discuss the general facts of SGEM.

Consider the following sine-Gordon equation [17,21,22]:

$$u_{xx} - u_{tt} = m^2 \sin(u), \quad (2)$$

where  $u = u(x, t)$  and  $m$  is a real constant.

Applying the wave transformation  $u = u(x, t) = U(\xi)$ ,  $\xi = \mu(x - ct)$  to Eq. (2), yields the following nonlinear ordinary differential equation (NODE):

$$U'' = \frac{m^2}{\mu^2(1-c^2)} \sin(U), \quad (3)$$

where  $U = U(\xi)$ ,  $\xi$  is the amplitude of the travelling wave and  $c$  is the velocity of the travelling wave. Reconsidering Eq. (3), we can write its full simplification as:

$$\left[ \left( \frac{U}{2} \right)' \right]^2 = \frac{m^2}{\mu^2(1-c^2)} \sin^2 \left( \frac{U}{2} \right) + K, \quad (4)$$

where  $K$  is the integration constant.

Substituting  $K = 0$ ,  $w(\xi) = \frac{U}{2}$  and  $a^2 = \frac{m^2}{\mu^2(1-c^2)}$  in Eq. (4), gives:

$$w' = a \sin(w), \quad (5)$$

Putting  $a = 1$  in Eq. (5), we have:

$$w' = \sin(w), \quad (6)$$

Equation (6) is variables separable equation, we obtain the following two significant equations from solving it:

$$\sin(w) = \sin(w(\xi)) = \frac{2pe^\xi}{p^2e^{2\xi} + 1} \Bigg|_{p=1} = \operatorname{sech}(\xi), \quad (7)$$

$$\cos(w) = \cos(w(\xi)) = \frac{p^2 e^{2\xi} - 1}{p^2 e^{2\xi} + 1} \Big|_{p=1} = \tanh(\xi), \tag{8}$$

where  $p$  is the integral constant.

For the solution of the following nonlinear partial differential equation;

$$P(u, u_x, u_t, u_{xx}, u_{tt}, u_{xt} \dots), \tag{9}$$

we consider,

$$U(\xi) = \sum_{i=1}^n \tanh^{i-1}(\xi) [B_i \operatorname{sech}(\xi) + A_i \tanh(\xi)] + A_0, \tag{10}$$

Equation (10) can be rewritten according to Eqs. (7) and (8) as follows:

$$U(w) = \sum_{i=1}^n \cos^{i-1}(w) [B_i \sin(w) + A_i \cos(w)] + A_0. \tag{11}$$

We determine the value  $n$  under the terms of NODE by the balance principle. Letting the coefficients of  $\sin^i(w)\cos^j(w)$  to be all zero, yields a system of equations. Solving this system by using Wolfram Mathematica 9 gives the values of  $A_i$ ,  $B_i$ ,  $\mu$  and  $c$ . Finally, substituting the values of  $A_i$ ,  $B_i$ ,  $\mu$  and  $c$  in Eq. (10), we obtain the new travelling wave solutions to Eq. (9)

### 3 Applications

Consider the Lonngren-wave equation given in Eq. (1);

Applying the transformation  $u = u(x, t) = U(\xi)$ ,  $\xi = \mu(x - ct)$  to Eq. (1), we have:

$$c^2 \mu^2 U'' + (1 - c^2 \alpha)U + c^2 \beta U^2 = 0 \tag{12}$$

We obtain  $n = 2$ , by applying the balance principle on Eq. (12). With  $n = 2$  and using Eq. (11), yields:

$$U(w) = B_1 \sin(w) + A_1 \cos(w) + B_2 \cos(w) \sin(w) + A_2 \cos^2(w) + A_0. \tag{13}$$

Differentiating Eq. (13) twice, gives:

$$U''(w) = B_1 \cos^2(w) \sin(w) - B_1 \sin^3(w) - 2A_1 \sin^2(w) \cos(w) + B_2 \cos^3(w) \sin(w) - 5B_2 \sin^3(w) \cos(w) - 4A_2 \cos^2(w) \sin^w(w) + 2A_2 \sin^4(w), \tag{14}$$

Putting Eqs. (13) and (14) in Eq. (12), yields;

$$\begin{aligned} & A_0 - c^2 \alpha A_0 + c^2 \beta A_0^2 + A_1 \cos(w) - c^2 \alpha A_1 \cos(w) - 2c^2 \mu^2 A_1 \sin^2(w) \cos(w) + 2c^2 \beta A_0 A_1 \cos(w) + \\ & c^2 \beta A_1^2 \cos^2(w) + A_2 \cos^2(w) - c^2 \alpha A_2 \cos^2(w) \\ & - 4c^2 \mu^2 A_2 \cos^2(w) \sin^2(w) + 2c^2 \mu^2 A_2 \sin^4(w) + 2c^2 \beta A_0 A_2 \cos^2(w) + \\ & 2c^2 \beta A_1 A_2 \cos^3(w) + c^2 \beta A_2^2 \cos^4(w) + B_1 \sin(w) - c^2 \alpha B_1 \sin(w) + \\ & c^2 \mu^2 B_1 \cos^2(w) \sin(w) - c^2 \mu^2 B_1 \sin^3(w) + 2c^2 \beta A_0 B_1 \sin(w) + \\ & 2c^2 \beta A_1 B_1 \cos(w) \sin(w) + 2c^2 \beta A_2 B_1 \cos^2(w) \sin(w) + c^2 \beta B_1^2 \sin^2(w) \\ & + B_2 \cos(w) \sin(w) - c^2 \alpha B_2 \cos(w) \sin(w) + c^2 \mu^2 B_2 \cos^3(w) \sin(w) - \\ & 5c^2 \mu^2 B_2 \sin^3(w) \cos(w) + 2c^2 \beta A_0 B_2 \cos(w) \sin(w) + 2c^2 \beta A_1 B_2 \cos^2 \sin(w) \\ & + 2c^2 \beta A_2 B_2 \cos^3 \sin(w) + 2c^2 \beta B_1 B_2 \sin^2(w) \cos(w) + c^2 \beta B_2^2 \cos^2(w) \sin^2(w) = 0. \end{aligned}$$

We collect a set of algebraic equations by equating each summation of the coefficients of the trigonometric terms of the same power to zero in the abovementioned equation. We solve the set of generated to obtained the values of the coefficients. To get the new solitary solutions,  $u(x, t)$  to Eq. (1), we substitute in each case the obtained results of the coefficients into Eq. (10) along with  $n = 2$ .

Case-1: When we consider following coefficients:

$$A_0 = \frac{2\mu^2}{\beta}, A_1 = 0, B_1 = 0, A_2 = -\frac{6\mu^2}{\beta}, B_2 = 0, \alpha = \frac{1}{c^2} - 4\mu^2,$$

these produce new dark solution as:

$$u_1(x, t) = \frac{2\mu^2}{\beta} (1 - 3 \tanh[\mu(x - ct)]^2). \quad (15)$$




Fig. 1 The 3D and 2D surfaces of Eq. (15).

Case-2: If it is taken as

$$A_0 = \frac{6\mu^2}{\beta}, A_1 = 0, B_1 = 0, A_2 = -\frac{6\mu^2}{\beta}, B_2 = 0, \alpha = \frac{1}{c^2} + 4\mu^2,$$

they produce a new singular solution as:

$$u_2(x, t) = \frac{6\mu^2}{\beta} \operatorname{sech}[\mu(x - ct)]^2. \quad (16)$$

Case-3: When we take

$$A_0 = \frac{2\mu^2}{\beta}, A_1 = 0, B_1 = 0, A_2 = -\frac{3\mu^2}{\beta}, B_2 = \frac{3i\mu^2}{\beta}, \alpha = \frac{1}{c^2} - \mu^2,$$

they give mixed complex singular solution as:

$$u_3(x, t) = \frac{\mu^2}{\beta} \left( -1 + \frac{3i}{i + \sinh(\mu(x - ct))} \right). \quad (17)$$




Fig. 2 The 3D and 2D surfaces of Eq. (16).




Fig. 3 The 3D and 2D surfaces of Eq. (17).

Case-4:

$$A_0 = \frac{3(c^2\alpha - 1)}{c^2\beta}, A_1 = 0, B_1 = 0, A_2 = \frac{3 - 3c^2\alpha}{c^2\beta}, B_2 = \frac{3i(c^2\alpha - 1)}{c^2\beta}, \mu = -\frac{\sqrt{c^2\alpha - 1}}{c},$$

give mixed complex rational solution as:

$$u_4(x, t) = \frac{3i(c^2\alpha - 1)}{c^2\beta \left( i - \sinh\left[\left(\frac{\sqrt{c^2\alpha - 1}}{c}\right)(x - ct)\right] \right)}. \tag{18}$$




Fig. 4 The 3D and 2D surfaces of Eq. (18).

Case-5:

$$A_0 = \frac{2 - 2c^2\alpha}{c^2\beta}, A_1 = 0, B_1 = 0, A_2 = \frac{3(c^2\alpha - 1)}{c^2\beta}, B_2 = -\frac{3i(c^2\alpha - 1)}{c^2\beta}, \mu = -\frac{i\sqrt{c^2\alpha - 1}}{c},$$

which produces the following trigonometric travelling wave solution as:

$$u_5(x, t) = \frac{1}{c^2\beta} \left( c^2\alpha - 1 + \frac{3(c^2\alpha - 1)}{-1 + \sin[(x - ct)\frac{\sqrt{c^2\alpha - 1}}{c}]} \right). \tag{19}$$




Fig. 5 The 3D and 2D surfaces of Eq. (19).

Case-6:

$$A_0 = \frac{3\mu^2}{\beta}, A_1 = 0, B_1 = 0, A_2 = -\frac{3\mu^2}{\beta}, B_2 = \frac{3i\mu^2}{\beta}, c = -\frac{1}{\sqrt{\alpha - \mu^2}},$$

which introduces the following complex mixed solution as:

$$u_7(x, t) = \frac{3\mu^2}{\beta} \operatorname{sech} \left[ \mu \left( x + \frac{t}{\sqrt{\alpha - \mu^2}} \right) \right] \left( \operatorname{sech} \left[ \mu \left( x + \frac{t}{\sqrt{\alpha - \mu^2}} \right) \right] + i \operatorname{tanh} \left[ \mu \left( x + \frac{t}{\sqrt{\alpha - \mu^2}} \right) \right] \right). \tag{20}$$




Fig. 6 The 3D and 2D surfaces of Eq. (20).

#### 4 Results and Discussion

The powerful SGEM as one of the prominent methods for obtaining the some new travelling wave solutions to the nonlinear partial differential equations has been used in this paper. This method is based on both important properties of the sine-Gordon equation such as Eqs. (7) and Eq.(8). The SGEM includes trigonometric functions, which will be used later for obtaining novel solutions in Eq.(11). Many new solutions can be obtained by using the properties of these trigonometric functions. This is one of the main properties of SGEM. Therefore, it gives many coefficients to the considered model such as complex, exponential and trigonometric.

## 5 Conclusions

In this manuscript, by selecting of some of them, we have obtained the same solution, Eq.(15); moreover, we have found some entirely new complex, exponential, dark and hyperbolic solutions to the model considered when we compared the solutions obtained with the help of  $\exp$  the  $(G'/G)$ -expansion method and the modified extended tanh method used in [14]. These solutions are new physical properties of model equation Eq.(1.1). The effectiveness and the simplicity of the method show that its powerful and reliable mathematical tool that can be applied in solving various NEEs. For computational calculations, we have used the packet programs for drawing graphical surfaces in this paper. To the best of our knowledge, the application of SGEM to the Lonngren-wave equation has not been submitted to the literature beforehand.

## References

- [1] P.J. Olver, Evolution equations possessing infinitely many symmetries, *J. Math. Physics*, 18 (1977) 1212-1217.
- [2] A.M. Wazwaz, New solitons and kinks solutions to the Sharma-Mukhtar equation, *Appl. Math. Computation* 188 (2007) 1205-1213.
- [3] C. Cattani, A review on Harmonic Wavelets and their fractional extension, *Journal of Advanced Eng. Comp.*, 2018, 2(4), 224-238.
- [4] D. Lu, A.R. Seadawy, M.M. A. Khater, Bifurcations of new multi soliton solutions of the van der Waals normal form for fluidized granular matter via six different methods, *Results in Physics*, 2017, 7, 2028-2035.
- [5] A. Ciancio, A. Quartarone, A Hybrid Model For Tumor-Immune Competition, *U.P.B. Sci. Bull. Series A*, 2013, 75(4), 125-136.
- [6] M. Arshad, A.R. Seadawy, D.Lu, Bright-dark solitary wave solutions of generalized higher-order nonlinear Schrödinger equation and its applications in optics, *J Electromag Waves and Appl*, 2017, 31(16), 1711-1722.
- [7] I. Cilingir, H. Demir, New Algorithm for the Lid-driven Cavity Flow Problem with Boussinesq-Stokes Suspension, *Karaelmas J. Science and Eng*, 8(2), (2018), 462-472.
- [8] T. Caraballo, M. Herrera-Cobos, P. Marín-Rubio, An iterative method for non-autonomous nonlocal reaction-diffusion equations, *Appl. Math. Nonlinear Sciences*, 2(1) (2017) 73-82.
- [9] Baskonus, H.M.; Bulut, H.; Emir, D.G. Regarding New Complex Analytical Solutions for the Nonlinear Partial Vakhnenko-Parkes Differential Equation via Bernoulli Sub-Equation Function Method, *Math. Letters*, 2015, 1(1), 1-9.
- [10] M. Askin, Effect of the Transition Metal Elements on the Relaxation Times in the Agar Solutions, *Asian Journal of Chemistry*, 19(4), (2007), 3191-3196.
- [11] C. Cattani, A. Ciancio, On the fractal distribution of primes and prime-indexed primes by the binary image analysis, *Physica A*, 2016, 460, 222-229
- [12] A. Ciancio, H.M. Baskonus, T.A. Sulaiman, H. Bulut, New structural dynamics of isolated waves via the coupled nonlinear Maccari's system with complex structure, *Indian J of Physics*, 92(10), (2018), 1281-1290.
- [13] S.M. El-Shaboury, M.K. Ammar, W.M. Yousef, Analytical solutions of the relative orbital motion in unperturbed and in J- perturbed elliptic orbits, *Appl. Math. Nonlinear Sciences*, 2(2), (2017), 403-414
- [14] S. Akcagil, T.T. Aydemir, Comparison Between the  $(G'/G)$ -Expansion Method and the Modified Extended tanh Method, *Open Physics* 14, 88 (2016)
- [15] T.A. Sulaiman, A.I. Aliyu, A. Yusuf, A Solution of Telegraph Equation by Natural Decomposition Method *International Conference on Mathematics and Mathematics Education (ICMME-2016)*, Elazig/Turkey, 12-14 May 2016.
- [16] K.E. Lonngren, H.C.S. Hsuan, W.F. Ames, On the Soliton, Invariant and Shock Solutions of a Fourth-Order Nonlinear Equation, *Journal of Mathematical Analysis and Applications*, 52, 538 (1975).
- [17] M I Rabinovich, D I Trubetskov, Introduction in Theory of Waves, Nauka, Moscow, (1984).
- [18] C. Yan, A Simple Transformation for Nonlinear Waves, *Physics Letters A* 22(4), 77 (1996)
- [19] H.M. Baskonus, New Acoustic Wave Behaviors to the Davey-Stewartson Equation with Power Nonlinearity Arising in Fluid Dynamics, *Nonlinear Dynamics* 86(1), 177 (2016)
- [20] Z. Yan, A New Sine-Gordon Equation Expansion Algorithm to Investigate Some Special Nonlinear Differential Equations, *Academia Sinica* 23, 300 (2004)
- [21] S. Liu, Z. Fu and S. Liu, Exact Solutions to Sine-Gordon-Type Equations, *Physics Letters A*, 351 (2006), 59-63.
- [22] Z. Yan and H. Zhang, New Explicit and exact Travelling Wave Solutions for a System of Variant Boussinesq equations in Mathematical Physics, *Physics Letters A*, 252 (1999), 291-296.
- [23] Y. Zhen-Ya, Z. Hong-ong and F. En-Gui, New Explicit and Travelling Wave Solutions for a Class of Nonlinear Evolution Equations, *Acta Physica Sinica*, 48(1) (1999), 1-5.
- [24] Araci, S.; Ozer, O. Extended q-Dedekind-type Daehee- Changhee sums associated with extended q-Euler polynomials,

- Adv. Differen. Eq, 2015, 2015(1), 272-276.
- [25] H.M. Baskonus and H. Bulut, Analytical Studies on the (1+1)-dimensional Nonlinear Dispersive Modified Benjamin-Bona-Mahony Equation Defined by Seismic Sea Waves, *Waves in Random and Complex Media*, 25(4), 1-13, 2015.
- [26] Aslan, I. Exact and explicit solutions to nonlinear evolution equations using the division theorem, *Appl. Math. and Comp.* 2011, 217, 8134-8139.
- [27] Sulaiman, T.A.; Bulut, H.; Yokus, A.; Baskonus, H.M. On the exact and numerical solutions to the coupled Boussinesq equation arising in ocean engineering, *Indian J of Physics* 2018, 1-10.
- [28] V.B. Awati, M. Jyoti, Homotopy analysis method for the solution of lubrication of a long porous slider, *Appl. Math. Nonlinear Sciences*, 1(2) (2016) 507-516
- [29] Cattani. C.; Sulaiman, T.A.; Baskonus, H.M.; Bulut, H. On the soliton solutions to the Nizhnik-Novikov-Veselov and the Drinfel'd-Sokolov systems, *Opt. and Quan. Elect.* 2018, 50(138), 1-11.
- [30] O. Ozer; Pekin A, An Algorithm For Explicit Form of Fundamental Units of Certain Real Quadratic Fields and Period Eight, *European J. Pure and Applied Mathematics*, 2015, 8(3), 343-356.
- [31] Cattani. C.; Sulaiman, T.A.; Baskonus, H.M.; Bulut, H. Solitons in an inhomogeneous Murnaghan's rod, *Europ Phys J Plus*, 2018, 133(228), 1-11.
- [32] Baskonus, H.M. New complex and hyperbolic function solutions to the generalized double combined Sinh-Cosh-Gordon equation, *AIP Conf. Proc.* 2017, 1798(020018), 1-10.
- [33] Seadawy, A. Three-dimensional nonlinear modified Zakharov-Kuznetsov equation of ion-acoustic waves in a magnetized plasma, *Comput. Math. Appl.*, 2016, 71, 201-206.
- [34] Unlukal, C.; Penel, M.; Penel, B. Risk Assessment with Failure Mode and Effect Analysis and Gray Relational Analysis Method in Plastic Enjection Proses, *ITM Web of Conf.*, 2018, 22(01023), 1-10
- [35] Sulaiman, T.A.; Yokus, A.; Gulluoglu, N.; Baskonus, H.M.; Bulut, H. Regarding the Numerical and Stability Analysis of the Sharma-Tosso-Olver Equation, *ITM Web of Conf.*, 2018, 22(01036), 1-9.
- [36] M.Dewasurendra, K.Vajravelu, On the Method of Inverse Mapping for Solutions of Coupled Systems of Nonlinear Differential Equations Arising in Nanofluid Flow, Heat and Mass Transfer, *Appl. Math. Nonlinear Sciences* 3(1) (2018) 1-14
- [37] H. Bulut, Classification of exact solutions for generalized form of K(m,n) equation, *Abstract and Applied Analysis*, 2013 (2013), 1-11.
- [38] H. M. Baskonus, H.Bulut, F. B. M Belgacem, Analytical Solutions for Nonlinear Long-Short Wave Interaction Systems with Highly Complex Structure, *J. Comput. Appl. Math.*, 312, 257 (2017)
- [39] Ilhan, O.A.; Sulaiman, T.A.; Bulut, H.; Baskonus, H.M. On the new wave Solutions to a Nonlinear Model Arising in Plasma Physics, *Europ Phys J Plus*, 2018, 133(27), 1-6.
- [40] Ozer, O. A Note On Structure of Certain Real Quadratic Number Fields, *Iranian Journal of Science and Technology*, 2017, 41(3), 759-769.
- [41] Baskonus, H.M.; Sulaiman, T.A.; Bulut, H.; Akturk, T. Investigations of dark, bright, combined dark-bright optical and other soliton solutions in the complex cubic nonlinear Schrödinger equation with  $-$ potential, Superlattice and Microstructures, 2018, 115, 19-29.
- [42] S. Duran; M. Askin, T.A. Sulaiman, New soliton properties to the ill-posed Boussinesq equation arising in nonlinear physical science, *An Int.J. Optimization and Control: Theories and Applications*, 2017, 7(3), 240-247.
- [43] Baskonus, H.M.; Askin, M., Travelling Wave Simulations to the Modified Zakharov-Kuznetsov Model Arising In Plasma Physics, 6th International Youth Science Forum "LITTERIS ET ARTIBUS", Computer Science and Engineering, Lviv, Ukraine, 24-26 November 2016.
- [44] C.M.Khalique, I.E.Mhlanga, Travelling waves and conservation laws of a (2+1)-dimensional coupling system with Korteweg-de Vries equation, *Appl. Math. Nonlinear Sciences* 3(1) (2018) 241-254
- [45] Baskonus, H.M.; Koc, D.A.; Bulut, H. New travelling wave prototypes to the nonlinear Zakharov-Kuznetsov equation with power law nonlinearity, *Nonlin. Sci. Letters A: Math., Phys. and Mech.* 2016, 7(2), 67-76.
- [46] W.X. Ma; J.Li; C.M. Khalique, A Study on Lump Solutions to a Generalized Hirota-Satsuma-Ito Equation in (2+1)-Dimensions, *Complexity*, 2018, 2018(9059858), 1-7.
- [47] Bulut, H.; Atas, S.S.; Baskonus, H.M. Some novel exponential function structures to the Cahn-Allen equation, *Cogent Physics*, 2016, 3(1240886), 1-8.
- [48] Yokus, A.; Sulaiman, T.A.; Gulluoglu, M.T.; Bulut, H. Stability Analysis, Numerical and Exact Solutions of the (1+1)-Dimensional NDMBBM Equation, *ITM Web of Conf.*, 2018, 22(01064), 1-10.
- [49] Yokus, A.; Sulaiman, T. A.; Baskonus, H. M.; Atmaca, S. P. On the exact and numerical solutions to a nonlinear model arising in mathematical biology, *ITM Web of Conf.*, 2018, 22(01061), 1-10.
- [50] F.T. Akyildiz, K. Vajravelu, Galerkin-Chebyshev Pseudo Spectral Method and a Split Step New Approach for a Class of Two dimensional Semi-linear Parabolic Equations of Second Order, *Appl. Math. Nonlinear Sciences* 3(1) (2018) 255-264
- [51] H.M. Baskonus, H. Bulut, On the Complex Structures of Kundu-Eckhaus Equation via Improved Bernoulli Sub-Equation Function Method, *Waves in Random and Complex Media*, 25(4), 720 (2015).

- [52] I. Cilingir; H. Demir, Application of the Hybrid Differential Transform Method to the nonlinear equations, *Applied Mathematics*, 2012, 3(3), 1-10.
- [53] A.R. Seadawy, Ionic acoustic solitary wave solutions of two-dimensional nonlinear Kadomtsev-Petviashvili-Burgers equations in quantum plasma, *Mathematical Methods and Applied Sciences*, 2017, 40, 1598-1607.
- [54] H.M. Baskonus, H. Bulut, New Complex Hyperbolic Function Solutions for the (2+1)-Dimensional Dispersive Long Water-Wave System, *Mathematical and Computational Applications*, 21(2), 1 (2016),
- [55] O. Ozer, Determination of Fundamental Units of Real Quadratic Number Fields Related with Specific Continued Fraction Expansions. *Egyptian Computer Science Journal*, 2018, 42(2), 1-12.
- [56] Bulut, H.; Sulaiman, T.A., Baskonus, H.M., Yazgan, T., Novel Hyperbolic Behaviors to Some Important Models Arising in Quantum Science, *Opt. and Quan. Elect.*, 2017, 49(349), 1-16.
- [57] M. Rosa, M.L. Gandarias, Multiplier method and exact solutions for a density dependent reaction-diffusion equation, *Appl. Mathematics and Nonlinear Sciences* 1(2), 2016 311-320.
- [58] H. M. Baskonus, H. Bulut, Analytical Studies on the (1+1)-dimensional Nonlinear Dispersive Modified Benjamin-Bona-Mahony Equation Defined by Seismic Sea Waves, *Waves in Random and Complex Media*, 25(4), 576-586, 2015.
- [59] H. M. Baskonus, Complex Soliton Solutions to the Gilson-Pickering Model, *Axioms*, 8(1), 18, 2019.
- [60] H. Demir, Temporal differential transform and spatial finite difference methods for unsteady heat conduction equations with anisotropic diffusivity, *Gazi University Journal of Science*, 2014, 27(4), 1063-1076.
- [61] H. Bulut, Comparison between The Alternating Group Explicit Method and Adomian Decomposition Method for Solution of Coupled Viscous Burgers Equation, *Nonlinear Science Letters A*, 1(2), 161-172, (2010).
- [62] V.B. Awati, M. Jyoti, Homotopy analysis method for the solution of lubrication of a long porous slider, *Appl. Math. Nonlinear Sciences*, 1(2) (2016) 507-516
- [63] H.M. Baskonus, New complex and hyperbolic function solutions to the generalized double combined Sinh-Cosh-Gordon equation, *AIP Conf. Proc.* 1798(020018) (2017) 1-10.
- [64] A. Biswas, M.O. Al-Amr, H. Rezazadeh, M. Mirzazadeh, M. Eslami, Q. Zhou, S.P. Moshokoa, M. Belic, Resonant optical solitons with dual-power law nonlinearity and fractional temporal evolution, *Optik* 165 (2018) 233-239.
- [65] A.F. Qasim, M.O. Al-Amr, Approximate solution of the Kersten-Krasil'shchik coupled Kdv-MKdV system via reduced differential transform method, *Eurasian Journal of Science and Engineering* 4 (2) (2018) 1-9.
- [66] M.O. Al-Amr, Exact solutions of the generalized (2+1)-dimensional nonlinear evolution equations via the modified simple equation method, *Comput. Math. Appl.* 69 (5) (2015) 390-397.
- [67] A.J. Al-Sawoor, M.O. Al-Amr, A new modification of variational iteration method for solving reaction-diffusion system with fast reversible reaction, *J. Egyptian Math. Soc.* (2014), Vol. 22, No. 3, pp. 396-401.
- [68] A.J. Al-Sawoor, M.O. Al-Amr, Numerical solution of a reaction-diffusion system with fast reversible reaction by using Adomian's decomposition method and He's variational iteration method, *Al-Rafidain J. Comput. Sci. Math.* 9 (2) (2012) 243-257.
- [69] M.O. Al-Amr, S. El-Ganaini, New exact traveling wave solutions of the (4+1)-dimensional Fokas equation, *Comput. Math. Appl.* 74 (2017) 1274-1287.
- [70] M.O. Al-Amr, Exact solutions of a family of higher-dimensional space-time fractional KdV type equations, *Computer Science and Information Technology* 8 (6) (2018) 131-141.
- [71] A.J. Al-Sawoor, M.O. Al-Amr, Reduced differential transform method for the generalized Ito system, *Int. J. Enhanc. Res. Sci. Tech. Eng.* 2 (11) (2013) 135-145.