Applied Mathematics and Nonlinear Sciences

<https://www.sciendo.com>

Numerical Solution of Abel's Integral Equations using Hermite Wavelet

R. A. Mundewadi^{1†}, S. Kumbinarasaiah²

1 Department of Mathematics, M.E.S Degree College, Malleswaram, Bangalore, India,
E-mail: rkmundewadi@gmail.com

2 Department of Mathematics, Karnatak University, Dharwad, India,
E-mail: kumbinarasaiahs@gmail.com

Submission Info

Communicated by Juan Luis García Guirao
Received March 21st 2019
Accepted May 8th 2019
Available online June 28th 2019

Abstract

A numerical method is developed for solving the Abel's integral equations is presented. The method is based upon Hermite wavelet approximations. Hermite wavelet method is then utilized to reduce the Abel's integral equations into the solution of algebraic equations. Illustrative examples are included to demonstrate the validity, efficiency and applicability of the proposed technique. Algorithm provides high accuracy and compared with other existing methods.

Keywords: Abel's integral equations, Hermite wavelets, collocation method.

MSC codes: 65T60, 65R20, 97N40

1 Introduction

Wavelets theory is a new emerging tool in applied mathematical research area. It is applicable in various fields, such as, signal analysis for waveform representation and segmentations, time-frequency analysis and Harmonic analysis. Wavelets permit the accurate representation of a variety of functions and operators. Moreover, wavelets establish a connection with fast numerical algorithms [1, 2]. Since 1991 the various types of wavelet methods have been applied for the numerical solution of different kinds of integral equations [3]. Namely, the Haar wavelets method [3], Legendre wavelets method [4], Rationalized haar wavelet [5], Hermite cubic splines [6], Coifman wavelet scaling functions [7], CAS wavelets [8], Bernoulli wavelets [9], wavelet preconditioned techniques [25-28]. Some of the papers are found for solving Abel's integral equations using the wavelet based methods, such as Legendre wavelets [10] and Chebyshev wavelets [11].

Abel's integral equations have applications in various fields of science and engineering. Such as microscopy,

[†]Corresponding author.

Email address: rkmundewadi@gmail.com

seismology, semiconductors, scattering theory, heat conduction, metallurgy, fluid flow, chemical reactions, plasma diagnostics, X-ray radiography, physical electronics, nuclear physics [12-14].

In 1823, Abel, when generalizing the tautochrone problem derived the equation:

$$\int_0^x \frac{y(t)}{\sqrt{(x-t)}} dt = f(x), \quad (1)$$

where $f(x)$ is a known function and $y(x)$ is an unknown function to be determined. This equation is a particular case of a linear Volterra integral equation of the first kind. For solving eq. (1) different numerical based methods have been developed over the past few years, such as product integration methods [15], collocation method [16, 17], homotopy analysis transform method [18]. The generalized Abel's integral equations on a finite segment appeared for the first time in the paper of Zeilon [19]. There are several numerical methods for approximating the solution of singular integral equations is known. Baker [20] studied the numerical treatment of integral equations. A numerical solution of weakly singular volterra integral equations was introduced in [21]. Babolian and Salimi [22] discussed an operational matrix method based on block-pulse functions for singular integral equations. In this paper, we introduced the Hermite wavelets based numerical method for solving Abel's integral equations.

The article is organized as follows: In Section 2, formulation of Hermite wavelets and function approximation is presented. Section 3 is devoted the method of solution. In section 4, numerical results are demonstrated the accuracy of the proposed method by some of the illustrative examples. Lastly, the conclusion is given in section 5.

2 Properties of Hermite Wavelets

Wavelets constitute a family of functions constructed from dilation and translation of a single function called mother wavelet. When the dilation parameter a and translation parameter b varies continuously, we have the following family of continuous wavelets:

$$\psi_{a,b}(x) = |a|^{-1/2} \psi\left(\frac{x-b}{a}\right), \forall a, b \in \mathbb{R}, a \neq 0.$$

If we restrict the parameters a and b to discrete values as $a = a_0^{-k}, b = nb_0 a_0^{-k}, a_0 > 1, b_0 > 0$. We have the following family of discrete wavelets

$$\psi_{k,n}(x) = |a|^{1/2} \psi(a_0^k x - nb_0), \forall a, b \in \mathbb{R}, a \neq 0,$$

where $\psi_{k,n}$ form a wavelet basis for $L^2(\mathbb{R})$. In particular, when $a_0 = 2$ and $b_0 = 1$, then $\psi_{k,n}(x)$ forms an orthonormal basis. Hermite wavelets are defined as [24]

$$\psi_{k,n}(x) = \begin{cases} \frac{2^{k+1/2}}{\sqrt{\pi}} H_m(2^k x - 2n + 1), & \frac{n-1}{2^{k-1}} \leq x < \frac{n}{2^{k-1}} \\ 0, & \text{otherwise} \end{cases} \quad (2)$$

where $m = 0, 1, \dots, M-1$. Here is Hermite polynomials of degree m with respect to weight function $W(x) = \sqrt{1-x^2}$ on the real line $\mathbb{R}$ and satisfies the following recurrence formula $H_0(x) = 1, H_1(x) = 2x$,

$$H_{m+2}(x) = 2xH_{m+1}(x) - 2(m+1)H_m(x) \quad (3)$$

where $m = 0, 1, 2, \dots$

Function approximation: Here we approximating the solution $y(x)$ of Abel's integral equations using Hermite wavelet basis as follows:

$$y(x) = \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} c_{n,m} \psi_{n,m}(x) \quad (4)$$

where $\psi_{n,m}(x)$ is given in eq.(2). We approximate $y(x)$ by truncating the series represented in eq.(4) as,

$$y(x) = \sum_{n=1}^{2^{k-1}} \sum_{m=0}^{M-1} c_{n,m} \psi_{n,m}(x) = C^T \Psi(x) \quad (5)$$

where C and $\Psi(x)$ are $2^{k-1}M \times 1$ matrix,

$$C^T = [c_{1,0}, \dots, c_{1,M-1}, c_{2,0}, \dots, c_{2,M-1}, \dots, c_{2^{k-1},0}, \dots, c_{2^{k-1},M-1}] \quad (6)$$

$$\Psi(x) = [\psi_{1,0}, \dots, \psi_{1,M-1}, \psi_{2,0}, \dots, \psi_{2,M-1}, \dots, \psi_{2^{k-1},0}, \dots, \psi_{2^{k-1},M-1}] \quad (7)$$

3 Convergence and Error Analysis

Theorem 3.1. The series solution of Hermite wavelet $y(x) = \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} c_{n,m} \psi_{n,m}(x)$ is converges to $y(x)$.

Proof: Let $L^2(R)$ be the infinite dimensional Hilbert space and $\psi_{n,m}$ is defined as eq.(2) forms an orthonormal basis.

Let $y(x) = \sum_{i=0}^{M-1} c_{n,i} \psi_{n,i}(x)$ where $c_{n,i} = \langle y(x), \psi_{n,i}(x) \rangle$ for a fixed n .

Let us denote the sequence of partial sums S_n of $\{c_{n,i} \psi_{n,i}(x)\}$,

Let S_n and S_m be the partial sums with $n \geq m$. Now we have to prove S_n is a Cauchy sequence in Hilbert space $L^2(R)$.

Choose, $S_n = \sum_{i=0}^n c_{n,i} \psi_{n,i}(x)$, Now $\langle y(x), S_n \rangle = \left\langle y(x), \sum_{i=0}^n c_{n,i} \psi_{n,i}(x) \right\rangle = \sum_{i=m+1}^n |c_{n,i}|^2$

We claim that $\|S_n - S_m\|^2 = \sum_{i=m+1}^n |c_{n,i}|^2, \forall n > m$

Now $\left\| \sum_{i=m+1}^n c_{n,i} \psi_{n,i}(x) \right\|^2 = \left\langle \sum_{i=m+1}^n c_{n,i} \psi_{n,i}(x), \sum_{i=m+1}^n c_{n,i} \psi_{n,i}(x) \right\rangle = \sum_{i=m+1}^n |c_{n,i}|^2, \forall n > m$

thus, $\left\| \sum_{i=m+1}^n c_{n,i} \psi_{n,i}(x) \right\|^2 = \sum_{i=m+1}^n |c_{n,i}|^2, \forall n > m$

Since, Bessel's inequality, we have $\sum_{i=m+1}^n |c_{n,i}|^2 \leq \|y(x)\|^2$ is bounded and convergent.

Hence, $\left\| \sum_{i=m+1}^n c_{n,i} \psi_{n,i}(x) \right\|^2 \rightarrow 0$ as $m, n \rightarrow \infty$.

This implies, $\left\| \sum_{i=m+1}^n c_{n,i} \psi_{n,i}(x) \right\| \rightarrow 0$. and

Therefore $\{S_p\}$ is a Cauchy sequence and it converges to s (say).

We assert that $y(x) = s$

Now $\langle s - y(x), \psi_{n,i}(x) \rangle = \langle s, \psi_{n,i}(x) \rangle - \langle y(x), \psi_{n,i}(x) \rangle = \langle s, \psi_{n,i}(x) \rangle - \left\langle \lim_{n \rightarrow \infty} S_n, \psi_{n,i}(x) \right\rangle = 0$,

This implies,

$$\langle s - y(x), \psi_{n,i}(x) \rangle = 0$$

Hence $y(x) = s$ and $\sum_{i=0}^n c_{n,i} \psi_{n,i}(x)$ converges to $y(x)$ as $n \rightarrow \infty$ and proved.

Theorem 3.2. Suppose that $y(x) \in C^m[0, 1]$ and $C^T \Psi(x)$ is the approximate solution using Hermite wavelet. Then the error bound would be given by,

$$\|E(x)\| \leq \left\| \frac{2}{m! 4^m 2^{m(k-1)}} \max_{x \in [0,1]} |y^{(m)}(x)| \right\|.$$

Proof: Applying the definition of norm in the inner product space, we have,

$$\|E(x)\|^2 = \int_0^1 [y(x) - C^T \Psi(x)]^2 dx.$$

Divide interval $[0, 1]$ into 2^{k-1} subintervals $I_n = \left[\frac{n-1}{2^{k-1}}, \frac{n}{2^{k-1}}\right]$, $n = 1, 2, 3, \dots, 2^{k-1}$.

$$\|E(x)\|^2 = \sum_{n=1}^{2^{k-1}} \int_{\frac{n-1}{2^{k-1}}}^{\frac{n}{2^{k-1}}} [y(x) - C^T \Psi(x)]^2 dx.$$

$$\|E(x)\|^2 \leq \sum_{n=1}^{2^{k-1}} \int_{\frac{n-1}{2^{k-1}}}^{\frac{n}{2^{k-1}}} [y(x) - P_m(x)]^2 dx.$$

where $P_m(x)$ is the interpolating polynomial of degree m which approximates $y(x)$ on I_n . By using the maximum error estimate for the polynomial on I_n , then

$$\|E(x)\|^2 \leq \sum_{n=1}^{2^{k-1}} \int_{\frac{n-1}{2^{k-1}}}^{\frac{n}{2^{k-1}}} \left[\frac{2}{m! 4^m 2^{m(k-1)}} \max_{x \in I_n} |y^{(m)}(x)| \right]^2 dx.$$

$$\|E(x)\|^2 \leq \sum_{n=1}^{2^{k-1}} \int_{\frac{n-1}{2^{k-1}}}^{\frac{n}{2^{k-1}}} \left[\frac{2}{m! 4^m 2^{m(k-1)}} \max_{x \in [0,1]} |y^{(m)}(x)| \right]^2 dx.$$

$$\|E(x)\|^2 = \int_0^1 \left[\frac{2}{m! 4^m 2^{m(k-1)}} \max_{x \in [0,1]} |y^{(m)}(x)| \right]^2 dx$$

$$\|E(x)\| \leq \left\| \frac{2}{m! 4^m 2^{m(k-1)}} \max_{x \in [0,1]} |y^{(m)}(x)| \right\|.$$

which, we have used the well-known maximum error bound for the interpolation.

4 Method of Solution

Consider the Abel's integral equation of the form,

$$\lambda y(x) = f(x) + \int_0^x \frac{y(t)}{\sqrt{x-t}} dt, \quad 0 \leq x, t \leq 1 \quad (8)$$

where $\lambda = 0$ or $\lambda = 1$. We first approximate $y(x)$ as truncated series defined in eq.(4). That is,

$$y(x) \approx \sum_{n=1}^{2^{k-1}} \sum_{m=0}^{M-1} c_{n,m} \psi_{n,m}(x) = C^T \Psi(x) \quad (9)$$

where C and $\Psi(x)$ are defined in eq.(6) and (7). Then substituting eq.(9) in eq.(8), we get

$$\lambda C^T \Psi(x) = f(x) + \int_0^x \frac{C^T \Psi(t)}{\sqrt{x-t}} dt \quad (10)$$

Next, assume eq.(10) is precise at following collocation points $x_i = \frac{2i-1}{2^k M}$, $i = 1, 2, \dots, 2^{k-1}M$. Then we obtain

$$\lambda C^T \Psi(x_i) = f(x_i) + \int_0^{x_i} \frac{C^T \Psi(t)}{\sqrt{x_i-t}} dt \quad (11)$$

Next, we obtain the system of algebraic equations with $2^{k-1}M$ unknown coefficients. By solving this system of equations, we get Hermite wavelet coefficients and then substituting these coefficients in eq.(9), we get the approximate solution of eq.(8).

5 Numerical Examples

In this section, we present Hermite wavelets method for the numerical solution of Abel's integral equation to demonstrate the capability of the present method.

$$\text{Errorfunction} = \|y_e(x_i) - y_a(x_i)\|_\infty = \sqrt{\sum_{i=1}^n (y_e(x_i) - y_a(x_i))^2}$$

where y_e and y_a are the exact and approximate solution respectively.

Example 1. Consider the Abel's integral equation of first kind [22],

$$\frac{2}{105} \sqrt{x} (105 - 56x^2 + 48x^3) = \int_0^x \frac{y(t)}{x-t} dt, \quad 0 \leq x \leq 1. \quad (12)$$

We apply the present method to solve eq.(12) with $k = 1$ and $M = 4$. Then we get truncating approximate solution with unknowns as,

$$y(x) \approx \sum_{m=0}^3 c_{1,m} \psi_{1,m}(x) = C^T \Psi(x) \quad (13)$$

Then applying the procedure discussed in the section 3. We get a system of four algebraic equations with four unknowns and solving this system, we obtain the Hermite wavelet coefficients as, $c_{1,0} = \frac{1257}{1513}$, $c_{1,1} = -\frac{69}{5000}$, $c_{1,2} = \frac{277}{10000}$, $c_{1,3} = \frac{69}{5000}$, and substituting in eq.(13), we obtain: $y(x) = \frac{1257}{1513} \psi_{10}(x) - \frac{69}{5000} \psi_{11}(x) + \frac{277}{10000} \psi_{12}(x) + \frac{69}{5000} \psi_{13}(x)$ On simplifying, we get $y(x) = x^3 - x^2 + 1$, which is exact solution of eq.(12). Numerical results with exact solutions are shown in table 1 and graphically shown in figure 1.

Example 2. Consider the Abel's integral equation of the first kind [22, 23],

$$x = \int_0^x \frac{y(t)}{\sqrt{x-t}} dt, \quad 0 \leq x \leq 1. \quad (14)$$

which has the exact solution $y(x) = \frac{2}{\pi} \sqrt{x}$. We solved the eq.(14) using the present method and obtained approximate solution is compared with exact and other existing methods which reflects in table 2 and figure 2. Error analysis is shown in table 3 and figure 3.

Table 1 Numerical results of example 1.

x	Exact solution	Present method ($k = 1, M = 8$)	Abs. Error
0.1	0.9910	0.9910	1.61e-12
0.2	0.9680	0.9680	4.71e-13
0.3	0.9370	0.9370	2.27e-12
0.4	0.9040	0.9040	3.04e-12
0.5	0.8750	0.8750	2.52e-12
0.6	0.8560	0.8560	9.67e-13
0.7	0.8530	0.8530	9.64e-13
0.8	0.8720	0.8720	2.36e-12
0.9	0.9190	0.9190	2.36e-12

Table 2 Numerical results of example 2.

x	Exact solution	Present method ($k = 1, M = 10$)	Method [22] ($k = 1, M = 8$)	Method [23] ($m=16$)
0.1	0.201317	0.200842	0.200128	0.200460
0.2	0.284705	0.284667	0.286092	0.297987
0.3	0.348691	0.348628	0.347394	0.337588
0.4	0.402634	0.402609	0.404161	0.405769
0.5	0.450158	0.450129	0.449568	0.464014
0.6	0.493124	0.493113	0.492704	0.490550
0.7	0.532634	0.532607	0.532315	0.539721
0.8	0.569410	0.569440	0.569156	0.562698
0.9	0.603951	0.603690	0.603742	0.606044

Table 3 Error analysis of example 2.

x	Present method ($k = 1, M = 10$)	Method [22] ($k = 1, M = 8$)	Method [23] ($m=16$)
0.1	4.73e-04	1.18e-03	8.57e-04
0.2	3.77e-05	1.38e-03	1.32e-02
0.3	6.21e-05	1.29e-03	1.11e-02
0.4	2.37e-05	1.52e-03	3.13e-03
0.5	2.85e-05	5.90e-04	1.38e-02
0.6	9.87e-06	4.19e-04	2.57e-03
0.7	2.72e-05	3.19e-04	7.08e-03
0.8	3.05e-05	2.54e-04	6.71e-03
0.9	2.59e-04	2.08e-04	2.09e-03

Table 4 Numerical results of example 3.

x	Exact solution	Present method ($k = 1, M = 10$)	Method [23] ($m=16$)
0.1	0.953462589245592	0.953462604453520	0.95646081381695
0.2	0.912870929175277	0.912870928482700	0.90601007037324
0.3	0.877058019307029	0.877058021105882	0.88361513925322
0.4	0.845154254728517	0.845154255308354	0.84340093819493
0.5	0.816496580927726	0.816496581976408	0.80822420481499
0.6	0.790569415042095	0.790569415421744	0.79221049469412
0.7	0.766964988847370	0.766964989996025	0.76284677221990
0.8	0.745355992499930	0.745355991505969	0.74933888037055
0.9	0.725476250110012	0.725476258975992	0.72434536240934

Table 5 Error analysis of example 3.

x	Present method ($k = 1, M = 10$)	Method [23] ($m=16$)
0.1	1.52e-08	2.99e-03
0.2	6.92e-10	6.86e-03
0.3	1.79e-09	6.55e-03
0.4	5.79e-10	1.75e-03
0.5	1.04e-09	8.27e-03
0.6	3.79e-10	1.64e-03
0.7	1.14e-09	4.11e-03
0.8	9.93e-10	3.98e-03
0.9	8.86e-09	1.13e-03

Example 3. Consider the Abel's integral equation of the second kind [22, 23],

$$4y(x) = \frac{4}{\sqrt{x+1}} - \arcsin\left(\frac{1-x}{1+x}\right) + \frac{\pi}{2} - \int_0^x \frac{y(t)}{\sqrt{x-t}} dt, \quad 0 \leq x \leq 1. \quad (15)$$

which has the exact solution $y(x) = \frac{1}{\sqrt{x+1}}$. Applying the Hermite wavelet method for solving eq.(15), then obtained approximate solution is compared with the exact solution and method[23] are shown in table 4 and figure 4. Error analysis is shown in table 5. **Example 4.** Consider the Abel's integral equations of the second kind [22, 23],

$$y(x) = 2\sqrt{x} - \int_0^x \frac{y(t)}{\sqrt{x-t}} dt, \quad 0 \leq x \leq 1. \quad (16)$$

which has the exact solution $y(x) = 1 - \exp(\pi t) \operatorname{erfc}(\sqrt{\pi t})$. We solved the eq.(16) by the present method, we obtain the approximate solution and is compared with exact and other existing methods as shown in table 6 and figure 5. Error analysis is shown in table 7 and figure 6.

6 Conclusion

The Hermite wavelet method is applied for the numerical solution of Abel's integral equations. The present method reduces an integral equation into a set of algebraic equations. Obtained results are higher accuracy with exact ones and existing methods [22, 23], which can be observed in section 5. The numerical results shows that

Table 6 Numerical results of example 4.

x	Exact solution	Present method ($k = 1, M = 10$)	Method [22] ($k = 0, M = 16$)	Method [23] ($m=16$)
0.1	0.414059	0.411229	0.415689	0.402472
0.2	0.508352	0.507572	0.505528	0.519751
0.3	0.564309	0.563685	0.566205	0.554755
0.4	0.603347	0.602926	0.601908	0.605031
0.5	0.632868	0.632521	0.634188	0.640487
0.6	0.656323	0.656059	0.655109	0.654785
0.7	0.675601	0.675358	0.676588	0.678700
0.8	0.691842	0.691690	0.691596	0.688860
0.9	0.705787	0.705398	0.704377	0.706495

Table 7 Error analysis of example 4.

x	Present method ($k = 1, M = 10$)	Method [22] ($k = 0, M = 16$)	Method [23] ($m=16$)
0.1	2.82e-03	1.62e-03	1.15e-02
0.2	7.79e-04	2.82e-03	1.13e-02
0.3	6.23e-04	1.89e-03	9.55e-03
0.4	4.21e-04	1.43e-03	1.68e-03
0.5	3.46e-04	1.32e-03	7.61e-03
0.6	2.63e-04	1.21e-03	1.53e-03
0.7	2.42e-04	9.86e-04	3.09e-03
0.8	1.50e-04	2.45e-04	2.98e-03
0.9	3.87e-04	1.40e-03	7.08e-04

Fig. 1 Comparison of numerical solutions with exact solutions of example 1.

Fig. 2 Comparison of numerical solutions, exact solutions and existing methods of example 2.

Fig. 3 Comparison of error analysis of example 2.

Fig. 4 Comparison of Numerical solutions of example 3.

Fig. 5 Comparison of Numerical solutions of example 4.

Fig. 6 Comparison of Error analysis of example 4.

the accuracy improves with increasing the values of M for better accuracy. Convergence theorem reveals that existence of solution.

References

- [1] C. K. Chui, Wavelets: A Mathematical Tool for Signal Analysis, SIAM, Philadelphia, PA., 1997.
- [2] G. Beylkin, R. Coifman, V. Rokhlin, Fast wavelet transforms and numerical algorithms I, Commun. Pure Appl. Math., 44 (1991), 141–183.
- [3] Ü. Lepik, E. Tamme, Application of the Haar wavelets for solution of linear integral Equations, Ant. Turk–Dynam. Sys. Appl. Proce., (2005), 395–407.
- [4] K. Maleknejad, M.T. Kajani, Y. Mahmoudi, Numerical solution of linear Fredholm and Volterra integral equation of the second kind by using Legendre wavelets, J. Kybernet., 32 (2003), 1530–1539.
- [5] K. Maleknejad, F. Mirzaee, Using rationalized haar wavelet for solving linear integral equations, App. Math. Comp., 160 (2005), 579–587.
- [6] K. Maleknejad, M. Yousefi, Numerical solution of the integral equation of the second kind by using wavelet bases of hermite cubic splines, App. Math. Comp., 183 (2006), 134–141.
- [7] K. Maleknejad, T. Lotfi, Y. Rostami, Numerical computational method in solving fredholm integral equations of the second kind by using coifman wavelet, App. Math. Comp., 186 (2007), 212–218.
- [8] S. Yousefi, A. Banifatemi, Numerical solution of Fredholm integral equations by using CAS wavelets, App. Math. Comp., 183 (2006), 458–463.
- [9] S. C. Shiralashetti, R. A. Mundewadi, Bernoulli Wavelet Based Numerical Method for Solving Fredholm Integral Equations of the Second Kind, J. Inform. Comp. Sci., 11(2) (2016), 111–119.
- [10] S. A. Yousefi, Numerical solution of Abel's integral equation by using Legendre wavelets, Appl. Math. Comp., 175 (2006), 574–80.
- [11] S. Sohrabi, Comparison Chebyshev wavelets method with BPFs method for solving Abel's integral equation, Ain Shams Eng. J., 2 (2011), 249–254.
- [12] A. M. Wazwaz, Linear and nonlinear integral equations: methods and applications, Berlin: Higher Education, Beijing, Springer, 2011.
- [13] A. M. Wazwaz, A first course in integral equations, Singapore: World Scientific Publishing, 1997.
- [14] R. Gorenflo, S. Vessella, Abel integral equations, analysis and applications, In: Lecture notes in mathematics, Heidelberg: Springer, 1991.
- [15] C.T.H. Baker, The numerical treatment of integral equations, Clarendon Press, Oxford, 1977.
- [16] S.C. Shiralashetti, S. Kumbinarasaiah, Theoretical study on continuous polynomial wavelet bases through wavelet

- series collocation method for nonlinear Lane–Emden type equations, *Appl. Math. Comput.*, 315 (2017), 591–602.
- [17] H. Brunner, *Collocation methods for Volterra integral and related functional differential equations*, Cambridge Monographs on Applied and Computational Mathematics, Cambridge Univ. Press, Cambridge, 2004.
 - [18] S. Noeiaghdam, E. Zarei, H. B. Kelishami, Homotopy analysis transform method for solving Abel’s integral equations of the first kind, *Ain Shams Eng. J.*, 7 (2016), 483–495.
 - [19] N. Zeilon, Sur quelques points de la theorie de l’equation integrale d’Abel, *Arkiv Mat Astr Fysik*, 18 (1924), 1–19.
 - [20] C.T.H. Baker, *The numerical treatment of integral equations*, Oxford: Clarendon Press, 1977.
 - [21] P. Baratella, A. P. Orsi, A new approach to the numerical solution of weakly singular Volterra integral equations, *J. Comput. Appl. Math.*, 163(2) (2004), 401–418.
 - [22] [22] E. Babolian, A. Salimi Shamloo, Numerical solution of Volterra integral and integro-differential equations of convolution type by using operational matrices of piecewise constant orthogonal functions, *J. Comp. Appl. Math.*, 214 (2008), 495–508.
 - [23] A. Shahsavaran, M. R. Moazami Goudarzi, O. Moradtalab, Solving Abel’s integral equation of the first kind using piecewise constant functions and Taylor expansion by collocation method, In: 40th Annual Iranian mathematics conference.
 - [24] A. Ali, M.A. Iqbal, S.T. Mohyud-Din, Hermite wavelets method for Boundary Value problems, *Inter. J. Modern Appl. Phys.*, 3(1) (2013), 38–47.
 - [25] M.H. Kantli, S.C. Shiralashetti, Finite difference Wavelet–Galerkin method for the numerical solution of elastohydrodynamic lubrication problems, *Journal of Analysis*, 26(2) (2018), 285–295.
 - [26] N.M. Bujurke, M.H. Kantli, B.M. Shettar, Jacobian free Newton-GMRES method for the solution of elastohydrodynamic grease lubrication in line contact using wavelet based pre-conditioners, *Proceedings of the National Academy of Sciences, India Section A: Physical Sciences*, 88(2) (2018), 247–265.
 - [27] N.M. Bujurke, M. H. Kantli, B.M. Shettar, Wavelet preconditioned Newton-Krylov method for elastohydrodynamic lubrication of line contact problems, *Applied Mathematical Modelling*, 46 (2017), 285–298.
 - [28] S.C. Shiralashetti, M.H. Kantli, A.B. Deshi, A new wavelet multigrid method for the numerical solution of elliptic type differential equations, *Alexandria Engineering Journal*, 57 (2018), 203–209.