



Sex estimation of upper long bones by selected measurements in a Radom (Poland) population from the 18th and 19th centuries AD

*Jacek Tomczyk¹, Joanna Nieczuja-Dwojacksa², Marta Zalewska³,
Wojciech Niemirow^{4,5}, Wioleta Olczyk²*

¹Department of Biological Anthropology, Institute of Ecology and Bioethics, Cardinal Stefan Wyszyński University in Warsaw, Poland

²Department of Human Ecology, Faculty of Biology and Environmental Sciences, Cardinal Stefan Wyszyński University in Warsaw, Poland

³Department of Environmental Hazard Prevention and Allergology, Medical University of Warsaw, Poland

⁴Faculty of Mathematics, Informatics and Mechanics, University of Warsaw, Poland

⁵Faculty of Mathematics and Computer Science, Nicolaus Copernicus University, Toruń, Poland

ABSTRACT: Several studies have shown that sex estimation methods based on measurements of the skeleton are specific to populations. Metric traits of the upper long bones have been reported as reliable indicators of sex. This study was designed to determine whether the four long bones can be used for the sex estimation of an historical skeletal population from Radom (Poland). The material used consists of the bones of 169 adult individuals (including 103 males and 66 females) from the 18th and 19th centuries. Twelve measurements were recovered from clavicle, humerus, radius and ulna. The initial comparison of males and females indicated significant differences in all measurements ($p < 0.0001$). The accuracy of sex estimation ranged from 68% to 84%. The best predictor for sex estimation of all the measurements in Radom's population was the maximum length of the radius (84%), and the ulna (83%), and the vertical diameter of the humeral head (83%). The Generalized Linear Model (GLM) detected the strongest significant relationship between referential sex and the vertical diameter of the humeral head ($p < 0.0001$), followed by the maximal length of the ulna ($p = 0.0117$). In other measurements of the upper long bones, GLM did not detect statistically significant differences.

KEY WORDS: bioarchaeology, sex estimation, upper extremities long bones

Introduction

Sex estimation is an important first step in the development of a biological profile in human osteology and is used in both bioarchaeological studies (González et al. 2007; Bašić et al. 2013) and forensic anthropological analysis (Kamdi et al. 2014; Peckmann et al. 2015). In these fields, it is important to recognize the sex of the skeleton, regardless of the state of preservation. Today, significant accuracy in sex determination is possible through the DNA analysis from bones. Unfortunately, these methods require sophisticated equipment, and can fail in cases of dry or seriously damaged bone samples (Faerman et al. 1998; Santos et al. 1998; Rösing et al. 2007). Moreover, aDNA extraction is also an invasive technique, and it cannot be often used in the case of valuable historical bone collections. Therefore, if the skeleton is reasonably complete and a reliable morphological diagnosis is possible, there is no need to apply molecular methods. In such cases, sex estimation can generally be detected by two different approaches: observation of morphological traits and/or measurement of different bones (Asala et al. 2004; Cowal and Pastor 2008; Robinson and Bidmos 2011; Spradley and Jantz 2011; Albanese 2013; Bašić et al. 2013).

It is widely accepted that the pelvic girdle is the most accurate area by which to estimate sex and methods using these elements tend to make successful predictions in 90% to 95% of individuals (Ferembach et al. 1980; Volk and Ubelaker 2002). Comparative studies show varying degrees of the accuracy assessment of sex depending on the area of the pelvic region (sacroiliac segment, ischio-

pubic segment and acetabular segment) (Bruzek and Murail 2006; González et al. 2007). The skull is somewhat less reliable for use in estimating sex, ranging from 77% to 92% (Krogman and Iscan 1986; Byers 2002; Bass 2005; Bruzek and Murail 2006). Such a large divergence in dimorphism assessment can be the result of the diversity of populations (Howells 1996; Bass 2005; Dayal et al. 2008; Spradley and Jantz 2011).

Sex estimation when the pelvic girdle and the skull are absent or destroyed is problematic. Therefore, sex estimation based on other bones is an important complement to widely used methods. Long bones have demonstrated their usefulness in sex assessment studies (e.g., Ruff and Hayes 1988; Liu 1989; González-Reimers et al. 2000). However, it is not just the morphologic data for long bones which is studied, but also the measurements for scapula (Spradley and Jantz 2011), ribs (Kubicka and Piontek 2016), carpal bones (Mastrangelo et al. 2011), calcaneus (Peckmann et al. 2015), talus (Otong et al. 2016), and even the internal auditory canal located on the temporal bone (Gonçalves et al. 2015).

Dimensions from the upper long bones (humerus, radius, ulna, and clavicle) have been used to successfully distinguish between the sexes in several European (Cowal and Pastor 2008; Bašić et al. 2013; Kranioti and Tzanakis 2015), Asian (Sakaue 2004; Lee et al. 2014) and African populations (Steyn and Işcan 1999; Barrier and Abbé 2008). In mentioned studies, different measurements have been shown to be good indicators of sex dimorphism. The ability to estimate the sex of each of these bones equips anthropologists with more options when dealing with cases in which the recovered remains are incomplete or destroyed.

The aim of this study is to estimate the degree of sex dimorphism in the 18th and 19th century population of Radom (Poland) and to develop metric standards for the estimation of sex. The upper long bones will be examined for this purpose (clavicle, humerus, ulna, and radius). In particular, the objective is to collect metric data that will generate formulae suitable for Radom's group. These formulae will probably produce the highest levels of accuracy possible for estimating sex in skeletal remains from this geographic region.

Materials and methods

The samples were selected from the Radom Cemetery collection curated at the Department of Biological Anthropology at Cardinal Stefan Wyszyński University (Warsaw, Poland). The present work utilized material obtained during the excavation seasons 2010 to 2013.

Radom is a medium-sized city located in east-central Poland, about 100 km south of Poland's capital, Warsaw.

Although the city of Radom has a long history dating back to the early Middle Ages, our study focused on individuals from the 18th and 19th centuries (Fuglewicz 2011). According to historical information, we know that the first urban municipal cemetery was founded in 1791, but due to the lack of space, a new cemetery was established in 1811 at another location, while the municipal cemetery at the citadel has been abandoned and forgotten (Piątkowski 2000; Zapłata 2011). This means that all the examined human remains come from the turn of the century, and they were buried within a span of 20 years.

Anthropological analysis of the skeletal remains was conducted at the Depart-

ment of Biological Anthropology of the Cardinal Stefan Wyszyński University in Warsaw.

During the archaeological excavations, a total of 275 burials were uncovered. However, in the present study, the analyses were narrowed to the adult individuals, whose state of bone preservation allowed further examination. Only the bones with complete ossification were included in the present study. Bones showing any deformity or degeneration were excluded from further analyses. According to the above criteria only 180 individuals were chosen. But the final analyses were narrowed only to those individuals whose assessment of sex, conducted by two independent investigators, were identical (JT, WO). Above criteria have been met by 169 individuals (103 male and 66 female), who were selected to the further studies.

The age of the individual was estimated on the basis of changes in the morphology of the pubic symphysis, using the Brooks and Suchey (1990) system and standards for changes in the topography of the auricular surface (Buikstra and Ubelaker 1994; White and Folkens 2000). The referential sex of the individuals was estimated by the Phenice (1969) method. It includes visual assessments of pelvic traits, such as the greater sciatic notch, ischiopubic ramus, subpubic concavity and ventral arc. Although cranial non-metric traits are not as accurate as pelvic traits for sex estimation, they can provide successful estimates, so they cannot be ignored in the analyses. Cranial features, such as the supra-orbital ridges, nuchal area, mastoids, and chin, were used in sex estimation (Buikstra and Ubelaker 1994; Konigsberg and Hens 1998; Walker 2008).

All measurements were made according to Martin (1957): i) clavicle: maximum length (M1), superior (vertical-inferior) diameter at midshaft (M4), anterior (sagittal-posterior) diameter at midshaft (M5); ii) humerus: maximum length (M1), transverse diameter of head (M9), vertical diameter of head (M10); iii) radius: maximum length (M1), medial-lateral diameter at midshaft (M4), anterior-posterior diameter (M5); iv) ulna: maximum length (M1), anterior-posterior diameter (M11), medial-lateral diameter (M12).

Measurements were taken with a Vernier caliper calibrated to 0.1 mm and an osteometric board. Wherever possible, the left side was used. Selected measurements (superior-inferior diameter of the clavicle, transverse diameter of the humeral head and medio-lateral diameter of the ulna) were dubbed twice by the same investigator (JT). The results of the measurements of these observations were compared with the referential sex as estimated from the skull and pelvis.

Statistical analyses were undertaken as follows. First, reproducibility of the selected measurements, made by this same investigator, was assessed by calculating Spearman's correlation coefficient. Second, the Cohen's Kappa coefficient was used to compare observations between investigators. Then, the one-sample Kolmogorov-Smirnov test was conducted for all variables in either sex in order to test for any deviations from a normal distribution. The F-test of equality of variance between sexes was used. Differences in mean values between the sexes were examined by the t-student test or the Welch's test. The t-student test is more reliable when the two-sample has equal variances, while the Welch's test is better when variance is unequal.

In our analysis we consider 'n' objects, among them nF belonging to class "F" (females), and nM belonging to class "M" (males). For each of the objects we also have the numeric values of some measurements. We wanted to find a measurement which the best discriminates between "F" and "M". For a given measurement we chose a cut-point C, and computed the following indicators:

TF = number of "F" among those objects which have the measurement $\leq C$

TM = number of "M" among those objects which have the measurement $> C$

$$ACC = (TF+TM)/n$$

ACC (accuracy of classification) is the fraction of correctly recognized objects.

Finally, the Generalized Linear Model (GLM) was used to assess which measurement was the most reliable method for assessing the sex of skeletons in the study population of Radom. Statistical analyses were performed using the R Project for Statistical Computing (2013). Differences showing $p \leq 0.05$ were considered statistically significant.

Results

Assessment of sex was done by two independent investigators. The first investigator identified 109 males and 71 females from 180 individuals, while the second investigator recognized 106 males and 77 females from 180 individuals. It means that eleven individuals were classified differently by two investigators, and those individuals were excluded from further studies. The value for Kappa is 0.8961 indicating a very good level of agreement. Such a result verifies the high repeatability of the observations

Table 1. Results of statistical analyses in both sexes

Measurement		Kolmogorov-Smirnov test		
		Total	Males	Females
Clavicle	Maximal length	0.274	0.277	0.281
	Antero-posterior diameter	0.274	0.276	0.280
	Superior-inferior diameter	0.274	0.276	0.280
Humerus	Maximal length	0.274	0.277	0.281
	Vertical diameter of head	0.274	0.276	0.280
	Transverse diameter of head	0.274	0.277	0.282
Radius	Maximal length	0.274	0.277	0.281
	Antero-posterior diameter	0.274	0.276	0.279
	Medio-lateral diameter	0.274	0.276	0.279
Ulna	Maximal length	0.275	0.277	0.283
	Antero-posterior diameter	0.274	0.276	0.279
	Maximal length	0.274	0.276	0.279

and it allowed us to estimate the referential sex. For intra examiner reproducibility the Spearman correlation values for the transverse diameter of the humeral head was 0.97, superior-inferior diameter of the clavicle was 0.96, and for the medio-lateral diameter of the ulna was 0.95. Such results suggest that all the measurements have high reproducibility.

The distribution of all variables of each sex was detected by the Kolmogorov-Smirnov test. This test did not detect any deviations from the normal distribu-

tion at the statistically significant level (Table 1).

Lack of equality of variance was detected by the F-test in three cases: the transverse diameter of the humeral head ($F= 0.5402$, $p= 0.0265$), and the antero-posterior ($F= 5.7712$, $p= 0.0001$) and medio-lateral ($F= 1.7339$, $p= 0.0195$) diameter of the radius (Table 2). In these cases, the Welch's test was used to calculate statistical significance between the mean of measurements.

Table 2. F test to compare two variances between sexes

Measurement		F	p
Clavicle	Maximal length	0.82	0.483
	Antero-posterior diameter	0.86	0.573
	Superior-inferior diameter	0.83	0.467
Humerus	Maximal length	0.02	0.071
	Vertical diameter of head	0.79	0.365
	Transverse diameter of head	0.54	0.026*
Radius	Maximal length	0.88	0.666
	Antero-posterior diameter	5.77	0.0001*
	Medio-lateral diameter	1.73	0.0195*
Ulna	Maximal length	0.96	0.908
	Antero-posterior diameter	1.79	0.737
	Maximal length	1.35	0.207

Table 3. Maximum, minimum, average and standard deviations of the measurements

Measurement	Sex	Min	Max	Mean	CI (95%)		SD	p	
					Lower bound	Upper bound			
Clavicle	Maximal length	M	109	172	142.5	140.1	143.7	10.6	0.0001
		F	109	163	134	132.2	137.7	9.7	
	Antero-posterior diameter	M	7	18	12.4	11.7	12.3	1.8	0.0001
		F	7	15	10.6	10.2	11.3	1.7	
	Superior-inferior diameter	M	7	16	10.8	10.6	11.4	1.9	0.0001
		F	7	15	9.3	8.5	9.4	1.8	
Humerus	Maximal length	M	270	366	325.1	323.0	330.9	24.6	0.0001
		F	231	360	303.9	293.9	305.0	19.3	
	Vertical diameter of head	M	37	52	45.7	45.3	46.6	3.1	0.0001
		F	32	47	41.3	40.8	42.1	2.7	
	Transverse diameter of head	M	32	48	42	41.2	42.7	3.2	0.0001*
		F	32	43	37.5	36.3	37.7	2.3	
Radius	Maximal length	M	215	280	242.7	238.6	245.3	14.3	0.0001
		F	185	252	218	216.5	223.4	13.5	
	Antero-posterior diameter	M	9	14	12.7	11.8	12.2	1.0	0.0009*
		F	8	23	11	9.7	10.2	2.4	
	Medio-lateral diameter	M	12	19.6	16.2	16.2	16.9	1.5	0.0001*
		F	9.6	20	14.4	14.2	15.4	1.9	
Ulna	Maximal length	M	225	300	260.4	256.6	263.4	14.5	0.0001
		F	200	273	237	232.7	241.2	14.9	
	Antero-posterior diameter	M	10	20	13.8	13.0	13.7	1.7	0.0001
		F	8	19	12	11.5	12.4	1.8	
	Medio-lateral diameter	M	12	21	16.5	16.5	17.4	1.9	0.0001
		F	8	20.1	14.7	14.5	15.4	2.2	

*Welch test sample.

All variables of measurements are shown in Table 3. Generally, all of the values were larger for males than for females, and these differences were statistically significant. The greatest differences between sexes in mean values were detected in the measurement of length: radial maximal length (males: 242 mm, females: 218 mm), ulnar maximal length (males: 260 mm, females: 237 mm), humeral maximal length (males: 325 mm, females: 303 mm), and clavicle maximal length (males: 142 mm, females: 134 mm). The differences between sexes in

mean values of transverse measurements were at a similar level.

The accuracy of sex estimation for the upper long bones size is represented in Table 4.

The accuracy in estimation of sex using different measurements ranged from 68% to 84%. In general, the values for the antero-posterior diameter and maximal length of clavicle were better predictors than the superior-inferior diameter. However, these received values against other analyzed bones, are very small. The measured values showed that vertical diameter of the humeral head was a better

Table 4. Results for classification of accuracy and demarking point

	Measurement	Sex	Accuracy (%)		Demarking point (mm)
Clavicle	Maximal length	M	60.5	73.3	F<140<M
		F	83.3		
	Antero-posterior diameter	M	69.8	73.5	
		F	75.8		
Superior-inferior diameter	M	55.1	68.5	F<10<M	
	F	79.2			
Humerus	Maximal length	M	74.3	83.3	F<316<M
		F	82.0		
	Vertical diameter of head	M	80.4	78.5	
		F	85.4		
Transverse diameter of head	M	78.4	84.5	F<39<M	
	F	78.8			
Radius	Maximal length	M	84.2	84.5	F<228<M
		F	85.1		
	Antero-posterior diameter	M	78.8	82.4	
		F	84.4		
Medio-lateral diameter	M	70.0	72.3	F<15.5<M	
	F	74.1			
Ulna	Maximal length	M	83.0	83.2	F<248<M
		F	83.5		
	Antero-posterior diameter	M	64.4	76.0	
		F	79.1		
Medio-lateral diameter	M	57.1	68.4	F<16.2<M	
	F	87.2			

predictor than the transverse diameter of the head or the maximal length. The values for the maximal length of both the radius and the ulna were better predictors than the transverse diameter (antero-posterior and medio-lateral).

The best predictor for sex estimation of all the measurements in Radom's population is the maximum length of radius (84.5%) and ulna (83.2%), and the vertical diameter of the humeral head (83.3%). The worst predictive values were the superior-inferior diameter of the clavicle (68.5%) and the medio-lateral diameter of the ulna (68.4%) (Fig. 1).

The discriminate function coefficient of the upper long bones dimension for

Radom's population is presented in Table 4. The functions are displayed based on a single variable. Females are indicated when the discriminate score is lower than the demarcation points and males are indicated when the discriminate score is higher. For example, a maximal length of clavicle size of 143 mm would be identified as a male because the diameter is greater than the 140 mm function coefficient.

In the final step, the GLM was used to examine the relationship between the following measurements, with the referential sex estimation from the skull and the pelvis. GLM, which ignores inter-individual variability, detected a weak sta-




Fig. 1. Boxplots with the main measurements (A – vertical diameter of humeral head, B – maximum length of radius, C – maximal length of ulna)

tistically significant correlation between the referential sex and maximal length of the ulna ($p = 0.0117$), while a stronger significant relationship was found between the referential sex and the vertical diameter of humeral head ($p < 0.0001$). In other measurements of the upper long bones, GLM did not detect any statistically significant differences.

Discussion

The skeleton plays a significant role in various sciences, such as forensic science, medicine and anthropology. The estimation of sex using metric methods has been explored by many researchers of both contemporary (Barrier and L'Abbé 2008; Celbis and Agritmis 2006) and historical populations (Bergmann et al. 1962; Cowal and Pastor 2008; Bašić et al. 2013). The humerus or clavicle have been studied intensively and standards have been obtained for several populations, from the Adriatic coast (Bašić et al. 2013), the southern India region (Patil et al. 2011), Iran (Akhlaghi et al. 2012), and even prehistoric New Zealand populations (Murphy 1994). Also, the ulna and radius have been the subject of many osteometric studies from different world regions, such as Crete

(Kranioti and Michalodimitrakis 2009), Turkey (Celbis and Agritmis 2006) and South Africa (Barrier and L'Abbé 2008). Although many Polish populations have been described through metric characteristics (Piontek 1969; Ćwirko-Godycki and Swedborg 1977; Psonak and Kwiatkowska 2011), unfortunately there are few studies of sex estimation using the measurement of the upper long bones (Bergmann et al. 1961; 1962).

It should be noted that all these studies have shown that populations have different metric manifestations in both sexes. These variations are related to body size and consequently to metric differences in the dimensions of individual skeletal elements. Observed differences can be attributed to genetic factors (Holden and Mace 1999; Nyati et al. 2006; Gustafsson et al. 2007), physical activity (Ruff 1987; 2003; Carlson et al. 2007; Cowgill and Hager 2007; Patil et al. 2011), and environmental and/or social conditions (Stini 1969; Stinson 1985; Celbis and Agritmis 2006). In this context, there is evidence that populations that have either a very low or very high protein intake demonstrate the least amount of sex dimorphic variation (Gray and Wolfe 1980). The multitude of combinations among these factors may partly

explain why it is so difficult to draw unambiguous conclusions on the etiology of the differences observed between the two sexes. Despite the above limitations, many studies, including our data, indicate that metric differences in the upper long bones are quite a good indicator of sex and they can be used in the diagnosis of the sex of individuals. The results have revealed that the mean values of the male measurements were significantly higher than those of females.

The clavicles of Radom's population show that the mean length and both transverse dimensions (antero-posterior and superior-inferior) of males were statistically significantly larger than the females' clavicles ($p < 0.0001$). However, in the literature, the measurement at the maximum/minimum mid-shaft diameters of the clavicles (Ray 1959; Shirley 2009) or the mid-shaft circumference is preferred over the sagittal-vertical diameters at mid-shaft (Akhlagi et al. 2012; Murphy 1994). This fact may be the result of the anatomical structure of this part of the clavicle, and the measurement technique in this anatomical region, which can be "twisted", so all the transverse measurements can be disrupted. Finding the maximum and minimum diameters is easier than approximating the sagittal-vertical diameters, because the latter require a rough estimation of the bone's anatomical orientation. Nevertheless, using our team transverse dimensions (antero-posterior and superior-inferior) also demonstrated that variation between the sexes is visible.

Three measurements (maximum length, vertical and transverse diameter of the head) were taken from each humerus of Radom's population. They show that the mean values of males were statistically significantly higher than

those of the females ($p < 0.0001$). This observation is consistent with the results noted by other researchers who conducted studies both on contemporary (Patil et al. 2011; Thakur et al. 2015) and historical populations (Bergmann et al. 1961; 1962; Piontek 1969; Bašić et al. 2013).

For the ulnar and radial dimensions the sex difference between maximal length was also statistically significant. The mean for males was 260 mm and for females 237 mm in the ulna length measurement, and 242 mm for male and 218 mm for females in the radius length measurement. Sex dissimilarities in these bones were observed not only in the European populations (Piontek 1969; Celbis and Agritmis 2006), but also in Asian (Sakaue 2004; Lee et al. 2014) and African (Barrier and L'Abbé 2008) populations. However, it is worth emphasizing that both sexes are characterized by similar differences between medio-lateral and antero-posterior diameters in the bones of the forearm. Similar results, a small difference between the measurements for both sexes, were obtained for contemporary populations from Africa (Barrier and L'Abbé 2008). Results from the Early Medieval population from Ostrów Lednicki (Poland) reported increased values for both measurements. Moreover, in this population from the Early Medieval period, the medio-lateral dimension of the males' ulna was clearly bigger than the females' ulna (23.7 mm and 15.3 mm, respectively) (Ćwirko-Godycki and Swedborg 1977). That may indicate the different types of physical activity between males and females living in Ostrów Lednicki. The similar dimensions between males and females from 18th and 19th century Radom in this anatomical region suggest

similar or identical types of physical activity in the upper limb in both sexes.

General studies prove that the maximum length of the humerus, the vertical diameter of the head, or the humeral midshaft circumference are the most significant of all the measurements of the humerus in estimating sex (Bergmann et al. 1962; Loth and İscan 2000; Thakur et al. 2015). Dimensions such as maximum ulnar length, or mid-shaft circumference, as well as radial length and circumference of the radial head are shown to be good indicators of sex differences, of all the measurements of forearm bones (Mall et al. 2001; Barrier and L'Abbé 2008). According to our studies, the maximal length of the radius and the vertical diameter of the humeral head are the most discerning for the upper limb measurements (84.5% and 83.3%, respectively), followed by ulnar length (83.2%). The medio-lateral diameter of the ulna and the superior-inferior diameter of the clavicle (68.4% and 68.5%, respectively) are less useful for sex estimation of all the analyzed measurements. A comparison of the results of the present study with those from other studies of skeletal long bones demonstrates that these same measurements are especially valuable. The vertical diameter of the humeral head, and the maximum length of radius and ulna are important sex indicators in many other populations (Kranioti and Michalodimitrakis 2009; Bašić et al. 2013; Lee et al. 2014). However, these same measurements can have a different value in the assessment of sex in the studies of populations. For the vertical diameter of the head, a contemporary population from Korea had 87% accuracy, while our study achieved 83% accuracy (Lee et al. 2014). In the maximal length of the radius, modern Cretans

had 91% accuracy, while the population from Radom achieved only 84% accuracy (Kranioti and Michalodimitrakis 2009). In the maximal length of the humerus, a German population had 80% accuracy, while the analysis of Radom's population achieved 77% accuracy (Mall et al. 2001).

In the analysed sample, the females were classified better than the males for all variables. The maximum length of the radius was considered a useful measurement for distinguishing sex, with a classification accuracy of 84% for males and 85% for females. The vertical diameter of the humeral head was characterized by an accuracy of 80.4% for males and 85.4% for females. The maximum length of the ulna was considered to have a classification accuracy of 83% for males and 83.5% for females. A similar phenomenon was also observed by some authors (Cowan and Pastor 2008; Barrier and L'Abbé 2008). The explanation of this fact can be a less pronounced robustness in the male group, which would have increased their chances of being classified as females. However, we cannot exclude the reason that males and females may differ in size variability within their own group.

The demarcation points of Radom's population were larger than those determined by other studies, such as from the Eastern Adriatic regions (Bašić et al. 2013), Korea (Lee et al. 2014), Crete (Kranioti and Michalodimitrakis 2009), and an Early Medieval population from Poland (the city of Strzelno) (Piontek 1969). The demarcation points of Radom's sample from the 18th and 19th centuries were smaller in comparison with the Early Medieval Polish population from Ostrow Lednicki (Bergmann et al. 1962; Ćwirko-Godycki and Swedborg 1979). It means that it is extremely diffi-

cult, or even impossible, to propose general standards for differences between the sexes that would be applicable to all populations.

GLM was used to examine the relationship between measurements with the referential sex estimated from the skull and the pelvis. The results of the GLM analysis indicated that the vertical diameter of the humeral head is the most valuable for sex diagnosis in this population, followed by maximal length of the ulna. The rest of the measurements were not statistically significant according to this model. Overall, the data generated by the present investigation suggest that this metric dimension for the determination of sex in the skeletons of the Radom population can be very useful. However, the application of this dimension is not recommended for geographically remote populations or archeological samples before further studies can confirm their suitability.

Conclusion

Where pelvis or skull bones are not available, long bones can be useful in determining sex. In this context, metric measurements are preferred due to their high levels of accuracy. For the purpose of the present study, the clavicle, humerus, radius, and ulna were selected to determine which measurements of these bones were the most useful for the determination of sex in a population from Radom. Our studies proved that the maximal length of the radius, the vertical diameter of the humeral head and the maximal length of the ulna are the most useful of the upper limb measurements.

Acknowledgements

The research was financed by the National Science Centre (Poland) during the years 2013-2017 (Grant No 2013/11/B/HS3/04117).

Authors' contributions

JT designed the research interpreted the results, wrote the paper; JND gave support and conceptual advice, collected the articles, commented the initial draft of the manuscript; WO collected the data; MZ and WN performed statistical analyses. The final version of paper was prepared by JT and approved by all authors.

Conflict of interest

The authors declare that there is no conflict of interests regarding publication of this paper.

Corresponding author

Jacek Tomczyk, Department of Biological Anthropology, Institute of Ecology and Bioethics, Cardinal Stefan Wyszyński University in Warsaw, Wóycickiego 1/3, 01-938 Warsaw, Poland
email address: j.tomczyk@uksw.edu.pl

References

- Akhlaghi M, Moradi B, Hajibeygi M. 2012. Sex determination using anthropometric dimensions of the clavicle in Iranian population. *J Forensic Leg Med* 19:381-85.
- Albanese J. 2013. A method for estimating sex using the clavicle, humerus, radius, and ulna. *J Forensic Sci* 58(6):1413-19.
- Asala SA, Bidmos MA, Dayal MR. 2004. Discriminant function sexing of fragmentary femur of South African blacks. *Forensic Sci Int* 145(1):25-9.

- Barrier IL, L'Abbé EN. 2008. Sex determination from the radius and ulna in a modern South African sample. *Forensic Sci Int* 179:1–7.
- Bass WM. 2005. *Human osteology: a laboratory and field manual*. Columbia, MO: Missouri Archaeological Society.
- Bašić Ž, Anterić I, Vilović K, Petaros A, Bosnar A, Madžar B, et al. 2013. Sex determination in skeletal remains from the medieval Eastern Adriatic coast – discriminant function analysis of humeri. *Croat Med J* 54:272–78.
- Bergmann P, Goździewski S, Weolon Z. 1961. Investigation into the asymmetry of arm bones in man. *Materiały i Prace Antropologiczne*. 59:87–98.
- Bergmann P, Goździewski S, Weolon Z. 1962. Sexual dimorphism in arm bones. *Materiały i Prace Antropologiczne* 59:99–106.
- Brooks ST, Suchey JM. 1990. Skeletal age determination based on the os pubis: a comparison of the Acsádi-Nemeskéri and Suchey-Brooks methods. *Hum Evol* 5:227–38.
- Bruzek J, Murail P. 2006. Methodology and reliability of sex determination from the skeleton. In: A Schmitt, E Cuhna, J Pinheiro, editors. *Forensic Anthropology and Medicine, Complementary Sciences From Recovery to Cause of Death*. New Jersey: Humana Press. 225–42.
- Buikstra J, Ubelaker DH. 1994. *Standards for Data Collection from Human Skeletal Remains*. Fayetteville, Arkansas Archaeological Survey.
- Byers SN. 2002. *Introduction to forensic anthropology: a textbook*. Boston, MA: Allyn and Bacon Publishers.
- Carlson KJ, Grine FE., Pearson OM. 2007. Robusticity and sexual dimorphism in the postcranium of modern hunter-gatherers from Australia. *Am J Phys Anthropol* 134:9–23.
- Celbis O, Agritmis H. 2006. Estimation of stature and determination of sex from radial and ulnar bone length in a Turkish corpse sample. *Forensic Sci Int* 158:135–39.
- Cowal LS, Pastor RF. 2008. Dimensional variation in the proximal ulna: Evaluation of a metric method for sex assessment. *Am J Phys Anthropol* 135:469–78.
- Cowgill LW, Hager LD. 2007. Variation in the development of postcranial robusticity: an example of Çatalhöyük, Turkey. *Int J Osteoarch* 17:235–52.
- Ćwirko-Godycki M, Swedborg I. 1977. Ludność pochowana na cmentarzu Ostrowa Lednickiego pod względem metrycznym z uwzględnieniem zmienności cech oraz objawów patologicznych na kościach. *Anthropol Rev* 43:2–36.
- Dayal MR, Spocter MA, Bidmos MA. 2008. An assessment of sex using the skull of black South Africans by discriminant function analysis. *HOMO* 59:209–21.
- Faerman M, Bar-Gal GK, Filon D, Greenblatt CL, Stager L, Oppenheim A, et al. 1998. Determining the sex of infanticide victims from the late Roman era through ancient DNA analysis. *J Archaeol Sci* 25:861–65.
- Ferembach D, Schwidetzky I, Stloukal M. 1980. Recommendations for age and sex diagnoses of skeleton. *J Hum Evol* 9:517–49.
- Fuglewicz B. 2011. Studia nad początkami radomskiego zespołu osadniczego w dolinie rzeki Mlecznej. In: A Buko, D Główska, M Trzeciecki, editors. *Radom: korzenie miasta i regionu. Tom 2*. Warszawa: Wydawnictwo Instytutu Archeologii i Etnologii PAN. 9–33.
- González PN, Bernal V, Perez SI, Barrientos G. 2007. Analysis of dimorphic structures of the human pelvis its implications for sex estimation in samples without reference collections. *J Archaeol Sci* 34:1720–30.
- González-Reimers E, Velasco-Vazquez J, Arnay-de-la-Rosa M, Santolaria-Fernandez F. 2000. Sex determination by discriminant function analysis of the right tibia in the prehispanic population of the Canary Islands. *Forensic Sci Int* 108:165–72.
- Gonçalves D, Thompson TJU, Cunha E. 2015. Sexual dimorphism of the lateral angle of the internal auditory canal and its potential for sex estimation of burned hu-

- man skeletal remains. *Int J Legal Med* DOI 10.1007/s00414-015-1154-x.
- Gray J, Wolfe L. 1980. Height and sexual dimorphism of stature among human societies. *Am J Phys Anthropol* 53:441–56.
- Gustafsson A, Werdelin L, Tullberg BS, Lindenfors P. 2007. Stature and sexual stature dimorphism in Sweden, from the 10th to the end of the 20th century. *Am J Hum Biol* 19: 861–70.
- Holden C, Mace R. 1999. Sexual dimorphism in stature and women's work: a phylogenetic cross-cultural analysis. *Am J Phys Anthropol* 110:27–45.
- Howells WW. 1996. Howells' craniometric data on the internet. *Am J Phys Anthropol* 101:441–42.
- Kamdi A, Gayatri Shereke AR, Krishnaiah M, Chaitanya K. 2014. Morphometric parameters and sex determination of clavicle in Telangana region. *J Dent Med Sci* 13:1–5.
- Konigsberg L, Hens S. 1998. Use of ordinal categorical variables in skeletal assessment of sex from the cranium. *Am J Phys Anthropol* 107:97–112.
- Kranioti EF, Michalodimitrakis M. 2009. Sexual dimorphism of the humerus in contemporary Cretans – a population-specific study and a review of the literature. *J Forensic Sci* 54:996–1000.
- Kranioti EF, Tzanakis N. 2015. Estimation of sex from the upper limb in modern Cretans with the Aid of ROC-Analysis: A Technical Report. *FRCIJ* 1:1–7. DOI: 10.15406/frcij.2015.01.00008.
- Krogman WM, Iscan MY. 1986. *The human skeleton in forensic medicine*. Springfield, IL: Charles C. Thomas.
- Kubicka AM, Piontek J. 2016. Sex estimation from measurements of the first rib in a contemporary Polish population. *Int J Legal Med* 130:265–72.
- Lee J-H, Kim Y-S, Lee U-Y, Park D-K, Jeong Y-G, Lee NS, et al. 2014. Sex determination using upper limb bones in Korean populations. *Anat Cell Biol* 47:196–201.
- Liu W. 1989. Sex discriminant analysis of long bones of upper limb. *Acta Anthropol Sinica* 8:231–39.
- Loth SR, İşcan MY. 2000. Sex determination. In: J Siegal, P Saukko, G Knupfer, editors. *Encyclopedia of Forensic Sciences*. London: Academic Press. 242–52.
- Mall G, Hubig M, Buttner A, Kuznik J, Penning R, Graw M. 2001. Sex determination and estimation of stature from the long bones of the arm. *Forensic Sci Int* 117:23–30.
- Martin R. 1957. *Lehrbuch der Anthropologie*. Stuttgart.
- Mastrangelo P, De Luca S, Sánchez-Mejorada G. 2011. Sex assessment from carpal bones: discriminant function analysis in a contemporary Mexican sample. *Forensic Sci Int* 209:1–15.
- Murphy AMC. 1994. Sex determination of prehistoric New Zealand Polynesian clavicles. *New Zealand J Archaeol* 16:85–91.
- Nyati LH, Norris SA, Cameron N, Pettifor JM. 2006. Effect of ethnicity and sex on the growth of the axial and appendicular skeleton of children living in a developing country. *Am J Phys Anthropol* 130:135–41.
- Otong ES, Rufai AA, Amaza DS, Mustapha Z, Jacks TW., Attah M. 2016. Estimation of sex using antero-posterior dimension of talus in North Eastern Nigerians. *Br J Med Res* 14(3):1–8.
- Patil G, Kolagi S, Ramadurg U. 2011. Sexual dimorphism in the humerus: a study on South Indians. *J Clin Diagnost Res* 5:538–41.
- Peckmann TR, Orr K, Meek S, Manolis SK. 2015. Sex determination from the talus in a contemporary Greek population using discriminant function analysis. *J Forensic Leg Med* 33:14–19.
- Phenice TW. 1969. A newly developed visual method of sexing os pubis. *Am J Phys Anthropol* 30:297–01.
- Piątkowski S. 2000. *Radom – zarys dziejów miasta*. Radom: Wydawnictwo Radom.
- Piontek J. 1969. Early Medieval human remains from ancient burial grounds in Strzelno, district Mogilno. *Anthropol Rev* 35:317–21.

- Psonak D, Kwiatkowska B. 2012. Selection of skull morphological traits for sexual dimorphism analysis. *HOMO* 63:258–74.
- R Development Core Team. 2013. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria, <http://www.R-project.org>.
- Ray LJ. 1959. Metrical and non-metrical features of the clavicle of the Australian Aboriginal. *Am J Phys Anthropol* 17:217–26.
- Robinson MS, Bidmos MA. 2011. An assessment of the accuracy of discriminant function equations for sex determination of the femur and tibia from a South African population. *Forensic Sci Int* 206(1-3):212 e1-5.
- Rösing FW, Graw M, Marré B, Ritz-Timme S, Rothschild MA, Rötzscher K, et al. 2007. Recommendation for the forensic diagnosis of sex and age from skeletons. *HOMO* 58:75–89.
- Ruff C. 1987. Sexual dimorphism in human lower limb bone structure: relationship to subsistence strategy and sexual division of labor. *J Human Evol* 16:391–416.
- Ruff CB, Hayes WC. 1988. Sex differences in age-related remodeling of the femur and tibia. *J Orthop Res* 6:886–96.
- Ruff C. 2003. Growth in bone strength, body size, and muscle size in a juvenile longitudinal sample. *Bone* 33:317–29.
- Sakaue K. 2004. Sexual determination of long bones in recent Japanese. *Anthropol Sci* 112:75–81.
- Santos FR, Pandya A, Tyler-Smith C. 1998. Reliability of DNA-based sex test. *Nat Genet* 18:103.
- Shirley NR. 2009. Age and sex estimation from the human clavicle: an investigation of traditional and novel methods. Ph.D. thesis, Knoxville: The University of Tennessee.
- Spradley MK, Jantz RL. 2011. Sex estimation in forensic anthropology: skull versus postcranial elements. *J Forensic Sci* 56:289–96.
- Steyn M, İşcan MY. 1999. Osteometric variation in the humerus: sexual dimorphism in South Africans. *Forensic Sci Int* 106:77–85.
- Stini WA. 1969. Nutritional stress and growth: sex difference in adaptive response. *Am J Phys Anthropol* 31:417–26.
- Stinson S. 1985. Sex differences in environmental sensitivity during growth and development. *Yearb Phys Anthropol* 28:123–47.
- Thakur SB, Deshmukh A, Joshi DS. 2015. Determination of sex from humerus in Marathawada Region. *Int J Res Trend Sci Technol* 15: 220–22.
- Volk C, Ubelaker DH. 2002. A test of the Phenice method for the estimation of sex. *J Forensic Sci* 47:19–24.
- Walker P. 2008. Sexing skulls using discriminant function analysis of visually assessed traits. *Am J Phys Anthropol* 136:39–50.
- White TD, Folkens PA. 2000. *Human osteology*. San Diego: Academic Press.
- Zapłata R. 2011. Wstępne wyniki badań archeologicznych w wykopie 1/2010 na stanowisku 1 „Majdan” i w wykopie I-2/2010 na stanowisku 2 „osada” w Radomiu. In: A Buko, D Główska, M Trzeciecki, editors. *Radom: korzenie miasta i regionu, Tom 2*. Warszawa: Wyd. Instytutu Archeologii i Etnologii PAN 89–112.