

EFFICACY OF PHYTOGENIC FEED ADDITIVE ON PERFORMANCE, PRODUCTION AND HEALTH STATUS OF MONOGASTRIC ANIMALS -**A REVIEW**

Santi Devi Upadhaya, In Ho Kim*

Department of Animal Resource and Science, Dankook University, No. 29 Anseodong, Cheonan, Choongnam 330-714 South Korea *Corresponding author: inhokim@dankook.ac.kr

Abstract

The growing concerns of consumers on the use of antibiotic as a growth promoter in livestock feed have fueled the interest in alternative products. In the recent years a group of natural products known as phytogenics has been a focus of several studies. Phytogenics are a heterogeneous group of feed additives originating from plants and consist of herbs, spices, fruit, and other plant parts. These feed additives are reported to have a wide range of activities including antimicrobial, anthelminthic, antioxidant, growth enhancer, and immune modulator. Besides these properties they are also reported to stimulate feed intake and endogenous secretion and enhance production. They include many different bio-active ingredients such as alkaloids, bitters, flavonoids, glycosides, mucilage, saponins, tannins phenolics, polyphenols, terpenoids, polypeptide, thymol, cineole, linalool, anethole, allicin, capsaicin, allylisothiocyanate, and piperine. These feed additives have been tested in the form of extracts, cold pressed oils, essential oils in a number of animals but the results are variable. Therefore, their application as feed additive has been limited, largely owing to their inconsistent efficacy and lack of full understanding of the modes of action. The future of these feed additives depend on the characteristics of herbs, the knowledge on their major and minor constituents, the in-depth knowledge on their mode of action and their value based on the safety to animal and their products. The aim of this review is to summarize on the current knowledge on the use of phytogenic as a feed additive in monogastric animals.

Key words: feed additive, monogastric animal, phytogenic

The use of antibiotic growth promoters (AGP) in livestock feed has been practiced for many decades. There were serious concerns on the resistance issue of antibiotics based on different research studies. The resistance to antimicrobial agents as a serious threat to human health was only emphasized in the 1990s (Neu, 1992; Kunin, 1993) due to the fact that there were great public concerns on the usage of antibiotics as AGP in food animals. This led to the ban on most of the antibiotic feed additives in the European Union in 1999. Eventually, its use as a growth promoter

in livestock feed has been fully banned in the European Union since January 2006 (Regulation 1831/2003/EC). The concerns about development of antibiotic resistance and the ban of antibiotics as AGP have initiated the surge of exploring alternatives to antibiotics with similar antimicrobial and growth-promoting effects. Such feed additive is expected not to induce resistance to bacteria and to have no potential side effects to animals. In the recent years, some feed additives such as probiotics (Musa et al., 2009), prebiotics (Gibson et al., 2004), organic acids (De Lange et al., 2010; Upadhaya et al., 2014, 2016 a), enzymes (Bedford and Cowieson, 2012) and phytogenics (Randrianarivelo et al., 2010; Gong et al., 2014; Dhama et al., 2015) are used as a replacement for AGP. Phytogenic feed additive has been reported to enhance performance, feed conversion ratio, carcass meat safety and quality in animals (Stanacev et al., 2011; Dhama et al., 2014, 2015). Besides enhancing performance, phytogenic also has anti-oxidant property, the effects of which are associated with essential oils (EOs) and their components (Lee and Shibamoto, 2001; Alagawamy et al., 2016). Phytogenic also has beneficial effects on nutrient utilization possibly by stimulating digestive enzymes such as lipase, amylase, or protease (Platel and Srinivasan, 2004) and improves gastrointestinal morphology (Jamroz et al., 2006; Upadhaya et al., 2016 b). Janz et al. (2007) indicated that oregano fed pigs had a tendency of lower oxidation in meat. Phytogenic supplementation may have a positive impact on nutrient digestibility (Hernandez et al., 2004). The supplementation of blends of EOs also demonstrated the stable egg yield and reproductive performance compared with that of an AGP (Bozkurt et al., 2009).

However, the lipophilic nature of phytogenic compounds limits the efficient delivery of these compounds to the gut. This problem could be resolved by microencapsulation and combination with other compounds (Yang et al., 2015). A study by Hafeez et al. (2016) demonstrated that matrix encapsulation of essential oils with active ingredients such as carvacrol, thymol and limonene at the dosage of 100 mg/kg feed, improved performance and apparent ideal digestibility of nutrients compared with control in broiler chickens.

Considering this collective information, this study sought to provide an overview of current knowledge on the efficacy of phytogenic feed additive on performance, production and health status of monogastric animals based on literature review from different studies.

Characteristics of phytogenic feed additive

The phytogenic feed additives (PFA) derived from herbs, spices or aromatic plants (Windisch et al., 2008) are relatively young classes of feed additive that have gained considerable attention in the recent years in feed industry (Hashemipour et al., 2013; Dhama et al., 2015; Abd El-Hack et al., 2016; Alagawany et al., 2016). These plant derived products are residue-free unlike synthetic antibiotics and are also generally considered safe to be used as the ingredients in the food industry as well as in animal diet as an ideal growth promoter (Varel, 2002; Hashemi et al., 2008; Li et al. 2016). The herbs and plant extracts used as feed additives include many different bio-active ingredients such as alkaloids, bitters, flavonoids, glycosides, mucilage, saponins, tannins phenolics, polyphenols, terpenoids, polypeptide, thymol, cineole,

linalool, anethole, allicin, capsaicin, allylisothiocyanate, and piperine (Wenk, 2006; Windisch et al., 2008; Grashorn, 2010; Kiczorowska et al., 2015; Al-Yasiry and Kiczorowska, 2016). Therefore, the effects expected of herbs and plant extracts are also various. Furthermore, the phytogenic feed additives have considerable variation in their chemical composition depending on their ingredients, influences of location, climatic conditions, harvest or storage conditions (Huyghebaert et al., 2011). Some other factors that influence the efficacy of the phytogenic mainly include the plant parts and their physical properties, the genetic variation of the plant, age of the plant, different dosage used, extraction method, harvest time, and compatibility with other ingredients (Yang et al., 2009). In addition, the beneficial effect of dietary phytogenic can be influenced by the nutritional status of animals, the infection, the diet composition and the environment condition (Giannenas et al., 2003).

The mode of action of most phytogenics is still not fully elucidated. Nevertheless, the antimicrobial, immunomodulatory, antioxidative, and growth-promoting effects in animals have been reported by various researchers (Windisch et al., 2008; Jacela et al., 2010; Abd El- Hack and Alagawany, 2015).

Phytogenic feed additive and animal health

Antimicrobial activity

The primary mode of action of PFA is the modulation of beneficial intestinal microbiota by controlling potential pathogens. The essential oils (EOs) present in PFA contain most of the bioactive substances of the plant which include carvacrol, eugenol, thymol, capsaicin, cineole and so on are well known for their antibacterial, antifungal, antiviral and anti-coccidial properties (Windisch et al., 2008; Giannenas and Kyriajakis, 2009; Dragan et al., 2014; Gopi et al., 2014; Oyuntsetseg et al., 2014; Farag et al., 2016; Patil and Patil, 2016). For instance, clove, which is rich in eugenol, is used as antibacterial in human and veterinary medicine (Rhayour et al., 2003). Likewise, the blend of EO of oregano and thyme (0.05%) with carvacrol and thymol as active compounds were reported to be effective against Salmonella located in the digestive tract of chicken (Koscova et al., 2006). In an in vitro study, cinnamaldehyde supplementation at the dose rate of 10 and 100 µg/ml reduced the viability of Eimeria tenella parasites as compared with control (Lee et al., 2011). The reduction in fecal E. coli count was also observed in our previous study when 1 g/kg of eugenol was supplemented in growing pigs' diet (Yan and Kim, 2012) and reduction in the proliferation of Clostridium perfringens and E. coli were also detected in the small and large intestine of *Clostridium perfrigens* challenged broilers when PFA containing 250 mg/kg EOs of thyme and star anise was added to the diet (Cho et al., 2014). In another challenge experiment, Wati et al. (2015) demonstrated that when Salmonella enteritidis and E.coli challenged broilers were fed diet containing 150 mg/kg PFA (fennel, lemon balm, peppermint. anise, oak, clove and thyme), lower counts of Salmonella, E. coli and Clostridium and higher counts of Lactobacillus were observed. The antimicrobial mode of action is supposed to be due to the hydrophobicity of these phenolic compounds and due to the fact that these compounds can enter the bacterial cell membrane leading to disintegration of cell membrane, leakage of ions and eventually cell death (Burt, 2004). However, limonene, a non-phenolic

compound is also reported to exhibit anti-bacterial effect (Burt, 2004). The antibacterial activity of EO from spices such as cinnamon, clove, garlic, mustard, onion and oregano was found to be slightly more against gram-positive than gram-negative bacteria (Ceylan and Fung, 2004). The effect of blends of EO consisting of clove, coriander and curcuma (50-500 ppm) on cell integrity, as measured using propidium iodide for gram-positive bacteria, was found to be dose dependent and the growth inhibition of gram-negative bacteria, in contrast, occurred mostly without cell integrity loss (Thapa et al., 2012). The additive effects of EOs such as thymol and carvacrol with organic acids such as lactic acid, citric and acetic acid or oregano essential oil with citric acid have been reported to reduce pathogenic bacteria thereby producing adequate antimicrobial effects (Zhou et al., 2007; Hulankova and Borilova, 2011). Our previous study also showed that when 0.2% Korean pine extract along with 2% citric acid was supplemented to the weaning pigs' diet, it significantly increased fecal lactobacillus counts (Zhang et al., 2012). Some other studies reported that blends of Eos (10 g/kg herb such as thyme, oregano, marjoram, rosemary or yarrow or 1 g/kg of essential oil) and (0.05%, 0.1% and 0.15% EOs of thymol and carvacrol) had no effect on the microbial population and composition in the digestive tract or fecal excretions of broilers (Cross et al., 2007) and pigs (Muhl and Liebert, 2007 a). In their review, Brenes and Roura (2010) suggested that minor components in EO may have a critical role as the bacteriostatic activity and may have synergistic effects. The EOs from peppermint and eucalyptus (250 ml/L drinking water) has been reported to inhibit respiratory pathogens such as Mycoplasma gallisepticum and protect against H9N2 virus infection (Barbour et al., 2011). Likewise, administration of eucalyptus and peppermint oil blends via oral (0.25 ml/L drinking water for 12 hours/day) and spray route (0.1 ml/20 ml water/10 birds) reduced Newcastle disease infection in broilers (Awaad et al., 2016). The findings of different studies warrant more detailed investigation to develop a blend of EOs having better antimicrobial properties.

Impact on animal immunity

For the alleviation of diseases, modulation of immune response has been of great interest to researchers (Ozek et al., 2011; Mahima et al., 2012). Nuclear factor "kappa light-chain-enhancer" of activated B cells (NFKB) represents a key transcription factor regulating the synthesis of genes involved in immune reactions and inflammatory response (Ghosh et al., 1998). Several herbs and EOs have been reported as having immunomodulatory effects such as lymphocyte expression, phagocytosis, modulation of cytokine and immunoglobulin secretion, histamine release and so on (Mahima et al., 2012). A blend of oregano, anise, and citrus peel (40 mg/kg diet) supplementation to piglets' diet has been demonstrated to evolve anti-inflammatory effect by reducing the gene expression of NFkB and TNF alpha (Kroismayr et al., 2008). The supplementation of poultry feed with anise was reported to improve lymphocyte counts (Soltan et al., 2008). The increase in IgG in broilers was noted with the inclusion of 0.1% of herb mixture consisting of Phlomis umbrosa Turez, Cynancum wilfordii Hem, Zingiber officinale Rosc and Platycodi Radix in broiler (Begum et al., 2014). Yan and Kim (2012) also indicated that eugenol and cinnamaldehyde (1 g/kg) exhibited lymphocyte enhancing activity in growing pigs. A study by Li et

al. (2012 a) demonstrated that when 0.01% of EO product containing 18% thymol and cinnmaldehyde was supplemented to weaning pigs' diet, it reduced inflammatory mediator, interleukin-6 concentration in the plasma indicating the positive effect of PFA on animal health. Administration of encapsulated EOs containing thymol and cinnamaldehyde (50, 100 or 150 g/ton of feed) has been reported to confer positive immune effects such as increase in lymphocyte proliferation rate, phagocytic rate as well as increase in immunoglobulins such as IgA and IgM in the blood of weaning pigs (Li et al., 2012 b). Some studies reported that when 10 or 20 g plant extract per liter water from herb such as Withania somnifera having bioactive compound withanolides (a steroidal lactone) was administered to broiler chick diets, it improved hemoglobin, packed cell volume, white blood cells count and antibody titre against viral disease such as infectious bursal disease and infectious bronchitis suggesting the improvement in hematological profile and immunological status of birds (Mushtaq et al., 2012; Pant et al., 2012). Studies conducted on the immunomodulatory potential of the herbs such as W. somnifera, T. cordifolia and A. indica against chicken infectious anemia, an immunosuppressive viral disease of young chicks, revealed that these herbs have excellent capacity in stimulating both the cellular and humoral immune responses in chicks against the causative virus and also ameliorating effects on viral pathogenesis were observed (Latheef et al., 2013). In another study, Alhajj et al. (2015) reported that inclusion of 6 g anise seed per kg diet showed higher antibody titer against Newcastle disease virus and infectious bronchitis in broiler chickens. The precise mechanism of phytogenic on immunomodulation is still not fully clear and therefore further studies are needed to clarify the role of PFA on immune modulation

Impact on gastrointestinal morphology and nutrient digestibility

Phytogenic feed additives originating from different herbs are known to have laxative, spasmolytic and anti-flatulence effects in gastro-intestinal tract. Essential oils such as 0.01% thymol and cinnamaldehyde (Li et al., 2012 a) and PFA such as menthol or cinnamaldehyde (Maenner et al., 2011) have been reported to positively affect nutrient digestibility in swine. Likewise, feeding broilers a diet supplemented with 200 mg/kg EO from peppermint led to the increase of crude protein digestibility (Emami et al., 2012). A study by Ahmed (2013) reported that the supplementation of blends of oregano, anise, orange peel and chicory essential oils (0.0125%) in the diet of weaned pig improved protein digestibility. Beneficial effects on nutrient digestibility using different PFA in some of our previous studies as summarized in Table 1 have also been observed in pigs and poultry (Yan et al., 2010; 2011 a, 2012 a, 2012 b; Zhang et al., 2012; Gheisar et al., 2015; Upadhaya et al., 2016 b). The reason for improvement in nutrient absorption may be partly explained due to stimulation in secretions of saliva, bile and enhanced enzyme activity (Platel and Srinivasan, 2004; Jang et al., 2007). The improved nutrient digestibility consequently enhances the health status of animals. In contrast, Muhl and Liebert (2007 b) failed to observe improved nutrient digestibility and enzymatic activity in weaned piglets fed diets containing a 0.05%, 0.10% or 0.15% phytogenic product having carvacrol, thymol and tannins as key constituents. The inconsistent results in apparent digestibility may be caused

L	able 1. Re:	sponse of phytogenic additive	by non-ruminant a	nimals as	per the re:	search stu	udies conducted at Dankook University	
				Overall g	rowth per.	formance		
Feed additive	Dose	Major compounds	Species		response		Other responses	Reference
				ADG	FI	G:F		
1	2	3	4	5	9	7	8	6
Essential oil blend	0.1 mg/k	g Thyme, rosemary, oregano	Growing-finish- ing pigs	Yes	N/A	N/A	Positive effect on nitrogen and energy digestibility, Improved longissimus area	Yan et al., 2010
Herb extract	1 g/kg	Saurus chinensis	Finishing pigs	N/A	N/A	N/A		Ao et al., 2011
	2 g/kg			N/A	N/A	N/A	Serum triglyceride and LDL/C were reduced, Ammonia levels reduced in slurry	
Herb extract	1 g/kg	Houttuynia cordata	Finishing pigs	Yes	Yes	N/A	Improved meat quality	Yan et al., 2011 a
	1 g/kg	Taraxacum officinale		N/A	Yes	N/A	Improved nitrogen digestibility, Im- proved longissimus muscle area	
Herb extract mixture	250 mg/k _i	g Buckwheat, thyme, curcuma black pepper and ginger	, Growing pigs	Yes	Yes	N/A	Both doses increased WBC and reduced fecal NH $_3$ and H $_2$ S gas emission	
	500 mg/ki	۵۵		Yes	Yes	N/A		Yan et al., 2011 b
Herb	1 g/kg	Date pits	Weaning pigs	N/A	N/A	N/A		Cho et al., 2012
	0.5 g/kg	Japenese honeysuckle		Yes	N/A	N/A		
	1 g/kg	Houttuynia cordata		Yes	N/A	N/A		
	1 g/kg	Laquer tree extract		Yes	N/A	N/A		
	1 g/kg	Yellow ginger	Growing pigs	Yes	N/A	N/A	Improved meat quality	
	1 g/kg	Hoantchy root		Yes	N/A	N/A		
Essential oil blend	1 g/kg	Eugenol	Growing pigs	N/A	N/A	N/A	Reduced fecal <i>E. coli</i> and reduced fe- cal NH ₃ and H ₂ S gas emission, higher lymphocyte concentration	Yang and Kim, 2012
	1 g/kg	Cinnalmaldehyde		N/A	N/A	N/A		
Herb extract mixture	250 mg/k _i	g Buckwheat, thyme, curcuma black pepper and ginger	, Weaning pigs	N/A	N/A	N/A	Improved nutrient digestibility and reduced fecal E. coli population	Yan et al., 2012 a

934

Herb extract	1 g/kg	Houttuynia cordata	Weaning pigs	N/A	N/A	N/A	Reduced fecal E. coli	Yan et al., 2012 b
	1 g/kg	Taraxacum officinale		Yes	N/A	Yes	Improved energy digestibility and reduced fecal <i>E. coli</i> counts	
Herb extract	2 g/kg	Korean pine (flavonoids and phenolic compounds)	Weaning pigs	N/A	N/A	N/A	Improved DM digestibility	Zhang et al., 2012
Herb extract	0.5 mg/kg	Coptis chinensis (berberine)	Finishing pigs	N/A	N/A	N/A		Zhou et al., 2013
	1 mg/kg			N/A	N/A	N/A	Improved meat quality	
Herb extract	0.5 mg/kg	Evening primrose	Weaning pigs					Lei et al., 2014
	1 mg/kg							
	2 mg/kg			Yes	N/A	Yes	Increased lactobacillus counts and reduced LDL/cholesterol, Increased DM digestibility	
Gel based additive	50 g/pig	Caraway (limonene and carvone)	Weaning pigs	Yes	N/A	Yes	Improved digestibility of nutrients and increased ileum and jejunum villi	Upadhaya et al., 2016
	100 g/pig			Yes	N/A	Yes	length	
Essential oil, herb and spices mix	250 mg/kg	Thyme and star anise	Broilers	N/A	N/A	Yes	Reduced serum total cholesterol, Inhibited <i>Clostridium</i> spp. and <i>E. coli</i> in GIT	Cho et al., 2014
Phytogenic blend	151 ppm	Quillaja, anise and thyme	Ducks	Yes	N/A	Yes	Improved nutrient digestibility and reduced drip loss and cooking loss	Gheisar et al., 2015
Fermented plants	0.5 mg/kg	Gynura procumbens, Rehmannia glutinosa, Scutellaria baicalensis fermented by L. plantarum, S. cerevisiae and B. licheni- formis respectively	Broilers	Yes	N/A	Yes	Linear reduction in ammonia gas emis- sion from excreta, Improved dry matter and nitrogen retention	Jeong and Kim, 2015
	1 mg/kg			Yes	N/A	Yes		
	2 mg/kg			Yes	N/A	Yes		

Use of phytogenics as a feed additive for monogastric animals

	6	Li et al., 2015		Begum et al., 2014		
	8	Reduced excreta gases emission			Improved IgG levels and reduced abdominal fat	
	7	Yes		N/A	N/A	N/A
	9	Yes		N/A	N/A	N/A
ole 1 contd.	5	Yes		N/A	Yes	N/A
Tal	4	Broilers		Broilers		
	3	Korean pine (Flavonoids and phenolic compounds)		Phlomis umbrosa Turez, Cynancum wilfordii Hem, Zingiber officinale Rosc and Platycodi Radix		
	2	0.5 mg/kg	1 mg/kg	0.5 mg/kg	1 mg/kg	1.5 mg/kg
	1	Herb extract		Herb extract		

by endogenous loss from mucus secretion induced by plant extracts (Jamroz et al., 2006). Besides imparting positive health benefit, the improved nutrient absorption may reduce feed cost by allowing appropriate modifications to nutrient density diets. Zeng et al. (2015) investigated the acceptance of commercial EO product at the dosage of 0.025% consisting of 4.5% cinnamaldehyde and 13.5% thymol in low energy density in weaned pig diets with wheat and extruded full-fat soybean as the major ingredients. The supplementation of EOs improved the apparent digestibility of dry matter; crude protein and energy compared with pigs fed the low energy density diet without EO indicating PFA may contribute to the reduction of feed cost. The ability of epithelial cells to regenerate villus can be improved though decreased numbers of pathogenic bacteria in the gut thereby enhancing intestinal absorptive capacity (Mourao et al., 2006). Inclusion of blend of plant extracts from sage, lemon, balm, nettle and coneflower (20, 30, 30, and 20%, respectively) at 500 mg/kg feed has been reported to improve the villus height in weaning pigs (Hanczakowska and Swiatkiewicz, 2012). The inclusion of 100 and 200 mg/kg thymol and carvacrol in broiler chickens' diet improved villus surface, villus height, villus height to crypt depth and muscular layer of jejunum and ileum (Hashemipour et al., 2013). The addition of Euphorbia hirta (7.5 g/kg) increased the villus height compared to the control birds (Hashemi et al., 2014). Our recent study about gel based feed additive (50 or 100 g/pig/day) supplementation to weaned pig with essential oils from caraway seed as main ingredient improved the villus length of ileum and jejunum (Upadhaya et al., 2016 b). The dietary supplementation with 2.0 and 2.5% of Boswellia serrata resin to broiler led to a significant increase in the length of the duodenum and total intestine (Kiczorowska et al., 2016). Such an effect of PFA is expected due to their well-documented inhibitory effects against pathogens. However, the use of PFA in relation to gut morphology is not consistent. For instance, Manzanilla et al. (2004) reported that weaning pigs under stress fed plant extract mixture (150 or 300 mg/kg) consisting of 5% carvacrol, 3% cinnamaldehyde and 2% capsicum oleoresin reduced villus length. Likewise, Oetting et al. (2006) indicated birds fed 700 ppm, 1,400 ppm and 2,100 ppm of herbal extract consisting of a mixture of essential oils of thyme, clove, oregano, eugenol and carvacrol did not have any significant change in gastrointestinal morphology. The negative or no influence of phytogenic in gut morphology was hypothesized due to irritation of intestinal tissues (Windisch et al., 2009).

Impact of phytogenic feed additive on animal performance

Several studies documented the use of PFA as a growth promoter (Srinivasan, 2006; Windisch, et al., 2008; Jacela et al., 2010; Kim et al., 2012; Elagib et al., 2013; Bartos et al., 2016; Li et al., 2016). Pastorelli et al. (2012) noted that 10 mg/ kg of *Lippia citriodora* led to faster growth in finishing pigs. However, Bruno et al. (2013) noted that there was no effect of 2000 ppm of herb extract consisting of *Rosmarinus officinalis, Mentha piperita, Lippia sidoides* and *Porophyllum ruderale* on growth performance in weaning pigs. Garlic, thyme and cornflower as feed additives in broilers have been reported to improve growth performance (Aji et al., 2011; Khan et al., 2011; Rahimi et al., 2011). The body weight was significantly improved when 50 or 100 mg of garlic and onion was administered to the diet of broiler chickens (Aji

et al., 2011). The supplementation of fenugreek seeds (1, 2 and 3%) significantly improved feed conversion ratio of broiler chickens (Mamoun et al., 2014). The dietary supplementation of 0.5 g/kg anise seed containing anothole as active ingredient significantly improved body weight gain and performance index but had no significant effect on feed intake and feed conversion ratio in broilers; moreover, a higher level of anise seed (1.5 g/kg) reduced growth performance compared with control (Soltan et al., 2008). Another study reported that supplementation of 1 or 2 g of anise seed in broilers diet improved body weight, daily weight gain and feed conversion ratio but had no effect on feed intake (Alhajj et al., 2015). The studies on PFA at Dankook University from 2010 to 2016 by our research groups have shown that the effects of different kinds of phytogenic additive on growth performances by swine and poultry were variable as presented in Table 1. The physiological difference in the digestive tract, the origin of the EOs or herb species, the dose of phytogenic supplemented to the feed and the environmental conditions used in the trial might have caused the difference in the results. Dietary supplementation of phytogenic compounds has the potential to increase feed intake by improving the palatability of diet possibly due to enhanced flavor and odor, especially with the use of EOs (Kroismay et al., 2006). The stability of EOs during feed processing has to be taken into consideration because processing may lead to loss of activity by the active ingredients present in the phytogenic compound. A considerable loss of activity of EOs has been reported when pelleting temperature of 58°C was applied (Maenner et al., 2011). Addition of a blend of EO from basil, caraway, laurel, lemon, oregano, sage, tea and thyme (100 g/t) in broiler diet has been demonstrated to improve the body weight gain with positive effects on feed to gain ratio (Khattak et al., 2014). The reason for increase in voluntary feed intake resulting in improved weight gain was reported to be due to the improvement in the flavor and palatability of feed by the aromatic EOs. However, feed intake was significantly reduced when feed was supplemented with fennel (100 mg/kg) or caraway oil (100 mg/kg) indicating that the palatability was reduced due to flavor in a choice feed experiment conducted in growing pigs by Schone et al. (2006). Our recent study by Upadhaya et al. (2016 b) on phytogenic feed additive (50 or 100 g/pig/day) containing caraway oil showed no significant effect on feed intake in weaning pigs. A study by Czech et al. (2009) indicated that inclusion of herbal extract (0.8 g/kg feed) from garlic bulbs, common liquorice roots and tillers, common thyme herb and caraway fruits stimulates growth performance and exerts beneficial actions in the gastro-intestinal tract by increasing the activity of digestive enzymes. Therefore, the properties of improved flavor and palatability are variable depending on the type and dose of PFA used. Akthar et al. (2003) reported that the inclusion of black cumin seeds into laying hen diets at a level of 1.5% led to a 12% increase of laying rate from 40 to 52 weeks of age. Consistently, diets supplemented with 3% black cumin seeds increased egg production and egg weight of 27-week-old laying hens during a 5-week experimental period in contrast to the supplementation of 1% (Aydin et al., 2008). On the other hand, Bolukbaşi et al. (2009) observed no improvement of laying hen performance with respect to black cumin seed oil (1, 2 or 3 ml/kg diet) supplementation. No effect was observed on laying rate and weight of settable eggs in response to an EO mixture (consisting of EO from oregano, laurel,

sage, myrtle, fennel and citrus) supplementation at 24, 36 or 48 mg/kg (Bozkurt et al., 2009). Likewise, Bolukbasi et al. (2008) observed no effect on laying rate but an increase in egg weight when diets were supplemented with 200 mg/kg EO of thyme, sage or rosemary over a period of 12 weeks. The supplementation of herb blends consisting of 0.05% garlic powder, 0.3% cinnamon powder and 0.03% of each of the following dried herbs: yarrow, rosemary, thyme, basil and oregano improved egg laying capacity of hen by 1.79% as compared with control (Gerzilov et al., 2015). The beneficial influence of the PFA on improved performance and feed conversion ratio could be also explained due to the antioxidant activity of bioactive compounds such as carvacrol, thymol, cineol and pinene (Faleiro et al., 2005; Hazzit et al., 2006) as well as from improved enzyme activity in the alimentary tract, stimulation of useful and inhibition of pathogenic microflora which eventually resulted in improved absorption and utilization of nutrients (Windisch et al., 2008; Frankic et al., 2009).

Other health benefits of phytogenic feed additive

Besides, immune enhancing, antimicrobial, and performance enhancing effects, phytogenics also have anti-oxidant property. The excellent plant derived antioxidants are obtained from rosemary, olive leaves, thyme, marjoram, sage, oregano, etc (Botsoglou et al., 2005, 2013; Rahal et al., 2014). Some other common herbs, spices and fruits that have antioxidant property are ginger, turmeric, garlic, plum, pine bark extract, berries, pomegranate, caraway, cinnamon, clove etc (Khan et al., 2012; Zhao et al., 2011; Botsoglou et al., 2013), the effects of which are associated with EOs and their components (Lee and Shibamoto, 2001; Miura et al., 2002). The demand for natural antioxidants in food is increasing due to their health benefits against oxidative stress and several diseases (Labaque et al., 2013; Perez-Barron et al., 2015; Juadjur et al., 2015; Kim et al., 2015). For instance, leaf powder from herbs such as Ocimum sanctum and Withania somnifera at the dose rate of 0.1% has been demonstrated to reverse the cadmium-induced oxidative stress in chicken due to the active ingredients (withanolides, methyl eugenol and flavonoids) of these herbs possessing antioxidants effect (Bharavi et al., 2010) and by neutralization of superoxide, hydrogen peroxide and nitric oxide by scavenging radicals or due to increased production of catalase, superoxide dismutase and glutathione peroxidase (Ali et al., 2006; Yarru et al., 2009). The supplementation of thymol (80 mg/animal/day) helped to reduce fear responses in quail when exposed to stressful situations (Labaque et al., 2013). A study by Pastorelli et al. (2012) demonstrated that the supplementation of Lippia citriodora containing verbascoside as active constituent at the dose rate of 10 mg/ kg for 56 days led to higher levels of serum IgA and lower levels of reactive oxygen metabolites in fattening pigs.

Impact of phytogenic FA on meat and egg quality

Meat quality

Plant oils used as feedstuffs in monogastric nutrition may readily alter fatty acid composition of all body lipid fractions by generally increasing the amount of polyunsaturated fatty acid and, therefore, their susceptibility to oxidation leading to the development of oxidative rancidity and associated increase in unpleasant odours. Concomitantly, plant oils usually contain natural antioxidants, which may contribute to an improved oxidative stability of meat and meat products, compensating for the increased degree of unsaturation. Herbs such as oregano and sage have been extensively studied for their antioxidant activity (Brenes and Roura, 2010). The supplementation of plant extract mix containing the combination of oregano essential oil and sweet chestnut wood extract (0.2%) to pigs' diet was reported to reduce the lipid oxidation of meat (Ranucci et al., 2014). The oxidative stability of meat obtained from broilers, hens or turkeys in a series of studies have been reported to increase with the use of dietary supplementation of EOs (Florou-Paneri et al., 2006; Giannenas et al., 2005). For instance, dietary supplementation of 100 mg/kg EO blends with 5% carvacrol, 3% cinnamaldehyde and 2% capsicum oleoresin as active constituents improved the concentration of anti-oxidants in the liver of broiler chicken (Karadas et al., 2014). Likewise, a study by Ghazaghi et al. (2014) noted that supplementation of Mentha spicata (1-4%) in the diet improved meat quality of Japanese quail. Al-Kassie et al. (2014) reported that the inclusion of mixture of hot red pepper and black pepper at a level of 0.75 and 1% in the diets significantly improved the dressing percentage of broilers. Our studies with some of the PFA have shown no adverse effect in meat quality in finishing pigs (Li et al., 2016) and ducks (Gheiser et al., 2015) or positive effect in meat quality aspects of pigs which could be attributed to antioxidative properties of PFA (Yan et al., 2010, 2011 a; Zhou et al., 2013). On the contrary, Simitzis et al. (2010) reported that dietary oregano EO (0.25, 0.5 and 1 ml/kg of fed diet) failed to improve the lipid oxidation status of pork. The inconsistency in findings may be contributed by the different fatty acid composition in the meat of poultry and swine. Although poultry meat contains low lipid content, its relative concentration of polyunsaturated fatty acids is higher than pork (Janz et al., 2007). Thus, poultry meat is highly susceptible to oxidative deterioration, which might contribute to an enhanced response on dietary EOs supplementation on the lipid oxidation status of poultry meat.

Egg quality

The study on the effects of PFA on egg quality is limited and variable. In a previous study, Bolukbaşi et al. (2008) reported that supplementing 200 mg/kg of thyme, sage or rosemary EO increased the proportion of egg shell. However, in another study, no benefit was determined in relative shell weight with respect to dietary supplementation with three different levels (1, 2 or 3 ml/kg) of black cumin EO (Bolukbaşi et al., 2009). Similarly, EO of oregano (50 or 100 mg/kg) did not influence egg quality characteristics such as yolk color score, Haugh unit or shell thickness when supplemented to the diet at the age of 32 weeks (Florou-Paneri et al., 2005). In another study, Abdel-Wareth and Lohakare (2014) noted that the inclusion of peppermint (*Mentha piperita*) leaves (0, 5, 10, 15, or 20 g/kg for 12 weeks) in HyLine brown laying hens (64-week-old) led to significantly increased egg weight, egg production and egg mass. More studies are needed to verify the effects of PFA on egg quality characteristics.

Conclusion

The search for alternatives to antibiotics is gaining grounds in recent years. Phytogenic compounds represent one of the most promising alternatives to antibiotics because they consist of a large variety of active ingredients. However, their application in food animal production has been limited, largely owing to their inconsistent efficacy and lack of full understanding of the modes of action. Differences between or within PFA depend significantly on several variables, which makes it necessary to define the exact composition of the supplements which have been added to diets. In addition, minor components present are critical to the activity of PFA and may have a synergistic influence. Sometimes the minor components may counteract the exerted effects. Therefore, in the future, the detailed study on the constituents of PFA including EOs is needed so as to evaluate their different biological effects, their safety with regards to animal health and the quality of animal products. To make the best use of phytogenic substances for economically effective and sustainable animal production, a better understanding of the effects of phytogenic compounds and their mode of action is a must.

References

- A b d El-Hack M.E., A lagawany M. (2015). Performance, egg quality, blood profile, immune function, and antioxidant enzyme activities in laying hens fed diets with thyme powder. J. Anim. Feed Sci., 24: 127–133.
- Abd El-Hack M., Alagawany M., Farag M., Tiwari R., Karthik K., Dhama K., Zorriehzahra J., Adel M. (2016). Beneficial impacts of thymol essential oil on health and production of animals, fish and poultry: A review. J. Essen. Oil Res., 28: 365–382.
- A b d e l W a r e t h A.A.A., L o h a k a r e J.D. (2014). Effect of dietary supplementation of peppermint on performance, egg quality, and serum metabolic profile of Hy-Line Brown hens during the late laying period. Anim. Feed Sci. Technol., 197: 114–120.
- Ahmed S.T., Hossain M.E., Kim G.M., Hwang J.A., Ji H., Yang C.J. (2013). Effect of resveratrol and essential oils on growth performance, immunity, digestibility and fecal microbial shedding in challenged piglets. Asian-Australas. J. Anim. Sci., 26: 683–690.
- Aji S.B., Ignatuius K., Ado A.Y., Nuhu J.B., Abdulkarim A. (2011). Effect of feeding onion (*Allium cepa*) and garlic (*Allium sativum*) on some performance characteristics of broiler chickens. Res. J. Poult. Sci., 4: 22–27.
- A k h t a r M.S., N a s i r Z., A b i d A.R. (2003). Effect of feeding powdered *Nigella sativa* L. seeds on poultry egg production on their suitability for human consumption. Vet Arch., 73: 181–190.
- A l a g a w a n y M., A l i A s h o u r E., R e d a F.M. (2016). Effect of dietary supplementation of garlic (*Allium sativum*) and turmeric (*Curcuma longa*) on growth performance, carcass traits, blood profile and oxidative status in growing rabbit. Ann. Anim. Sci., 16: 489–505.
- Alhajj M.S., Alhobaishi M., Ger El Nabi A.R., Al-Mufarrej S.I. (2015). Immune responsiveness and performance of broiler chickens fed a diet supplemented with high levels of Chinese star anise fruit (*Illicium verum* Hook. F). J. Anim. Vet. Adv., 14: 36–42.
- Al-Kassie Galib A.M., Al-Nasrawi Mamdooh A.M., Ajeena Saba J. (2011). Use of black pepper (*Piper nigrum*) as feed additive in broilers diet. Roavs, 1: 169–173.
- Al Yasiry A.R.M., Kiczorowska B. (2016). Frankincense therapeutic properties. Postep. Hig. Med. Dosw., 70: 380–391.
- A o X., Yan L., Meng Q.W., Zhou T.X., Wang J.P., Kim H.J., Cho J.H., Kim I.H. (2011). Effects of *Saururus chinensis* extract supplementation on growth performance, meat quality and slurry noxious gas emission in finishing pigs. Livest. Sci., 138: 187–192.

- Applegate T.J., Klose V., Steiner T., Ganner A., Schatzmayr G. (2010). Probiotics and phytogenics for poultry: myth or reality? J. Appl. Poult. Res., 19: 194–210.
- Awaad M.H.H., Afify M.A.A., Zoulfekar S.A., Mohammed F.F., Elmenawy M.A., Hafez H.M. (2016). Modulating effect of peppermint and eucalyptus essential oils on vVND infected chickens. Pak. Vet. J., 36: 350–355.
- Ay din R., K ar a m an M., C i c e k T., Y ar d i b i H. (2008). Black cumin (*Nigella sativa* L.) supplementation into the diet of the laying hen positively influences egg yield parameters, shell quality and decreases egg cholesterol. Poultry Sci., 87: 2590–2595.
- Barbour E.K., Saade M., Abdel Nour A., Kayali G., Kidess S., Bou Ghannam R., Harakeh S., Shaib H. (2011). Evaluation of essential oils in the treatment of broilers coinfected with multiple respiratory etiologic agents. Int. J. App. Res. Vet. Med., 9: 317–323.
- Bartos P., Dolan A., Smutny L., Sístkova M., Celja I., Soch M., Havelka Z. (2016). Effects of phytogenic feed additives on growth performance and on ammonia and greenhouse gases emissions in growing-finishing pigs. Anim. Feed Sci. Technol., 212: 143–148.
- B e d f o r d M.R., C o w i e s o n A.J. (2012). Exogenous enzymes and their effects on intestinal microbiology. Anim. Feed Sci. Technol., 173: 76–85.
- B e g u m M., H o s a i n M.M., K i m I.H. (2014). Effects of the plant extract YGF251 on growth performance, meat quality, relative organ weight, nutrient digestibility and blood profiles in broiler chickens: possible role of insulin-like growth factor 1. Vet. Medicina, 59: 415–423.
- Bharavi K., Gopala Reddy A., Rao G.S., Ravikumar P., Rajasekhar Reddy A., Rama Rao S.V. (2010). Reversal of cadmium induced oxidative stress and its bio-accumulation by culinary herbs *Murraya koenigii* and *Allium sativum*. Res. J. Pharmacol., 4: 60–65.
- Bolukbasi S.C., Erhan M.K., Kaynar O. (2008). The effect of feeding thyme, sage and rosemary on laying hen performance, cholesterol and some proteins ratio of egg yolk and *Escherichia coli* count in feces. Arch Geflugelkd., 72: 231–237.
- Bolukbasi S.C., Kaynar O., Erhan M.K., Uruthan H. (2009). Effect of feeding *Nigella sativa* oil on laying hen performance, cholesterol and some proteins ratio of egg yolk and *Escherichia coli* count in feces. Arch Geflugelkd., 73: 167–172.
- Bozkurt M., Alcicek A., Çabuk M., Kucukyilmaz K., Çatli A.U. (2009). Effect of an herbal essential oil mixture on growth, laying traits, and egg hatching characteristics of broiler breeders. Poultry Sci., 88: 2368–2374.
- Brenes A., Roura E. (2010). Essential oils in poultry nutrition: Main effects and modes of action. Anim. Feed Sci. Tech., 158: 1–14.
- Bruno D.G., Martins S.M.M.K., Parazzi L.J., Afonso E.R., Del Santo T.A., Teixeira S.D.N., Moreno A.M., Moretti A.D. (2013). Phytogenic feed additives in piglets challenged with Salmonella typhimurium. Rev. Bras. Zootecn., 42: 137–143.
- Burt S. (2004). Essential oils: their antibacterial properties and potential applications in foods a review. Inter. J. Food Microbiol., 94: 223–253.
- Ceylan E., Fung D.Y.C. (2004). Antimicrobial activity of spices. J. Rapid Meth. Aut. Mic., 12: 1-55.
- Cho J.H., Kim H.J., Kim I.H. (2014). Effects of phytogenic feed additive on growth performance, digestibility, blood metabolites, intestinal microbiota, meat color and relative organ weight after oral challenge with *Clostridium perfringens* in broilers. Livest. Sci., 160: 82–88.
- C h o J.H., Z h a n g S., K i m I.H. (2012). Effects of anti-diarrhoeal herbs on growth performance, nutrient digestibility, and meat quality in pigs. Asian-Australas. J. Anim. Sci., 25: 1595–1604.
- Cross D.E., McDevitt R.M., Hillman K., Acamovic T. (2007). The effect of herbs and their associated essential oils on performance, dietary digestibility and gut microflora in chickens from 7 to 28 days of age. Brit. Poultry Sci., 48: 496–506.
- C z e c h A., K o w a l c z u k E., G r e l a E.R. (2009). The effect of a herbal extract used in pig fattening on the animals' performance and blood components. Ann. UMCS-EE, 27: 25–33.
- De Lange C.F.M., Pluske J.R., Gong J., Nyachoti C.M. (2010). Strategic use of feed ingredients and feed additives to stimulate gut health and development in young pigs. Livest. Sci., 134: 124–134.
- Dhama K., Chakraborty S., Tiwari R., Verma A.K., Saminathan M., Amarpal Malik Y.S., Nikousefat Z., Javdani M., Khan R.U. (2014). A concept paper on novel

technologies boosting production and safeguarding health of humans and animals. Res. Opin. Anim. Vet. Sci., 4: 353–370.

- Dhama K., Latheef S.K., Mani S., Samad H. A., Kartik K., Tiwari R., Khan R.U., Alagawany M., Farag M.R., Alam G.M., Laudadio V., Tufarelli V. (2015). Multiple beneficial applications and modes of action of herbs in poultry health and production – A review. Int. J. Pharmacol., 11: 152–176.
- Dragan L., Gyorke A., Ferreira J.F., Pop I. A., Dunca I., Dragan M., Mircean V., Dan I., Cozma V. (2014). Effects of Artemisia annua and Foeniculum vulgare on chickens highly infected with Eimeria tenella (Phylum Apicomplexa). Acta Vet Scand., 56: 22, DOI: 10.1186/1751-0147-56-22.
- Elagib H.A.A., El-Amin W.I.A., Elamin K.M., Malik H.E.E. (2013). Effect of dietary garlic (*Allium sativum*) supplementation as feed additive on broiler performance and blood profile. J. Anim. Sci. Adv., 3: 58–64.
- Emami Khodambashi N., Samie A., Rahmani H.R., Ruiz-Feria C.A. (2012). The effect of peppermint essential oil and fructooligosaccharides, as alternatives to virginiamycin, on growth performance, digestibility, gut morphology and immune response of male broilers. Anim. Feed Sci. Technol., 175: 57–64.
- Faleiro L., Miguel G., Gomes S., Costa L., Venancio Teixeira F., Figueiredo A., Barroso A.C., Pedro J.G. (2005). Antibacterial and antioxidant activities of essential oils isolated from *Thymbra capitata* L. (Cav.) and *Origanum vulgare* L.J. Agric. Food Chem., 53: 8162–8168.
- F a r a g M.R., A l a g a w a n y M., T u f a r e 11 i V. (2016). *In vitro* antioxidant activities of resveratrol, cinnamaldehyde and their synergistic effect against cyadox-induced cytotoxicity in rabbit erythrocytes. Drug Chem. Toxicol., 17: 1–10.
- Florou Paneri P., Giannenas I., Christaki E., Govaris A., Botsoglou N. (2006). Performance of chickens and oxidative stability of the produced meat as affected by feed supplementation with oregano, vitamin C, vitamin E and their combinations. Arch. Geflugelkd., 70: 232–239.
- Frankic T., Voljc M., Salobir J., Rezar V. (2009). Use of herbs and spices and their extracts in animal nutrition. Acta Agr. Slov., 94: 95–102.
- Gerzilov V., Nikolov A., Petrov P., Bozakova N., Penchev G., Bochukov A. (2015). Effect of a dietary herbal mixture supplement on the growth performance, egg production and health status in chickens. J. Cent. Eur. Agric., 16: 10–27.
- Ghazaghi M., Mehri M., Kasmani F.B. (2014). Effects of dietary *Mentha spicata* on performance, blood metabolites, meat quality and microbial ecosystem of small intestine in growing Japanese quail. Anim. Feed Sci. Technol., 194: 89–98.
- G h e i s a r M.M., I m Y.W., L e e H.H., C h o i Y.I., K i m I.H. (2015). Inclusion of phytogenic blends in different nutrient density diets of meat-type ducks. Poultry Sci., Doi: org/10.3382/ps/pev301.
- Ghosh S., May M.J., Kopp E.B. (1998). NF-κB and rel proteins: evolutionarily conserved mediators of immune responses. Ann. Rev. Immunol., 16: 225–260.
- G i a n n e n a s I.A., K y r i a z a k i s I. (2009). Phytobased products for the control of intestinal diseases in chickens in the post antibiotic era. In: Phytogenics in animal nutrition: natural concepts to optimize gut health and performance, Steiner T. (ed.). Nottingham University Press, UK, pp. 61–85.
- Giannenas I., Florou-Paneri P., Papazahariadou M., Christaki E., Botsoglou N.A., Spais A.B. (2003). Effect of dietary supplementation with oregano essential oil on performance of broilers after experimental infection with *Eimeria tenella*. Arch Tierernahr., 57: 99–106.
- Giannenas I.A., Florou-Paneri P., Botsoglou N.A., Christaki E., Spais A.B. (2005). Effect of supplementing feed with oregano and/or alpha-tocopheryl acetate on growth of broiler chickens and oxidative stability of meat. J. Anim. Feed Sci., 14: 521–535.
- Gibson G.R., Probert H.M., Loo J.V., Rastall R.A., Roberfroid M.B. (2004). Dietary modulation of the human colonic microbiota: Updating the concept of prebiotics. Nutr. Res. Rev., 17: 259–275.
- Gong J., Yin F., Hou Y., Yin Y. (2014). Chinese herbs as alternatives to antibiotics in feed for swine and poultry production: Potential and challenges in application. Can. J. Anim. Sci., 94: 223–241.

- Gopi M., Karthik K., Manjunathachar H.V., Tamilmahan P., Kesavan M., Dashprakash M., Balaraju B.L., Purushothaman M.R. (2014). Essential oils as a feed additive in poultry nutrition. Adv. Anim. Vet. Sci., 2: 1–7.
- G r a s h o r n M.A. (2010). Use of phytobiotics in broiler nutrition an alternative to in feed antibiotics? J. Anim. Feed Sci., 19: 338–347.
- Hafeez A., Manner K., Schieder C., Zentek J. (2016). Effect of supplementation of phytogenic feed additives (powdered vs. encapsulated) on performance and nutrient digestibility in broiler chickens. Poultry Sci., 95: 622–629.
- H a n c z a k o w s k a E., S w i a t k i e w i c z M. (2012). Effect of herbal extracts on piglet performance and small intestinal epithelial villi. Czech J. Anim. Sci., 57: 420–429.
- H a s h e m i S.R., Z u l k i f l i I., Z u n i t a Z., S o m c h i t M.N. (2008). The effect of selected sterilization methods on antibacterial activity of aqueous extract of herbal plants. J. Biol. Sci., 8: 1072–1076.
- Hashemi S.R., Zulkifli I., Davoodi H., HairBejo M., Loh T.C. (2014). Intestinal histomorphology changes and serum biochemistry responses of broiler chickens fed herbal plant (*Euphorbia hirta*) and mix of acidifier. Iranian J. Appl. Anim. Sci., 4: 95–103.
- Hashemipour H., Kermanshahi H., Golian A., Veldkamp T. (2013). Effect of thymol and carvacrol feed supplementation on performance, antioxidant enzyme activities, fatty acid composition, digestive enzyme activities, and immune response in broiler chickens. Poultry Sci., 92: 2059–2069.
- Hazzit M., Baaliouamer A., Faleiro M.L., Miguel M.G. (2006). Composition of the essential oils of *Thymus* and *Origanum* species from Algeria and their antioxidant and antimicrobial activities. J. Agric. Food Chem., 54: 6314–6321.
- Hernandez F., Madrid J., Garcia V., Orengo J., Megias M.D. (2004). Influence of two plant extracts on broiler performance, digestibility, and digestive organ size. Poultry Sci., 83: 169–174.
- Hulankova R., Borilova G. (2011). *In vitro* combined effect of oregano essential oil and caprylic acid against *Salmonella* serovars, *Escherichia coli* O157:H7, *Staphylococcus aureus* and *Listeria monocytogenes*. Acta Vet. Brno, 80: 343–348.
- Huyghebaert G., Ducatelle R., Van Immerseel F. (2011). An update on alternatives to antimicrobial growth promoters for broilers. Vet. J., 187: 182–188.
- Jacela J.Y., DeRouchey J.M., Tokach M.D., Goodband R.D., Nelssen J.L., Renter D.G., Dritz S.S. (2010). Feed additives for swine: Fact sheets – prebiotics and probiotics, and phytogenics. J. Swine Health Prod., 18: 132–136.
- Jamroz D., Wertelecki T., Houszka M., Kamel C. (2006). Influence of diet type on the inclusion of plant origin active substances on morphological and histochemical characteristics of the stomach and jejunum walls in chicken. J. Anim. Physiol. Anim. Nutr., 90: 255–268.
- Janz J.A.M., Morel P.C.H., Wilkinson B.H.P., Purchas R.W. (2007). Preliminary investigation of the effects of low-level dietary inclusion of fragrant essential oils and oleoresins on pig performance and pork quality. Meat Sci., 75: 350–355.
- Jeong J.S., K im I.H. (2015). Effect of fermented medicinal plants (*Gynura Procumbens, Rehmannia Glutinosa, Scutellaria Baicalensis*) as alternative performance enhancers in broilers. J. Poultry Sci., 52: 119–126.
- Juadjur A., Mohn C., Schantz M., Baum M., Winterhalter P., Richling E. (2015). Fractionation of an anthocyanin-rich bilberry extract and *in vitro* antioxidative activity testing. Food Chem., 167: 418–424.
- Karadas F., Pirgozliev V., Rose S.P., Dimitrov D., Oduguwa O., Bravo D. (2014). Dietary essential oils improve the hepatic antioxidative status of broiler chickens. Brit. Poultry Sci., 55: 329–334.
- Khan R.U., Nikousefat Z., Tufarelli V., Naz S., Javdani M., Laudadio V. (2012). Garlic (*Allium sativum*) supplementation in poultry diets: Effect on production and physiology. World Poultry Sci. J., 68: 417-424.
- Khattak F., Ronchi A., Castelli P., Sparks N. (2014). Effects of natural blend of essential oil on growth performance, blood biochemistry, cecal morphology, and carcass quality of broiler chickens. Poultry Sci., 93: 132–137.
- Kiczorowska B., Klebaniuk R., Bąkowski M., Al-Yasiry A.R.M. (2015). Culinary herbs nutritive value and content of minerals J. Elem., 20: 599–608.

- Kiczorowska B., Samolińska W., Al-Yasiry A.R.M. Kowalczyk-Pecka D. (2016). Effect of supplementation of mixtures for broiler chickens with *Boswellia serrata* on the condition of the gastrointestinal tract and rearing efficiency. Ann. Anim. Sci., 16: 1–15.
- Kim Y.S., Hwang J.W., Sung S.H., Jeon Y.J., Jeong J.H., Jeon B.T., Moon S.H., Park P.J. (2015). Antioxidant activity and protective effect of extract of *Celosia cristata* L. flower on tert-butyl hydroperoxide-induced oxidative hepatotoxicity. Food Chem., 68: 572–579.
- Koscova J., Nemcova R., Gancarcikova S., Jonecova Z., Scirankova L., Bomba A., Buleca V. (2006). Effect of two plant extracts and *Lactobacillus fermentum* on colonization of gastrointestinal tract by *Salmonella enteric* var. Dusseldorf in chicks. Biol. Brat., 61: 775–778.
- Kroisma A., Steiner T., Zhang C. (2006). Influence of a phytogenic feed additive on performance of weaner piglets (Abstract). J. Anim. Sci., 84: 329.
- Kroismayr A., Sehm J., Pfaffl M.W., Schedle K., Plitzner C., Windisch W. (2008). Effects of avilamycin and essential oils on mRNA expression of apoptotic and inflammatory markers and gut morphology of piglets. Czech J. Anim. Sci., 53: 377–387.
- K u n i n C.M. (1993). Resistance to antimicrobial drugs a worldwide calamity. Ann. Int. Med., 118: 557–561.
- Labaque M.C., Kembro J.M., Luna A., Marin R.H. (2013). Effects of thymol feed supplementation on female Japanese quail (*Coturnix coturnix*) behavioral fear response. Anim. Feed Sci. Technol., 183: 67–72.
- Latheef S.K., Dhama K., Wani M.Y., Samad H.A., Barathidasan R., Tiwari R., Singh S.D., Rai R.B. (2013). Ameliorative effects of four herbs (*Withania somnifera, Tinospora cordifolia, Azadirachta indica* and E care SE herbal) on the pathogenesis of chicken infectious anaemia virus. Int. J. Curr. Res., 5: 2327–2331.
- L e e K.G., S h i b a m o t o T. (2001). Inhibition of malonaldehyde formation from blood plasma oxidation by aroma extracts and aroma components isolated from clove and eucalyptus. Food Chem. Toxicol., 39: 1199–1204.
- Lee S.H., Lillehoj H.S., Jang S.I., Lee K.W., Bravo D., Lillehoj E.P. (2011). Effects of dietary supplementation with phytonutrients on vaccine-stimulated immunity against infection with *Eimeria tenella*. Vet. Parasitol., 181: 97–105.
- L e i Y., K i m D.W., Y i m Y.H., K i m I.H. (2014). The effect of evening primrose extract (*Oenotherabiennis*) on growth performance, nutrient digestibility, blood characteristics, faecal microbiota and fecal score in weaned pigs. Int. J. Agric. Innov. Res., 3: 307–314.
- Li H.L., Z h a o P.Y., L e i Y., H o s s a i n M.M., K i m I.H. (2015). Phytoncide, phytogenic feed additive as an alternative to conventional antibiotics, improved growth performance and decreased excreta gas emission without adverse effect on meat quality in broiler chickens. Livest. Sci., 181: 1–6.
- Li H.N., Zhao P.Y., Yan L., Hossain M.M., Kang J., Kim I.H. (2016). Dietary phytoncide supplementation improved growth performance and meat quality of finishing pigs. Asian Australas. J. Anim. Sci., 9: 1314–1321.
- Li P., Piao X., Ru Y., Han X., Xue L., Zhang H. (2012 a). Effects of adding essential oil to the diet of weaned pigs on performance, nutrient utilization, immune response and intestinal health. Asian-Australas. J. Anim. Sci., 25: 1617–1626.
- Li S.Y., Ru Y.J., Liu M., Xu B., Peron A., Shi X.G. (2012 b). The effect of essential oils on performance, immunity and gut microbial population in weaner pigs. Livest Sci., 145: 119–123.
- Maenner K., Vahjen W., Simon O. (2011). Studies on the effects of essential-oil-based feed additives on performance, ileal nutrient digestibility, and selected bacterial groups in the gastrointestinal tract of piglets. J. Anim. Sci., 89: 2106–2112.
- Mahima Rahal A., Deb R., Latheef S.K., Abdul Samad H., Tiwari R., Verma A.K., Kumar A., Dhama K. (2012). Immunomodulatory and therapeutic potentials of herbal, traditional/indigenous and ethnoveterinary medicines. Pak. J. Biol. Sci., 15: 754–774.
- M a m o u n T., M u k h t a r A., T a b i d i M.H. (2014). Effect of fenugreek seed powder on the performance, carcass characteristics and some blood serum attributes. Adv. Res. Agri. Vet. Sci., 1: 6–11.
- Manzanilla E.G., Perez J.F., Martin M., Kamel C., Baucells F., Gasa J. (2004). Effect of plant extracts and formic acid on the intestinal equilibrium of early-weaned pigs. J. Anim. Sci., 82: 3210–3218.

- Miura K., Kikuzaki H., Nakatani N. (2002). Anti-oxidant activity of chemical components from sage (*Salvia officinalis*) and thyme (*Thymus vulgaris*) measured by oil stability index method. J. Agric. Food Chem., 50: 1841–1851.
- Mourao J.L., Victor P., Alves A., Guedes C., de Lurdes Pinto M., Saavedra M.J., Spring P., Kocher A. (2006). Effect of mannan oligosaccharides on the performance, intestinal morphology and cecal fermentation of fattening rabbits. Anim. Feed Sci. Technol., 126: 107–120.
- M u h l A., L i e b e r t F. (2007 a). Growth and parameters of microflora in intestinal and fecal samples of piglets due to application of a phytogenic feed additive. J. Anim. Physiol. Anim. Nutr., 91: 411–418.
- M u h l A., L i e b e r t F. (2007 b). No impact of a phytogenic feed additive on digestion and unspecific immune reaction in piglets. J. Anim. Physiol. Anim. Nutr., 91: 426–431.
- Musa H.H., Wu S.L., Zhu C.H., Seri H.I., Zhu G.Q. (2009). The potential benefits of probiotics in animal production and health. J. Anim. Vet. Adv., 8: 313–321.
- Mushtaq M., Durrani F.R., Imtiaz N., Sadique U., Hafeez A., Akhtar S., Ahmad S. (2012). Effect of administration of *Withania somnifera* on some hematological and immunological profile of broiler chicks. Pak Vet. J., 32: 70–72.
- Neu H.C. (1992). The crisis in antibiotic resistance. Science, 257: 1064-1073.
- Oetting L.L., Utiyama C.E., Giani P.A., Ruiz U.D., Miyada V.S. (2006). Effects of herbal extracts and antimicrobials on apparent digestibility, performance, organs morphometry and intestinal histology of weanling pigs. Braz. J. Anim. Sci., 35: 1389–1397.
- Oyuntsetseg N., Khasnatinov M.A., Molor-Erdene P., Oyunbileg J., Liapunov A.V., Danchinova G.A., Oldokh S., Baigalmaa J., Chimedragchaa C. (2014). Evaluation of direct antiviral activity of the Deva-5 herb formulation and extracts of five Asian plants against influenza A virus H3N8. BMC Compl. Alt. Med., 14: 235.
- O z e k K., Wellmann K.T., Ertekin B., Tarim B. (2011). Effects of dietary herbal essential oil mixture and organic acid preparation on laying traits, gastrointestinal tract characteristics, blood parameters and immune response of laying hens in a hot summer season. J. Anim. Feed Sci., 20: 575–586.
- P a n t M., A m b w a n i T., U m a p a t h i V. (2012). Antiviral activity of Ashwagandha extract on infectious bursal disease virus replication. Indian J. Sci. Technol., 5: 2750–2751.
- Pastorelli G., Rossi R., Corino C. (2012). Influence of *Lippia citriodora* verbascoside on growth performance, antioxidant status, and serum immunoglobulins content in piglets. Czech J. Anim. Sci., 57: 312–322.
- Patil K.R., Patil C.R. (2016). Anti-inflammatory activity of bartogenic acid containing fraction of *Barringtonia racemosa* Roxb in acute and chronic animal models of inflammation. J. Trad. Compl. Med., dx.doi.org/10.1016/j.jtcme.2016.02.001. Accessed 11th Nov 2016.
- Perez-Barron G., Avila-Acevedo J.G., Garcia-Bores A.M., Montes S., Garcia-Jimenez S., Leon-Rivera I., Rubio-Osornio M., Monroy-Noyola A. (2015). Neuroprotective effect of *Buddleja cordata* methanolic extract in the 1-methyl- 4-phenylpyridinium Parkinson's disease rat model. J. Nat. Med., 69: 86–93
- Platel K., Srinivasan K. (2004). Digestive stimulant action of spices: A myth or reality? Indian J. Med. Res., 119: 167–179.
- R a h i m i A., M i r z a A g h a z a d e h A., D a n e s h y a M.R. (2011). Growth performance and some carcass characteristics in broiler chickens supplemented with Thymus extract (*Thymus vulgaris*) in drinking water. J. Am. Sci., 7: 400–405.
- Randrianarivelo R., Danthu P., Benoit C., Ruez P., Raherimandimby M., Starter S. (2010). Novel alternative to antibiotics in shrimp hatchery: Effects of the essential oil of *Cinnamosma fragrans* on survival and bacterial concentration of *Penaeus monodon* larvae. J. Appl. Microbiol., 109: 642–650.
- Ranucci D., Beghelli D., Trabalza-Marinucci M., Branciari R., Forte C., Olivieri O., Badillo Pazmay G.V., Cavallucci C., Acuti G. (2014). Dietary effects of a mix derived from oregano (*Origanum vulgare* L.) essential oil and sweet chestnut (*Castanea sativa* Mill.) wood extract on pig performance, oxidative status and pork quality traits. Meat Sci., http://dx.doi.org/10.1016/j.meatsci.2014.09.149.
- Regulation (EC) No 1831/2003 of the European Parliament and of the Council of 22 September 2003 on additives for use in animal nutrition. Off. J. European Union. L 574 268: 29–43.

- Rhayour K., Bouchikhi T., Tantaoui Elaraki A., Sendide K., Remmal A. (2003). The mechanism of bactericidal action of oregano and clove essential oils and of their phenolic major components on *Escherichia coli* and *Bacillus subtilis*. J. Essent. Oil Res., 15: 286–292.
- Schone F., Vetter A., Hartung H., Bergmann H., Biertümpfel A., Richter G., Müller S., Breitschuh G. (2006). Effects of essential oils from fennel (*Foeniculi aetheroleum*) and caraway (*Carvi aetheroleum*) in pigs. J. Anim. Physiol. Anim. Nutr., 90: 500–510.
- Simitzis P.E., Symeon G.K., Charismiadou M.A., Bizelis J.A., Deligeorgis S.G. (2010). The effects of dietary oregano oil supplementation on pig meat characteristics. Meat Sci., 84: 670–676.
- Soltan M.A., Shewita R.S., El-Katcha M.I. (2008). Effects of dietary anise seeds supplementation on growth performance, immune response, carcass traits and some blood parameters of broiler chickens. Int. J. Poult. Sci., 7: 1078–1088.
- Srinivasan K. (2006). Fenugreek (*Trigonella foenum-graecum*): A review of health beneficial physiological effects. Food Rev. Int., 22: 203–224.
- Stanacev V., Glamocic D., Milosevic N., Puvaca N., Stanacev V., Plavsa N. (2011). Effect of garlic (*Allium sativum* L.) in fattening chicks nutrition. Afr. J. Agric. Res., 6: 943–948.
- Thapa D., Losa R., Zweifel B., Wallace R.J. (2012). Sensitivity of pathogenic and commensal bacteria from the human colon to essential oils. Microbiol., 158: 2870–2877.
- U p a d h a y a S.D., L e e K.Y., K i m I.H. (2014). Protected organic acid blends as an alternative to antibiotics in finishing pigs. Asian-Australas. J. Anim. Sci., 27: 1600–1607.
- U p a d h a y a S.D., L e e K.Y., K i m I.H. (2016 a). Effect of protected organic acid blends on growth performance, nutrient digestibility and faecal micro flora in growing pigs. J. Appl. Anim. Res., 44: 232–242.
- U p a d h a y a S.D., K i m S.J., K i m I.H. (2016 b). Effects of gel-based phytogenic feed supplement on growth performance, nutrient digestibility, blood characteristics and intestinal morphology in weanling pigs. J. Appl. Anim. Res., 44: 384–389.
- Varel V.H. (2002). Livestock manure odor abatement with plant-derived oils and nitrogen conservation with urease inhibitors: A review. J. Anim. Sci., 80: E1–E7.
- Wati T., Ghosh T.K., Syed B., Haldar S. (2015). Comparative efficacy of a phytogenic feed additive and an antibiotic growth promoter on production performance, caecal microbial population and humoral immune response of broiler chickens inoculated with enteric pathogens. Anim. Nutr., 3: 213–219.
- We n k C. (2006). Are herbs, botanicals and other related substances adequate replacements for antimicrobial growth promoters? In: Antimicrobial Growth Promoters, Barug D., Jong J. de., Kies A.K., Verste M.W.A. (eds). Wageningen Academic Publishers, The Netherlands, pp. 329–340.
- Windisch W., Schedle K., Plitzner C., Kroismayr A. (2008). Use of phytogenic products as feed additives for swine and poultry. J. Anim. Sci., 86: E140–E148.
- Windisch W., Rohrer E., Schedle K. (2009). Phytogenic feed additives to young piglets and poultry: Mechanisms and application. In: Phytogenics in Animal Nutrition: Natural concepts to optimize gut health and performance, Steiner T. (ed). Nottingham University Press, pp.19–39.
- Yan L., K im I.H. (2012). Effect of eugenol and cinnamaldehyde on the growth performance, nutrient digestibility, blood characteristics, fecal microbial shedding and fecal noxious gas content in growing pigs. Asian-Aust. J. Anim. Sci., 25: 1178–1183.
- Yan L., Wang J.P., Kim H.J., Meng Q.W., Ao X., Hong S.M., Kim I.H. (2010). Influence of essential oil supplementation and diets with different nutrient densities on growth performance, nutrient digestibility, blood characteristics, meat quality and fecal noxious gas content in growerfinisher pigs. Livest. Sci., 128: 115–122.
- Yan L., Meng Q.W., Kim I. H. (2011 a). The effects of dietary *Houttuynia cordata* and *Taraxacum officinale* extract powder on growth performance, nutrient digestibility, blood characteristics and meat quality in finishing pigs. Livest. Sci., 141: 188–193.
- Yan L., Meng Q.W., K im I.H. (2011 b). The effect of an herb extract mixture on growth performance, nutrient digestibility, blood characteristics and fecal noxious gas content in growing pigs. Livest. Sci., 141: 143–147.
- Yan L., Zhang Z.F., Park J.C., Kim I.H. (2012 a). Evaluation of *Houttuynia cordata* and *Taraxa-cum officinale* on growth performance, nutrient digestibility, blood characteristics and fecal microbial shedding in weaning pigs. Asian-Australas. J. Anim. Sci., 25: 1439–1444.

- Yan L., Meng Q.W., Kim I.H. (2012 b). Effect of an herb extract mixture on growth performance, nutrient digestibility, blood characteristics, and fecal microbial shedding in weanling pigs. Livest. Sci., 145: 189–195.
- Yang C., Chowdhury M.A.K., Hou Y., Gong J. (2015). Phytogenic compounds as alternatives to in-feed antibiotics: potentials and challenges in application. Pathogens, 4: 137–156.
- Yang Y., Iji P.A., Choct M. (2009). Dietary modulation of gut microflora in broiler chickens: a review of the role of six kinds of alternatives to in-feed antibiotics. World's Poultry Sci. J., 65: 97–114.
- Zeng Z., Xu X., Zhang Q., Li P., Zhao P., Li Q., Liu J., Piao X. (2015). Effects of essential oil supplementation of a low-energy diet on performance, intestinal morphology and microflora, immune properties and antioxidant activities in weaned pigs. Anim. Sci. J., 86: 279–285.
- Zhang H.Y., Piao X.S., Zhang Q., Li P., Yi J.Q., Liu J.D., Li Q.Y., Wang G.Q. (2013). The effects of *Forsythia suspensa* extract and berberine on growth performance, immunity, antioxidant activities, and intestinal microbiota in broilers under high stocking density. Poultry Sci., 92: 1981–1988.
- Zhang S., Jung J.H., Kim H.S., Kim B.Y., Kim I.H. (2012). Influences of phytoncide supplementation on growth performance, nutrient digestibility, blood profiles, diarrhea scores and fecal microflora shedding in weaning pigs. Asian-Aust. J. Anim. Sci., 25: 1309–1315.
- Zhou F., Ji B., Zhang H., Jiang H., Yang Z., Li J., Li J., Ren Y., Yan W. (2007). Synergistic effect of thymol and carvacrol combined with chelators and organic acids against *Salmonella* Typhimurium. J. Food Prot., 70: 1704–1709.
- Zhou T.X., Zhang Z.F., Kim I.H. (2013). Effects of dietary *Coptis chinensis* herb extract on growth performance, nutrient digestibility, blood characteristics and meat quality in growing-finishing pigs. Asian-Aust. J. Anim. Sci., 26: 108–115.

Received: 10 VI 2016 Accepted: 13 XII 2016