



THE EFFECT OF DIFFERENT CONCENTRATIONS OF CAFFEINE, PENTOXIFYLLINE AND 2'-DEOXYADENOSINE ON THE BIOLOGICAL PROPERTIES OF FROZEN-THAWED CANINE SEMEN*

Marek Lecewicz*, Rafał Strzeżek, Anna M. Majewska, Piotr S. Purpurowicz, Władysław Kordan

Department of Animal Biochemistry and Biotechnology,
University of Warmia and Mazury in Olsztyn, Oczapowskiego 5, 10-718 Olsztyn, Poland

*Corresponding author: mlecew@uwm.edu.pl

Abstract

Artificial insemination (AI) and semen cryopreservation are the most accessible and commonly used techniques for breeding domestic animals. Among many parameters, such as plasma membrane integrity and acrosome structure, one of the key factors that determine the quality of frozen-thawed samples for artificial insemination is sperm motility. Sperm motility is one of the key parameters that determine the quality of frozen-thawed samples for AI. The total number of progressively motile spermatozoa in thawed canine semen is correlated with fertility. A variety of substances were used to compare sperm motility with the control. The aim of this study was to determine the effect of semen extender supplementation with motility stimulants, pentoxifylline (PTX), caffeine (CAF) and 2'-deoxyadenosine (DX), after different post-thaw incubation times (30, 60, 120 min) on the motility, selected kinematic parameters, plasma membrane integrity and mitochondrial membrane potential of cryopreserved canine spermatozoa. During attempts to improve the quality of cryopreserved semen, the applied substances exerted beneficial effects at a concentration of 10 mM. We demonstrated that both phosphodiesterase inhibitors, caffeine and pentoxifylline, as well as 2'-deoxyadenosine increased the motility and selected kinematic parameters of thawed canine spermatozoa.

Key words: canine, cryopreservation, motility stimulants, kinematic motility

Artificial insemination (AI) and semen cryopreservation are the most accessible and commonly used techniques for breeding domestic animals. However, fewer assisted reproduction techniques have been developed for dogs than farm animals (Jewgenow et al., 2017; Lucio et al., 2017). Despite the above, there is a growing interest in frozen dog semen. The freezing-thawing procedure supports the preservation and propagation of genetic material. Cryopreservation also enables long-dis-

* This work was supported by the University of Warmia and Mazury in Olsztyn (No. 11.610.003-300).

tance transport of semen, reduces animal transport costs and increases the genetic diversity of species (Comizzoli *et al.*, 2009; Belala *et al.*, 2016; Lucio *et al.*, 2017). Canine semen has low resistance to freezing due to the alternations in phospholipids/cholesterol ratio triggered by freezing process, which makes cells vulnerable with the spermatozoa plasma membrane destabilization and consequent reduced sperm fertility (Inanc *et al.*, 2018). Cryopreservation can contribute to a decrease in sperm motility, loss of membrane integrity and acrosome structure, as well as damage to sperm DNA (Kim *et al.*, 2010; Karger *et al.*, 2017). The above risks necessitate new solutions to enhance the post-thaw parameters of canine spermatozoa, optimize oocyte fertilization and increase contraception rates.

Sperm motility is one of the key parameters that determine the quality of frozen-thawed samples for AI (Van den Berghe *et al.*, 2018). The total number of progressively motile spermatozoa in thawed canine semen is correlated with fertility (Mason *et al.*, 2017).

Sperm motility can decrease as a result of changes in plasma membrane permeability in the tail region of spermatozoa, and the formation of ice crystals in mitochondria and axonemes during cryopreservation (Rasul *et al.*, 2001; Yu *et al.*, 2002; Prapaiwan *et al.*, 2016). In the past, light microscopy was routinely used to evaluate the main semen parameters. Today, spermatozoa are accurately and rapidly assessed with the use of computer-assisted sperm analysis systems (CASA). These systems are capable of detecting even minor changes in sperm motility, and they are highly useful tools for improving the quality of cryopreserved canine semen (Rijsselaere *et al.*, 2012). In addition, it is possible to visualize the trajectory of movement of individual spermatozoa, enabling a detailed description of parameters related to motility and type of movement (Antończyk *et al.*, 2010). Furthermore, evaluation of kinematic parameters of sperm after cryopreservation process could be insufficient to determine the structural and biochemical changes in gametes. The maintenance of the sperm fertilization potential depends also on plasma membrane integrity and mitochondrial membrane potential of spermatozoa (Fraser *et al.*, 2002; Dziekońska *et al.*, 2009).

Various substances have been used to improve the quality of cryopreserved semen. The most popular motility stimulants include methylxanthines such as caffeine (CAF) and pentoxifylline (PTX) which are competitive and non-selective inhibitors of cyclic adenosine monophosphate (cAMP) phosphodiesterase that increase intracellular cAMP concentration and enhance tyrosine phosphorylation in the sperm tail region (Yunes *et al.*, 2005; Esteves *et al.*, 2007; Brie *et al.*, 2016). It is known that PTX inhibits tumour necrosis factor- α which is responsible for DNA fragmentation and programmed cell death. It also reduces superoxide anions and lipid peroxidation associated with sperm membrane damage (Pecker *et al.*, 1997; Mundle *et al.*, 1999; Zhang *et al.*, 2005).

2'-deoxyadenosine (DX), an adenosine analogue with a modified ribose ring, delivers similar effects. 2'-deoxyadenosine stimulates motility via a different molecular mechanism which probably involves the activation of A2-receptor-mediated adenylate cyclase which increases the intracellular concentration of cAMP (Milani *et al.*, 2010).

Motility enhancers exerted positive effects in fresh and frozen-thawed semen of several animal species, including humans (Carrel and Aston, 2013), bulls (Zhang et al., 2014), stallions (Stephens et al., 2013), rams (Maxwell et al., 1995), dogs (Milani et al., 2010) and turkeys (Slanina et al., 2018).

The aim of this study was to determine the effect of semen extender supplementation with motility stimulants, caffeine (CAF), pentoxifylline (PTX) and 2'-deoxyadenosine (DX), after different post-thaw incubation times (30, 60, 120 min) on the motility, selected kinematic parameters, plasma membrane integrity and mitochondrial membrane potential of cryopreserved canine spermatozoa.

Material and methods

Chemicals

The chemicals were purchased from: Sigma-Aldrich Chemical Company (Saint Louis, USA) – caffeine, pentoxifylline, 2'-deoxyadenosine, Tris; POCh (Gliwice, Poland) – citrate; Chempur (Piekary Śląskie, Poland) – fructose; Minitüb (Tiefenbach, Germany) – Orvus Es Paste (Equex STM).

Semen collection

Semen was obtained from four mixed-breed dogs. The animals were housed at the laboratory of the Department of Animal Biochemistry and Biotechnology. They were kept individually, fed a commercial dry food (Purina Dog Chow®) with mineral-vitamin supplementation (Can-Vit®) and had access to water *ad libitum*. Ejaculates were collected once weekly by digital manipulations (Linde-Forsberg, 1991) and only the second fraction (the sperm-rich fraction) was used in the experiments. Sperm-rich fractions of ejaculates were collected once weekly over a period of 10 weeks. The dogs were housed in individual pens and fed commercial canine food. Water was available *ad libitum*. Permission to conduct this study was granted by the Local Ethics Committee for Animal Experiments (LKE.065.01.2017).

A preliminary assessment of the ejaculates involved an evaluation of sperm motility and determination of sperm cell concentrations by the cytometric method in a hemocytometer. Semen samples with total motility higher than 80% and sperm counts higher than 200×10^6 cells per ml of semen were qualified for further analyses.

Freezing-thawing procedure

The procedure of cryopreservation was conducted according to Nizanski et al. (2001) with some modifications (Strzeżek et al., 2015). Briefly, after individual evaluation, semen samples were pooled and centrifuged ($700 \times g$, 5 min). Seminal plasma was removed, semen sediments were diluted with standard Tris-citrate-fructose (TCF) extender (Rota et al., 1995) to a concentration of 2×10^8 spermatozoa/ml and left to stand for 30 minutes at room temperature (20°C). The samples were then cooled to 5°C for 60 minutes. After chilling, a second dilution (1:1) was performed

with an extender containing 5.0 ml of TCF, 4.0 ml of chicken egg yolk, 0.8 ml of glycerol and 0.2 ml of Orvus Es Paste to obtain a concentration of 1×10^8 spermatozoa/ml. The final content of chicken egg yolk, glycerol and Orvus Es Paste in diluted semen samples was determined at 20%, 4% and 1%, respectively. Equilibration time at 5°C was 15 minutes. Semen was loaded into 0.25 ml plastic straws (Minitüb, Germany) and frozen in a closed styrofoam box by placing filled straws on a rack suspended 4 cm above the surface of liquid nitrogen. Freezing time was 10 minutes. Frozen straws and tubes were placed in a container with liquid nitrogen. Straws were thawed in a water bath at 70°C for 5 s (Nöthling et al., 2005).

Application of motility stimulants

Thawed semen was aliquoted and mixed with: 1) TCF extender plus caffeine at final concentrations of 5 mM, 10 mM, 20 mM; 2) TCF extender plus pentoxifylline at final concentrations of 5 mM, 10 mM, 20 mM; 3) TCF extender plus 2'-deoxyadenosine at final concentrations of 5 mM, 10 mM, 20 mM.

Sperm viability (motility, plasma membrane integrity and mitochondrial function) was examined at different time intervals of 0, 60 and 120 min for different variants of the extender relative to the control. The incubation temperature was 22°C (Kordan et al., 2010).

Motility

Sperm motility characteristics were evaluated using the Hamilton-Thorne Sperm Analyser IVOS version 12.3 (Hamilton-Thorne Biosciences, USA). Software settings recommended by the manufacturer for canine sperm analyses were used: frame acquired – 30, frame rate – 60 Hz, minimum cell contrast – 75, minimum cell size – 6 pixels, straightness threshold – 75%, path velocity threshold – 100 µm/s, low velocity average pathway VAP (velocity straight line) cut-off – 9.9 µm/s, low VSL (velocity straight line) cut-off – 20 µm/s, static size gates – 0.80–4.93, static intensity gates – 0.49–1.68, static elongation gates – 22–84. Total motility (%), progressive motility (%), velocity average pathway (VAP, µm/s), velocity straight line (VSL, µm/s), curvilinear velocity (VCL, µm/s), amplitude lateral head (ALH, µm), beat cross frequency (BCF), linearity (LIN, %) and straightness (STR, %) were determined by the IVOS analyser.

Plasma membrane integrity

Sperm plasma membrane integrity was assessed by dual fluorescent staining with SYBR-14 and PI (Live/Dead Sperm Viability Kit; Molecular Probes, Eugene, USA), as described by Garner and Johnson (1995), with minor modifications. Briefly, aliquots of sperm samples (20×10^6 spermatozoa/ml) were incubated with SYBR-14 (1 mM SYBR-14 in DMSO) and PI solutions (2.4 µM PI in Tyrode's salt solution) for 10 min at 37°C. After incubation, stained sperm cells were placed on microscopic slides and examined under a fluorescence microscope (Olympus CH 30 RF-200, Tokyo, Japan) at 600 × magnification. Sperm cells displaying only bright green fluorescence were considered as viable spermatozoa with an undamaged plasma membrane. A minimum of 200 cells per slide was examined in random fields of each aliquot.

Mitochondrial function assessed by fluorescent microscopy

Sperm mitochondrial function was assessed by dual staining with fluorescent probes, 5,5',6,6'-tetrachloro-1,1',3,3'-tetraethylbenzimidazolylcarbocyanine iodide, JC-1 (Molecular Probes, Eugene, USA) with propidium iodide (PI, Sigma Chemical Co., St. Louis, MO, USA), according to previously described methods (Garner and Thomas, 1999) with some modifications (Dziekońska et al., 2009). Aliquots of sperm samples (20×10^6 spermatozoa/ ml) were incubated with JC-1 solution (1 mg of JC-1/ml dimethyl sulfoxide, DMSO) for 15 min at 37°C. After incubation, sperm samples were stained with PI (10 µl of PI solution in 0.5 mg/ml phosphate buffered solution) for 10 min at 37°C, washed ($600 \times g$, 5 min at room temperature), and sperm pellets were resuspended in a HEPES buffered solution (10 mM HEPES, 0.85% NaCl, 0.1% bovine serum albumin, pH 7.4). Stained sperm samples were placed on microscopic slides, covered with coverslips (22×22 mm) and examined under a fluorescence microscope (Olympus CH 30 RF-200, Tokyo, Japan). Viable spermatozoa with functional mitochondria emitted orange-red fluorescence. Two slides were evaluated per sample, and 200 spermatozoa were counted per slide.

Statistical analysis

Data were expressed as means \pm SEM and analysed by one-way analysis of variance (ANOVA). The mean values were compared between the control group and treatment groups using Duncan's multiple range test in the Statistica software package (StatSoft Inc., Tulsa, OK, USA). Differences were considered significant at $P \leq 0.05$.

Results

Total motility

The influence of motility stimulants on total sperm motility in samples incubated over different times is shown in Figure 1A. Total motility was significantly higher in thawed semen samples supplemented with motility stimulants at a concentration of 10 mM, regardless of stimulant type and incubation time, relative to control. Caffeine and 2'-deoxyadenosine applied at a concentration of 5 mM improved total motility after 0 and 60 minutes of incubation.

Progressive motility

Progressive motility was significantly higher in all semen samples treated with CAF, PTX and 2 DX at concentrations of 5 and 10 mM, regardless of incubation time (Figure 1B).

Kinematic parameters of sperm motility

When applied at a concentration of 10 mM, all of the tested stimulants significantly influenced VAP after 0, 60 and 120 minutes of incubation. A similar result was noted in VSL values, excluding semen samples treated with PTX. VCL was

most influenced by CAF at a concentration of 10 mM, however, CAF did not exert a significant influence on VCL values immediately after supplementation. DX and PTX did not influence the analysed parameter, but in non-incubated samples, VCL was significantly higher under exposure to pentoxifylline at a concentration of 10 mM. ALH increased significantly after treatment with 10 mM of CAF and 10 mM of PTX, regardless of incubation time. BCF remained constant through incubation with no significant differences (Figure 2). The highest percentage of spermatozoa characterized by linear movement was observed in semen samples treated with 10 mM of CAF after 60 minutes of incubation. The highest STR values were noted after the application of CF, DX and PTX at a concentration of 10 mM immediately after supplementation, and after 120 minutes of incubation with 10 mM of CAF and 10 mM of DX.




Figure 1. Percentage of total (A) and progressive motility (B) of frozen-thawed canine spermatozoa supplemented with different concentrations of caffeine, pentoxifylline and 2'-deoxyadenosine. The results are expressed as the mean values \pm SEM of 20 ejaculates from 4 dogs. Values (a, b) with different letters are significant at $P \leq 0.05$ for every incubation time




Figure 2. The mean values \pm SEM of VAP, VSL, VCL, ALH, BCF, LIN and STR in thawed semen supplemented with different concentrations of motility stimulants during 120 minutes of incubation, determined by the CASA method. The results are expressed as mean values \pm SEM. Values (a, b) with different letters are significant at $P \leq 0.05$ for every incubation time




Figure 3. The percentage of canine spermatozoa with integral plasma membrane determined in the SYBR-14/PI assay (A) and the percentage of canine spermatozoa with functional mitochondria determined in the JC-1/PI assay (B) after the addition of different concentrations of caffeine, pentoxifylline and 2'-deoxyadenosine. The results are expressed as mean values \pm SEM

Plasma membrane integrity and mitochondrial membrane potential

The influence of different concentrations of motility stimulants after 0, 60 and 120 minutes of incubation on plasma membrane integrity and mitochondrial activity is shown in Figure 3A and Figure 3B, respectively. Significant differences in the percentage of spermatozoa with intact plasma membrane and higher mitochondrial membrane potential were not observed relative to control.

Discussion

Despite the availability of sophisticated assisted reproductive techniques, frozen-thawed semen is characterized by lower motility and viability than freshly ejaculated

sperm. Cryopreservation induces crystallization and osmotic changes which can cause cellular damage (Stănescu and Bîrțoiu, 2011; Setyawan et al., 2015). An intact plasma membrane is essential for the maintenance of sperm cell integrity and the events associated with oocyte fertilization. However, most freezing-induced injuries in spermatozoa affect the plasma membrane (Bailey et al., 2000; Lucio et al., 2017).

A variety of extenders and cryopreservation protocols have been developed over the years, but the whelping rates of frozen dog semen are still unsatisfactory in comparison with fresh semen (Uchoa et al., 2012; Rodenas et al., 2014; Johnson et al., 2014). In dogs, the success of AI with cryopreserved semen is considerably limited due to problems associated with insemination timing (Alhaider and Watson, 2009). The selection of the appropriate extender, the freezing process and the thawing protocol are critical for the success of AI and the quality of cryopreserved canine sperm (Silva et al., 2006; Karger et al., 2017).

The purpose of our study was to verify whether post-thaw motility parameters, plasma membrane integrity and mitochondrial membrane potential can be improved by adding motility stimulants at different concentrations to the thawing medium. The results indicate that both phosphodiesterase inhibitors, caffeine and pentoxifylline, as well as 2'-deoxyadenosine increased the motility and selected kinematic parameters (VAP, VSL, VCL, ALH, BCF, LIN, STR) of thawed canine spermatozoa relative to control. Nevertheless, we did not observe any significant influence of used substances on plasma membrane integrity and mitochondrial membrane potential. We found that the tested substances exerted beneficial effects at a concentration of 10 mM. Similar results were reported by López and Alvariño (2000) in rabbits. The authors found that sperm motility increased at a higher concentration of caffeine (10 mM), but was not affected at lower caffeine concentrations (2.5 or 5 mM). In other studies, high caffeine concentrations exerted opposite effects on sperm motility. Stephens et al. (2013) demonstrated a non-significant increase in motility parameters after the addition of CAF at concentrations of 1, 2 and 3.5 mmol/L to thawed equine semen. The addition of 5 mM of caffeine also significantly enhanced the motility of bovine semen (Barakat et al., 2015). Caffeine also improved the qualitative parameters of spermatozoa in boars (Yamaguchi et al., 2013) and laboratory mice (Nabavi et al., 2013). These findings suggest that caffeine's effect on sperm motility parameters could be species-specific.

In the present study, PTX exerted a similar effect on sperm motility parameters. Semen samples supplemented with 10 mM of pentoxifylline were characterized by significantly higher motility parameters relative to control. Similar findings were reported by Koutsarova et al. (1997). The authors found that the addition of pentoxifylline at a concentration of 0.0036 mol/L significantly increased the percentage of progressively motile spermatozoa in fresh semen, and that the addition of PTX at a concentration of 0.0072 mol/L to semen samples at thawing improved the quality of frozen-thawed sperm.

The addition of PTX to semen samples before cryopreservation significantly decreased the velocity, linearity and motility of spermatozoa, which could result from the depletion of the available energy substrate (Yovich et al., 1990). However, various authors demonstrated that the addition of PTX to thawed semen significantly

increased sperm parameters. In a study by Gradil and Ball (2010), the addition of pentoxifylline (3.5 or 7.0 mM) to cryopreserved equine semen immediately after thawing significantly increased total and progressive motility relative to control. Furthermore, Stephens *et al.* (2013) observed that use of pentoxifylline increased sperm VCL, VAP and VSL in equine semen. Similar results were noted in previous studies where PTX treatment increased the percentage of motile spermatozoa in frozen-thawed human, feline and boar semen (Hammitt *et al.*, 1989; Stachecki *et al.*, 1995). In many previous studies, pentoxifylline was used at a concentration of 3.6 mM of (Tesarik *et al.*, 1992; Kay *et al.*, 1993; Lewis *et al.*, 1993) but the beneficial effects of this motility stimulant were observed at both lower and higher concentrations (Kay *et al.*, 1993; Lewis *et al.*, 1993; Centola *et al.*, 1995; Sharma *et al.*, 1996). Esteves *et al.* (2007) demonstrated that the addition of pentoxifylline to low-quality human sperm before freezing did not prevent acrosomal loss during the cryopreservation process and did not improve post-thaw motility or viability. Despite the above, pentoxifylline improved the ability of thawed spermatozoa to undergo the acrosome reaction in response to calcium ionophore.

Bani-Hani *et al.* (2011) suggested that the stimulatory effect of PTX on semen motility could be modulated by higher energy production. When added at a concentration of 5 mM, PTX increased the activity of seminal creatine kinase, an enzyme which catalyses the formation of ATP by the creatine-creatine phosphate shuttle (Banihani and Abu-Alhayjaa, 2016).

There is a general scarcity of published studies investigating the influence of 2'-deoxyadenosine on frozen-thawed spermatozoa. This motility stimulant can increase the proportion of motile spermatozoa by approximately 21–39% and induce changes in motility characteristics, linear velocity and the frequency of sperm head rotation in human semen. The percentage of penetrated oocytes was determined at 80.8% in the hamster zona-free ovum test (Mbizvo *et al.*, 1993; Lanzafame *et al.*, 1994) demonstrated that DX exerted a prolonged effect on hyperactivated motility. Similarly to our findings, Milani *et al.* (2010) reported that the addition of 5 mM and 7.5 mM of DX had a positive effect on the motility of thawed canine semen. Imoedemhe *et al.* (1992) observed a significant improvement in sperm motility and the number of recoverable human spermatozoa relative to unexposed control samples. In the present study, sperm membrane integrity was nearly equally enhanced by the addition of pentoxifylline and 2-deoxyadenosine to sperm prepared with the two-layer Percoll gradient method.

There are limited reports concerning the influence of pentoxifylline and 2-deoxyadenosine on sperm membrane integrity. A positive correlation between the percentage of sperm tail swelling patterns and motile spermatozoa was found in a study investigating membrane functional integrity in mouse spermatozoa supplemented with pentoxifylline (Ponce *et al.*, 1999). Laokirkkiat *et al.* (2007) demonstrated that the addition of pentoxifylline or 2-deoxyadenosine to human semen prepared with the two-layer Percoll gradient method led to nearly equal enhancement of sperm membrane integrity.

In theory, spermatozoa are the most suitable candidates for cryopreservation on account of their general properties. However, freezing-thawing can severely impair

the function of male gametes. Cryoinjury can be caused by various factors, including membrane disturbances and changes in osmotic pressure which lead to changes in cell volume (Pena, 2007). The motility of frozen-thawed spermatozoa can be compromised by active transport and changes in plasma membrane permeability in the tail region of sperm cells (Yu et al., 2002).

In conclusion, the motility enhancers used in our study (CAF, PTX, DX) improved post-thaw motility, selected kinematic parameters of motility (VAP, VSL, VCL, ALH, BCF, LIN, STR) and the overall quality of frozen-thawed canine semen. These effects were concentration-dependent, in particular in CAF and PTX. All of the tested motility enhancers produced the most satisfactory results when administered at a concentration of 10 mM. However, the applied treatments did not exert a significant influence on plasma membrane integrity or mitochondrial membrane potential in cryopreserved dog semen.

References

- Alhaider A.K., Watson P.F. (1999). The effects of hCG and growth factors on in vitro nuclear maturation of dog oocytes obtained during anoestrus. *Reprod. Fertil. Dev.*, 21: 538–548.
- Antończyk A., Nizański W., Twardoń J., Kozdrowski R., Ochota M., Błasiak K., Mikołajewska N., Stańczyk E. (2010). Current views on computer assisted sperm analysis. *Med. Weter.*, 66: 663–667.
- Bailey J.L., Bilodeau J.F., Cormier N. (2000). Semen cryopreservation in domestic animals: a damaging and capacitating phenomenon. *J. Androl.*, 21: 1–7.
- Bani-Hani S.A. (2011). Semen quality and chemical oxidative stress; quantification and remediation. *ETD Archive*, 24, <https://engagedscholarship.csuohio.edu/etdarchive/24>.
- Banihani S.A., Abu-Alhayja R.F. (2016). The activity of seminal creatine kinase is increased in the presence of pentoxifylline. *Andrologia*, 48: 603–604.
- Barakat I.A., Danfour M.A., Galewan F.A., Dkhil M.A. (2015). Effect of various concentrations of caffeine, pentoxifylline, and kallikrein on hyperactivation of frozen bovine semen. *Biomed. Res. Int.*, 2015: 1–7.
- Belala R., Fatmi S., Kaidi R., Iguer-Ouada M. (2016). Benefits of cholesterol and α -tocopherol loaded cyclodextrins in dog semen cryopreservation. *Revue Méd. Vét.*, 167: 22–27.
- Brie D., Sahebkar A., Penson P.E., Dinca M., Ursoniu S., Serban M.C., Zanchetti A., Howard G., Ahmed A., Aronow W.S., Muntner P., Lip G.Y., Wong N.D., Rysz J., Banach M. (2016). Effects of pentoxifylline on inflammatory markers and blood pressure: a systematic review and meta-analysis of randomized controlled trials. *J. Hypertens.*, 34: 2318–2329.
- Carrel D.T., Aston K.I. (2013). *Spermatogenesis: Methods and Protocols*. New York, USA, Humana Press.
- Centola G.M., Cartie R.J., Cox C. (1995). Differential responses of human sperm to varying concentrations of pentoxifylline with demonstration of toxicity. *J. Androl.*, 16: 136–142.
- Comizzoli P., Crosier A.E., Songsasen N., Gunther M.S., Howard J.G., Wildt D.E. (2009). Advances in reproductive science for wild carnivore conservation. *Reprod. Domest. Anim.*, 44: 47–52.
- Dziekońska A., Fraser L., Strzeżek J. (2009). Effect of different storage temperatures on the metabolic activity of spermatozoa following liquid storage. *J. Anim. Feed Sci.*, 18: 638–649.
- Esteves S.C., Spaine D.M., Cedenho A.P. (2007). Effects of pentoxifylline treatment before freezing on motility, viability and acrosome status of poor quality human spermatozoa cryopreserved by the liquid nitrogen vapor method. *Braz. J. Med. Biol. Res.*, 40: 985–992.
- Fraser L., Leciewicz M., Strzeżek J. (2002). Fluorometric assessments of viability and mitochondrial status of boar spermatozoa following liquid storage. *Pol. J. Vet. Sci.*, 5: 85–92.

- Garner D.L., Johnson L.A. (1995). Viability assessment of mammalian sperm using SYBR-14 and propidium iodide. *Biol. Reprod.*, 53: 276–284.
- Garner D.L., Thomas C.A. (1999). Organelle-specific probe JC-1 identifies membrane potential differences in the mitochondrial function of bovine sperm. *Mol. Reprod. Dev.*, 53: 222–229.
- Gradil C.M., Ball B.A. (2000). The use of pentoxifylline to improve motility of cryopreserved equine spermatozoa. *Theriogenology*, 54: 1041–1047.
- Hammit D.G., Bedia E., Rogers P.R., Syrop C.H., Donovan J.F., Williamson R.A. (1989). Comparison of motility stimulants for cryopreserved human semen. *Fertil. Steril.*, 52: 495–502.
- Imoedemhe D.A., Sigue A.B., Pacpaco E.A., Olazo A.B. (1992). Successful use of the sperm motility enhancer 2-deoxyadenosine in previously failed human *in vitro* fertilization. *J. Assist. Reprod. Genet.*, 9: 53–56.
- Inanc M.E., Tekin K., Olgac K.T., Yilmaz B., Cil B., Tasdemir U., Tuncer P.B., Buyukleblebici S., Durmaz E., Uysal O. (2018). Effect of cholesterol loaded cyclodextrin on semen cryopreservation of Aksaray Malakli shepherd dogs of different ages. *Anim. Reprod. Sci.*, 193:191–200.
- Jewgenow K., Braun B.C., Dehnhard M., Zahmel J., Goeritz F. (2017). Research on reproduction is essential for captive breeding of endangered carnivore species. *Reprod. Domest. Anim.*, 52: 18–23.
- Johnson A.E., Freeman E.W., Wildt D.E., Songsasen N. (2014). Spermatozoa from the maned wolf (*Chrysocyon brachyurus*) display typical canid hyper-sensitivity to osmotic and freezing-induced injury, but respond favorably to dimethyl sulfoxide. *Cryobiology*, 68: 361–370.
- Karger S., Geiser B., Grau M., Heuwieser W., Arlt S.P. (2017). Short communication: Progressive motility of frozen-thawed canine semen is highest five minutes after thawing. *Reprod. Domest. Anim.*, 52: 350–352.
- Kay V.J., Coutts J.R.T., Robertson L. (1993). Pentoxifylline stimulates hyperactivation in human spermatozoa. *Hum. Reprod.*, 8: 727–731.
- Kim S.H., Yu D.H., Kim Y.J. (2010). Effects of cryopreservation on phosphatidylserine translocation, intracellular hydrogen peroxide, and DNA integrity in canine sperm. *Theriogenology*, 73: 282–292.
- Kordan W., Lecewicz M., Strzezek R., Dziekońska A., Fraser L. (2010). Effect of platelet activating factor (PAF) supplementation in semen extender on viability and ATP content of cryopreserved canine spermatozoa. *Pol. J. Vet. Sci.*, 13: 571–579.
- Koutsarova N., Todorov P., Koutsarov G. (1997). Effect of pentoxifylline on motility and longevity of fresh and thawed dog spermatozoa. *J. Reprod. Fertil.*, 51: 117–121.
- Lanzafame F., Chapman M.G., Guglielmino A., Gearon C.M., Forman R.G. (1994). Pharmacological stimulation of sperm motility. *Hum. Reprod.*, 9: 192–199.
- Laokirkkiat P., Kunathikom S., Choavaratana R., Petyim S., Prechapanich J. (2007). Comparison between sperm treated with pentoxifylline and 2-deoxyadenosine using hypo-osmotic swelling test. *J. Med. Assoc. Thai.*, 90: 211–215.
- Lewis S.E., Moohan J.M., Thompson W. (1993). Effects of pentoxifylline on human sperm motility in normozoospermic individuals using computer assisted analysis. *Fertil. Steril.*, 59: 418–423.
- Linde-Forsberg C. (1991). Achieving canine pregnancy by using frozen or chilled extended semen. *Vet. Clin. North Am. Small Anim. Pract.*, 21: 457–485.
- López F.J., Alvarino J.M. (2000). Effects of added caffeine on results following artificial insemination with fresh and refrigerated rabbit semen. *Anim. Reprod. Sci.*, 58: 147–154.
- Lucio C.F., Angrimani D.S.R., Brito M.M., Vannucchi C.I. (2017). Oxidative stress challenges during the sperm cryopreservation in dogs. *J. Vet. Androl.*, 2: 1–7.
- Mason S.J. (2017). A retrospective clinical study of endoscopic-assisted transcervical insemination in the bitch with frozen-thawed dog semen. *Reprod. Domest. Anim.*, 52: 275–280.
- Maxwell W.M., Robinson S.J., Roca J., Molinia F.C., Sanchez-Partida L.G., Evans G. (1995). Motility, acrosome integrity and fertility of frozen ram spermatozoa treated with caffeine, pentoxifylline, cAMP, 2-deoxyadenosine and kallikrein. *Reprod. Fertil. Dev.*, 7: 1081–1087.
- Mbizvo M.T., Johnston R.C., Baker G.H.W. (1993). The effect of the motility stimulants, caf-

- feine, pentoxifylline, and 2-deoxyadenosine on hyperactivation of cryopreserved human sperm. *Fertil. Steril.*, 59: 1112–1117.
- Milani C., Fontbonne A., Sellem E., Stelletta C., Gérard O., Romagnoli S. (2010). Effect of post-thaw dilution with caffeine, pentoxifylline, 2'-deoxyadenosine and prostatic fluid on motility of frozen-thawed dog semen. *Theriogenology*, 74: 153–164.
- Mundle S.D., Reza S., Ali A., Mativi Y., Shetty V., Venugopal P., Gregory S.A., Raza A. (1999). Correlation of tumor necrosis factor alpha (TNF alpha) with high Caspase 3-like activity in myelodysplastic syndromes. *Cancer Lett.*, 140: 201–207.
- Nabavi N., Todehdeghhan F., Shiravi A. (2013). Effect of caffeine on motility and vitality of sperm and *in vitro* fertilization of outbred mouse in T6 and M16 media. *Iran. J. Reprod. Med.*, 11: 741–746.
- Nizanski W., Dubiel A., Bielas W., Dejneka G.J. (2001). Effects of three cryopreservation methods and two semen extenders on the quality of dog semen after thawing. *J. Reprod. Fertil.*, 57: 365–369.
- Nöthling J.O., Shuttleworth R. (2005). The effect of straw size, freezing rate, and thawing rate upon post-thaw quality of dog semen. *Theriogenology*, 63: 1469–1480.
- Peeker R., Abramsson L., Marklund S.L. (1997). Superoxide dismutase isoenzymes in human seminal plasma and spermatozoa. *Mol. Hum. Reprod.*, 3: 1061–1066.
- Pena F.J. (2007). Detecting subtle changes in sperm membranes in veterinary andrology. *Asian J. Androl.*, 9: 731–737.
- Ponce A.A., Fiol de Cuneo M., Ruiz R.D., Vincenti L.M., Santillá M.E., Stutz G., Lacuara J.L. (1999). Influence of pentoxifylline on sperm membrane functional integrity. *Arch. Androl.*, 43: 77–84.
- Prapaiwan N., Tharasanit T., Punjachaipornpol S., Yamtang D., Roongsitthichai A., Moonarmart W., Kaeoket K., Manee-In S. (2016). Low-density lipoprotein improves motility and plasma membrane integrity of cryopreserved canine epididymal spermatozoa. *Asian-Australas. J. Anim. Sci.*, 29: 646–651.
- Rasul Z., Ahmad N., Anzar M. (2001). Changes in motion characteristics, plasma membrane integrity, and acrosome morphology during cryopreservation of buffalo spermatozoa. *J. Androl.*, 22: 278–283.
- Rijsselaere T., Van Soom A., Maes D., Nizanski W. (2012). Computer-assisted sperm analysis in dogs and cats: an update after 20 years. *Reprod. Domest. Anim.*, 47: 204–207.
- Rodenas C., Parrilla I., Roca J., Martinez E.A., Lucas X. (2014). Effects of rapid cooling prior to freezing on the quality of canine cryopreserved spermatozoa. *J. Reprod. Dev.*, 60: 355–361.
- Rota A., Ström B., Linde-Forsberg C. (1995). Effects of seminal plasma and three extenders on canine semen stored at 4 degrees C. *Theriogenology*, 44: 885–900.
- Setyawan E.M., Kim M.J., Oh H.J., Kim G.A., Jo Y.K., Lee S.H., Choi Y.B., Lee B.C. (2015). Maintaining canine sperm function and osmolyte content with multistep freezing protocol and different cryoprotective agents. *Cryobiology*, 71: 344–349.
- Sharma R.K., Tolentino M.V. Jr, Thomas A.J. Jr, Agarwal A. (1996). Optimal dose and duration of exposure to artificial stimulants in cryopreserved human spermatozoa. *J. Urol.*, 155: 568–573.
- Silva A.R., Cardoso R.C., Silva L.D. (2006). Influence of temperature during glycerol addition and post-thaw dilution on the quality of canine frozen semen. *Reprod. Domest. Anim.*, 41: 74–78.
- Slanina T., Miškeje M., Tirpák F., Błaszczyk M., Formicki G., Massányi P. (2018). Caffeine strongly improves motility parameters of turkey spermatozoa with no effect on cell viability. *Acta Vet. Hung.*, 66: 137–150.
- Stachecki J.J., Dresser B.L., Pope C.E., Armant D.R. (1995). Stimulation of ejaculated domestic cat spermatozoa motility with caffeine, pentoxifylline, and 2'-deoxyadenosine. *Arch. Androl.*, 34: 63–68.
- Stănescu (Pascal) M., Bîrțoiu A.I. (2012). Freezing of dog's sperm: a review. *Rom. Biotechnol. Lett.*, 17: 7709–7716.
- Stephens T.D., Brooks R.M., Carrington J.L., Cheng L., Carrington A.C., Porr C.A., Splan R.K. (2013). Effects of pentoxifylline, caffeine, and taurine on post-thaw motility and longevity of equine frozen semen. *J. Equine Vet. Sci.*, 33: 615–621.

- Strzeżek R., Polakiewicz W., Kordan W. (2015). The effect of two packaging systems on the post-thaw characteristics of canine sperm. *Pol. J. Vet. Sci.*, 18: 249–253.
- Tesarik J., Thébault A., Testart J. (1992). Effect of pentoxifylline on sperm movement characteristics in normozoospermic and asthenozoospermic specimens. *Hum. Reprod.*, 7: 1257–1263.
- Uchoa D.C., Silva T.F., Mota Filho A.C., Silva L.D. (2012). Intravaginal artificial insemination in bitches using frozen/thawed semen after dilution in powdered coconut water (ACP-106c). *Reprod. Domest. Anim.*, 47: 289–292.
- Van den Berghe F., Paris M.C.J., Briggs M.B., Farstad W.K., Paris D.B.B.P. (2018). A two-step dilution tris-egg yolk extender containing Equex STM significantly improves sperm cryopreservation in the African wild dog (*Lycaon pictus*). *Cryobiology*, 80: 18–25.
- Yamaguchi S., Suzuki C., Noguchi M., Kasa S., Mori M., Isozaki Y., Ueda S., Funahashi H., Kikuchi K., Nagai T., Yoshioka K. (2013). Effects of caffeine on sperm characteristics after thawing and inflammatory response in the uterus after artificial insemination with frozen-thawed boar semen. *Theriogenology*, 79: 87–93.
- Yovich J.M., Edirisinghe W.R., Cummins J.M., Yovich J.L. (1990). Influence of pentoxifylline in severe male factor infertility. *Fertil. Steril.*, 53: 715–722.
- Yu I., Songsasen N., Godke R.A., Leibo S.P. (2002). Differences among dogs in response of their spermatozoa to cryopreservation using various cooling and warming rates. *Cryobiology*, 44: 62–78.
- Yunes R., Fernández P., Doncel G.F., Acosta A.A. (2005). Cyclic nucleotide phosphodiesterase inhibition increases tyrosine phosphorylation and hyper motility in normal and pathological human spermatozoa. *Biocell.*, 29: 287–293.
- Zhang J., Guo H., Su J., Zhao L., Li Y., Sun W., Han H., Hu S., Zhao G., Li Y., Dai Y., Li X. (2014). A comparison of the effects of pentoxifylline on quality of fresh and frozen-thawed bull spermatozoa. *Agriculture Biotechnology*, 3: 20–29.
- Zhang X., Sharma R.K., Agarwal A., Falcone T. (2005). Effect of pentoxifylline in reducing oxidative stress-induced embryotoxicity. *J. Assist. Reprod. Genet.*, 22: 415–417.

Received: 6 I 2019

Accepted: 14 III 2019