On partial sums of normalized Mittag-Leffler functions

Dorina Răducanu

Abstract

This article deals with the ratio of normalized Mittag-Leffler function $\mathbb{E}_{\alpha,\beta}(z)$ and its sequence of partial sums $(\mathbb{E}_{\alpha,\beta})_m(z)$. Several examples which illustrate the validity of our results are also given.

1 Introduction

Let $\mathcal{A}$ be the class of functions f normalized by

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (1.1)$$

which are analytic in the open unit disk $\mathcal{U} = \{z \in \mathbb{C} : |z| < 1\}$.

Denote by $\mathcal{S}$ the subclass of $\mathcal{A}$ which consists of univalent functions in $\mathcal{U}$.

Consider the function $E_\alpha(z)$ defined by

$$E_\alpha(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\alpha n + 1)}, \quad \alpha > 0, \quad z \in \mathcal{U}, \quad (1.2)$$

where $\Gamma(s)$ denotes the familiar Gamma function. This function was introduced by Mittag-Leffler in 1903 [9] and is therefore known as the Mittag-Leffler function.

Key Words: Analytic functions, partial sums, Mittag-Leffler function, univalent function.
2010 Mathematics Subject Classification: 33E12, 30A10.
Received: 01.07.2016
Revised: 15.09.2016
Accepted: 21.09.2016

Another function $E_{\alpha,\beta}(z)$, having similar properties to those of Mittag-Leffler function, was introduced by Wiman [20], [21] and is defined by

$$E_{\alpha,\beta}(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\alpha n + \beta)}, \quad \alpha > 0, \beta > 0, z \in \mathcal{U}. \quad (1.3)$$

During the last years the interest in Mittag-Leffler type functions has considerably increased due to their vast potential of applications in applied problems such as fluid flow, electric networks, probability, statistical distribution theory etc. For a detailed account of properties, generalizations and applications of functions (1.2) - (1.3) one may refer to [6], [7], [12], [16].

Geometric properties including starlikeness, convexity and close-to-convexity for the Mittag-Leffler function $E_{\alpha,\beta}(z)$ were recently investigated by Bansal and Prajapat in [1]. Differential subordination results associated with generalized Mittag-Leffler function were also obtained in [14].

The function defined by (1.3) does not belong to the class $\mathcal{A}$. Therefore, we consider the following normalization of the Mittag-Leffler function $E_{\alpha,\beta}(z)$:

$$\mathbb{E}_{\alpha,\beta}(z) = \Gamma(\beta)zE_{\alpha,\beta}(z) = z + \sum_{n=1}^{\infty} \frac{\Gamma(\beta)}{\Gamma(\alpha n + \beta)} z^{n+1}, \quad \alpha > 0, \beta > 0, z \in \mathcal{U}. \quad (1.4)$$

Note that some special cases of $\mathbb{E}_{\alpha,\beta}(z)$ are:

$$\begin{cases} \mathbb{E}_{2,1}(z) = z \cosh \sqrt{z} \\ \mathbb{E}_{2,2}(z) = \sqrt{z} \sinh(\sqrt{z}) \\ \mathbb{E}_{2,3}(z) = 2[\cosh(\sqrt{z}) - 1] \\ \mathbb{E}_{2,4}(z) = 6[\sinh(\sqrt{z}) - \sqrt{z}]/\sqrt{z}. \end{cases} \quad (1.5)$$

Recently, several results related to partial sums of special functions, such as Bessel [10], Struve [22], Lommel [2] and Wright functions [3] were obtained.

Motivated by the work of Bansal and Prajapat [1] and also by the above mentioned results, in this paper we investigate the ratio of normalized Mittag-Leffler function $\mathbb{E}_{\alpha,\beta}(z)$ defined by (1.4) to its sequence of partial sums

$$\begin{cases} (\mathbb{E}_{\alpha,\beta})_0(z) = z \\ (\mathbb{E}_{\alpha,\beta})_m(z) = z + \sum_{n=1}^m A_n z^{n+1}, \quad m \in \mathbb{N} = \{1, 2, \dots\}, \end{cases} \quad (1.6)$$

where

$$A_n = \frac{\Gamma(\beta)}{\Gamma(\alpha n + \beta)}, \quad \alpha > 0, \beta > 0, n \in \mathbb{N}.$$

We obtain lower bounds on ratios like

$$\Re \left\{ \frac{\mathbb{E}_{\alpha,\beta}(z)}{(\mathbb{E}_{\alpha,\beta})_m(z)} \right\}, \Re \left\{ \frac{(\mathbb{E}_{\alpha,\beta})_m(z)}{\mathbb{E}_{\alpha,\beta}(z)} \right\}, \Re \left\{ \frac{\mathbb{E}'_{\alpha,\beta}(z)}{(\mathbb{E}_{\alpha,\beta})'_m(z)} \right\}, \Re \left\{ \frac{(\mathbb{E}_{\alpha,\beta})'_m(z)}{\mathbb{E}'_{\alpha,\beta}(z)} \right\}.$$

Several examples will be also given.

Results concerning partial sums of analytic functions may be found in [4], [8], [11], [13], [17], [18], [19].

2 Main results

In order to obtain our results we need the following lemma.

Lemma 2.1. *Let $\alpha \geq 1$ and $\beta \geq 1$. Then the function $\mathbb{E}_{\alpha,\beta}(z)$ satisfies the next two inequalities:*

$$|\mathbb{E}_{\alpha,\beta}(z)| \leq \frac{\beta^2 + \beta + 1}{\beta^2}, \quad z \in \mathcal{U} \quad (2.1)$$

$$|\mathbb{E}'_{\alpha,\beta}(z)| \leq \frac{\beta^2 + 3\beta + 2}{\beta^2}, \quad z \in \mathcal{U}. \quad (2.2)$$

Proof. Under the hypothesis we have $\Gamma(n + \beta) \leq \Gamma(\alpha n + \beta)$ and thus

$$\frac{\Gamma(\beta)}{\Gamma(\alpha n + \beta)} \leq \frac{\Gamma(\beta)}{\Gamma(n + \beta)} = \frac{1}{(\beta)_n}, \quad n \in \mathbb{N}, \quad (2.3)$$

where

$$(x)_n = \begin{cases} 1 & , n = 0 \\ x(x+1) \dots (x+n-1) & , n \in \mathbb{N} \end{cases}$$

is the well-known Pochhammer symbol.

Note that

$$(x)_n = x(x+1)_{n-1}, \quad n \in \mathbb{N} \quad (2.4)$$

and

$$(x)_n \geq x^n, \quad n \in \mathbb{N}. \quad (2.5)$$

Making use of (2.3) - (2.5) and also of the well-known triangle inequality, for $z \in \mathcal{U}$, we obtain

$$|\mathbb{E}_{\alpha,\beta}(z)| = \left| z + \sum_{n=1}^{\infty} \frac{\Gamma(\beta)}{\Gamma(\alpha n + \beta)} z^{n+1} \right| \leq 1 + \sum_{n=1}^{\infty} \frac{\Gamma(\beta)}{\Gamma(\alpha n + \beta)} \leq 1 + \sum_{n=1}^{\infty} \frac{1}{(\beta)_n}$$

$$\begin{aligned}
 &= 1 + \frac{1}{\beta} \sum_{n=1}^{\infty} \frac{1}{(\beta+1)_{n-1}} \leq 1 + \frac{1}{\beta} \sum_{n=1}^{\infty} \frac{1}{(\beta+1)^{n-1}} = 1 + \frac{1}{\beta} \sum_{n=0}^{\infty} \left(\frac{1}{\beta+1} \right)^n \\
 &= \frac{\beta^2 + \beta + 1}{\beta^2}
 \end{aligned}$$

and thus, inequality (2.1) is proved.

Using once more the triangle inequality, for $z \in \mathcal{U}$, we obtain

$$|\mathbb{E}'_{\alpha,\beta}(z)| = \left| 1 + \sum_{n=1}^{\infty} \frac{(n+1)\Gamma(\beta)}{\Gamma(\alpha n + \beta)} z^n \right| \leq 1 + \sum_{n=1}^{\infty} \frac{n\Gamma(\beta)}{\Gamma(\alpha n + \beta)} + \sum_{n=1}^{\infty} \frac{\Gamma(\beta)}{\Gamma(\alpha n + \beta)}. \quad (2.6)$$

For $\beta \geq 1$ we have

$$\frac{n}{(\beta)_n} = \frac{n}{\beta(\beta+1)_{n-1}} = \frac{n}{\beta(\beta+1)_{n-2}(\beta+n-1)} \leq \frac{1}{\beta(\beta+1)_{n-2}}. \quad (2.7)$$

Taking into account inequalities (2.3) - (2.5) and (2.7), from (2.6), we obtain

$$\begin{aligned}
 |\mathbb{E}'_{\alpha,\beta}(z)| &\leq 1 + \sum_{n=1}^{\infty} \frac{n}{(\beta)_n} + \sum_{n=1}^{\infty} \frac{1}{(\beta)_n} \\
 &\leq 1 + \frac{1}{\beta} + \frac{1}{\beta} \sum_{n=2}^{\infty} \frac{1}{(\beta+1)_{n-2}} + \frac{1}{\beta} \sum_{n=1}^{\infty} \frac{1}{(\beta+1)_{n-1}} \\
 &\leq 1 + \frac{1}{\beta} + \frac{1}{\beta} \sum_{n=0}^{\infty} \left(\frac{1}{\beta+1} \right)^n + \frac{1}{\beta} \sum_{n=0}^{\infty} \left(\frac{1}{\beta+1} \right)^n = \frac{\beta^2 + 3\beta + 2}{\beta^2}
 \end{aligned}$$

and thus, inequality (2.2) is also proved. $\square$

Let $w(z)$ be an analytic function in $\mathcal{U}$. In the sequel, we will frequently use the following well-known result:

$$\Re \left\{ \frac{1+w(z)}{1-w(z)} \right\} > 0, \quad z \in \mathcal{U} \quad \text{if and only if} \quad |w(z)| < 1, \quad z \in \mathcal{U}.$$

Theorem 2.1. *Let $\alpha \geq 1$ and $\beta \geq \frac{1+\sqrt{5}}{2}$. Then*

$$\Re \left\{ \frac{\mathbb{E}_{\alpha,\beta}(z)}{(\mathbb{E}_{\alpha,\beta})_m(z)} \right\} \geq \frac{\beta^2 - \beta - 1}{\beta^2}, \quad z \in \mathcal{U} \quad (2.8)$$

and

$$\Re \left\{ \frac{(\mathbb{E}_{\alpha,\beta})_m(z)}{\mathbb{E}_{\alpha,\beta}(z)} \right\} \geq \frac{\beta^2}{\beta^2 + \beta + 1}, \quad z \in \mathcal{U}. \quad (2.9)$$

Proof. From inequality (2.1) we get

$$1 + \sum_{n=1}^{\infty} A_n \leq \frac{\beta^2 + \beta + 1}{\beta^2}, \text{ where } A_n = \frac{\Gamma(\beta)}{\Gamma(\alpha n + \beta)}, \quad n \in \mathbb{N}.$$

The last inequality is equivalent to

$$\frac{\beta^2}{\beta + 1} \sum_{n=1}^{\infty} A_n \leq 1.$$

In order to prove the inequality (2.8), we consider the function $w(z)$ defined by

$$\frac{1 + w(z)}{1 - w(z)} = \frac{\beta^2}{\beta + 1} \frac{\mathbb{E}_{\alpha, \beta}(z)}{(\mathbb{E}_{\alpha, \beta})_m(z)} - \frac{\beta^2 - \beta - 1}{\beta + 1}$$

or

$$\frac{1 + w(z)}{1 - w(z)} = \frac{1 + \sum_{n=1}^m A_n z^n + \frac{\beta^2}{\beta + 1} \sum_{n=m+1}^{\infty} A_n z^n}{1 + \sum_{n=1}^m A_n z^n}. \quad (2.10)$$

From (2.10), we obtain

$$w(z) = \frac{\frac{\beta^2}{\beta + 1} \sum_{n=m+1}^{\infty} A_n z^n}{2 + 2 \sum_{n=1}^m A_n z^n + \frac{\beta^2}{\beta + 1} \sum_{n=m+1}^{\infty} A_n z^n}$$

and

$$|w(z)| < \frac{\frac{\beta^2}{\beta + 1} \sum_{n=m+1}^{\infty} A_n}{2 - 2 \sum_{n=1}^m A_n - \frac{\beta^2}{\beta + 1} \sum_{n=m+1}^{\infty} A_n}.$$

The inequality $|w(z)| < 1$ holds true if and only if

$$\frac{2\beta^2}{\beta + 1} \sum_{n=m+1}^{\infty} A_n \leq 2 - 2 \sum_{n=1}^m A_n$$

which is equivalent to

$$\sum_{n=1}^m A_n + \frac{\beta^2}{\beta + 1} \sum_{n=m+1}^{\infty} A_n \leq 1. \quad (2.11)$$

To prove (2.11), it suffices to show that its left-hand side is bounded above by

$$\frac{\beta^2}{\beta+1} \sum_{n=1}^{\infty} A_n$$

which is equivalent to

$$\frac{\beta^2 - \beta - 1}{\beta + 1} \sum_{n=1}^m A_n \geq 0.$$

The last inequality holds true for $\beta \geq \frac{1 + \sqrt{5}}{2}$.

We use the same method to prove inequality (2.9). Consider the function $w(z)$ given by

$$\frac{1 + w(z)}{1 - w(z)} = \frac{\beta^2 + \beta + 1}{\beta + 1} \frac{(\mathbb{E}_{\alpha, \beta})_m(z)}{\mathbb{E}_{\alpha, \beta}(z)} - \frac{\beta^2}{\beta + 1}.$$

From the last equality we obtain

$$w(z) = \frac{-\frac{\beta^2 + \beta + 1}{\beta + 1} \sum_{n=m+1}^{\infty} A_n z^n}{2 + 2 \sum_{n=1}^m A_n z^n - \frac{\beta^2 - \beta - 1}{\beta + 1} \sum_{n=m+1}^{\infty} A_n z^n}$$

and

$$|w(z)| < \frac{\frac{\beta^2 + \beta + 1}{\beta + 1} \sum_{n=m+1}^{\infty} A_n}{2 - 2 \sum_{n=1}^m A_n - \frac{\beta^2 - \beta - 1}{\beta + 1} \sum_{n=m+1}^{\infty} A_n}.$$

Then, $|w(z)| < 1$ if and only if

$$\frac{\beta^2}{\beta + 1} \sum_{n=m+1}^{\infty} A_n + \sum_{n=1}^m A_n \leq 1. \quad (2.12)$$

Since the left-hand side of (2.12) is bounded above by

$$\frac{\beta^2}{\beta + 1} \sum_{n=1}^{\infty} A_n$$

we have that the inequality (2.9) holds true. Now, the proof of our theorem is completed. $\square$

In the next theorem we consider ratios involving derivatives.

Theorem 2.2. *Let $\alpha \geq 1$ and let $\beta \geq \frac{3 + \sqrt{17}}{2}$. Then*

$$\Re \left\{ \frac{\mathbb{E}'_{\alpha, \beta}(z)}{(\mathbb{E}_{\alpha, \beta})'_m(z)} \right\} \geq \frac{\beta^2 - 3\beta - 2}{\beta^2}, \quad z \in \mathcal{U} \quad (2.13)$$

and

$$\Re \left\{ \frac{(\mathbb{E}_{\alpha, \beta})'_m(z)}{\mathbb{E}'_{\alpha, \beta}(z)} \right\} \geq \frac{\beta^2}{\beta^2 + 3\beta + 2}, \quad z \in \mathcal{U}. \quad (2.14)$$

Proof. From (2.2) we have

$$1 + \sum_{n=1}^{\infty} (n+1)A_n \leq \frac{\beta^2 + 3\beta + 2}{\beta^2}, \quad \text{where } A_n = \frac{\Gamma(\beta)}{\Gamma(\alpha n + \beta)}, \quad n \in \mathbb{N}.$$

The above inequality is equivalent to

$$\frac{\beta^2}{3\beta + 2} \sum_{n=1}^{\infty} (n+1)A_n \leq 1.$$

To prove (2.13), define the function $w(z)$ by

$$\frac{1 + w(z)}{1 - w(z)} = \frac{\beta^2}{3\beta + 2} \frac{\mathbb{E}'_{\alpha, \beta}(z)}{(\mathbb{E}_{\alpha, \beta})'_m(z)} - \frac{\beta^2 - 3\beta - 2}{3\beta + 2}$$

which gives

$$w(z) = \frac{\frac{\beta^2}{3\beta + 2} \sum_{n=m+1}^{\infty} (n+1)A_n z^n}{2 + 2 \sum_{n=1}^m (n+1)A_n z^n + \frac{\beta^2}{3\beta + 2} \sum_{n=m+1}^{\infty} (n+1)A_n z^n}$$

and

$$|w(z)| < \frac{\frac{\beta^2}{3\beta + 2} \sum_{n=m+1}^{\infty} (n+1)A_n}{2 - 2 \sum_{n=1}^m (n+1)A_n - \frac{\beta^2}{3\beta + 2} \sum_{n=m+1}^{\infty} (n+1)A_n}.$$

The condition $|w(z)| < 1$ holds true if and only if

$$\sum_{n=1}^m (n+1)A_n + \frac{\beta^2}{3\beta + 2} \sum_{n=m+1}^{\infty} (n+1)A_n \leq 1. \quad (2.15)$$

The left-hand side of (2.15) is bounded above by

$$\frac{\beta^2}{3\beta+2} \sum_{n=1}^{\infty} (n+1)A_n \quad \text{if} \quad \frac{\beta^2-3\beta-2}{3\beta+2} \sum_{n=1}^m (n+1)A_n \geq 0$$

which holds true for $\beta \geq \frac{3+\sqrt{17}}{2}$.

The proof of (2.14) follows the same pattern. Consider the function $w(z)$ given by

$$\begin{aligned} \frac{1+w(z)}{1-w(z)} &= \frac{\beta^2+3\beta+2}{3\beta+2} \frac{(\mathbb{E}_{\alpha,\beta})'_m(z)}{\mathbb{E}'_{\alpha,\beta}(z)} - \frac{\beta^2}{3\beta+2} \\ &= \frac{1 + \sum_{n=1}^m (n+1)A_n z^n - \frac{\beta^2}{3\beta+2} \sum_{n=m+1}^{\infty} (n+1)A_n z^n}{1 + \sum_{n=1}^{\infty} (n+1)A_n z^n}. \end{aligned} \quad (2.16)$$

From (2.16), we can write

$$w(z) = \frac{-\frac{\beta^2+3\beta+2}{3\beta+2} \sum_{n=m+1}^{\infty} (n+1)A_n z^n}{2 + 2 \sum_{n=1}^m (n+1)A_n z^n - \frac{\beta^2-3\beta-2}{3\beta+2} \sum_{n=m+1}^{\infty} (n+1)A_n z^n}$$

and

$$|w(z)| < \frac{\frac{\beta^2+3\beta+2}{3\beta+2} \sum_{n=m+1}^{\infty} (n+1)A_n}{2 - 2 \sum_{n=1}^m (n+1)A_n - \frac{\beta^2-3\beta-2}{3\beta+2} \sum_{n=m+1}^{\infty} (n+1)A_n}.$$

The last inequality implies that $|w(z)| < 1$ if and only if

$$\frac{2\beta^2}{3\beta+2} \sum_{n=m+1}^{\infty} (n+1)A_n \leq 2 - 2 \sum_{n=1}^m (n+1)A_n$$

or equivalently

$$\sum_{n=1}^m (n+1)A_n + \frac{\beta^2}{3\beta+2} \sum_{n=m+1}^{\infty} (n+1)A_n \leq 1. \quad (2.17)$$

It remains to show that the left-hand side of (2.17) is bounded above by

$$\frac{\beta^2}{3\beta + 2} \sum_{n=1}^{\infty} (n+1)A_n.$$

This is equivalent to

$$\frac{\beta^2 - 3\beta - 2}{3\beta + 2} \sum_{n=1}^m (n+1)A_n \geq 0 \text{ which holds true for } \beta \geq \frac{3 + \sqrt{17}}{2}.$$

Now, the proof of our theorem is completed. $\square$

3 Examples

In this section we give several examples which illustrate our theorems.

A result involving the functions $\mathbb{E}_{2,2}(z)$ and $\mathbb{E}_{2,3}(z)$, defined by (1.5), can be obtained from Theorem 2.1 by taking $m = 0, \alpha = 2, \beta = 2$ and $m = 0, \alpha = 2, \beta = 3$, respectively.

Corollary 3.1. *The following inequalities hold true:*

$$\Re \left\{ \frac{\sinh(\sqrt{z})}{\sqrt{z}} \right\} \geq \frac{1}{4} = 0,25, \quad \Re \left\{ \frac{\sqrt{z}}{\sinh(\sqrt{z})} \right\} \geq \frac{4}{7} \approx 0,57$$

and

$$\Re \left\{ \frac{\cosh(\sqrt{z}) - 1}{z} \right\} \geq \frac{5}{18} \approx 0,28, \quad \Re \left\{ \frac{z}{\cosh(\sqrt{z}) - 1} \right\} \geq \frac{18}{13} \approx 1,38.$$

Setting $m = 0, \alpha = 2$ and $\beta = 4$ in Theorem 2.1 and Theorem 2.2 respectively, we obtain the next result involving the function $\mathbb{E}_{2,4}(z)$, defined by (1.5), and its derivative.

Corollary 3.2. *The following inequalities hold true:*

$$\Re \left\{ \frac{\sinh(\sqrt{z}) - \sqrt{z}}{z\sqrt{z}} \right\} \geq \frac{11}{96} \approx 0,11, \quad \Re \left\{ \frac{z\sqrt{z}}{\sinh(\sqrt{z}) - \sqrt{z}} \right\} \geq \frac{32}{7} \approx 4,57$$

and

$$\Re \left\{ \frac{\sqrt{z} \cosh(\sqrt{z}) - \sinh(\sqrt{z})}{z\sqrt{z}} \right\} \geq \frac{1}{24} \approx 0,04,$$

$$\Re \left\{ \frac{z\sqrt{z}}{\sqrt{z} \cosh(\sqrt{z}) - \sinh(\sqrt{z})} \right\} \geq \frac{8}{5} = 1,6.$$

Remark 3.1. If we consider $m = 0$ in inequality (2.13), we obtain $\Re \{E'_{\alpha,\beta}(z)\} > 0$. In view of Noshiro-Warschawski Theorem (see [5]), we have that the normalized Mittag-Leffler function is univalent in $\mathcal{U}$ for $\alpha \geq 1$ and $\beta \geq \frac{3 + \sqrt{17}}{2}$.

References

- [1] D. Bansal, J. K. Prajapat, *Certain geometric properties of the Mittag-Leffler functions*, Complex Var. Elliptic Equ., **61**(3)(2016), 338-350.
- [2] M. Çağlar, E. Deniz, *Partial sums of the normalized Lommel functions*, Math. Inequal. Appl., **18**(3)(2015), 1189-1199.
- [3] M. Din, M. Raza, N. Yağmur, *Partial sums of normalized Wright functions*, arXiv:1606.02750v1 [math.CV], 2016.
- [4] B. A. Frasin, *Generalization of partial sums of certain analytic and univalent functions*, Appl. Math. Lett., **21**(2008), 735-741.
- [5] A. W. Goodman, *Univalent functions, vol. I*, Mariner Publishing Company, Inc., 1983.
- [6] R. Gorenflo, F. Mainardi, S. V. Rogosin, *On the generalized Mittag-Leffler type function*, Integral Transform. Spec. Funct., **7**(1998), 215-224.
- [7] I. S. Gupta, L. Debnath, *Some properties of the Mittag-Leffler functions*, Integral Transform. Spec. Funct., **18**(5)(2007), 329-336.
- [8] L. J. Liu, S. Owa, *On partial sums of the Libera integral operator*, J. Math. Anal. Appl., **213**(2)(1997), 444-454.
- [9] G. M. Mittag-Leffler, *Sur la nouvelle fonction $E_\alpha(x)$* , C. R. Acad. Sci. Paris, **137**(1903), 554-558.
- [10] H. Orhan, N. Yağmur, *Partial sums of generalized Bessel functions*, J. Math. Inequal., **8**(4)(2014), 863-877.
- [11] S. Owa, H. M. Srivastava, N. Saitoh, *Partial sums of certain classes of analytic functions*, Int. J. Comput. Math., **81**(10)(2004), 1239-1256.
- [12] T. R. Prabhakar, *A singular integral equation with a generalized Mittag-Leffler function in the kernel*, Yokohama Math. J., **19**(1997), 7-15.
- [13] V. Ravichandran, *Geometric properties of partial sums of univalent functions*, Math. Newslett., **22**(3)(2012), 208-221.

- [14] D. Răducanu, *Differential subordinations associated with generalized Mittag-Leffler functions* (submitted).
- [15] T. O. Salim, *Some properties relating to the generalized Mittag-Leffler function*, Advances Appl. Math. Anal., **4**(1)(2009), 21-30.
- [16] A. K. Shukla, J. C. Prajapati, *On a generalization of Mittag-Leffler function and its properties*, J. Math. Anal. Appl., **337**(2007), 797-811.
- [17] H. Silverman, *Partial sums of starlike and convex functions*, J. Math. Anal. Appl., **209**(1997), 221-227.
- [18] E. M. Silvia, *On partial sums of convex functions of order α* , Houston J. Math., **11**(1985), 397-404.
- [19] A. E. Tudor, D. Răducanu, *On a subclass of analytic functions involving harmonic means*, An. Șt. Univ. Ovidius Constanța, **23**(1)(2015), 267-275.
- [20] A. Wiman, *Über den Fundamental satz in der Theorie der Functionen $E_\alpha(x)$* , Acta Math., **29**(1905), 191-201.
- [21] A. Wiman, *Über die Nullstellun der Functionen $E_\alpha(x)$* , Acta Math., **29**(1905), 217-134.
- [22] N. Yağmur, H. Orhan, *Partial sums of generalized Struve functions*, Miskolc. Math. Notes (accepted.)

Dorina RĂDUCANU,
Faculty of Mathematics and Computer Science,
Transilvania University of Brașov,
Bdul Iuliu Maniu 50, 500091 Brașov, Romania.
Email: draducanu@unitbv.ro