

DOI: 10.1515/ausm-2016-0021

Some Hermite-Hadamard type integral inequalities for operator AG-preinvex functions

Ali Taghavi

Department of Mathematics, Faculty of Mathematical Sciences, University of Mazandaran, Iran email: taghavi@umz.ac.ir

Haji Mohammad Nazari

Department of Mathematics, Faculty of Mathematical Sciences, University of Mazandaran, Iran email: m.nazari@stu.umz.ac.ir Vahid Darvish

Department of Mathematics, Faculty of Mathematical Sciences, University of Mazandaran, Iran email: vahid.darvish@mail.com (vdarvish@wordpress.com)

Abstract. In this paper, we introduce the concept of operator AGpreinvex functions and prove some Hermite-Hadamard type inequalities for these functions. As application, we obtain some unitarily invariant norm inequalities for operators.

1 Introduction and preliminaries

The following Hermite-Hadamard inequality holds for any convex function f defined on $\mathbb R$

$$(b-a)f\left(\frac{a+b}{2}\right) \le \int_{a}^{b} f(x)dx \le (b-a)\frac{f(a)+f(b)}{2}, \quad a,b \in \mathbb{R}.$$
 (1)

2010 Mathematics Subject Classification: 47A63, 15A60, 47B05, 47B10, 26D15 Key words and phrases: Hermite-Hadamard inequality, operator AG-preinvex function, log-convex function, positive linear operator It was firstly discovered by Hermite in 1881 in the journal Mathesis (see [8]). But this result was nowhere mentioned in the mathematical literature and was not widely known as Hermite's result [10].

Beckenbach, a leading expert on the history and the theory of convex functions, wrote that this inequality was proven by Hadamard in 1893 [2]. In 1974, Mitrinovič found Hermites note in Mathesis [8]. Since (1) was known as Hadamards inequality, the inequality is now commonly referred as the Hermite-Hadamard inequality [10].

Definition 1 [13] A continuous function $f : I \subset \mathbb{R} \to \mathbb{R}^+$ is said to be an AG-convex function (arithmetic-geometrically or log-convex function) on the interval I if

$$f(\lambda a + (1 - \lambda)b) \le f(a)^{\lambda} f(b)^{1 - \lambda}.$$
 (2)

for $a, b \in I$ and $\lambda \in [0, 1]$, *i.e.*, log f is convex.

Theorem 1 [13] Let f be an AG-convex function defined on [a, b]. Then, we have

$$f\left(\frac{a+b}{2}\right) \leq \sqrt{f\left(\frac{3a+b}{4}\right)f\left(\frac{a+3b}{4}\right)}$$
$$\leq \exp\left(\frac{1}{b-a}\int_{a}^{b}\log(f(u))du\right)$$
$$\leq \sqrt{f\left(\frac{a+b}{2}\right)} \cdot \sqrt[4]{f(a)} \cdot \sqrt[4]{f(b)}$$
$$\leq \sqrt{f(a)f(b)}, \qquad (3)$$

where $\mathfrak{u} = \log \mathfrak{t}$.

Let B(H) stands for the C^{*}-algebra of all bounded linear operators on a complex Hilbert space H with inner product $\langle \cdot, \cdot \rangle$. An operator $A \in B(H)$ is positive and write $A \ge 0$ if $\langle Ax, x \rangle \ge 0$ for all $x \in H$. Let $B(H)_{sa}$ stand for the set of all self-adjoint elements of B(H).

Let A be a self-adjoint operator in B(H). The Gelfand map establishes a *-isometrically isomorphism Φ between the set C(Sp(A)) of all continuous functions defined on the spectrum of A, denoted by Sp(A), and the C*-algebra C*(A) generated by A and the identity operator 1_H on H as follows:

for any $f, g \in C(Sp(A)))$ and any $\alpha, \beta \in \mathbb{C}$ we have:

• $\Phi(\alpha f + \beta g) = \alpha \Phi(f) + \beta \Phi(g);$

- $\Phi(fg) = \Phi(f)\Phi(g)$ and $\Phi(\bar{f}) = \Phi(f)^*$;
- $\|\Phi(f)\| = \|f\| := \sup_{t \in \operatorname{Sp}(A)} |f(t)|;$
- $\Phi(f_0) = 1_H$ and $\Phi(f_1) = A$, where $f_0(t) = 1$ and $f_1(t) = t$, for $t \in \operatorname{Sp}(A)$.

With this notation we define

$$f(A) = \Phi(f)$$
 for all $f \in C(Sp(A))$

and we call it the continuous functional calculus for a self-adjoint operator A.

If A is a self-adjoint operator and f is a real valued continuous function on Sp(A), then $f(t) \ge 0$ for any $t \in Sp(A)$ implies that $f(A) \ge 0$, i.e. f(A) is a positive operator on H. Moreover, if both f and g are real valued functions on Sp(A) then the following important property holds:

 $f(t) \ge g(t)$ for any $t \in \operatorname{Sp}(A)$ implies that $f(A) \ge g(A)$, (4)

in the operator order of B(H), see [14].

Definition 2 A real valued continuous function f on an interval I is said to be operator convex function if

$$f(\lambda A + (1 - \lambda)B) \le \lambda f(A) + (1 - \lambda)f(B),$$

in the operator order, for all $\lambda \in [0, 1]$ and self-adjoint operators A and B in B(H) whose spectra are contained in I.

In [4] Dragomir investigated the operator version of the Hermite-Hadamard inequality for operator convex functions. Let $f: I \to \mathbb{R}$ be an operator convex function on the interval I then, for any self-adjoint operators A and B with spectra in I, the following inequalities holds

$$\begin{split} f\left(\frac{A+B}{2}\right) &\leq 2\int_{\frac{1}{4}}^{\frac{3}{4}} f(tA+(1-t)B)dt \\ &\leq \frac{1}{2}\left[f\left(\frac{3A+B}{4}\right)+f\left(\frac{A+3B}{4}\right)\right] \\ &\leq \int_{0}^{1} f\left((1-t)A+tB\right)dt \\ &\leq \frac{1}{2}\left[f\left(\frac{A+B}{2}\right)+\frac{f(A)+f(B)}{2}\right] \\ &\leq \frac{f(A)+f(B)}{2}, \end{split}$$

for the first inequality in above, see [12].

In [5], Ghazanfari et al. gave the concept of operator preinvex function and obtained Hermite-Hadamard type inequality for operator preinvex function.

Definition 3 [5] Let X be a real vector space, a set $S \subseteq X$ is said to be invex with respect to the map $\eta : S \times S \to X$, if for every $x, y \in S$ and $t \in [0, 1]$,

$$x + t\eta(x, y) \in S.$$

It is obvious that every convex set is invex with respect to the map $\eta(x, y) = x - y$, but there exist invex sets which are not convex (see [1]).

Let $S \subseteq X$ be an invex set with respect to η . For every $x, y \in S$. the η -path $P_{x\nu}$ joining the points x and $\nu := x + \eta(y, x)$ is defined as follows

$$P_{xv} := \{z : z = x + t\eta(y, x), t \in [0, 1]\}.$$

The mapping η is said to satisfy the condition (C) if for every $x,y\in S$ and $t\in [0,1],$

$$\eta(\mathbf{y},\mathbf{y}+t\eta(\mathbf{y},\mathbf{x}))=-t\eta(\mathbf{x},\mathbf{y}),\quad \eta(\mathbf{x},\mathbf{y}+t\eta(\mathbf{x},\mathbf{y}))=(1-t)\eta(\mathbf{x},\mathbf{y}).$$

Note that for every $x, y \in S$ and every $t_1, t_2 \in [0, 1]$, from conditions in (C), we have

$$\eta(y + t_2\eta(x, y), y + t_1\eta(x, y)) = (t_2 - t_1)\eta(x, y),$$
(5)

see [9] for details.

Definition 4 Let $S \subseteq B(H)_{sa}$ be an invex set with respect to $\eta : S \times S \rightarrow B(H)_{sa}$. Then, the continuous $f : \mathbb{R} \rightarrow \mathbb{R}$ is said to be operator preinvex with respect to η on S, if for every $A, B \in S$ and $t \in [0, 1]$,

$$f(A + t\eta(B, A)) \le (1 - t)f(A) + tf(B), \tag{6}$$

in the operator order in B(H).

Every operator convex function is operator preinvex with respect to the map $\eta(A, B) = A - B$, but the converse does not hold (see [5]).

Theorem 2 [5] Let $S \subseteq B(H)_{sa}$ be an invex set with respect to $\eta : S \times S \rightarrow B(H)_{sa}$ and η satisfies condition (C). If for every $A, B \in S$ and $V = A + \eta(B, A)$ the function $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ is operator preinvex with respect to η on η -path P_{AV} with spectra of A and spectra of V in the interval I. Then we have the inequality

$$f\left(\frac{A+V}{2}\right) \le \int_0^1 f((A+t\eta(B,A))dt \le \frac{f(A)+f(B))}{2}$$

Throughout this paper, we introduce the concept of operator AG-preinvex functions and obtain some Hermite-Hadamard type inequalities for these class of functions. These results lead us to obtain some inequalities unitarily invariant norm inequalities for operators.

2 Some inequalities for operator AG-preinvex functions

In this section, we prove some Hermite-Hadamard type inequalities for operator AG-preinvex functions.

Definition 5 [13] A continuous function $f: I \subseteq \mathbb{R} \to \mathbb{R}^+$ is said to be operator AG-convex (concave) if

$$f(\lambda A + (1 - \lambda)B) \le (\ge) f(A)^{\lambda} f(B)^{1-\lambda}$$

for $0\leq\lambda\leq 1$ and self-adjoint operators A and B in B(H) whose spectra are contained in I.

Example 1 [6, Corollary 7.6.8] Let A and B be to positive definite $n \times n$ complex matrices. For $0 < \alpha < 1$, we have

$$|\alpha A + (1 - \alpha)B| \ge |A|^{\alpha}|B|^{1 - \alpha} \tag{7}$$

where $|\cdot|$ denotes determinant of a matrix.

Let f be an operator AG-convex function, for commutative positive operators $A, B \in B(H)$ whose spectra are contained in I, then we have

$$f\left(\frac{A+B}{2}\right) \leq \int_{0}^{1} \sqrt{f(\alpha A + (1-\alpha)B)f((1-\alpha)A + \alpha B)} d\alpha$$

$$\leq \sqrt{f(A)f(B)}, \qquad (8)$$

(see [13] for more inequalities).

Definition 6 Let $S \subseteq B(H)_{sa}$ be an invex set with respect to $\eta : S \times S \rightarrow B(H)_{sa}$. A continuous function $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^+$ is called operator AG-preinvex with respect to η on S if

$$f(A + t\eta(B, A)) \le f(A)^{1-t}f(B)^t$$

for $t \in [0, 1]$ such that spectra of A and B are contained in I.

Remark 1 Let f be an operator AG-preinvex function, in a commutative case, we then get

$$\begin{array}{rcl} f(A+t\eta(B,A)) &\leq & f(A)^{1-t}f(B)^t \\ &\leq & (1-t)f(A)+tf(B) \\ &\leq & \max\{f(A),f(B)\} \end{array}$$

It means that f is operator quasi preinvex i.e., $f(A+t\eta(B,A)) \le \max\{f(A), f(B)\}$.

We need the following lemma for giving Hermite-Hadamard type inequalities for operator preinvex function.

Lemma 1 Let $S \subseteq B(H)_{sa}$ be an invex set with respect to $\eta : S \times S \to B(H)_{sa}$ and $f : I \subseteq \mathbb{R} \to \mathbb{R}^+$ be a continuous function on the interval I. Suppose that η satisfies condition (C). Then for every $A, B \in S$ and $V = A + \eta(B, A)$ and for some fixed $s \in (0, 1]$ the function f is operator AG-preinvex with respect to η on η -path P_{AV} with spectra of A and V in the interval I if and only if the function $\phi_{A,B}$ defined by

$$\varphi_{A,B}(t) = f(A + t\eta(B, A)) \tag{9}$$

is a log-convex function on [0, 1].

Proof. Let φ be a log-convex function on [0, 1], we should prove that f is operator AG-preinvex with respect to η .

For every $C_1 := A + t_1 \eta(B, A) \in P_{AV}$, $C_2 := A + t_2 \eta(B, A) \in P_{AV}$, fixed $\lambda \in [0, 1]$, by (9) we have

$$\begin{split} f(C_1 + \lambda \eta(C_2, C_1)) &= f(A + t_1 \eta(B, A) + \lambda \eta(A + t_2 \eta(B, A), A + t_1 \eta(B, A))) \\ &= f(A + t_1 \eta(B, A) + \lambda (t_2 - t_1) \eta(B, A)) \\ &= f(A + (t_1 + \lambda t_2 - \lambda t_1) \eta(B, A)) \\ &= f(A + ((1 - \lambda)t_1 + \lambda t_2) \eta(B, A)) \\ &= \phi((1 - \lambda)t_1 + \lambda t_2) \\ &\leq \phi(t_1)^{1 - \lambda} \phi(t_2)^{\lambda} \\ &= (f(A + t_1 \eta(B, A)))^{1 - \lambda} (f(A + t_2 \eta(B, A)))^{\lambda}. \end{split}$$

Conversely, let f be operator AG-preinvex, then, by (6)

$$\begin{split} \phi((1-\lambda)t_1 + \lambda t_2) &= f(A + ((1-\lambda)t_1 + \lambda t_2)\eta(B,A)) \\ &= f(A + t_1\eta(B,A) + \lambda(t_2 - t_1)\eta(B,A)) \\ &= f(A + t_1\eta(B,A) + \lambda\eta(A + t_2\eta(B,A),A + t_1\eta(B,A))) \\ &\leq f(A + t_1\eta(B,A))^{1-\lambda}f(A + t_2\eta(B,A))^{\lambda} \\ &= \phi(t_1)^{1-\lambda}\phi(t_2)^{\lambda}. \end{split}$$

Theorem 3 Let $S \subseteq B(H)_{sa}$ be an invex set with respect to $\eta : S \times S \to B(H)_{sa}$ and $f : I \subseteq \mathbb{R} \to \mathbb{R}^+$ be a continuous function on the interval I. Suppose that η satisfies condition (C). Then for the operator AG-preinvex function f with respect to η on η -path P_{AV} such that spectra of A and V are in I, we have

$$\begin{split} f\left(\frac{A+V}{2}\right) &\leq \sqrt{f\left(\frac{3A+V}{4}\right)f\left(\frac{A+3V}{4}\right)} \\ &\leq \exp\left(\int_{0}^{1}\log(f(A+t\eta(B,A)))dt\right) \\ &\leq \sqrt{f\left(\frac{A+V}{2}\right)}\sqrt[4]{f(A)}\sqrt[4]{f(V)} \\ &\leq \sqrt{f(A)f(V)} \\ &\leq \frac{f(A)+f(V)}{2} \end{split}$$

where $A,B\in S$ and $V=A+\eta(B,A)$ and for some fixed $s\in(0,1]$

Proof. Since f is an operator AG-preinvex function, so by Lemma 1 we have $\varphi(t) = f(A + t\eta(B, A))$ is log-convex on [0, 1].

On the other hand, in [11] we obtained the following inequalities for logconvex function φ on [0, 1]:

$$\varphi\left(\frac{1}{2}\right) \leq \sqrt{\varphi\left(\frac{1}{4}\right)\varphi\left(\frac{3}{4}\right)}$$

$$\leq \exp\left(\int_{0}^{1}\log(\varphi(\mathfrak{u}))d\mathfrak{u}\right)$$

$$\leq \sqrt{\varphi\left(\frac{1}{2}\right)} \cdot \sqrt[4]{\varphi(0)} \cdot \sqrt[4]{\varphi(1)}$$

$$\leq \sqrt{\varphi(0)\varphi(1)}.$$
(10)

By knowing that

$$\begin{split} \phi(0) &= f(A) \\ \phi\left(\frac{1}{4}\right) &= f\left(A + \frac{1}{4}\eta(B,A)\right) = f\left(\frac{3A + V}{4}\right) \\ \phi\left(\frac{1}{2}\right) &= f\left(A + \frac{1}{2}\eta(B,A)\right) = f\left(\frac{A + V}{2}\right) \\ \phi(1) &= f(V), \end{split}$$

we obtain

$$\begin{split} f\left(\frac{A+V}{2}\right) &\leq \sqrt{f\left(\frac{3A+V}{4}\right)f\left(\frac{A+3V}{4}\right)} \\ &\leq \exp\left(\int_{0}^{1}\log(f(A+t\eta(B,A)))dt\right) \\ &\leq \sqrt{f\left(\frac{A+V}{2}\right)}\sqrt[4]{f(A)}\sqrt[4]{f(V)} \\ &\leq \sqrt{f(A)f(V)}. \end{split}$$

3 Some unitarily invariant norm inequalities for operator AG-preinvex functions

In this section we prove some unitarily invariant norm inequalities for operators.

We consider the wide class of unitarily invariant norms $||| \cdot |||$. Each of these norms is defined on an ideal in B(H) and it will be implicitly understood that when we talk of |||T|||, then the operator T belongs to the norm ideal associated with $||| \cdot |||$. Each unitarily invariant norm $||| \cdot |||$ is characterized by the invariance property |||UTV||| = |||T||| for all operators T in the norm ideal associated with $||| \cdot |||$ and for all unitary operators U and V in B(H).

For $1 \leq p < \infty$, the Schatten p-norm of a compact operator A is defined by $\|A\|_p = (\operatorname{Tr} |A|^p)^{1/p}$, where Tr is the usual trace functional. Note that for compact operator A we have, $\|A\| = s_1(A)$, and if A is a Hilbert-Schmidt operator, then $\|A\|_2 = (\sum_{j=1}^{\infty} s_j^2(A))^{1/2}$. These norms are special examples of the more general class of the Schatten p-norms which are unitarily invariant [3].

Remark 2 The author of [7] proved that if $A, B, X \in B(H)$ such that A, B are positive operators, then for $0 \le \nu \le 1$ we have

$$|||A^{\nu}XB^{1-\nu}||| \le |||AX|||^{\nu}|||XB|||^{1-\nu}.$$
(11)

Let X = I in above inequality, we then get

$$|||A^{\nu}B^{1-\nu}||| \le |||A|||^{\nu}|||B|||^{1-\nu}.$$
(12)

Lemma 2 Let f be an operator AG-preinvex function and η satisfies the condition (C). Then the function $\varphi_{A,B} : [0,1] \to \mathbb{R}$ defined as follows

$$\varphi(t) = |||f(A + t\eta(B, A))|||$$

is log-convex.

Proof. Let $t_1, t_2 \in [0, 1]$, we have

$$\begin{split} \phi((1-\lambda)t_1+\lambda t_2) &= & |||f(A+((1-\lambda)t_1+\lambda t_2)\eta(B,A))||| \\ &= & |||f(A+t_1\eta(B,A)+\lambda(t_2-t_1)\eta(B,A))||| \\ &= & |||f(A+t_1\eta(B,A)+\lambda\eta(A+t_2\eta(B,A),A+t_1\eta(B,A)))||| \\ &\leq & |||f(A+t_1\eta(B,A))^{1-\lambda}f(A+t_2\eta(B,A))^{\lambda}||| \\ &\leq & |||f(A+t_1\eta(B,A))|||^{1-\lambda}|||f(A+t_2\eta(B,A))|||^{\lambda} \text{ by } (12) \\ &= & \phi(t_1)^{1-\lambda}\phi(t_2)^{\lambda}. \end{split}$$

Theorem 4 Let $S \subseteq B(H)_{sa}$ be an invex set with respect to $\eta : S \times S \to B(H)_{sa}$ and $f : I \subseteq \mathbb{R} \to \mathbb{R}^+$ be a continuous function on the interval I. Suppose that η satisfies condition (C). Then for the operator AG-preinvex function f with respect to η on $\eta\mbox{-path}\ P_{AV}$ such that spectra of A and V are in I, we have

$$\begin{split} \left| \left| \left| \left| f\left(\frac{A+V}{2}\right) \right| \right| &\leq \sqrt{\left| \left| \left| f\left(\frac{3A+V}{4}\right) \right| \right| \right| \left| \left| \left| \left| \left| f\left(\frac{A+3V}{4}\right) \right| \right| \right| \right|} \\ &\leq \exp\left(\int_{0}^{1} \log(|||f(A+t\eta(B,A))|||) dt \right) \\ &\leq \sqrt{\left| \left| \left| f\left(\frac{A+V}{2}\right) \right| \right| \right|} \sqrt[4]{|||f(A)|||} \sqrt[4]{|||f(V)||} \\ &\leq \sqrt{|||f(A)||| |||f(V)|||} \\ &\leq \frac{|||f(A)||| + |||f(V)|||}{2}. \end{split}$$

where $A,B\in S$ and $V=A+\eta(B,A)$ and for some fixed $s\in(0,1]$

Proof. Since f is an operator AG-preinvex function, so by Lemma 2 we have $\varphi(t) = |||f(A + t\eta(B, A))|||$ is log-convex on [0, 1].

On the other hand, in [11] we obtained the following inequalities for logconvex function φ on [0, 1]:

$$\begin{aligned} \varphi\left(\frac{1}{2}\right) &\leq \sqrt{\varphi\left(\frac{1}{4}\right)\varphi\left(\frac{3}{4}\right)} \\ &\leq \exp\left(\int_{0}^{1}\log(\varphi(u))du\right) \\ &\leq \sqrt{\varphi\left(\frac{1}{2}\right)} \cdot \sqrt[4]{\varphi(0)} \cdot \sqrt[4]{\varphi(1)} \\ &\leq \sqrt{\varphi(0)\varphi(1)}. \end{aligned} \tag{13}$$

By knowing that

$$\begin{split} \varphi(0) &= \||f(A)\|| \\ \varphi\left(\frac{1}{4}\right) &= \left\| \left\| f\left(A + \frac{1}{4}\eta(B, A)\right) \right\| \right\| = \left\| \left\| f\left(\frac{3A + V}{4}\right) \right\| \\ \varphi\left(\frac{1}{2}\right) &= \left\| \left\| f\left(A + \frac{1}{2}\eta(B, A)\right) \right\| \right\| = \left\| \left\| f\left(\frac{A + V}{2}\right) \right\| \\ \varphi(1) &= \left\| \|f(V)\| \right\|, \end{split}$$

we obtain

$$\begin{split} \left| \left| \left| f\left(\frac{A+V}{2}\right) \right| \right| &\leq \sqrt{\left| \left| \left| f\left(\frac{3A+V}{4}\right) \right| \right| \right| \left| \left| \left| \left| \left| f\left(\frac{A+3V}{4}\right) \right| \right| \right| \right|} \\ &\leq \exp\left(\int_{0}^{1} \log(\left| \left| \left| f(A+t\eta(B,A)) \right| \right| \right|) dt \right) \\ &\leq \sqrt{\left| \left| \left| f\left(\frac{A+V}{2}\right) \right| \right| \right|} \sqrt[4]{\left| \left| \left| f(A) \right| \right| \left| \frac{4}{\sqrt{\left| \left| \left| f(V) \right| \right| \right|}} \right|} \\ &\leq \sqrt{\left| \left| \left| f(A) \right| \left| \left| \left| \left| \left| f(V) \right| \right| \right| \right|}. \end{split}$$

Let $\eta(B,A)=B-A$ in the above theorem, then we obtain the following inequalities:

$$\begin{aligned} \left| \left| \left| f\left(\frac{A+B}{2}\right) \right| \right| &\leq \sqrt{\left| \left| \left| f\left(\frac{3A+B}{4}\right) \right| \right| \right| \left| \left| \left| \left| f\left(\frac{A+3B}{4}\right) \right| \right| \right| \right|} \\ &\leq \exp\left(\int_{0}^{1} \log(|||f((1-t)A+tB)||| dt \right) \\ &\leq \sqrt{\left| \left| \left| f\left(\frac{A+B}{2}\right) \right| \right| \right|} \sqrt[4]{\left| \left| \left| f(A) \right| \right| \left| \left| \left| f(B) \right| \right| \right|} \\ &\leq \sqrt{\left| \left| \left| f(A) \right| \right| \left| \left| \left| \left| f(B) \right| \right| \right| } \\ &\leq \frac{\left| \left| \left| f(A) \right| \right| \left| \left| \left| \left| f(B) \right| \right| \right| }{2}. \end{aligned} \right|$$
(14)

References

- T. Antczak, Mean value in invexity analysis, Nonlinear Anal., 60 (2005), 1473–1484
- [2] E. F. Beckenbach, Convex functions, Bull. Amer. Math. Soc., 54 (1948), 439–460.
- [3] R. Bhatia, *Matrix Analysis*, GTM 169, Springer-Verlag, New York, 1997.
- [4] S. S. Dragomir, Hermite-Hadamards type inequalities for operator convex functions, Applied Mathematics and Computation., 218 (2011), 766–772.

- [5] A. G. Ghazanfari, M. Shakoori, A. Barani, S. S. Dragomir, Hermite-Hadamard type inequality for operator preinvex functions, arXiv:1306.0730v1
- [6] R. A. Horn, C. R. Johnson, *Matrix Analysis*, Cambridge University Press, 2012.
- [7] F. Kittaneh, Norm inequalities for fractional powers of positive operators, *Lett. Math. Phys.*, 27 (1993), 279–285.
- [8] D. S. Mitrinović, I. B. Lacković, Hermite and convexity, Aequationes Math., 28 (1985), 229–232.
- [9] S. R. Mohan, S. K. Neogy, On invex sets and preinvex function, J. Math. Anal. Appl., 189 (1995), 901–908.
- [10] J. E. Pečarić, F. Proschan, Y. L. Tong, Convex Functions, Partial Orderings, and Statistical Applications, Academic Press Inc., San Diego, 1992.
- [11] A. Taghavi, V. Darvish, H. M. Nazari, S. S. Dragomir, Hermite-Hadamard type inequalities for operator geometrically convex functions, *Monatsh. Math.* 10.1007/s00605-015-0816-6.
- [12] A. Taghavi, V. Darvish, H. M. Nazari, S. S. Dragomir, Some inequalities associated with the Hermite-Hadamard inequalities for operator hconvex functions, Accepted for publishing by J. Adv. Res. Pure Math., (http://rgmia.org/papers/v18/v18a22.pdf)
- [13] A. Taghavi, V. Darvish, H. M. Nazari, Some integral inequalities for operator arithmetic-geometrically convex functions, arXiv:1511.06587v1
- [14] K. Zhu, An introduction to operator algebras, CRC Press, 1993.

Received: December 18, 2015