

Checklist of the vascular flora of Wielkopolska (Poland): casual alien species

Bogdan Jackowiak*, Zbigniew Celka, Julian Chmiel, Karol Latowski
 & Waldemar Żukowski

Department of Plant Taxonomy, Faculty of Biology, Adam Mickiewicz University, Umultowska 89, 61-614 Poznań, Poland
 * corresponding author (e-mail: bogjack@amu.edu.pl)

Abstract. The list of alien vascular plant species only temporarily occurring in Wielkopolska refers to the previously published list of native and permanently established plants. Together, these two lists document the vascular flora of this region at the beginning of the 21st century. The current list, like the previous one, is a result of critical analysis of both contemporary and historical data, collected since the beginning of the 19th century. All information accessible in herbarium collections, publications and unpublished materials was used. A critical analysis was conducted at the taxonomic, nomenclatural, chorological and habitat levels, based on the verification of negative information not supported by sufficient arguments. The list is presented in an alphabetical order. Information on each species includes: family affinity, life form, geographic and historic status. In cases particularly disputable, the standard characteristic of a species was supplemented with an additional commentary.

Key words: regional biodiversity assessment, flora, vascular plants, alien species, synanthropisation, Great Poland, Central Europe

1. Introduction

The appearance of alien species in the flora has been a subject of intensive research for a long time. This phenomenon was noticed already in the mid-nineteenth century by De Candolle (1855), who introduced the term of an adventive plant to literature. The classification regarding the time of introduction and the stability of occurrence of alien species in a new region was of key importance for the development of research in this area (Thellung 1918/1919). With reference to the concept of A. Thellung, many variants of the classification of alien plants and definitions of particular groups of anthropophytes were created in the twentieth century (e.g., Holub & Jirásek 1967; Kornaś 1968, 1977, 1981; Schroeder 1969; Jackowiak 1990; Protopopova 1991). Finally, an attempt was made to put in order the classification system and definitions (Pyšek *et al.* 2004), because research in this field is developing very dynamically.

In many countries and regions, the appearance of alien species is carefully analyzed and evaluated in

terms of biodiversity changes and the invasion process. Alien plant species were commonly recognized in the floras of many countries (Crawley *et al.* 1996; Celesti-Grapow *et al.* 2009; Pyšek *et al.* 2012; Protopopova & Shevera 2014; Elvisto *et al.* 2016). This phenomenon has also been summed up on a European scale (Lambdon *et al.* 2008). However, it is still necessary to monitor this phenomenon both on a regional and local scale, and to pay even greater attention to casual alien species.

The subject of this paper are alien species occurring accidentally in Wielkopolska (western part of Poland) over the whole period of geobotanical research conducted in this region. The aim of the study was to evaluate the changes in regional flora resulting from the presence of accidentally introduced alien species. The checklist of casual alien species supplements the list of native and permanently established species of Wielkopolska (Jackowiak *et al.* 2013b).

The observations of alien plant species appearing in Wielkopolska have a long history (Jackowiak *et al.* 2013a). The first scientific work on this region was

published at the end of the 19th century: “Immigration of some plants in the city of Poznań after 1850 year ...” (Pfuhl 1896). In the first half of the twentieth century, two works are primarily noteworthy: (*i*) “The newcomers and waifs in the vegetation of Poznań” (Szulczewski 1931) and (*ii*) “Geographical analysis of the synanthropic flora of the city of Poznań” (Krawiecka 1951). Both in the second half of the twentieth century and presently, foreign plant species have been as standard included in local floristic studies. Particular attention has been paid to alien plant species in geobotanical monographs devoted to anthropogenic changes of the flora of Poznań (Jackowiak 1990, 1993), Gniezno Lake District (Chmiel 1993) and archeological objects (Celka 1999, 2004). A lot of data from Wielkopolska has been included in works concerning the entire country (Rostański & Sowa 1986/1987; Tokarska-Guzik 2005; Urbisz 2012).

2. Material and methods

2.1. Limits of the research area and material

The area of study almost entirely coincides with the Wielkopolska-Kujawy Lowland, thus, it corresponds to a second-level unit in the geobotanical division of Poland (Szafer 1972). The research area also included fragments of two other geobotanical units: the Northern Marginal Plateau Region and Trzebnicko-Ostrzeszowskie Hills.

Similarly as in our previous publications (e.g., Żukowski & Jackowiak 1995; Jackowiak *et al.* 2013a, 2013b), Wielkopolska is defined in this study within broad limits, covering both Kujawy (in the east) and Ziemia Lubuska (Central Nadodrza) in the west (Fig. 1). In total, the area of research covered about 50 thousand km².

The list of plant species found in the Wielkopolska region includes data from all available literature sources, as well as materials collected in the herbarium of Adam Mickiewicz University (POZ), and verified own and other authors' data not yet published. The main list of sources is included in the work on the history of floristic studies in Wielkopolska (Jackowiak *et al.* 2013a). Currently, it has been supplemented with publications that appeared in 2013–2017.

2.2. Taxonomy, nomenclature and basis for species classification

The alphabetical list of alien species not established in the vascular flora of Wielkopolska includes: name of the species and its author, in some cases also synonyms, family name, life form, information on native range, classification according to the source of introduction, and data regarding the observation period. Additional notes have been added to several dozen of species.

The taxonomy, nomenclature and biological form of species, as well as the region of origin, were determined primarily on the basis of the flora of Rothmaler *et al.* (2005), Jäger *et al.* (2008) and the Plant List (2013). In




Fig. 1. Location and borders of the Wielkopolska Region (according to Jackowiak *et al.* 2007)

general, species names are given, only in justified cases, subspecies names were used (e.g. *Lathyrus japonicus* Willd. subsp. *maritimus* (L.) P. W. Ball). The list also includes hybrid species (e.g., *Mentha ×piperita* L.).

The following categories of plants are included in the classification of biological forms: (1) annual plants, including summer annual plants that sprout, flower, produce seeds, and die during the warmer months of the year, as well as annual plants that germinate in autumn or winter, live through the winter, then bloom in winter or spring; (2) biennial plants that take two years to complete their biological lifecycle. In the first year, the plant grows leaves, stems, and roots (vegetative structures), then it enters a period of dormancy over the colder months. Usually, the stem remains very short and leaves are low to the ground, forming a rosette; (3) perennial, monocarpic plants that live more than two years, but they die after the first fruiting; (4) perennial, polycarpic plants that live, bloom and bear fruit for more than two years; (5) dwarf shrubs and sub-shrubs; (6) shrubs and trees.

In terms of the source of the introduction, casual alien species (diaphytes sensu Kornaś 1981) were divided into two subgroups: ephemeralophytes and ergasiophygophytes (Thellung 1918/19). The first sub-group includes species not cultivated in Wielkopolska and neighboring regions, but brought from other areas ("occasional escapees" sensu Munz 1968, and "ephemeral taxa" sensu Elven & Elvebak 1996). They correspond to the term "adventive" introduced

by De Candolle (1855), later often used both in this and a much broader sense (Muhlenbach 1979; Burda 1991; Provost 1998). Only for species belonging to this subgroup, the region of origin was given. The second subgroup includes casual alien species spreading from cultivation ("waifs" sensu Hickman 1993).

Important information, especially in the case of ephemeral species, is their observation date. Taking into account the history of floristic research in Wielkopolska (Jackowiak *et al.* 2013a), data on the appearance of species were divided into four periods: (1) until 1901, (2) in the years 1901-1951; (3) in the years 1952-2001; (4) after 2001. It was also indicated whether the species was observed once or repeatedly.

3. Results

3.1. Systematic structure of casual alien plant flora

The list of alien plants appearing ephemerally in Wielkopolska from the beginning of the 19th century to the present includes 481 species (Appendices 1-2). They represent 297 genera and 80 families.

The distribution of species within families is very diverse. Close to 60% of the flora of casual alien plants consists of species belonging to 10 families that are the most numerous in Wielkopolska (Table 1). On the other hand, worth of noting is the high number of families represented by a single species (33 families), and two species (14 families).

Table 1. Species diversity of families represented in the flora of casual alien plants of Wielkopolska

Family name	Number of species in the family	Number of families	Total number of species	%
Asteraceae	62	1	62	12.9
Poaceae	46	1	46	9.6
Brassicaceae	37	1	37	7.7
Rosaceae	34	1	34	7.1
Fabaceae	30	1	30	6.2
Liliaceae	21	1	21	4.4
Lamiaceae	19	1	19	4.0
Caryophyllaceae	15	1	15	3.1
Apiaceae	14	1	14	2.9
Chenopodiaceae, Solanaceae	12	2	24	5.0
Onagraceae, Polygonaceae	9	2	18	3.7
Malvaceae	8	1	8	1.7
Boraginaceae, Papaveraceae, Pinaceae	7	3	21	4.4
Amaranthaceae, Cucurbitaceae, Ranunculaceae, Salicaceae	6	4	24	5.0
Campanulaceae, Cuscutaceae, Scrophulariaceae	5	3	15	3.1
Caprifoliaceae, Euphorbiaceae, Linaceae, Resedaceae, Rubiaceae	4	5	20	4.2
Convolvulaceae, Fumariaceae, Plantaginaceae, Violaceae	3	4	12	2.5
see Appendix 1 and 2	2	14	28	5.8
see Appendix 1 and 2	1	33	33	6.9
Total		80	481	100.0

Table 2. Species diversity of genera represented in the flora of casual alien plants of Wielkopolska

Names of genera	Number of species in genus	Number of genera	Total number of species	%
<i>Allium</i>	9	1	9	1.9
<i>Oenothera</i>	8	1	8	1.7
<i>Bromus, Chenopodium, Rosa</i>	7	3	21	4.4
<i>Amaranthus, Artemisia, Centaurea, Spiraea, Vicia</i>	6	5	30	6.2
<i>Cuscuta, Lathyrus, Silene</i>	5	3	15	3.1
<i>Linum, Pinus, Prunus, Rumex, Salix, Solanum</i>	4	6	24	5.0
see Appendix 1 and 2	3	21	63	13.1
see Appendix 1 and 2	2	54	108	22.5
see Appendix	1	203	203	42.2
Total	297	481	100.0	

The distribution of species in genera is very characteristic. The most numerous group consists of genera represented by single or two species. These genera constitute together nearly 65% of the analyzed flora (Table 2).

3.2 Biological spectrum of casual alien plant flora

In the flora of casual alien plants of Wielkopolska dominate annual species, whose share exceeds 42% (Fig. 2). A significant share have also perennial and polycarpic herbaceous plants, followed by trees and shrubs. On the other hand, biennial plants and shrubs and semi-shrubs are of minor importance. The group "other" includes species that may occur in various

biological forms, but the lack of information from the region prevents their unambiguous determination.

3.3. Geographical origin, sources of dispersion and the frequency of records of casual alien species

Among casual alien species, escapees from crops predominate, constituting 60.3% of the analyzed flora (290 species). Thus, the share of foreign plants, introduced directly from other geographical areas, is 39.7% (191 species). In many cases, it was not easy to classify a species into a proper group, due to the limited knowledge of crops grown within the contemporary borders of Wielkopolska, especially in the 19th and the first half of the 20th century. For this reason, the



**Fig. 2.** Share of biological forms in the flora of casual alien plants of Wielkopolska

Table 3. The number of ephemeral species representing different continents and range groups in the flora of Wielkopolska

status of certain species may prove to be controversial.

The frequency of records of casual alien species is very low. Over 70% of them were observed only once (338 species), while about 30% were recorded from two to several times (136 species). It is worth noting that there is a significant difference between the two subgroups of casual alien species. Over 81% of ephemeral species (155 species) were recorded only once. However, ergasiophytes are significantly more frequently observed, of which about 63.1% of species have been observed only once (183 species).

The phytogeographic analysis of ephemeral species shows that in the flora of Wielkopolska, plants from all continents are represented (Table 3). The most numerous group consists of alien species whose native ranges cover Europe (66.5%). They include species with very different primary ranges covering: only Europe (30 species), southern and southwestern Europe and northern Africa (39 species), Eurasia (34 species) and finally Europe, Asia and North Africa (22 species). The second place is occupied by a group of species covering different parts of the African continent (39.3%). It is worth noting here that next to the taxa with a wider intercontinental range, there are also plants whose native range is

limited to Africa (9 species). The third place is taken by a group of plants distributed in Asia (18.3%), including 8 species found naturally only on this continent. The participation of elements representing North America and South America, including Central America is the same (11.5% each). Among the ephemeral species, one species originating from Australia was recorded.

4. Conclusion

The full list of vascular plants found in the Wielkopolska region during nearly 200 years includes more than 2,230 species (Jackowiak *et al.* 2013b). A significant part of the flora is composed of non-established species (481). This means that every fifth species (21.6%) occurs only temporarily. As the history of the invasion of many plant species shows, it is possible that at least some of casual alien taxa may become permanently established in the future. Therefore, regular monitoring of casual alien species is required. This applies to both escapees from crops, which prevail on the checklist of flora of Wielkopolska, and ephemeral species that spread regardless of human activity.

References

- ANTKOWIAK W., CZARNA A. & WAWRZYNIAK M. 2008. *Pyrus ×myoslavensis* (*P. communis* L. × *P. salicifolia* Pall.) – a new spontaneous pear hybrid. *Dendrobiology* 60: 45-49.
- BURDA R. I. 1991. *Antropogennaia Transformacia Flory*. Naukova Dumka, Kiev.
- CELESTI-GRAPOW L., ALESSANDRINI, A., ARRIGONI P. V., ASSINI S., BANFI E., BARNI E., BOVIO M., BRUNDU G., CAGIOTTI M. R., CAMARDA I., CARLI E., CONTI F., DEL GUACCHIO E., DOMINA G., FASCETTI S., GALASSO G., GUBELLINI L., LA VALVA V., LUCCHESE F., MARCHIORI S., MAZZOLA P., PECCENINI S., POLDINI L., PRETTO F., PROSSER F., SINISCALCO C., VILLANI M. C., VIEGI L., WILHALM T. & BLASI C. 2009. Inventory of the non-native flora of Italy. *Plant Biosystems* 143 (2): 386-430.
- CELKA Z. 1999. Rośliny naczyniowe grodzisk Wielkopolski. Prace Zakładu Taksonomii Roślin UAM w Poznaniu 9: 1-159. Bogucki Wyd. Nauk., Poznań.
- CELKA Z. 2004. Distribution Atlas of Vascular Plants on the Earthworks of Wielkopolska. Publications of the Department of Plant Taxonomy of Adam Mickiewicz University in Poznań 13: 1-447. Bogucki Wyd. Nauk., Poznań.
- CHMIEL J. 1993. Flora roślin naczyniowych wschodniej części Pojezierza Gnieźnieńskiego i jej antropogeniczne przeobrażenia w wieku XIX i XX, cz. 1 i 2. Prace Zakładu Taksonomii Roślin UAM w Poznaniu 1; 1: 1-202, 2: 1-212. Wyd. Sorus, Poznań.
- CHMIEL J. 2006. Zróżnicowanie przestrzenne flory jako podstawa ochrony przyrody w krajobrazie rolniczym. Prace Zakładu Taksonomii Roślin UAM w Poznaniu 14: 1-250. Bogucki Wyd. Nauk., Poznań.
- CRAWLEY M. J., HARVEY P. H. & PURVIS A. 1996. Comparative ecology of the native and alien floras of the British Isles. *Phil. Trans. R. Soc. Lond. B* 351(1345): 1251-1259. DOI: 10.1098/rstb.1996.0108
- CZARNA A. 2003. *Cicerbita macrophylla* (Willd.) Wallr. w Wielkopolsce. *Rocznik AR Pozn.* 354, ser. Bot. 6: 15-16.
- CZARNA A. 2005. Vascular flora of the railway station in Jarocin (Western Poland). *Rocznik AR Pozn.* 373, ser. Bot.-Stec. 9: 39-46.
- CZARNA A. 2007. Current flora of herbaceous plants in castle park in Goluchów (Wielkopolska region, Poland). *Rocznik AR Pozn.* 384, ser. Bot.-Stec. 11: 35-47.
- CZARNA A. 2009. Rośliny naczyniowe środkowej Wielkopolski. 184 pp. Wyd. Uniwersytetu Przyrodniczego w Poznaniu.
- CZARNA A. & MIELCARSKI Cz. 2004. *Salvia glutinosa* L. na terenie miasta Poznania. *Rocznik AR Pozn.* CCCLXIII, ser. Bot. 7: 47-51.
- DANIELEWICZ W. & MALIŃSKI T. 1994. Obce gatunki drzew i krzewów w Wielkopolskim Parku Narodowym. *Przegl. Przyrodn.* 5(3/4): 181-190.
- DANIELEWICZ W. & MALIŃSKI T. 1995. Materiały do znajomości dendroflory Wielkopolskiego Parku Narodowego. *Morena* 3: 7-27.
- DANIELEWICZ W. & MALIŃSKI T. 1997. Drzewa i krzewy obcego pochodzenia w lasach Wielkopolskiego Parku Narodowego. *Rocznik Dendrol.* 45: 65-81
- De Candolle A. P. 1855. *Géographie Botanique Raisonné*, vol. 2. V. Masson, Paris.
- DOLATOWSKI J. & SENETA W. 2008. *Dendrologia*. 544 pp. Wyd. Nauk. PWN, Warszawa.
- ELVEN R. & ELVEBAKK A. 1996. Part 1. Vascular plants. In: A. ELVEBAKK & P. PRESTRUÐ (eds.). *A Catalogue of Svalbard Plants, Fungi, Algae, and Cyanobacteria*. Norsk Polarinstitut Skrifter 198: 9-55.
- ELVISTO T., MARGUS PENSA M. & PALUOJA E. 2016. Indigenous and alien vascular plant species in a northern European urban setting (Tallinn, Estonia). *Plant Geography. Proceedings of the Estonian Academy of Sciences* 64 (4): 431-441.
- GĄBKA M. & OWIANNY P. M. 2009. First records of the *Hygrophila polysperma* (Roxb.) T. Anderson (Acanthaceae) in Poland. *Roczniki Akademii Rolniczej w Poznaniu. Botanika-Steciana* 13: 9-14.
- GOLENIA A. & BAYEROWA H. 1956. Kanianka polna (*Cuscuta arvensis* Breyr. var. *calycina* Engelm.; *C. pentagona*, *calycina* Engelm.) i jej nowe stanowiska w Polsce. *Fragm. Flor. Geobot.* 2(2): 5-9.
- HICKMAN J. C. (ed.). 1993. *The Jepson Manual*. Univ. California Press, Berkeley.
- HOLUB J. & JIRÁSEK V. 1967. Zur Vereinheitlichung der Terminologie in der Phytogeographie. *Folia Geobot. Phytotax.* 2: 69-113.
- JACKOWIAK B. 1990. Antropogenicne przemiany flory roślin naczyniowych Poznania. Wyd. Nauk. UAM, seria Biologia, 42, 232 pp. Poznań.
- JACKOWIAK B. 1993. Atlas of distribution of vascular plants in Poznań. Publications of the Department of Plant Taxonomy of the Adam Mickiewicz University in Poznań 3: 1-409.
- JACKOWIAK B., CELKA Z., CHMIEL J., LATOWSKI K. & ŻUKOWSKI W. 2007. Red list of vascular flora of Wielkopolska (Poland). *Biodiv. Res. Conserv.* 5-8: 95-127.
- JACKOWIAK B., CELKA Z., CHMIEL J., LATOWSKI K. & ŻUKOWSKI W. 2013a. The history of research on the vascular flora of Wielkopolska (Poland). *Biodiv. Res. Conserv.* 31: 1-8.
- JACKOWIAK B., CELKA Z., CHMIEL J., LATOWSKI K. & ŻUKOWSKI W. 2013b. Checklist of the vascular flora of Wielkopolska (Poland): native species and naturalized alien species. *Biodiv. Res. Conserv.* 31: 9-96.
- JÄGER E. J., EBEL F., HANELT P. & MÜLLER G. K. (eds.). 2008. *Exkursionsflora von Deutschland*. Band 5, Krautige Zier- und Nutzpflanzen. 880 pp. Spektrum Akademischer Verlag, Springer-Verlag Berlin, Heidelberg.
- KAŻMIERCZAKOWA R. 2014a. CR *Artemisia pontica* L. In: R. KAŻMIERCZAKOWA, K. ZARZYCKI & Z. MIREK (eds.). *Polska Czerwona Księga Roślin. Paprotniki i rośliny kwiatowe*, wyd. 3, pp. 521-522. PAN, Instytut Ochrony Przyrody, Kraków.
- KAŻMIERCZAKOWA R. 2014b. VU *Reseda phytœuma* L. In: R. KAŻMIERCZAKOWA, K. ZARZYCKI & Z. MIREK (eds.). *Polska Czerwona Księga Roślin. Paprotniki i rośliny kwiatowe*, wyd. 3, pp. 234-236. PAN, Instytut Ochrony Przyrody, Kraków.

- KORNAŚ J. 1968. Geograficzno-historyczna klasyfikacja roślin synantropijnych. Mater. Zakł. Fitosoc. Stos. U.W. 25: 33-41.
- KORNAŚ J. 1977. Analiza flor synantropijnych. Wiad. Bot. 21(2): 85-91.
- KORNAŚ J. 1981. Oddziaływanie człowieka na florę: mechanizmy i konsekwencje. Wiad. Bot. 25(3): 165-182.
- KRAWIECOWA A. 1951. Analiza geograficzna flory synantropijnej miasta Poznania. PTPN, Prace Kom. Biol. 13(1): 1-131.
- LADEMANN O. 1937. Adventivpflanzen der östlichen Niedersachsen. Verh. Bot. Ver. Prov. Brandenburg 77: 1-24.
- LAMBDON P. W., PYŠEK P., BASNOU C., HEJDA M., ARIANOUTSOU M., ESSL F., JAROŠÍK V., PERGL J., WINTER M., ANASTASIU P., ANDRIOPoulos P., BAZOS I., BRUNDU G., CELESTI-GRAPOW L., CHASSOT P., DELIPETROU P., JOSEFSSON M., KARK S., KLOTZ S., KOKKORIS Y., KÜHN I., MARCHANTE H., PERGLOVÁ I., PINO J., VILÀ M., ZIKOS A., ROY D. & HULME P. E. 2008. Alien flora of Europe: species diversity, temporal trends, geographical patterns and research needs. Preslia 80(2): 101-149.
- LATOWSKI K. 1981. Dalsze materiały florystyczne z terenów kolejowych Wielkopolski. Bad. Fizjogr. Pol. Zach., seria B-Botanika 32: 207-211.
- LATOWSKI K. 2009. Nowe i osobliwe antropofity we florze naczyniowej Wielkopolskiego Parku Narodowego. Cz. I. Szkarlatka jagodowa [*Phytolacca acinosa* Roxburgh]. Morena 14: 215-222.
- MEUSEL H., JÄGER E., RAUSCHERT S. & WEINERT E. 1978. Vergleichende Chorologie der zentraleuropäischen Flora. II. Text xi+418 pp., Karten pp. 259-421. Gustav Fischer Verlag, Jena.
- MIREK Z., PIĘKOŚ-MIRKOWA H., ZAJĄC A. & ZAJĄC M. 2002. Flowering plants and pteridophytes of Poland. A checklist. In: Z. MIREK (ed.). Biodiversity of Poland, 1, 442 pp. W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków.
- MUHLENBACH V. 1979. Contributions to the synanthropic (adventive) flora of the railroads in St. Louis, Missouri, U.S.A. Ann. Missouri Bot. Garden 66: 1-108.
- MUNZ P. A. 1968. A California Flora and Supplement. Univ. California Press, Berkeley.
- OKLEJEWICZ K., WOLANIN M. & WOLANIN M. N. 2015. Czerwona księga roślin województwa podkarpackiego. Zagrożone gatunki roślin. Zbiorowiska roślinne. 299 pp. Stowarzyszenie na Rzecz Rozwoju i Promocji Podkarpacia "Pro Carpathia", Rzeszów.
- PAMPUCH A. 1840. Flora Tremesensis. 70 pp. Trzemeszno.
- PFUHL F. 1896. Die bisher in der Provinz Posen nachgewiesenen Gefäßpflanzen. Zeitschr. Botan. Abteil. Naturwiss. Verein. 3(1): 1-70.
- PROTOPOPOVA V. V. 1991. The Synanthropic Flora of Ukraine and its Development. 204 pp. Naukova Dumka Press, Kiev.
- PROTOPOPOVA V. V. & SHEVERA M. V. 2014. Ergasiophytes of the Ukrainian flora. Biodiv. Res. Conserv. 35: 31-46.
- PROVOST M. 1998. Flore Vasculaire de Basse-Normandie, vol. 2. Université de Caen, Caen.
- PURCEL A. 2007. Obce gatunki drzew i krzewów na terenie Wielkopolskiego Parku Narodowego – ich udział i rola w biocenozach parku. Ph. D. Thesis, Katedra Przyrodniczych Podstaw Leśnictwa Akademia Rolnicza im. Augusta Cieszkowskiego w Poznaniu, Poznań.
- PURCEL A. 2009. Obce gatunki drzew i krzewów w Wielkopolskim Parku Narodowym – ich występowanie i rola w biocenozach parku. Morena 14: 35-191.
- PYŠEK P., DANIELKA J., SÁDLO J., CHRTEK J. JR., CHYTRÝ M., JAROŠÍK V., KAPLAN Z. & KRAHULEC F. 2012. Catalogue of alien plants of the Czech Republic (2nd edition): checklist update, taxonomic diversity and invasion patterns. Preslia 84(2): 155-255.
- PYŠEK P., RICHARDSON D. M., REJMÁNEK M., WEBSTER G. L., WILLIAMSON M. & KIRSCHNER J. 2004. Alien plants in checklists and floras: towards better communication between taxonomists and ecologists. Taxon 53(1): 131-143.
- RITSCHL G. 1850. Flora des Großherzogthums Posen, im Auftrage des naturhistorischen Vereins zu Posen. 291 pp. Druck und Verlag von E. S. Mittler und Sohn, Berlin.
- ROTHMALER W., JÄGER E. J. & WERNER K. 2005. Exkursionsflora von Deutschland. Band 4, Gefäßpflanzen: Kritischer Band. 980 pp. Spektrum Akademischer Verlag, Elsevier GmbH, München.
- ROSTAŃSKI K. & SOWA R. 1986/1987. Alfabetyczny wykaz efemerofitów Polski. Fragm. Flor. Geobot. 31-32 (1-2): 151-205.
- SCHROEDER F. G. 1969. Zur Klassifizierung der Anthropochoren. Vegetatio 16: 225-238.
- SOTEK Z. 2014. CR *Plantago coronopus* L. In: R. KAŻMIERCZAKOWA, K. ZARZYCKI & Z. MIREK (eds.). Polska Czerwona Księga Roślin. Paprotniki i rośliny kwiatowe, wyd. 3, pp. 485-487. PAN, Instytut Ochrony Przyrody, Kraków.
- SZULCZEWSKI J. W. 1931. Przybysze i przybłędy w roślinności Poznania. PTPN, Prace Kom. Mat.-Przyr., seria B, 5: 59-74.
- SZULCZEWSKI J. W. 1951. Wykaz roślin naczyniowych w Wielkopolsce dotąd stwierdzonych. PTPN, Prace Kom. Biol. 12(6): 1-128.
- THE PLANT LIST 2013. <http://www.theplantlist.org/>
- THELLUNG A. 1918/1919. Zur Terminologie der Adventiv- und Ruderalfloristik. Allg. Bot. Zeitschr. 24/25: 36-42.
- TOKARSKA-GUZIK B. 2005. The Establishment and Spread of Alien Plant Species (Kenophytes) in the Flora of Poland. Prace naukowe Uniw. Śląskiego w Katowicach 2372: 1-192.
- URBISZ A. 2012. Occurrence of temporarily-introduced alien plant species (ephemeralophytes) in Poland – scale and assessment of the phenomenon. Prace naukowe Uniw. Śląskiego w Katowicach 2897: 1-200.
- ZAJĄC A. 1979. Pochodzenie archeofitów występujących w Polsce. Rozpr. habil. Uniw. Jagiell., 29: 1-213. Druk UJ w Krakowie.
- ZAJĄC A. & ZAJĄC M. (eds.). 2001. Distribution Atlas of Vascular Plants in Poland. xii+714 pp. Edited by Laboratory of Computer Chorology, Institute of Botany, Jagiellonian University, Cracow.
- ZAJĄC M., ZAJĄC A. & TOKARSKA-GUZIK B. 2009. Extinct and endangered archaeophytes and the dynamics of their diversity in Poland. Biodiv. Res. Conserv. 13: 17-24.

- ZIELIŃSKI J. 2016. Różowokwiatowe mieszańce robinii (*Robinia* L., Fabaceae) w uprawie. Propozycja utworzenia grup kultywarów: Ambigua i Margaretae. Rocznik Polskiego Towarzystwa Dendrologicznego 64: 21-33.
- ZIELIŃSKI J., BIEL G., DANIELEWICZ W., TOMASZEWSKI D. & GAWLAK M. 2015. Różowokwiatowe robinie (*Robinia* L., Fabaceae) dziczejące w Polsce. Rocznik Polskiego Towarzystwa Dendrologicznego 63: 9-33.
- ŻUKOWSKI W. & JACKOWIAK B. 1995. List of endangered and threatened vascular plants in Western Pomerania and Wielkopolska (Great Poland). In: W. ŻUKOWSKI & B. JACKOWIAK (eds.). *Endangered and threatened vascular plants of Western Pomerania and Wielkopolska*. Publications of the Department of Plant Taxonomy of the Adam Mickiewicz University in Poznań 3: 9-96. Bogucki Wyd. Nauk., Poznań.
- ŻUKOWSKI W. 1959. Nowe i rzadkie gatunki synantropijne dla flory miasta Poznania. Przyr. Polski Zach. 3: 151-153.
- ŻUKOWSKI W., LATOWSKI K., JACKOWIAK B. & CHMIEL J. 1995. Rośliny naczyniowe Wielkopolskiego Parku Narodowego. Prace Zakładu Taksonomii Roślin UAM w Poznaniu 4: 1-229. Bogucki Wyd. Nauk. Poznań.
- ŻUKOWSKI W. & PIASZYK M. 1971. Rozmieszczenie gatunków z rodzaju *Artemisia* L. w Polsce. Bad. Fizjogr. Pol. Zach. 24: 107-129.

Appendix 1. Checklist of casual alien species of Wielkopolska – ephemeral species

Name of species	Name of family	BF	OP	Native range
Abutilon theophrasti Med.	Malvaceae	A	4b	Asia (N China)
Achillea nobilis L.	Asteraceae	P	2a	Europe S, C; Asia W
Acroptilon repens (L.) DC.	Asteraceae	P	3b	Asia W, C
Aegilops cylindrica Host	Poaceae	A	3b, 4b	Europe S, Russia
Aegilops speltoides Tausch subsp. <i>ligustica</i> (Savign.) Zhuk. [= <i>Aegilops ligustica</i> Asch. & Graebn.]	Poaceae	A	2a	Medit
Amaranthus deflexus L.	Amaranthaceae	AP	4a?	S America
Amaranthus dinteri Schinz	Amaranthaceae	A	3b	Africa S
Amaranthus palmeri S. Watson	Amaranthaceae	A	3b	N America S, W
Amaranthus powellii S. Watson [= <i>Amaranthus bouchonii</i> Thell.]	Amaranthaceae	A	3a	N America S; S America N
Ambrosia trifida L.	Asteraceae	A	2a	N America
Ammi majus L.	Apiaceae	A	2a	Medit, Asia S, W
Amsinckia calycina (Moris) Chater	Boraginaceae	A	4a	S America S
Anacyclus radiatus Loisel.	Asteraceae	A	1a	Medit W
Anthemis austriaca Jacq.	Asteraceae	A	4a	Asia W; Europe S
Artemisia verlotiorum Lamotte	Asteraceae	P	?	Asia E (China)
Asperula arvensis L.	Rubiaceae	A	2a	Asia S, W; Africa N
Azolla filiculoides Lam. ¹	Azollaceae	AP	4a	N and S America (warm and moderate regions)
Bidens pilosa L.	Asteraceae	A	2a	S America ?
Bifora radians M. Bieb. ²	Apiaceae	A	4a	Asia W; Europe S, E
Brachypodium distachyon (L.) P. Beauv.	Poaceae	A	2a	Medit
Briza minor L.	Poaceae	A	2a	Medit
Bromus commutatus Schrad.	Poaceae	A	2b	Europe W, S, E; Africa N; Asia W
Bromus grossus Desf. ex DC.	Poaceae	A	1a, 2a	Europe (Austria, Belgium, Italy, Luxembourg, Czech Republic, Liechtenstein)
Bromus lanceolatus Roth	Poaceae	A	2a	Medit
Bromus rigidus Roth	Poaceae	A	2a	Medit
Bromus squarrosus L.	Poaceae	A	?	Europe S; Asia S, W
Bromus briziformis Fisch. et. C. A. Mey.	Poaceae	A	1a	Asia S, W
Bromus catharticus Vahl [= <i>Bromus willdenowii</i> Kunth]	Poaceae	AP	4a	S America
Bunium bulbocastanum L.	Apiaceae	P	2a	Europe S, W
Bupleurum fontanesii Caruel	Apiaceae	A	2a	Medit
Calamintha menthifolia Host [= <i>Calamintha sylvatica</i> Bromf.]	Lamiaceae	P	2a	Medit
Camelina rumelica Velen. ³	Brassicaceae	A	3a	Europe E, S; Asia W
Carthamus tinctorius L.	Asteraceae	A	2a	Asia (Middle East, India)
Centaurea calcitrapa L. ⁴	Asteraceae	B	1a, 2a	Medit
Centaurea diluta Aiton	Asteraceae	AP	2a	Medit W
Centaurea solstitialis L.	Asteraceae	A	4a	Medit
Centaurea stenolepis A. Kern.	Asteraceae	P	4a	Europe S, E (Anatolia N-W)

¹ Oxbow lakes in the Odra valley in the vicinity of Bytom Odrzański (leg S. Rosadziński 2008, det. K. Latowski, POZ).

² At the railway station in Jarocin (Czarna 2005).

³ On the railway embankment between Drawski Mlyn and Krzyż Wielkopolski (Latowski 1981).

⁴ The following species were also reported from Wielkopolska: *Centaurea decipiens* Thuill. and *C. pratensis* Thuill. (Szulczeński (1951). These species were not included in the checklist due to the lack of comparative materials and subsequent confirmations, as well as some taxonomic doubts.

Name of species	Name of family	BF	OP	Native range
<i>Chenopodium pratericola</i> Rydb.	Chenopodiaceae	A	2a	N America
<i>Chenopodium quinoa</i> L.	Chenopodiaceae	A	2a	S America (Ecuador, Peru, Bolivia)
<i>Chenopodium ambrosioides</i> L.	Chenopodiaceae	A	2a	S America; N America S
<i>Chenopodium berlandieri</i> Moq.	Chenopodiaceae	A	2a	N America
<i>Chenopodium hircinum</i> Schrader	Chenopodiaceae	A	2a	S America
<i>Chenopodium schraderanum</i> Schult.	Chenopodiaceae	A	?	Africa N
<i>Chloris barbata</i> Sw.	Poaceae	A	2a	Asia S, E, tropic
<i>Chloris truncata</i> R. Br.	Poaceae	B	2a	Australia
<i>Chloris virgata</i> Sw.	Poaceae		A	Eurasia; Africa; and the Americas (the warmer temperate, subtropical, and tropical regions)
<i>Chorispora tenella</i> (Pall.) DC.	Brassicaceae	A	4a	Europe S, E; Asia
<i>Citrullus lanatus</i> (Thunb.) Matsum. et Nakai	Cucurbitaceae	A	3a	Africa S (Namibia)
<i>Cladanthus mixtus</i> (L.) Oberpr. & Vogt [= <i>Chamaemelum mixtum</i> (L.) All.]	Asteraceae	A	2a	Medit
<i>Claytonia perfoliata</i> Donn ex Willd.	Portulacaceae	A	4a	N America
<i>Cochlearia officinalis</i> L.	Brassicaceae	BP	2a	Europe N, W, C
<i>Coincya monensis</i> (L.) Greuter et Burdet s. l.	Brassicaceae	AP	2a	Europe S, W
<i>Coleostephus myconis</i> (L.) Rchb. f.	Asteraceae	A	2a	Medit
<i>Commelinia coelestis</i> Willd.	Commelinaceae	P	2a	N America S
<i>Conopodium majus</i> (Gouan) Lorent	Apiaceae	P	2a	Europe W, S
<i>Consolida orientalis</i> (J. Gay) Schrödinger [= <i>Consolida hispanica</i> (Costa) Greuter et Burdet]	Ranunculaceae	A	4a	Medit
<i>Corispernum nitidum</i> Kit. ex Schult.	Chenopodiaceae	A	3a	Europe S, E, W; Asia W
<i>Coronopus didymus</i> (L.) Sm.	Brassicaceae	A	2a	S America
<i>Cotula anthemoides</i> L.	Asteraceae	A	2a	Africa S; Asia S, W
<i>Crambe abyssinica</i> Hochst. ex R. E. Fr.	Brassicaceae	A	?	Africa S (Abisinia)
<i>Crepis setosa</i> Haller f.	Asteraceae	A	1a	Europe C, S
<i>Crepis sprengeriana</i> (L.) All. [= <i>Picris sprengeriana</i> (L.) Poir.]	Asteraceae	A	2a	Medit
<i>Crupina vulgaris</i> Pers. ex Cass.	Asteraceae	A	1a	Europe S
<i>Cucurbita ficifolia</i> Bouché	Cucurbitaceae	A	2a	S America (Andes)
<i>Cuscuta australis</i> R. Br.	Cuscutaceae	A	3a	Australia; Asia; Europe S
<i>Cuscuta campestris</i> Yunck. ⁵	Cuscutaceae	A	3a	S America
<i>Cuscuta cesatiana</i> Bertol. ⁶ [= <i>Cuscuta arvensis</i> Beyr. var. <i>calycina</i> Engelm.]	Cuscutaceae	A	3a	Europe S, C to Asia S, W (India)
<i>Cuscuta gronovii</i> Willd. ex Schult.	Cuscutaceae	A	1a, 2a	N America
<i>Cuscuta trifolii</i> Bab. et Gibson [= <i>Cuscuta epithymum</i> (L.) Nathh. subsp. <i>trifolii</i> (Bab. & Gibson) Beger]	Cuscutaceae	A	?	Europe W, S
<i>Cynosurus echinatus</i> L.	Poaceae	A	2a	Medit
<i>Cyperus congestus</i> Vahl	Cyperaceae	P	2a	Africa S
<i>Cytisus elongatus</i> Waldst. & Kit. [= <i>Chamaecytisus glaber</i> (L.f.) Rothm.]	Fabaceae	T	2a	Europe S (Serbia, Montenegro, Bulgaria, Romania)

⁵ Railway station in Zbąszynek (leg. et det. K. Latowski 1983, POZ).⁶ Sieraków near Rawicz, on *Origanum majorana* L. and *Capsicum annuum* L. (Golenia & Bayerowa 1956).

Name of species	Name of family	BF	OP	Native range
Dactyloctenium aegyptium (L.) Willd.	Poaceae	A	2a	Africa tropic, Asia S
Dasypyrum villosum (L.) Borbás	Poaceae	A	2a	Europe E, S; Asia W
Dimorphotheca pluvialis (L.) Moench	Asteraceae	AP	2a	Africa S
Draba muralis L.	Brassicaceae	A	2a	Europe; Africa N, W; Asia W
Dracocephalum moldavica L.	Lamiaceae	A	2a	Asia (China, Sibirien, Central Asia), Europe E
Dracocephalum thymiflorum L.	Lamiaceae	A	3a	Asia W, C; Europe E
Echium plantagineum L.	Boraginaceae	AB	2a	Europe S, W; Africa N; Asia S, E
Elymus canadensis L.	Poaceae	P	4a	N America
Eragrostis mexicana (Hornem.) Link	Poaceae	A	2a	S America; N America
Eragrostis multicaulis Steud.	Poaceae	A	2a	Asia E; Europe S; Africa N
Eragrostis ciliaris (All.) Vignolo ex Janch.	Poaceae	A	4a	Eurasia; Africa
Erucastrum nasturtiifolium (Poir.) O. E. Schulz	Brassicaceae	BP	3a	Europe S, W
Erysimum crepidifolium Rchb.	Brassicaceae	BP	2a	Europe C, S
Erysimum diffusum Ehrh.	Brassicaceae	B	?	Europe S, E, C; Asia W
Euphorbia platyphyllos L.	Euphorbiaceae	A	2a	Europe (medit and subatl)
Fumaria capreolata L.	Fumariaceae	A	2a	Medit
Fumaria densiflora DC.	Fumariaceae	A	2a	Europe S (Medit); Asia C, W
Galeopsis segetum Neck.	Lamiaceae	A	2a, 4a	Europe S (Medit)
Galium parisiense L.	Rubiaceae	A	2a	Europe S, W, Africa N, Asia Minor
Galium tricornutum Dandy ⁷	Rubiaceae	A	1a, 2a, 3a	Europe N, W; Africa N; Asia W
Gaudinia fragilis (L.) P. Beauv.	Poaceae	A	2a	Medit
Geranium rotundifolium L.	Geraniaceae	A	3a?	Europe S, W; Asia S; Africa N
Glaucium flavum Crantz	Papaveraceae	A	2a	Medit
Glaucium corniculatum (L.) Rudolph	Papaveraceae	A	2a, 3a	Medit
Guizotia abyssinica (L. f.) Cass.	Asteraceae	A	2a	Africa E (Ethiopia)
Gypsophila perfoliata L.	Caryophyllaceae	P	4a	Europe E; Asia W
Heliotropium europaeum L.	Boraginaceae	A	2a, 4a	Medit
Herniaria incana Lam.	Caryophyllaceae	P	4b	Europe (Medit) to Asia W
Hirschfeldia incana (L.) Lagr.-Foss. [= Hirschfeldia adpressa Moench]	Brassicaceae	A	4a	Medit
Hordeum secalinum Schreb.	Poaceae	P	2a	Europe W, S to Asia Minor; Africa N
Humulus scandens (Lour.) Merr.	Cannabaceae	A	2a	Asia E
Hygrophila polysperma (Roxb.) T. Anderson ⁸	Acanthaceae	P	4a	Asia S, E (India, Malaysia)
Hypericum mutilum L.	Hypericaceae	A	1a	N America
Hypericum gymnanthum Engelm. & A. Gray	Hypericaceae	A	1a	N America
Juncus anthelatus (Wiegand) R. E. Brooks & Whittem.	Juncaceae	P	4a	N America E; C America (Mexico)
Lactuca tatarica (L.) C. A. Mey	Asteraceae	P	?	Europe E; Asia E to N. & W. America N
Lactuca virosa L.	Asteraceae	AB	1a, 3b	Medit
Lathyrus aphaca L.	Fabaceae	A	4a	Europe W, C, S; Asia W; Africa N
Legousia speculum-veneris (L.) Chaix	Campanulaceae	A	2a	Medit
Lepidium latifolium L.	Brassicaceae	P	4b	Medit, Asia S, W
Lepidium perfoliatum L.	Brassicaceae	A	4a	Europe E; Asia
Linaria repens (L.) Mill.	Scrophulariaceae	P	4a	Europe (Medit-Atl)

⁷ In the highland belt and in the Lower Silesia, this species occurs as an archaeophyte Zająć 1979; Zająć et al. 2009.). In Wielkopolska, it appeared only ephemeral in Poznań (Żukowski 1959).

Name of species	Name of family	BF	OP	Native range
Linum austriacum L.	Linaceae	P	?	Europe E, S, C; Africa N; Asia W (Turkey)
Lolium rigidum Gaudin	Poaceae	A	2a	Medit, Asia S, W
Lythrum junceum Banks & Sol.	Lythraceae	P	2a	Medit
Malva nicaeensis All. ⁹	Malvaceae	A	2a	Medit
Marrubium peregrinum L.	Lamiaceae	P	2a	Europe S, E; Asia Minor
Medicago arabica (L.) Huds.	Fabaceae	A	1a	Medit
Medicago polymorpha L.	Fabaceae	A	?	Medit
Melica picta K. Koch ¹⁰	Poaceae	P	3a	Europe C, E
Melilotus indicus (L.) All.	Fabaceae	A	2a	Medit, Asia S, W
Mimulus moschatus Douglas ex Lindl.	Scrophulariaceae	P	?	N America W
Minuartia hybrida (Vill.) Schischk.	Caryophyllaceae	A	1a	Medit, Asia S, W
Moenchia mantica (L.) Bartl.	Caryophyllaceae	A	2a	Europe S, E
Myagrum perfoliatum L.	Brassicaceae	A	4a	Europe S; Asia W
Nicotiana rustica L.	Solanaceae	A	2a	America C, S
Nicotiana tabacum L.	Solanaceae	A	2a	America C, S ?
Oenothera ×tacikii Rostański	Onagraceae	B	4a	Europe
Oenothera cruciata Nutt. ex Don	Onagraceae	B	4a	N America
Oenothera fallax Renner em. Rostański	Onagraceae	B	4a	America N
Oenothera flæmingina Hudziok	Onagraceae	B	4a	Europe
Oenothera parviflora L.	Onagraceae	B	4a	N America
Oenothera pycnocarpa G. F. Atk. et Bartlett	Onagraceae	B	4a	N America
Orobanche ramosa L.	Orobanchaceae	A	2a, 3a	Africa, Asia W, Europe
Oxalis tetraphylla Cav. [= Oxalis deppei Lodd. ex Sweet]	Oxalidaceae	P	4b	America C (Mexico)
Panicum capillare L.	Poaceae	A	4b	N America
Panicum dichotomiflorum Michx.	Poaceae	A	?	S America; N America
Papaver albiflorum (Boiss.) Paczoski ¹¹ [= Papaver dubium L. subsp. albiflorum]	Papaveraceae	A	4a	Europe S, W (medit)
Paspalum racemosum Lam.	Poaceae	A	2a	S America W
Petrorhagia saxifraga (L.) Link	Caryophyllaceae	P	2a	Europe S, C; Asia W
Phalaris canariensis L.	Poaceae	A	4a	Medit
Phalaris coerulescens Desf.	Poaceae	P	2a	Medit
Plantago afra L. ¹²	Plantaginaceae	A	4a	Africa C, N; Asia S, W; Europe S, E
Plantago lagopus L.	Plantaginaceae	A	2a	Medit
Polycarpon tetraphyllum (L.) L.	Caryophyllaceae	A	2a	Europe (medit)
Polygonum patulum M. Bieb.	Polygonaceae	A	2a	Europe W, Medit, Asia C (Caucas, Siberia, Mongolia, China)
Polypogon monspeliensis (L.) Desf.	Poaceae	A	2a	Medit, Asia S, W
Potentilla intermedia L.	Rosaceae	BP	2a	Europe N, W; Africa N; Asia (Siberia)
Rapistrum perenne (L.) All.	Brassicaceae	P	1a, 3a	Europe E, S; Asia W
Rapistrum rugosum (L.) All.	Brassicaceae	A	1a, 2a, 3a	Medit
Reseda alba L.	Resedaceae	AP	1a, 2a	Medit
Ricinus communis L.	Euphorbiaceae	A	3a?	Africa tropic, Middle East

⁸ Reported for the first time from the vicinity of Konin (Gąbka & Owiśniak 2009).⁹ *Malva mauritiana* L. – reported from Wielkopolska (Szulczeński 1951; Czarna 2009, Celka 2014 POZ), in this work it was included in *M. sylvestris* L.¹⁰ Native to Poland, occurs only in a single locality in the south-eastern part of the country (Zajac & Zajac 2001, Oklejewicz *et al.* 2015).¹¹ Railway tracks at the railway station in Kościan (Czarna 2009).¹² Observed in Kórnik near Poznań in the years 2014–2015; in total, a few individuals occurred (leg. A. Wysocki, det. K. Latowski; POZ); in 2016 – not found.

Name of species	Name of family	BF	OP	Native range
<i>Roemeria hybrida</i> (L.) DC.	Papaveraceae	A	4a	Medit, Asia W
<i>Rosa gorenkensis</i> Besser	Rosaceae	T	4a	Europe S, W; Asia W,
<i>Rumex bucephalophorus</i> L.	Polygonaceae	A	2a	Medit
<i>Rumex longifolius</i> DC.	Polygonaceae	P	?	Asia W, C
<i>Rumex salicifolius</i> Weinm. [= <i>Rumex triangulivalvis</i> (Danser) Rech. f.]	Polygonaceae	P	3a	N America
<i>Rumex stenophyllus</i> Ledeb.¹³	Polygonaceae	P	4a	Europe S, W, Azja C (Siberia)
<i>Salsola collina</i> Pall.	Chenopodiaceae	A	3a	Asia (Korea, Mongolia, China, Pakistan, Russia, Central Asia); Europe E
<i>Salsola soda</i> L.	Chenopodiaceae	A	3a	Medit
<i>Salvia dumetorum</i> Andrz. ex Besser [= <i>Salvia dumetorum</i> Andrz.]	Lamiaceae	P	2a	Europe S, E; Asia C
<i>Scirpus georgianus</i> R. M. Harper	Cyperaceae	P	4a	N America
<i>Senecio inaequidens</i> DC.	Asteraceae	P	4a	Africa (South Africa, Lesotho, Swaziland)
<i>Sesamoides interrupta</i> (Bureau) G. López [= <i>Sesamoides canescens</i> (L.) Kuntze]	Resedaceae	P	3a	Medit W
<i>Sideritis montana</i> L.	Lamiaceae	A	4a	Medit, Asia W
<i>Silene gallin yi</i> Heuff. ex Rchb. [= <i>Silene trinervia</i> Sebast. & Mauri]	Caryophyllaceae	P	2a	Europe (Balkans)
<i>Silene linicola</i> C. C. Gmel.	Caryophyllaceae	A	2a	Europe W (France)
<i>Silene scabriflora</i> Brot.	Caryophyllaceae	A	2a	Europe S, W
<i>Sisymbrium austriacum</i> Jacq.	Brassicaceae	BP	2a	Europe C, S, W
<i>Sisymbrium orientale</i> L.	Brassicaceae	ABP	2a, 3a	Europe S; Asia W
<i>Solanum alatum</i> Moench	Solanaceae	A	3a	Medit, Asia Minor
<i>Solanum angustifolium</i> Mill. [= <i>Solanum cornutum</i> Lam.]	Solanaceae	A	3a	N America S
<i>Solanum triflorum</i> Nutt.	Solanaceae	A	2a	S America; N America
<i>Sorghum bicolor</i> (L.) Moench	Poaceae	A	3a	Africa tropic
<i>Sorghum halepense</i> (L.) Pers.	Poaceae	P	3a	Europe (Medit) to Asia C
<i>Thlaspi alliaceum</i> L.	Brassicaceae	A	?	Africa N, E; Asia W; Europe S, E
<i>Thlaspi caerulescens</i> J. Presl et. C. Presl [= <i>Thlaspi alpestre</i> L.]	Brassicaceae	BP	2a	Europe W, C
<i>Thlaspi perfoliatum</i> L.	Brassicaceae	A	4a	Europe S; Asia W
<i>Urtica cannabina</i> L.	Urticaceae	P	3a	Europe E (Russia), Asia W, C (to China)
<i>Valerianella carinata</i> Loisel.	Valerianaceae	A	1a	Medit ?
<i>Veronica peregrina</i> L.	Scrophulariaceae	AB	4a	N America S
<i>Vicia articulata</i> Hornem.	Fabaceae	A	2a	Medit
<i>Vicia bithynica</i> (L.) L.	Fabaceae	A	2a	Medit
<i>Vicia lutea</i> L.	Fabaceae	A	3a	Europe; Asia W; Africa N
<i>Vicia pannonica</i> Crantz	Fabaceae	A	4a	Europe S, C; Asia W
<i>Viola ambigua</i> Waldst. & Kit.	Violaceae	P	3a?	Europe C, S, E (Caucasus)
<i>Vulpia geniculata</i> (L.) Link	Poaceae	A	2a	Europe S; Africa N
<i>Xanthium orientale</i> L. [= <i>Xanthium macrocarpum</i> DC.]	Asteraceae	A	2a	N America ?
<i>Xanthium spinosum</i> L.	Asteraceae	A	2a	S America

Explanations: BF – biological form, A – annual plants, B – biennial plants, P – perennial, herbaceous plants, S – dwarf shrubs and sub-shrubs, T – shrubs and trees; OP – observation period, 1 – to 1901, 2 – 1901-1951, 3 – 1951-2001, 4 – after 2001, a – species was observed once, b – species was observed repeatedly; Native range: N – north, W – west, E – east, S – south, C – central, Medit – Mediterranean, Atl – Atlantic; ? – no data information uncertain

¹³ Railway station in Jarocin (Czarna 2005, 2009).

Appendix 2. Checklist of casual alien species of Wielkopolska – ergasiophygophytes

Name of species	Name of family	BF	OP
Abies concolor (Gord. et Glend.) Lindl. ex Hildebr.	Pinaceae	T	4b
Acer saccharinum L.	Aceraceae	T	4b
Acer tataricum L. ¹	Aceraceae	T	4b
Achillea tomentosa L.	Asteraceae	P	?
Adonis annua L. em. Huds.	Ranunculaceae	A	2a
Aesculus ×carnea Hayne	Hippocastanaceae	T	4b
Aesculus flava Sol.	Hippocastanaceae	T	3a
Alcea rosea L.	Malvaceae	BP	4b
Allium carinatum L.	Liliaceae	P	2a
Allium cepa L.	Liliaceae	P	4a
Allium fistulosum L.	Liliaceae	P	2a
Allium giganteum Regel	Liliaceae	P	4a
Allium moly L.	Liliaceae	P	4a
Allium sativum L.	Liliaceae	P	2a
Allium schoenoprasum L.	Liliaceae	P	4a
Allium sphaerocephalon L.	Liliaceae	P	2a
Allium ampeloprasum L. s. l. [= Allium porrum L.]	Liliaceae	P	2a
Althaea officinalis L. ²	Malvaceae	P	4a
Alyssum saxatile L. ³	Brassicaceae	S	4a
Amaranthus caudatus L.	Amaranthaceae	A	4a
Amaranthus cruentus L. [= Amaranthus paniculatus L.]	Amaranthaceae	A	4a
Amelanchier lamarckii F. G. Schroed.	Rosaceae	T	4a
Ammophila arenaria (L.) Link ⁴	Poaceae	P	4a
Amorpha fruticosa L.	Fabaceae	T	4a
Anethum graveolens L.	Apiaceae	A	4a
Antirrhinum majus L.	Scrophulariaceae	AP	4a
Apium graveolens L.	Apiaceae	A	4a
Aquilegia caerulea E. James [= Aquilegia ×hybrida hort.]	Ranunculaceae	P	4a
Arabis alpina L. ⁵	Brassicaceae	P	4a
Arabis caucasica Willd.	Brassicaceae	P	4a
Aristolochia macrophylla Lam.	Aristolochiaceae	T	4a
Artemisia abrotanum L.	Asteraceae	S	2b
Artemisia dracunculus L.	Asteraceae	P	4b
Artemisia ludoviciana Nutt. ⁶ [= Artemisia platyphylla Rydb.]	Asteraceae	P	3a
Artemisia maritima L.	Asteraceae	S	1a
Artemisia pontica L. ⁷	Asteraceae	P	1a
Aster ×salignus Willd.	Asteraceae	P	4a
Avena sativa L.	Poaceae	A	4b

¹ Some information about *Acer tataricum* L. was attributed to *A. ginnala* Maxim. [*Acer tataricum* L. subsp. *ginnala* (Maxim.) Wesm.].

² Probably a synanthropic species. There are some doubts on its status on saline soils (see Zająć & Zająć 2001; Mirek *et al.* 2002).

³ In Poland, it is a native species in the Pieniny Mountains (Zająć & Zająć 2001). In lowlands, this species is cultivated and running wild from cultivation.

⁴ In Poland, it is a native species only on the Baltic Sea (Zająć & Zająć 2001). In addition, this species is cultivated in lowlands and running wild from cultivation.

⁵ In Poland, it is a native species in the Tatras and on the Babia Góra Mt. (Zająć & Zająć 2001). In addition, this species is cultivated in lowlands and running wild from cultivation.

⁶ The species was reported for the first time for Poland from the area of the Wielkopolski National Park. The specimens collected there belong to two varieties: var. *ludoviciana* and var. *incompta* (Nutt.) Cronq. (Żukowski *et al.* 1995).

⁷ In Poland, it is a native species only in the south-eastern part of Nidziańska Basin (Kaźmierczakowa 2014a). In other parts of Poland, only anthropogenic populations occur (Żukowski & Piaszyk 1971; Zająć & Zająć 2001; Kaźmierczakowa 2014a). In Wielkopolska, it was collected by Spribile (1892, POZ) on the Lake Tuczno near Inowrocław (Żukowski & Piaszyk 1971).

Name of species	Name of family	BF	OP
Berberis thunbergii DC. ⁸	Berberidaceae	T	4b
Bergenia crassifolia (L.) Fritsch	Saxifragaceae	P	4a
Beta vulgaris L.	Chenopodiaceae	AP	4a
Betula lenta L. ⁹	Betulaceae	T	4a
Bidens ferulifolia (Jacq.) DC.	Asteraceae	AB	4a
Borago officinalis L.	Boraginaceae	A	4a
Brassica napus L. subsp. napus	Brassicaceae	AB	4b
Buddleja davidii Franch.	Buddlejaceae	T	4a
Calandrinia pilosiuscula DC.	Portulacaceae	AP	1a
Calendula arvensis L.	Asteraceae	A	?
Calendula officinalis L.	Asteraceae	AP	4a
Callistephus chinensis (L.) Nees	Asteraceae	A	?
Campanula carpatica Jacq.	Campanulaceae	P	4a
Campanula rapunculus L.	Campanulaceae	B	2b
Cannabis sativa L.	Cannabaceae	A	4a
Capsicum annuum L.	Solanaceae	A	2a
Caragana arborescens Lam.	Fabaceae	T	4b
Carthamus lanatus L.	Asteraceae	A	2a
Carya ovata (Mill.) K. Koch	Juglandaceae	T	4a
Celastrus orbiculatus Thunb. ¹⁰	Celastraceae	P	4a
Centaurea dealbata Willd.	Asteraceae	P	4a
Centaurea montana L.	Asteraceae	P	2a
Cerastium tomentosum L.	Caryophyllaceae	P	4b
Chaenomeles japonica (Thunb.) Lindl. ex Spach	Rosaceae	T	4a
Chamaemelum nobile (L.) All. [= <i>Anthemis nobilis</i> L.]	Asteraceae	P	2a
Chenopodium capitatum (L.) Asch.	Chenopodiaceae	A	2a
Cicerbita macrophylla (Willd.) Wallr. ¹¹	Asteraceae	P	4a
Clarkia amoena (Lehm.) A. Nelson & J. F. Macbr. subsp. lindleyi (Douglas) H. F. Lewis & M. R. Lewis [= <i>Godetia grandiflora</i> Lindl.]	Onagraceae	A	4a?
Collomia grandiflora Douglas	Polemoniaceae	A	2a
Colutea arborescens L.	Fabaceae	T	4a
Colutea media Willd. ¹²	Fabaceae	T	4a
Consolida ajacis (L.) Schur	Ranunculaceae	A	4a
Convolvulus tricolor L.	Convolvulaceae	A	2a
Coriandrum sativum L.	Apiaceae	A	4a
Cornus mas L.	Cornaceae	T	4a
Cosmos bipinnatus Cav.	Asteraceae	A	4b
Cotoneaster divaricatus Rehder et E. H. Wilson	Rosaceae	T	4a
Cotoneaster integrerrimus Medik.	Rosaceae	T	?
Cotoneaster villosulus (Rehder & E. H. Wilson) Flinck & B. Hydm	Rosaceae	T	4a
Crataegus pedicellata Sarg.	Rosaceae	T	4a
Crataegus coccinea L.	Rosaceae	T	4a
Cucumis sativus L.	Cucurbitaceae	A	4b

⁸ Reported from the Wielkopolski National Park (Purcel 2009).⁹ Planted and running wild in the acidic oak forest in the Gołębki Forest District (leg. and det. Chmiel 1987, POZ).¹⁰ Wielkopolski National Park (Danielewicz & Maliński 1995; Purcel 2009).¹¹ Found in the village Dobrzyca (Czarna 2003, 2007).¹² Wielkopolski National Park (Purcel 2009).

Name of species	Name of family	BF	OP
Cucurbita moschata Duchesne	Cucurbitaceae	A	4a
Cucurbita pepo L.	Cucurbitaceae	A	4b
Dianthus barbatus L.	Caryophyllaceae	P	4b
Dicentra spectabilis L.	Fumariaceae	P	4b
Dipsacus sativus (L.) Honck.	Dipsacaceae	BP	4a
Doronicum columnae Ten.	Asteraceae	P	4a
Doronicum pardalianches L.	Asteraceae	P	4a
Echinacea purpurea (L.) Moench	Asteraceae	P	4a
Echinops exaltatus Schrad.	Asteraceae	P	4a
Elaeagnus commutata Bernh. ex Rydb.	Eleagnaceae	T	4a
Eruca sativa Mill. [= <i>Eruca vesicaria</i> (L.) Cav. subsp. <i>sativa</i> (Mill.) Thell.]	Brassicaceae	A	2a
Erysimum cheiri (L.) Crantz [= <i>Cheiranthus</i> × <i>cheiri</i> L.]	Brassicaceae	S	2a
Eschscholzia californica Cham.	Papaveraceae	AP	4a
Euphorbia lathyris L.	Euphorbiaceae	AB	4a
Euphorbia marginata Pursh	Euphorbiaceae	A	4b
Fagopyrum esculentum Moench	Polygonaceae	A	4b
Fagopyrum tataricum (L.) Gaertn.	Polygonaceae	A	4a
Fallopia baldschuanica (Regel) Holub [= <i>Fallopia aubertii</i> (L. Henry) Holub]	Polygonaceae	P	4a
Foeniculum vulgare Mill.	Apiaceae	P	4a
Fragaria ×ananassa Duchesne ex Rozier	Rosaceae	P	4b
Fraxinus angustifolia Vahl.	Oleaceae	T	4a
Gaillardia pulchella Foug.	Asteraceae	A	4a
Galium anisophyllum Vill. ¹³	Rubiaceae	B	2a
Gypsophila repens L. ¹⁴	Caryophyllaceae	P	4a
Helianthus annuus L.	Asteraceae	A	4a
Helianthus decapetalus L.	Asteraceae	P	?
Helianthus ×laetiflorus Pers.	Asteraceae	P	?
Helleborus viridis L.	Ranunculaceae	P	4a
Hemerocallis fulva (L.) L.	Liliaceae	P	4b
Hesperis sylvestris Crantz	Brassicaceae	BP	4a
Hibiscus trionum L.	Malvaceae	A	2a
Hieracium aurantiacum L. ¹⁵	Asteraceae	P	2a, 4a
Hordeum jubatum L.	Poaceae	P	4a
Hordeum vulgare L. em. Alef. ¹⁶	Poaceae	A	4b
Hyssopus officinalis L.	Lamiaceae	S	4a
Iberis amara L.	Brassicaceae	A	4a
Iberis umbellata L.	Brassicaceae	A	4b
Impatiens balsamina L.	Balsaminaceae	A	4a
Impatiens stenosepala E. Pritz. [= <i>Impatiens punctata</i> Franch. ex Hook. f.]	Balsaminaceae	A	4a
Ipomoea purpurea Roth	Convolvulaceae	A	4a
Ipomoea nil (L.) Roth [= <i>Ipomoea hederacea</i> (L.) Jacq.]	Convolvulaceae	A	4a
Iris germanica L.	Iridaceae	P	4b
Isatis tinctoria L.	Brassicaceae	BP	4a

¹³ In Poland, it is a native species in the Carpathians (Zajac & Zajac 2001). In lowlands, it appears only ephemerally (Szulczeński 1951).¹⁴ In Poland, it is a native species in the Tatras and Pieniny Mts (Zajac & Zajac 2001). In lowlands, this species occurs only as an escapee from cultivation.¹⁵ In Poland, it is a native species in the mountains and in the foothills (Zajac & Zajac 2001). In lowlands, this species occurs only as an escapee from cultivation.¹⁶ *Hordeum distichon* L., a species reported from Wielkopolska has been included here in *Hordeum vulgare* L. em. Alef.

Name of species	Name of family	BF	OP
<i>Larix ×eurolepis</i> A. Henry ¹⁷	Pinaceae	T	4b
<i>Lathyrus hirsutus</i> L. ¹⁸	Fabaceae	A	4a?
<i>Lathyrus japonicus</i> Willd. subsp. <i>maritimus</i> (L.) P. W. Ball ¹⁹	Fabaceae	P	2a
<i>Lathyrus latifolius</i> L.	Fabaceae	P	4a
<i>Lathyrus nissolia</i> L.	Fabaceae	A	?
<i>Lavandula angustifolia</i> Mill.	Lamiaceae	S	4a
<i>Lavatera trimestris</i> L.	Malvaceae	A	4a
<i>Lens culinaris</i> Medik.	Fabaceae	A	2a
<i>Lepidium sativum</i> L.	Brassicaceae	A	4b
<i>Leucanthemum maximum</i> (Ramond) DC.	Asteraceae	P	4a
<i>Levisticum officinale</i> W. D. J. Koch	Apiaceae	P	4b
<i>Lilium bulbiferum</i> L. ²⁰	Liliaceae	P	4b
<i>Lilium candidum</i> L.	Liliaceae	P	4a
<i>Linum grandiflorum</i> Desf.	Linaceae	A	4a
<i>Linum perenne</i> L.	Linaceae	P	2a
<i>Linum usitatissimum</i> L.	Linaceae	A	4b
<i>Lobelia erinus</i> L.	Campanulaceae	A	?
<i>Lobularia maritima</i> (L.) Desv.	Brassicaceae	AP	4a
<i>Lonicera caprifolium</i> L.	Caprifoliaceae	T	4a
<i>Lonicera involucrata</i> (Richardson) Banks ex Spreng.	Caprifoliaceae	T	4a
<i>Lunaria annua</i> L.	Brassicaceae	AB	4b
<i>Lupinus angustifolius</i> L.	Fabaceae	A	4a
<i>Lupinus luteus</i> L.	Fabaceae	A	4b
<i>Lychnis coronaria</i> (L.) Desr.	Caryophyllaceae	P	4b
<i>Lysimachia punctata</i> L.	Primulaceae	P	4a
<i>Malva verticillata</i> L.	Malvaceae	A	4b
<i>Mathiola incana</i> (L.) W. T. Aiton	Brassicaceae	A	4a?
<i>Melica altissima</i> L.	Poaceae	P	?
<i>Melica ciliata</i> L. ²¹	Poaceae	P	?
<i>Melissa officinalis</i> L.	Lamiaceae	P	4a
<i>Mentha ×piperita</i> L. [= <i>Mentha ×citrata</i> Ehrh.]	Lamiaceae	P	4a
<i>Mentha ×rotundifolia</i> (L.) Huds.	Lamiaceae	P	4a
<i>Mentha spicata</i> L. em. L.	Lamiaceae	P	2a
<i>Misanthus sacchariflorus</i> (Maxim.) Hack.	Poaceae	B	4b
<i>Morus alba</i> L.	Moraceae	T	4b
<i>Muscari botryoides</i> (L.) Mill.	Liliaceae	P	4a
<i>Muscari comosum</i> (L.) Mill. ²²	Liliaceae	P	2a, 4a
<i>Muscari neglectum</i> Guss. ex Ten.	Liliaceae	P	4a
<i>Myrrhis odorata</i> (L.) Scop.	Apiaceae	P	?
<i>Narcissus poeticus</i> L.	Amaryllidaceae	P	4b
<i>Narcissus pseudonarcissus</i> L.	Amaryllidaceae	P	4b
<i>Nepeta grandiflora</i> M. Bieb.	Lamiaceae	P	2a
<i>Nepeta racemosa</i> Lam. [= <i>Nepeta mussinii</i> Spreng. ex Henckel]	Lamiaceae	P	2a

¹⁷ Hybrid of *Larix decidua* Mill. subsp. *decidua* and *L. kaempferi* (Lamb.) Carrière. It is cultivated and running wild from cultivation (Danielewicz & Maliński 1994, 1997; Purcel 2009).

¹⁸ Only some localities in southern Poland are native (Zajac & Zajac 2001).

¹⁹ Only some localities on the Baltic Sea are native in Poland (Zajac & Zajac 2001). The species was reported from Poznań by Szulczewski (1951).

²⁰ Only certain mountain localities are native in Poland (Zajac & Zajac 2001).

²¹ Only the localities in the south-eastern part of the country are native in Poland (Zajac & Zajac 2001). In Wielkopolska: Trzemeszno surroundings (Pampuch 1840; Ritschl 1850).

²² Native in southern Poland (Zajac & Zajac 2001). In Wielkopolska, this species runs wild from cultivation.

Name of species	Name of family	BF	OP
Nicandra physalodes (L.) Gaertn.	Solanaceae	A	4a
Nigella damascena L.	Ranunculaceae	A	4b
Nonea erecta Bernh. ²³ [= <i>Nonea pulla</i> (L.) DC.]	Boraginaceae	P	2a
Oenothera fruticosa L.	Onagraceae	P	4a
Oenothera glazioviana Micheli	Onagraceae	B	4a
Omphalodes verna Moench	Boraginaceae	P	4b
Ornithogalum nutans L.	Liliaceae	P	4b
Ornithopus sativus Brot.	Fabaceae	A	4b
Panicum miliaceum L.	Poaceae	A	4b
Papaver orientale L.	Papaveraceae	P	4a
Papaver somniferum L.	Papaveraceae	A	4b
Petroselinum crispum (Mill.) A. W. Hill	Apiaceae	B	4a
Petunia ×hybrida (Hook.) Vilm.	Solanaceae	A	4a
Petunia integrifolia (Hook.) Schinz et Thell. [= <i>Petunia violacea</i> Lindl.]	Solanaceae	A	2a, 4a
Peucedanum ostruthium (L.) W. D. J. Koch ²⁴	Apiaceae	P	3a?
Phacelia tanacetifolia Benth.	Hydrophyllaceae	A	4b
Phaseolus vulgaris L.	Fabaceae	A	4a
Philadelphus coronarius L.	Hydrangeaceae	T	4b
Physalis alkekengi L.	Solanaceae	P	4b
Physalis ixocarpa Brot. ex Hornem.	Solanaceae	A	?
Physocarpus opulifolius (L.) Maxim.	Rosaceae	T	4b
Phyteuma nigrum F. W. Schmidt	Campanulaceae	P	?
Phytolacca americana L. ²⁵	Phytolaccaceae	P	2a
Phytolacca esculenta van Houtte ²⁶ [= <i>Phytolacca acinosa</i> Roxb.]	Phytolaccaceae	P	4a
Pimpinella anisum L.	Apiaceae	A	?
Pinus banksiana Lamb.	Pinaceae	T	4a
Pinus nigra J. F. Arnold	Pinaceae	T	4a
Pinus rigida Mill.	Pinaceae	T	4b
Pinus strobus L.	Pinaceae	T	4b
Pisum sativum L.	Fabaceae	A	4b
Plantago coronopus L. ²⁷	Plantaginaceae	ABP	4b
Populus balsamifera L. ²⁸	Salicaceae	T	4a
Populus suaveolens Fisch. ex Loudon ²⁹ [= <i>Populus maximowiczii</i> Henry]	Salicaceae	T	4a
Prunus domestica L. ³⁰	Rosaceae	T	4b
Prunus persica (L.) Batsch [= <i>Persica vulgaris</i> Mill.]	Rosaceae	T	4b
Prunus armeniaca L. [= <i>Armeniaca vulgaris</i> Lam.]	Rosaceae	T	4b
Prunus cerasus L. [= <i>Cerasus vulgaris</i> Mill.]	Rosaceae	T	4b

²³ Native in the south-eastern part of Poland (Zajac & Zajac 2001).²⁴ In Poland, native localities are located in the Sudetes (Zajac & Zajac 2001). In Wielkopolska, this species is running wild from cultivation.²⁵ Reported from Gubin (Lademann 1937).²⁶ Reported from the Wielkopolski National Park (Latowski 2009).²⁷ The only native locality in Poland is on Kępa Karsiborska (Sotek 2014). The species was noted at the roadside near the Botanical Garden in Poznań. Perhaps, this population originated from the conservative collection of this species located in the Botanical Garden.²⁸ In Wielkopolska, the balsam poplar section (*Tacamahaca*) is represented by, among others: *Populus ×berolinensis* (K. Koch) Dippel, *P. candicans* Aiton and *P. simonii* Carrière.²⁹ *Populus suaveolens* Fisch. ex Loudon (*P. maximowiczii* Henry) has some varieties, e.g., 'NE42' (= 'Hybrida 275'), which reproduce vegetatively by suckers, e.g., in excavations (Chmiel 1993, 2006; Dolatowski & Seneta 2008).³⁰ The species is diversified into several subspecies with varying degrees of spread and establishment in the region, such as, among others: subsp. *domestica*; subsp. *insititia* (L.) Bonnier et Layens and subsp. *syriaca* (Borkh.) Janch. ex Mansf.

Name of species	Name of family	BF	OP
Pseudotsuga menziesii (Mirbel) Franco	Pinaceae	T	4a
Ptelea trifoliata L.	Rutaceae	T	4a
Pyrus communis L. ³¹	Rosaceae	T	4b
Raphanus sativus L.	Brassicaceae	A	4b
Reseda odorata L.	Resedaceae	A	4a
Reseda phytisma L. ³²	Resedaceae	A	2a
Rheum rhabarbarum L.	Polygonaceae	P	4b
Rhus hirta (L.) Sudw. [= <i>Rhus typhina</i> L.]	Anacardiaceae	T	4b
Ribes aureum Pursh	Grossulariaceae	T	4b
Ribes rubrum L.	Grossulariaceae	T	4b
Robinia neomexicana A. Gray ³³	Fabaceae	T	4a
Robinia viscosa Vent. ³⁴	Fabaceae	T	4a
Rosa blanda Aiton	Rosaceae	T	4a
Rosa glauca Pourr.	Rosaceae	T	4a
Rosa multiflora Thunb. ex Murray	Rosaceae	T	4a
Rosa rugosa Thunb.	Rosaceae	T	4b
Rosa spinosissima L.	Rosaceae	T	4a
Rosa virginiana Herrm.	Rosaceae	T	4a
Rubus occidentalis L.	Rosaceae	T	4a
Rudbeckia hirta L.	Asteraceae	ABP	4b
Salix acutifolia Willd.	Salicaceae	T	4a
Salix cordata Michx	Salicaceae	T	4a
Salix daphnoides Vill. ³⁵	Salicaceae	T	?
Salix eleagnos Scop. ³⁶	Salicaceae	T	?
Salvia glutinosa L. ³⁷	Lamiaceae	P	4a
Sambucus ebulus L. ³⁸	Caprifoliaceae	P	4a
Satureja hortensis L.	Lamiaceae	A	?
Scilla bifolia L. ³⁹	Liliaceae	P	2a, 4a
Scilla sibirica Haw.	Liliaceae	P	4b
Scorzonera hispanica L.	Asteraceae	P	2a
Scrophularia vernalis L. ⁴⁰	Scrophulariaceae	BP	4a
Secale cereale L. ⁴¹	Poaceae	A	4a
Sempervivum tectorum L.	Crassulaceae	P	4b
Setaria italica (L.) P. Beauv.	Poaceae	A	4a
Sicyos angulatus L.	Cucurbitaceae	A	4a
Sida hermaphrodita (L.) Rusby	Malvaceae	P	4a
Silene armeria L. ⁴²	Caryophyllaceae	AP	2a
Silene chalcedonica (L.) E. Krause [= <i>Lychnis chalcedonica</i> L.]	Caryophyllaceae	P	4a

³¹ Very variable and often hybridizing species, e.g., *P. × myoslavensis* A. Czarna & W. Antkowiak = *P. communis* L. × *P. salicifolia* Pall. (Antkowiak et al. 2008). The checklist does not include *Mespilus germanica* L. (= *Pyrus germanica* Hook. F.), a species previously reported by Szulczewski (1951).

³² Only localities in the Nidzianka Basin are native in Poland (Kaźmierczakowa 2014b). The species was reported from Łopuchowo near Oborniki (Wielkopolska) by Szulczewski (1951).

³³ *Robinia neomexicana* A. Gray hybridizes with *R. pseudoacacia* L. = *R. × holdtii* Beissn. (Zieliński et al. 2015; Zieliński 2016).

³⁴ *Robinia viscosa* Vent. hybridizes with *R. pseudoacacia* L. = *R. × ambigua* Poir. (see Zieliński et al. 2015, Zieliński 2016).

³⁵ A native species on the coast and in the mountains (Zajac & Zajac 2001), besides, planted and running wild in lowlands.

³⁶ Native in the mountains and in the foothills (Zajac & Zajac 2001), besides, planted and running wild in lowlands.

³⁷ Native in southern Poland (Zajac & Zajac 2001). In addition, it was reported from Wielkopolska: Poznań (Czarna & Mielcarski 2004) and in the vicinity of Konin (leg. et det. J. Chmiel 2011, POZ).

³⁸ Native in southern Poland (Zajac & Zajac 2001); in Wielkopolska, running wild from cultivation.

³⁹ Native in south-eastern Poland and the Upper Silesia (Zajac & Zajac 2001); in Wielkopolska, running wild from cultivation (Szulczewski 1951).

⁴⁰ Running wild from cultivation in the Botanical Garden in Poznań (W. Rakowski, P. Górska, D. Tomaszewski, pers. comm., 1992).

⁴¹ In Wielkopolska, an intergeneric wild hybrid also occurs ×*Triticosecale blaringhemii* A. Camus = ×*Triticale rimpauii* Wittm. (*Secale cereale* L. × *Triticum aestivum* L. em. Fiori et Paol.), which is often cultivated.

⁴² Szulczewski (1951) also reports *Silene nemoralis* Waldst et Kit. [*Silene italica* subsp. *nemoralis* (Waldst. et Kit.) Nyman] from Wielkopolska, which is a native species in the mountains.

Name of species	Name of family	BF	OP
<i>Silybum marianum</i> (L.) Gaertn.	Asteraceae	AB	4b
<i>Sinapis alba</i> L.	Brassicaceae	A	4b
<i>Solanum tuberosum</i> L.	Solanaceae	P	4a
<i>Solidago graminifolia</i> (L.) Salisb.	Asteraceae	P	4b
<i>Solidago patula</i> Muhl. ex Willd. ⁴³	Asteraceae	P	2a
<i>Sorbaria kirilowii</i> (Regel & Tiling) Maxim.	Rosaceae	T	4a
<i>Sorbaria sorbifolia</i> (L.) A. Braun	Rosaceae	T	4a
<i>Sorbus aria</i> (L.) Crantz s. str.	Rosaceae	T	4a
<i>Sorbus hybrida</i> L. ⁴⁴	Rosaceae	T	4a
<i>Sorbus intermedia</i> (Ehrh.) Pers. ⁴⁵	Rosaceae	T	4b
<i>Spinacia oleracea</i> L.	Chenopodiaceae	A	2a
<i>Spiraea ×billardii</i> Hérincq	Rosaceae	T	4a
<i>Spiraea ×pseudosalicifolia</i> Silverside ⁴⁶	Rosaceae	T	4a
<i>Spiraea ×vanhouttei</i> (Briot) Zabel	Rosaceae	T	4a
<i>Spiraea chamaedryfolia</i> L.	Rosaceae	T	4b
<i>Spiraea media</i> Schmidt	Rosaceae	T	4b
<i>Spiraea salicifolia</i> L. ⁴⁷	Rosaceae	T	4a
<i>Stachys byzantina</i> K. Koch	Lamiaceae	P	4a
<i>Staphylea pinnata</i> L. ⁴⁸	Staphyleaceae	T	2a
<i>Symphtum cordatum</i> Waldst. et Kit. ⁴⁹	Boraginaceae	P	2a
<i>Syringa vulgaris</i> L.	Oleaceae	T	4a
<i>Tagetes erecta</i> L.	Asteraceae	A	4a
<i>Tagetes tenuifolia</i> Cav.	Asteraceae	A	4a
<i>Tanacetum balsamita</i> L.	Asteraceae	P	4a
<i>Telekia speciosa</i> (Schreb.) Baumg.	Asteraceae	P	4a
<i>Thuja occidentalis</i> L.	Cupressaceae	T	4b
<i>Thuja plicata</i> Donn ex D. Don	Cupressaceae	T	4b
<i>Thymus vulgaris</i> L.	Lamiaceae	S	4a
<i>Tragopogon porrifolius</i> L.	Asteraceae	ABP	2a
<i>Trifolium incarnatum</i> L.	Fabaceae	A	4b
<i>Trifolium resupinatum</i> L.	Fabaceae	A	4b
<i>Trigonella coerulea</i> (L.) Ser.	Fabaceae	A	2a
<i>Triticum aestivum</i> L. em. Fiori et Paol. ⁵⁰ [= <i>Triticum vulgare</i> Vill.]	Poaceae	A	4b
<i>Triticum spelta</i> L.	Poaceae	A	?
<i>Tropaeolum majus</i> L.	Tropaeolaceae	A	4b
<i>Tulipa gesneriana</i> L.	Liliaceae	P	4b
<i>Tulipa sylvestris</i> L.	Liliaceae	P	2a
<i>Tulipa tarda</i> Stapf	Liliaceae	P	4a
<i>Typha laxmanii</i> Lepech. ⁵¹	Typhaceae	P	4b
<i>Typha minima</i> Hoppe	Typhaceae	P	?
<i>Verbena rigida</i> Spreng.	Verbenaceae	A	3a?
<i>Verbena urticifolia</i> L.	Verbenaceae	ABP	3a?
<i>Viburnum lantana</i> L.	Caprifoliaceae	T	4a
<i>Vicia faba</i> L.	Fabaceae	A	4a

⁴³ Reported from Gniezno (Szulczewski 1951).⁴⁴ Reported from the Wielkopolski National Park (Purcel 2009).⁴⁵ In Wielkopolska, the species is planted and running wild from cultivation.⁴⁶ Sometimes confused with *Spiraea ×billardii* auct. non Hérincq (Purcel 2007).⁴⁷ Native only in the south-eastern part of Poland (Zajac & Zajac 2001); in Wielkopolska, running wild from cultivation.⁴⁸ Native only in the south part of Poland (Zajac & Zajac 2001); in Wielkopolska, running wild from cultivation (Szulczewski 1951).⁴⁹ Native only in the south-eastern part of Poland (Zajac & Zajac 2001); in Wielkopolska, running wild from cultivation (Szulczewski 1951).⁵⁰ See note 41.⁵¹ Specimens of this species were collected near Konin (leg. and det. J. Chmiel 2008, POZ) and Barcin (leg. et det. J. Chmiel 2016, POZ).

Name of species	Name of family	BF	OP
Vicia sativa L.	Fabaceae	A	4b
Vinca minor L. ⁵²	Apocynaceae	S	4b
Viola suavis M. Bieb.	Violaceae	P	3a?
Viola wittrockiana Gams	Violaceae	AP	4b
Vitis vinifera L.	Vitaceae	T	4b

Explanations: BF – biological form, A – annual plants, B – biennial plants, P – perennial, herbaceous plants, S – dwarf shrubs and sub-shrubs, T – shrubs and trees; OP – observation period, 1 – to 1901, 2 – 1901-1951, 3 – 1951-2001, 4 – after 2001, a – species was observed once, b – species was observed repeatedly; ? – no data information uncertain

⁵² The range boundary runs across Poland (Meusel *et al.* 1978). In central and northern Poland, this species is running wild from cultivation. In Wielkopolska, it is found in courtyards, cemeteries (especially evangelical), old gardens, places of former settlement and near sacred buildings.