REVIEW

Understanding red blood cell rheology in sepsis and its role in clinical practice. From biomolecular aspects to possible therapeutic interventions

Emanuel Moisă¹, Silviu Negoită^{1,2}, Dan Corneci^{1,2}

¹ Anaesthesiology and Critical Care Department, Elias Clinical Emergency Hospital, Bucharest, Romania

² Carol Davila University of Medicine and Pharmacy, Bucharest, Romania

Correspondence to:

Emanuel Moisă, MD, Department of Anaesthesia and Critical Care, Elias Clinical Emergency Hospital, 17 Marasti Bd, Bucharest, Romania
E-mail: emanuelmoisa@gmail.com

Conflicts of interests

Nothing to declare

Acknowledgements

None

Keywords: RBC rheology, nitric oxide, band 3 phosphorylation, reactive species, phosphatidylserine redistribution, deformability

These authors take responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation.

Central Eur J Clin Res 2018;1(1):40-58

Received: 1.01.2018, Accepted: 30.08.2018, Published: 3.09.2018

Copyright © 2018 Central European Journal of Clinical Research. This is an open-access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Erythrocyte rheology is of interest in understanding microcirculation and oxygen delivery and consumption alterations induced by sepsis and septic shock. Several mechanisms are proposed: (i) direct or indirect RBC membrane alterations, (ii) abnormal intraerythrocytic homeostasis, (iii) RBCs interaction with other cells and extracellular molecules, (iiii) increased reactive species production and altered redox homeostasis. In this review, we describe in part these mechanisms and what's the impact of these hemorheological disturbances on the outcome

and mortality rate. Also, we outline the possible therapeutic interventions and further perspectives regarding sepsis and septic shock management.

Introduction

In 2016 The Third International Consensus for Sepsis and Septic shock defined sepsis as "a life-threatening organ dysfunction caused by a dysregulated host response to infection" [1]. The concept of severe sepsis is not actual anymore [1], but in this review we also collected data about red blood cell rheology in context of severe sepsis (as it was defined by the previous definitions in 2001) [2]. Sepsis and septic shock still represent the most common cause of death

and critical illness worldwide [1]. Pathophysiology of sepsis is complex and involves alterations particularly observed in microcirculation and blood constituents homeostasis [1]. Of interest in this review is the way sepsis alters erythrocyte rheology and function, the actual possible therapeutic tools and future perspectives regarding the treatment of these alterations.

Basic principles of hemorheology. Sepsis and septic shock induced altered red blood cell (RBC) rheology

Hemorheology is a branch of fluid dynamics studying and describing flow and deformation behaviour of blood and its cells (i.e., erythrocytes, leukocytes, platelets) [3]. Blood rheology still represents an area of interest, after decades of studies and, yet, there are still many aspects that need to be elucidated, like aggregation's (one major determinant of erythrocyte rheology) implications in pathophysiology of several diseases [4, 5]. But, in this review, we will focus on RBC rheology and how sepsis and septic shock influences it.

Blood viscosity

Blood is a non-Newtonian two-phase fluid. Plasma represents the suspending phase for blood cells and is considered a Newtonian fluid, thus alterations of its viscosity will lead to changes in whole blood viscosity independent of blood cells properties (changed or not) [3,6]. Plasma viscosity can be increased in acute phase of inflammatory diseases (increased mostly by plasma proteins, like fibrinogen and other acute phase reactants) and by extension, in sepsis [3,7].

Added to plasma, blood cells make whole blood viscosity higher than plasma viscosity and this has implications in blood flow behaviour (fluid transition from Newtonian to non-Newtonian) [3]. Viscosity can be expressed as the ratio between shear stress and shear rate [8]. In terms of viscosity, shear rate determines RBC behaviour, at high rates causing RBC dispersion and preventing aggregation, and at low rates increasing their chance to aggregate [8]. Microvascular stasis (low flow state) in sepsis may lead to low shear rates and microthrombosis (capillary-stopped flow) [8,9].

Hematocrit can seriously alter blood viscosity as seen in hyperviscosity syndromes which can associate severely decreased tissue perfusion pressures [10]. On the other hand, a low hematocrit (anemia) leads to hyperdynamic circulation [11]. In sepsis, development of a hyperviscosity syndrome is controversial and less

probable [7]. Systemic inflammatory response syndrome (SIRS) is frequently associated with increased acute phase reactants which, in theory, leads to increased plasma viscosity [3], sepsis and septic shock associate seriously altered microcirculation (vasodilation) [1] and by extension, a hyperdynamic status, and also, in time, a progressive anemia [1,7,12] which, in turn, reduces blood viscosity [3,8]. Blood viscosity in sepsis can maybe be characterized by an increased variability [7] caused first and foremost by the high volumes of fluids following aggressive volemic resuscitation [12]. Given this, a hyperviscosity syndrome is unlikely to develop and or it can be at least, prevented [13]. Hematocrit is the only hemorheological parameter included in the Acute Physiology and Chronic Health Evaluation (APACHE II) predictive score [14]. In sepsis, hematocrit value can be falsely increased with anemia [12], and this can modify studies results in blood rheology and also lead to false lower APACHE II scores in ICU patients assessment. Moreover, blood viscosity can be influenced by blood transfusions in sepsis [7]. Finally, knowing that dialysis increases blood viscosity and alters RBC rheology [15,16] patients who require renal replacement therapy may present further altered hemorheology. *Piagnerelli et al* concluded in their RBC rheology in sepsis review that blood viscosity is influenced mainly by plasma viscosity, shear rate, deformability and aggregation [8].

Aggregation

Capillary-stopped flow is specific for sepsis [9], and RBC rheologic changes like increased aggregability and adhesion are incriminated as possible mechanisms [8,9,17]. Of course, this phenomenon is much complex and must be integrated in the overall pathophysiological mechanisms of sepsis.

Aggregation can be described as the equilibrium between aggregating and disaggregating forces [3]. The disaggregating forces (electrostatic repulsion between cells, RBC membrane elastic energy and fluid shear forces) are dependent of RBC integrity and describe the concept of RBC aggregability, while aggregation is determined by large macromolecules (i.e., fibrinogen) [3,4,6,18]. The way this imbalance between aggregating and disaggregating forces may appear and how it can be ameliorated will be discussed later in this review.

Deformability

The normal erythrocyte is a biconcave disc shaped-like anucleated cell, approximately

7 μm in diameter and 2 μm thick [9]. Its primary function (considered the only one for a long time) is to transport oxygen from lungs to the tissues, but in the past years, erythrocyte was described not only as an oxygen carrier, but also as an oxygen sensor [19,20]. Of course, with recent findings there is much more we know about RBCs function and it will be described in this review. Deformability represents erythrocytes ability to deform when exposed to blood flow and this is of utter importance for physiologic capillary circulation, where RBCs can undergo extreme deformability in order to pass and maintain homeostasis [8,9,21,22]. RBC deformability depends on (i) membrane surface area-to-volume ratio also known as erythrocyte geometry, (ii) elements forming cytoskeletal proteins (spectrin, ankyrin, band 3 protein domain, 4.1 protein and 4.2 protein) affecting RBC membrane mechanical features and RBC shape and (iii) intraerythrocytic water and ion homeostasis. [21,22,23,24]. Red blood cell deformability was recently reviewed by Huisjes et al [21].

RBCs deformability can be an early marker of sepsis [7,25,26,27] and also a marker of severity [28]. Furthermore, altered deformability can be a prognostic factor being associated with outcome [29] and progression to organ dysfunction [30]. Finally, less probable, it can be an element of differential diagnosis between septic and non-septic critically ill patients [7,31]. RBC deformability is altered through several proposed mechanisms we will discuss later in this review along with how therapeutic interventions ameliorated deformability and changed outcome. Erythrocytes increased sphericity, variable volume and deformability alterations are factors which may alter red blood cells distribution width (RDW, a basic paraclinical tool studied routinely with every complete blood count laboratory examination) along with decreased hematocrit and hemoglobin [9,11]. In fact, RDW is a measure of RBCs grade of anisocytosis [9,11], and it was found to be changed in septic patients by several studies [?]. Increased RDW was associated with poor immediate [32,33] and short term prognosis: increased 28-day [34,35], 30-day [36] and 90-day mortality [34]. It is also considered, that RDW value can be helpful in differentiating the severity of sepsis [33,37] and is associated with MDA and TNF- α levels during the first week of sepsis [33].

Mechanisms explaining RBC rheology in sepsis Erythrocyte membrane alterations in sepsis RBC membrane sialic acid (SA) content and bacterial compounds action on RBC membrane

The way RBC content of sialic acid in septic patients changes RBC shape was described by Piagnerelli et al. including in their studies non-septic critically ill patients, septic patients and healthy volunteers as control group [38,39]. Piagnerelli et al aim was to observe if there is any relationship between RBC membrane sialic acid content and RBC shape using flow cytometry. They reported significantly lower values of SA content in septic patients compared with non-septic patients and healthy volunteers, and this low values have been correlated with RBC shape; RBCs were more spherical in septic patients and also had a decreased capacity of sphericity in hypo-osmolar solution. [38,39]. Just recently, Qadri et al proved that artificial desialation of RBCs with *Clostridium perfringens* derived neuraminidase induced increased phosphatidylserine exposure and increased intraerythrocytic calcium levels [40]. Changes in RBC shape due to sialic acid content and sialidase activity was also described in patients with diabetes. Diabetic patients presented lower quantities of sialic acid and an increased sialidase activity [41,42] and also, a higher percent of senescent erythrocytes [42]. RBCs in diabetic patients present a lower deformability [42] and increased aggregability [41]. Moutzouri et al [43] reported that sepsis and diabetes have an additive effect on red blood cell deformability with significantly higher rigidity indices in patients with diabetes and sepsis compared with septic but not diabetic patients, diabetic but not septic patients and healthy volunteers [43].

Another proposed mechanism was a direct effect of bacterial virulence factors on RBC membrane sialic acid content. [8]. Milligan et al [44] and later, Mattingly et al [45], reported that serotype III isolates of group B *Streptococci* from infants having the disease produced more extracellular neuraminidase than serotype III isolates of group B *Streptococci* in those colonized and asymptotically. Neuraminidase production was not an absolute condition for infants to develop the disease accordingly to Mattingly et al [45]. These findings are of big importance in understanding the pathologic molecular mechanisms given the fact that sialic acid is attached to A glycoprotein (the most abundant sialoglycoprotein on erythrocytes) and is also a monosaccharide derivative of

neuraminic acid which makes it susceptible to glycoside hydrolysis action of neuraminidase, which leads to a reduction of membrane surface negative charge [9]. As described above, smaller RBC sialic acid content leads to a decreased deformability [38,39]. A low negative charge in membrane surface will also reduce the repelling forces between RBCs and vascular endothelium, therefore increasing their susceptibility to aggregation [3,4,6]. Liukkonen et al [46] studied *Streptococcus suis* (a bacterial pathogen causing severe infections in young piglets and meningitis in humans) erythrocyte binding specificity of sialic acid. In presence of sialidase, hemagglutination of erythrocytes exposed to *Streptococcus suis* was abolished [46]. It was observed that resialylation of desialylated RBC with beta 1-3(4)GlcNAc alpha 2-3-sialyltransferase caused a strong hemagglutination. Also, they found that *Streptococcus suis* binds to Band 3, Band 4.5 proteins and glycophorin A. [46] Parkkinen et al [47] findings about S-fimbriated *Escherichia coli* effects on erythrocytes are comparable with Liukkonen et al, reporting *Escherichia coli*'s specificity of binding to a proteic sequence of the O-linked oligosaccharide of glycophorin A; hemagglutination of bacteria was abolished when erythrocytes were treated with neuraminidase and trypsin [47].

Other bacteria have the capacity of producing neuraminidase: *Pseudomonas aeruginosa* [48], *Clostridium perfringens* [49], *Streptococcus pneumoniae* [50], *Vibrio cholerae* [51], leading to rheological changes in RBCs. Piagnerelli et al studied neuraminidase activity in non-septic critically ill ICU patients and septic patients, reporting increased levels of free sialic acid in plasma and increased blood neuraminidase activity [38,49]. RBCs incubation with *Clostridium perfringens* neuraminidase using different concentrations led to shape alterations of RBCs that became more spherical and this change was correlated with higher sialic acid hydrolysis rates [49]. Also, authors reported increased RBC levels of 2,3-DPG known to alter RBCs shape [52,53] and increased RBC levels of lactate, but it was not possible to determine if these shape changes were due to increased 2,3-DPG or the spherical shape determined an abnormal RBC glycolysis leading to increased 2,3-DPG levels (2,3-DPG was measured after at least 10 hours of incubation and RBCs shape was already changed) [49].

The importance of RBC sialoglycoproteins in sepsis can be correlated with Lizcano et al recent findings about erythrocytes importance in suppressing neutrophil activation and

apoptosis in blood [54]. The authors reported that sialoglycoproteins from RBC surface are important in neutrophil suppression through Siglec-9 (a receptor known to recognize sialic acid and to decrease innate immune cell activation), both ex vivo and in vitro [54].

Phosphatidylserine redistribution on the outer RBC membrane leaflet is associated with increased eryptosis and erythrocyte clearance in sepsis

Dinkla et al [55] observed that once SIRS is developed in experimental endotoxemia (LPS administered IV), membrane lipid remodeling appears in both human and animal experiments, leading to increased phosphatidylserine exposure. An interesting observation was that in experimental endotoxemia are not observed the exactly same effects as seen in sepsis [55]. In sepsis, bacterial virulence factors, others than neuraminidase were associated with membrane externalization which is intercepted as an erythrocyte senescence signal in the reticuloendothelial system leading to increased eryptosis and RBC clearance, explaining, in part, sepsis induced anemia [56]. Different bacterial compounds like pyocyanin [48], α -hemolysin [57], listeriolysin [58], peptidoglycans [59], lipopeptides [60] and sphingomyelinase [61] were associated with membrane externalization through exposure and redistribution of phosphatidylserine on the outer RBC membrane leaflet. Eryptotic RBCs adhesion is increased by bacterial peptidoglycans and it seems to follow a CXCL16-dependent fashion as it was reported by Abed et al [62]. Qadri et al exposed in vitro human RBCs to the virulence factor pyocyanin released by *Pseudomonas aeruginosa* and reported altered intraerythrocytic homeostasis of water (dehydration with cell shrinkage) and calcium (increased RBC cytosolic calcium), increased RBC reactive oxygen species (ROS) concentration and increased phosphatidylserine redistribution on the outer RBC membrane [48]. Pyocyanin is redox-active metabolite capable of producing ROS and also, can inhibit catalase, which, in turn can lead to increased hydrogen peroxide levels [9] (its effects are described later in the sepsis induced oxidative stress chapter).

Phosphatidylserine redistribution is influenced by other factors too, like altered intraerythrocytic calcium homeostasis and sepsis induced oxidative stress [9,48,63,64,65,66,67]. Calcium is important in regulating RBC membrane phosphatidylserine distribution [63] and calcium homeostasis is profoundly disturbed in sepsis [64,65]. Oxidative stress imbalances

have an important contribution on membrane damage too, through lipid peroxidation and protein oxidation which determines increased RBCs inner leaflet membrane exposure to the extracellular space [9,48,66,67]. Eryptosis is also associated with increased ceramide formation in sepsis [68].

Sepsis induces RBC membrane proteins alterations Band 3 protein phosphorylation

Phosphorylation of band 3 protein is of importance in sepsis because it plays a significant role in intraerythrocytic metabolic alterations of glycolysis and oxygen delivery, and by extension influences RBC deformability (through several mechanisms) and RBC NO release behaviour [9,69,70,71,72]. Band 3 protein phosphorylation and dephosphorylation was reported in both in vitro [70,71,72] and in vivo [72] studies, using tyrosine kinase inhibitors. Recently, Lin et al in a murine cecal ligation puncture (CLP) sepsis model reported increased cytoplasmic domain of Band 3 protein (cdB3) proteolysis via Western Blot testing. Furthermore, cdB3 proteolysis was higher in mice undergoing severe CLP compared with those undergoing sham CLP [73]. Spolarics et al [74] and Condon et al [72] observed that band 3 phosphorylation was associated not only with reduced RBC deformability but also with altered anion activity.

Band 3 phosphorylation and dephosphorylation depends on several intraerythrocytic alterations: calcium concentration [75], oxidative damage on RBC membrane [76], low oxygen partial pressure leading to increased deoxyhemoglobin levels [77]. These alterations are already described in sepsis and can, in part, explain this phenomenon. Altered calcium homeostasis in erythrocyte leads to phosphotyrosine phosphatase dissociation from band 3 allowing its phosphorylation in a reaction catalyzed by the enzyme phosphotyrosine kinase [75]. Band 3 phosphorylation then determines increased cytosol concentration of glyceraldehyde dehydrogenase, phosphofructokinase and aldolase (glycolytic enzymes anteriorly inactive and attached to cdb3) and enhanced glycolysis [75]. Band 3 phosphorylation is also determined by deoxyhemoglobin which through binding to cdb3 enhances glycolysis via the same mechanism described above [77]. Sepsis hypoxic status reduces drastically the hexose monophosphate pathway and further, NADPH concentration, which is vital in producing reductants in order to protect RBCs from oxidative

damage [78,79]. Thus, ROS produced from the increased hemoglobin autooxidation processes along with other sources of ROS, in a low oxidative defense state, will lead to increased glycolysis via hydrogen peroxide [76] in the same fashion with calcium and deoxyhemoglobin. Enhanced glycolysis will lead to increased ATP efflux [80]. ATP will bind to the P2Y receptors, inducing NO synthesis, which will further alter RBC deformability [81]. Moreover, glycolytic pathway stimulation, decreased levels of intraerythrocytic reductants (altered NO-glutathione interaction), band 3 phosphorylation and dephosphorylation influences intraerythrocytic NO chemical reactions and modifies its bioavailability and release from RBCs [69]. Finally, band 3 protein phosphorylation in presence of high fibrinogen levels determines a higher NO release from RBCs [82,83].

Piagnerelli et al suggested that membrane protein content alterations are not correlated with abnormal RBC rheology seen in sepsis [84]. 25 patients were included in study, 15 with sepsis, 10 non-septic critically ill and 10 healthy volunteers as control group. No significant differences were observed in protein content between septic and non-septic patients, but in spectrin content between control group and non-septic critically ill patients, with decreased spectrin levels for the latter. The authors reported decreased levels of band 3, band 4.1 and 4.2, ankyrin, spectrin and increased band 3/spectrin, Protein 4.2/band 3 ratios, in all critically ill patients, but as long as differences between the two groups were not found, the authors suggested their conclusion presented earlier. [84]

Altered intraerythrocytic homeostasis 2,3-diphosphoglycerate (2,3-DPG)

The erythrocyte's fundamental role is oxygen delivery from lungs to the tissues by binding the oxygen molecule to hemoglobin. 2,3-DPG represents one of the most important organic phosphates in the RBC and is produced through the Rapoport-Lubering shuttle – the side path of glycolysis taking place in RBC. 2,3 DPG is an allosteric regulator of hemoglobin and has the role of reducing hemoglobin's affinity for oxygen by forming a reversible complex with deoxyhemoglobin [9,85]. 2,3-DPG synthesis is increased in the presence of hypoxaemia and acidemia resulting in right shifting of the oxyhemoglobin dissociation curve (ODC), decreased oxygen affinity to hemoglobin and increased release of oxygen to the tissues (Bohr effect). Studies results may seem paradoxical as a

Calcium homeostasis is profoundly disturbed in septic patients [64]. Totson et al reported that hypocalcemia in septic patients can predict mortality [7] and Desai et al found that septic nonsurvivors had lower plasma calcium compared with survivors [65]. This alteration is observed both in extracellular [65,91] and intracellular [64,92] calcium physiology. Intracellular calcium has structural (erythrocyte morphology, rheology, adhesion, cellular volume) and biochemical (proteolysis, transglutaminase activity, redox status) effects on RBC [9,63]. Intracellular calcium homeostasis alterations in septic patients were described for liver cells [93], cardiomyocytes [94], lymphocytes [95] and of interest in this review, erythrocytes [64,92,96,97]. Ruef et al reported increased intraerythrocytic calcium concentration and decreased erythrocyte deformability after RBCs (from healthy donors) were incubated with *E. coli* lipid A [92]. Erythrocyte deformability was found to be inhibited by administration of verapamil (a calcium channel blocker) and protein kinase inhibitors like staurosporine [92]. Todd and Mollit observed that intracellular calcium homeostasis is altered in vivo (septic patients) and in vitro (whole blood incubated with *E. coli* endotoxin), respectively [64]. They found that increased intraerythrocytic calcium levels (as free cytosolic calcium measured by fluorescent spectroscopy) is specific for septic patients. These in vivo results were studied also in vitro by incubating whole blood (pretreated with dantrolene, or verapamil, or ATP and posttreated or not with ATP) with *E. coli* endotoxin in the presence or absence of extracellular calcium [64]. Todd and Mollit reported the same increase in free cytosolic calcium for the in vitro experiment and also found out that intracellular calcium concentration was not influenced by pretreatment with verapamil or dantrolene (a ryanodine receptor blocker known to decrease intracellular calcium), transmembrane calcium channels, intracellular calcium deposits and extracellular calcium concentration [64]. An interesting finding was that pretreatment with ATP led to a smaller increase in intraerythrocytic calcium and this phenomenon may rise the suspicion of an energy deficiency being responsible for the cytosolic calcium increase. Nowadays, RBCs ATP deficit in sepsis is described through many mechanisms we will discuss later and can support Todd and Mollit theory.

WBCs modulation and increased oxidative stress role in RBCs intracellular calcium homeostasis was reported by Todd and Mollit which incubated anticoagulated whole blood samples from healthy donors with LPS and observed an increase in erythrocyte cytosolic

calcium concentration, but this increase was dependent on WBCs presence; in absence of WBCs this change was not reported [98]. WBCs implication in these findings was demonstrated when incubation of whole blood pretreated with xanthine oxidase inhibitors (allopurinol), leukocytes modulators (pentoxifylline) and free radical scavengers (superoxide dismutase) with LPS resulted in smaller increases in cytosolic calcium concentration [98].

Adenosine triphosphate

In conditions of impaired oxygenation as seen in sepsis, erythrocytes are capable of releasing vasoactive molecules like nitric oxide, nitrosothiols and ATP [99]. In conditions of hypoxia ATP can be released [100,101] through erythrocyte Pannexin-1 channel [9,102]. The released ATP binds to P2Y receptors and causes vasodilation [9,29,103,104]. Mathie et al reported that nitric oxide hepatic vasodilation was ATP-induced in rabbits perfused in vitro with Krebs-Bülbring buffer via hepatic artery and portal vein. The study results suggested that ATP binding to P2Y receptors (which are abundant in endothelial cells) induces NO and causes vasodilation in hepatic artery vascular bed. ATP-induced NO release via endothelial cells was reported also by Busse et al [105]. ATP was found to be an important mediator of NO release in isolated perfused rabbit lungs [107,106] and was reported in dogs models too [106]. An interesting and contradictory hypothesis to what was presented until now comes from Serroukh et al [?] suggesting that the decreased functional capillary density can be caused by an ATP-induced vasoconstriction in microvascular bed, secondary to increased ATP release from abnormal erythrocytes, but further investigations are needed to establish this theory.

RBC reduced deformability determines ATP release [27,81,106,109] and, in turn, ATP deposits are low in sepsis [100,101]. Decreased intraerythrocytic ATP impairs PMCA activity leading to an increase in intracellular calcium and decreased RBC deformability [8]. Bateman et al [81] and Rozier et al [110] observed ATP release dependent on RBC oxygenation is impaired in presence reduced pH (in a 6 h rat hypotensive sepsis model) [81] and inhibited, at high lactate concentrations, respectively [110].

Another possible mechanism for increased RBC ATP release in sepsis can be related to pore-forming virulence of some bacteria. Recently, Greve et al, in an in vitro study on knock-out mice, reported pore forming in RBC membrane and immediate RBC ATP re-

lease after α -hemolysin is inserted into RBC membrane via binding to P2X1 receptors [111]. This phenomenon was described in vivo in humans and is associated with P2X7 receptors [112]. Hemolysin *E. coli* producing strains are observed mostly in patients with severe urinary tract infections and Greve et al speculated that P2X receptor antagonists maybe will be useful in urosepsis [111]. Other bacteria can induce the same cellular changes as described above: *Staphylococcus aureus* via α -toxin [113] and *Clostridium perfringens* [114] via β -toxin.

Sepsis induced oxidative stress

Sepsis induced oxidative stress represents an important area of interest nowadays for scientific researchers. Molecular studies regarding oxidative stress were eye-opening in understanding sepsis pathophysiology and a more actual concern for scientists is how we can therapeutically intervene to correct all these imbalances [115]. Sepsis mechanisms are complex and multifactorial leading to a vicious circle in which disease creates disease. It's effects are of utter importance in terms of microcirculation disturbances and oxygen physiology, from uptake and delivery in (normal/abnormal) RBCs to (abnormal) utilization and consumption in mitochondrion. Several studies emphasize the oxidative imbalance in sepsis and critically ill patients describing a diminished antioxidant activity [9,115-121] and an increased oxidative status [1,8,9,115-117,121]. Oxygen physiology is altered by transformation of oxygen into reactive oxygen species like superoxide ($O_2^{\cdot-}$), hydroxyl radical ($\cdot OH$), hydrogen peroxide (H_2O_2), hypochlorous acid ($HOCl$) and peroxynitrite ($ONOO^-$) [115], oxygen free radicals and oxidizing species [9]. In ROS producing and redox homeostasis alteration are implicated many factors and of interest are: neutrophils [115, 122], endothelial cells [9, 115], the erythrocyte itself [123] and plasma oxidants (nitric oxide, xanthine oxidase, malonyldialdehyde) [9].

For this review, ROS and altered redox homeostasis is important because several studies reported altered RBC deformability [124-135] and increased RBC aggregability [124,135] both related to oxidative imbalance in septic patients. Baskurt et al found that intracellular superoxide anion production is responsible for altered RBC deformability and extracellular superoxide anion increased RBC aggregation, when RBCs were exposed to this ROS [124]. RBC exposure to hydrogen peroxide [125,126,132], determines spectrin-hemoglobin cross-linking [126,132] decreased deformability

[126,132], ekynocytes formation [125,126] and increased RBC adhesion [126], while exposure of RBCs to lipid peroxidation products like MDA determines decreased RBC deformability [67]. ROS influence on band 3 phosphorylation was already discussed earlier in this review. (see band 3 phosphorylation).

Of importance is also RBCs capacity to produce ROS under certain conditions [9]. Erythrocyte autooxidation is determined mainly by H_2O_2 and superoxide and is prevented by catalase and superoxide dismutase (SOD) [136]. The most important factor determining RBC autooxidation is superoxide anion production [123] and this it's production is favoured by high temperature, low pH, increased 2,3-DPG [137,138] and a partial oxygenation of hemoglobin [137]. High temperature as a manifestation of systemic inflammation, hypoxia determined by microcirculation alteration and abnormal oxygen physiology, low pH and in turn, increased 2,3-DPG are frequent in sepsis, making RBCs more susceptible to autooxidation, mainly in areas of capillary stopped-flow where oxygen hemoglobin saturations are lower [81,139].

In sepsis, the antioxidant capacity is overwhelmed by the excessive oxidative species production [115]. This hypothesis is sustained by the studies in which administration of antioxidant molecules in septic patients or critically ill patients improved the outcome [116,121, 128,140-146]. Administration of N-acetylcysteine (NAC) alone was found to transiently improve tissue oxygenation [140,141], increase hepatic blood flow and liver function [142], improve static lung compliance and lower plasma IL-8 levels [141] and decrease lipid peroxidation with a reduction of organ failure at day 10 [143]. Selenium administration alone was found to reduce mortality rate by 25%, decrease inflammatory response [144] and reduce the incidence of acute kidney injury requiring renal replacement therapy [145]. Gadek et al [147] and Pontes-Aruda et al [148] administered a combination of antioxidant molecules (vitamin c, vitamin E, β -carotene, selenium) enterally in approximately same concentrations and number of days (4 days [147] versus 4-7 days [148]). They reported mostly the same results: improved oxygenation, reduced length of ventilatory support and ICU stay, less new organ failure [147,148] and reduced mortality [148]. The pitfall of this combination of antioxidant molecules is the impossibility of associating the benefits with a single or specific molecule, as was reported by Marik et al as well [116] who found that co-administration of vitamin C, thiamine and hydrocortisone

prevented organ dysfunction and decreased mortality.

Moreover, several studies reported an inversely proportional relationship between plasmatic levels of antioxidants and oxidants [117,118,120]. Decreased antioxidants like vitamin E [117,118,120], ascorbate [116,117,118], β -carotene [117,118] and retinol [117] were found in septic patients associating important lipid peroxidation activity (increased MDA [118] or other evidence supporting lipid peroxidation [117]. In support of required proper antioxidant levels and in terms of RBC rheology comes evidence suggesting that supplementation in septic patients with vitamin C improved outcome by preventing organ failure [116], determined inhibition of erythrocyte destruction through eryptosis [146] and vitamin E supplementation decreased mortality rate and ameliorated RBC deformability [128].

Nitric oxide

NO release and effects in sepsis and septic shock were described in several studies [27,81,149-156]. Under physiologic conditions, NO is synthesised in endothelial cells from L-arginine catalyzed by e isoform nitric oxide synthase (eNOS), known also as c isoform (constitutive, cNOS) [157]. Produced in small amounts, NO is vital in capillary tone regulation [8]. Of interest in sepsis, is the i isoforme, which is the inducible NOS (iNOS) found to be stimulated excessively in sepsis by different mediators like LPS, cytokines (IL1, IL6, TNF- α , IFN γ) and ATP [8,56]. Recently, Takatani et al reported that iNOS is associated with immune cell migration and hypothermia in the late phase of sepsis [158]. Pentraxin 3 is known to be overexpressed in sepsis and septic shock [159] and is also thought it can blunt endothelial NO production [160]. Another NOS isoform is the neuronal isoform (nNOS) thought to take part in SIRS pathophysiology [161] and more important, it seems to have serious implications in vascular alterations seen in sepsis [162]. Another important aspect is gram-positive bacteria possession of bacterial isoform NOS (bNOS) which was reported to have multiple roles: protection against oxidative stress, antibiotic resistance and host immune response [163,164].

Hemoglobin and NO interaction is important in sepsis. NO affinity for hemoglobin is high [99] (increased affinity for thiol groups or for the iron ion), but NO binding at Hb is dependent on oxygen partial pressure [69, 99]. At arterial oxygen partial pressure (high-pressure, PaO₂) NO and oxy-Hb form S-nitrosohemoglo-

bin (SNO-HBO₂) through NO binding at the thiol group of cysteine β 93 [69], but at low oxygen partial pressures (as it happens in sepsis), NO interacts with the iron ion binding to it and resulting in nitrosylhemoglobin molecules formation [165]. In 1996, Jubelin et al [166] suggested that erythrocytes can be able to produce NO. In 2004 Carvalho et al [167] demonstrated in vitro NO erythrocyte production using the amperometric method. One year later, Kleinbongard et al reported that RBCs possess a functional eNOS [168]. Recently, Lin et al in a murine CLP sepsis model, reported decreased levels of RBC NO content (total RBC NO, FeNO, and SNO content) [73]. RBC NO content was measured using chemiluminescence and was significantly decreased, with lower values in severe CLP mice versus sham CLP mice [73]. These results can be correlated with Herdade et al observations on NO efflux behaviour related to acute inflammatory phase [169]. The authors created an in vivo mice model of acute inflammation by injecting platelet-activator factor (PAF) in mice scrotum and reported an inversely proportional relationship between NO efflux and development of acute-phase response, with decreasing rates of NO efflux as the inflammatory response was developed and approximately normal NO values in the recovery phase. An important observation authors made was that RBC deformability was not completely recovered and this can be, in part, attributed to PAF's ability to breakdown RBCs sphingomyelin [169].

In sepsis, RBC NO and nitrosothiols production and release can be increased via band 3 phosphorylation and NO interaction with reactive species resulted from hemoglobin autooxidation [69,99]. These mechanisms were described in more detail earlier (see Band 3 phosphorylation). Simplyfied, it's important to keep in mind that NO is released from RBCs in conditions of hypoxia, which in sepsis has a significant impact on erythrocyte rheology and microcirculation [81].

Figure 2 summarizes many of the mechanisms presented in this review.

Abbreviations: superoxide dismutase (SOD), catalase (CAT), peroxiredoxin (Prx), glutathione peroxidase, vitamin C, E, A (vit. C, E, A), NADPH O (NADPH oxidase), xanthine oxidase (XO), reactive oxygen species (ROS), superoxide ($O_2^{\cdot-}$), hydroxyl radical ($\cdot OH$), hydrogen peroxide (H_2O_2), hypochlorous acid (HOCl) and peroxynitrite (ONOO $^-$), interleukin-1, -6 (IL-1, -6), lipopolysaccharides (LPS), nitric oxide (NO), inducible nitric oxide synthetase (iNOS), adenosine triphosphate (ATP), white blood cell (WBC), cytoplasmic domain of band 3 (cdB3),

phosphatidylserine (PS), purinergic receptors (P₂YR), glycolytic enzymes (GE), 2,3-diphosphoglycerate (2,3-DPG), oxygen partial pressure (pO₂), tumor necrosis factor- α (TNF- α), interferon- γ (IFN- γ), deoxygenated haemoglobin (DeoxyHb), plasma membrane calcium channel (PMCCa²⁺).

Pharmacologic interventions regarding modulation of NO activity in sepsis is of great concern nowadays. In 1993, Korbut and Gryglewski [181], reported decreased RBC (from rabbit) deformability in presence of L-NAME (N-nitro-L-arginine methylester, a NO synthase

49

in sepsis are not completely elucidated [188]. Increases in mortality rate were reported, or no effects at all on microcirculation [189], but, in one study authors reported an improvement in microcirculation when nitroglycerin was administered after fluid resuscitation [190]. Actually, nitric oxide pharmacologic interventions are not included in guidelines because there is no strong evidence supporting the use of any of the described agents [12].

Conclusions

The erythrocyte is not seen anymore only as an oxygen carrier, but also as an oxygen sensor and vasoactive molecules producer, playing a key-role in microcirculation alterations seen in sepsis and septic shock. RBCs deformability and aggregation can be influenced through several mechanisms that still need to be completely established, but many of them seem to be interconnected, to augment or inhibit one another, having different trends. In the future, RBCs maybe will be possible targets in sepsis and septic shock management, but further studies are required in order to improve the unfavorable outcome and mortality rate seen in these patients.

References

1. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *Jama* 2016;315(8):801
2. Levy MM, Fink MP, Marshall JC, Abraham E, Angus D, Cook D, et al. 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Intensive Care Medicine* 2003;29(4):530–8
3. Baskurt OK, Meiselman HJ. Blood rheology and hemodynamics. *Seminars in Thrombosis and Hemostasis* 2003;29(5):435–50
4. Baskurt OK, Neu B, Meiselman HJ. Red Blood Cell Aggregation. *Boca Raton. J Lab Clin Med* 1997;130(2):183–90
5. Nemeth N, Deak A, Szentkereszty Z, Peto K. Effects and influencing factors on hemorheological variables taken into consideration in surgical pathophysiology research. *Clinical Hemorheology and Microcirculation* 2018;69(1-2):133–40
6. Baskurt OK. In vivo correlates of altered blood rheology. *Biorheology* 2008;45(6):629–38
7. Totsimon K, Biro K, Szabo ZE, Toth K, Kenyeres P, Marton Z. The relationship between hemorheological parameters and mortality in critically ill patients with and without sepsis. *Clinical Hemorheology and Microcirculation* 2017;65(2):119–129
8. Piagnerelli M, Zouaoui Boudjeltia K, Vanhaeverbeek M, Vincent JL. Red blood cell rheology in sepsis. *Intensive Care Med* 2003;29:1052–1061
9. Bateman R, Sharpe M, Singer M, Ellis C. The Effect of Sepsis on the Erythrocyte. *International Journal of Molecular Sciences* 2017;18(9):1932
10. Lenz C, Rebel A, Waschke K, Koehler R, Frietsch T. Blood viscosity modulates tissue perfusion – sometimes and somewhere. *Transfusion Alternatives in Transfusion Medicine* 2007;9(4):265–72
11. Beck N. Anemia: General Considerations. *Diagnostic Hematology*. London: Springer 2009;199–218
12. Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R et al. Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock: 2016. *Intensive Care Medicine* 2017;43(3):304–377
13. Toth K, Kesmarky G. Clinical significance of hemorheological alterations In: Baskurt OK, Hardeman MR, Rampling MW, Meiselman HJ, (eds) *Handbook of Hemorheology and Hemodynamics*, Amsterdam 2007:392–404
14. Knaus WA, Draper EA, Wagner DP, Zimmerman JE. APACHE II: a severity of disease classification system. *Crit Care Medicine* 1985;13:818–829
15. Reinhart WH, Cagienard F, Schulzki T, Venzin RM. The passage of a hemodialysis filter affects hemorheology, red cell shape, and platelet aggregation. *Clinical Hemorheology and Microcirculation* 2014;57:49–62
16. Hasler C, Owen G, Brunner W, Reinhart W. Echinocytosis induced by haemodialysis. *Nephrology Dialysis Transplantation* 1998;13(12):3132–3137
17. Eichelbronner O, Sibbald WJ, Chin-Yee IH. Intermittent flow increases endotoxin-induced adhesion of human erythrocytes to vascular endothelial cells. *Intensive Care Medicine* 2003;29:709–714
18. Simmonds MJ, Meiselman HJ, Baskurt OK. Blood rheology and aging. *Journal of Geriatric Cardiology* 2013;10(3):291–301
19. Ellsworth ML. The red blood cell as an oxygen sensor: what is the evidence? *Acta Physiologica Scandinavica* 2000;168:551–559
20. Jensen FB. The dual roles of red blood cells

- in tissue oxygen delivery: oxygen carriers and regulators of local blood flow. *Journal of Experimental Biology* 2009;212:3387–3393
21. Huisjes R, Bogdanova A, Solinge WW, Schiffelers RM, Kaestner L, Wijk RV. Squeezing for Life – Properties of Red Blood Cell Deformability. *Frontiers in Physiology* 2018;9:656
 22. Bor-Kucukatay M, Wenby RB, Meiselman HJ, Baskurt OK. Effects of nitric oxide on red blood cell deformability. *American Journal of Physiology-Heart and Circulatory Physiology* 2003;284(5)
 23. Nans A, Mohandas N, Stokes DL. Native Ultrastructure of the Red Cell Cytoskeleton by Cryo-Electron Tomography. *Biophysical Journal* 2011;101(10):2341–50
 24. Gallagher PG. Hereditary elliptocytosis: spectrin and protein 4.1R. *Seminars in Hematology* 2004;41(2):142–64
 25. Reggiori G, Occhipinti G, De Gasperi A, Vincent JL, Piagnerelli M. Early alterations of red blood cell rheology in critically ill patients. *Critical Care Medicine* 2009;37:3041–3046
 26. Nemeth N, Berhes M, Kiss F, Hajdu E, Deak A, Molnar A, Szabo J, Fulesdi B. Early hemorheological changes in a porcine model of intravenously given *E. coli* induced fulminant sepsis. *Clinical Hemorheology and Microcirculation* 2015;61:479–496
 27. Bateman RM, Jagger JE, Sharpe MD, Ellsworth ML, Mehta S, Ellis CG. Erythrocyte deformability is a nitric oxide-mediated factor in decreased capillary density during sepsis. *American Journal of Physiology-Heart and Circulatory Physiology* 2001;280(6):H2848–56
 28. Moutzouri AG, Skoutelis AT, Gogos CG, Missirlis YM, Athanassiou GM. Red blood cell deformability in patients with sepsis: A marker for prognosis and monitoring of severity. *Clinical Hemorheology and Microcirculation* 2007;36:291–299
 29. Donadello K, Piagnerelli M, Reggiori G, Gottin L, Scolletta S, Occhipinti G, et al. Reduced red blood cell deformability over time is associated with a poor outcome in septic patients. *Microvascular Research* 2015;101:8–14
 30. Machiedo GW. The Incidence of Decreased Red Blood Cell Deformability in Sepsis and the Association With Oxygen Free Radical Damage and Multiple-System Organ Failure. *Archives of Surgery* 1989;124(12):1386
 31. Langenfeld JE, Livingston DH, Machiedo GW. Red cell deformability is an early indicator of infection. *Surgery* 1991;110:398–403
 32. Sadaka F, O'Brien J, Prakash S. Red Cell Distribution Width and Outcome in Patients With Septic Shock. *Journal of Intensive Care Medicine* 2012;28(5):307–13
 33. Lorente L, Martín MM, Abreu-González P, Solé-Violán J, Ferreres J, Labarta L, et al. Red Blood Cell Distribution Width during the First Week Is Associated with Severity and Mortality in Septic Patients. *PLoS ONE* 2014;9(8):e105436
 34. Kim CH, Park JT, Kim EJ, Han JH, Han JS, Choi JY, et al. An increase in red blood cell distribution width from baseline predicts mortality in patients with severe sepsis or septic shock. *Critical Care* 2013;17(6):R282
 35. Jo YH, Kim K, Lee JH, Kang C, Kim T, Park H-M, et al. Red cell distribution width is a prognostic factor in severe sepsis and septic shock. *The American Journal of Emergency Medicine* 2013;31(3):545–8
 36. Kumar S, Jandial A, Bhalla A, Sharma N, Varma N, Varma S. Elevated red cell distribution width as a prognostic marker in severe sepsis: A prospective observational study. *Indian Journal of Critical Care Medicine* 2017;21(9):552
 37. Debari V, Mahmood N, Mathew J, Kang B, Khan M. Broadening of the red blood cell distribution width is associated with increased severity of illness in patients with sepsis. *International Journal of Critical Illness and Injury Science* 2014;4(4):278
 38. Piagnerelli M, Boudjeltia KZ, Brohee D, Piro P, Carlier E, Vincent J-L, et al. Alterations of red blood cell shape and sialic acid membrane content in septic patients. *Critical Care Medicine* 2003;31(8):2156–62
 39. Piagnerelli M, Vincent J-L, Boudjeltia KZ, Brohee D, Vanhaeverbeek M. Modifications of Red Blood Cell Shape and Glycoproteins Membrane Content in Septic Patients. *Advances in Experimental Medicine and Biology Oxygen Transport To Tissue XXIII* 2003;109–14
 40. Qadri SM, Donkor DA, Nazy I, Branch DR, Sheffield WP. Bacterial neuraminidase-mediated erythrocyte desialylation provokes cell surface aminophospholipid exposure. *European Journal of Haematology* 2018;100(5):502–10
 41. Rogers ME, Williams DT, Niththyananthan R, Rampling MW, Heslop KE, Johnston DG. Decrease in erythrocyte glycophorin sialic acid content is associated with increased erythrocyte aggregation in human diabetes. *Clinical Science* 1992;82(3):309–13

42. Venerando B, Fiorilli A, Croci G, Tringali C, Goi G, Mazzanti L et al. Acidic and neutral sialidase in the erythrocyte membrane of type 2 diabetic patients. *Blood* 2002;99:1064-1070
43. Moutzouri AG, Athanassiou GA, Dimitropoulou D, Skoutelis AT, Gogos CA. Severe sepsis and diabetes mellitus have additive effects on red blood cell deformability. *Journal of Infection* 2008;57(2):147-51
44. Milligan TW, Baker CJ, Straus DC, Mattingly SJ. Association of elevated levels of extracellular neuraminidase with clinical isolates of type III group B streptococci. *Infection and Immunity* 1978;21(3):738-746
45. Mattingly SJ, Milligan TW, Pierpont AA, Straus DC. Extracellular neuraminidase production by clinical isolates of group B streptococci from infected neonates. *Journal of Clinical Microbiology* 1980;12(4):633-635
46. Liukkonen J, Haataja S, Tikkanen K, Kelm S, Finne J. Identification of N-acetylneuraminyl alpha 2->3 poly-N-acetyl-lactosamine glycans as the receptors of sialic acid-binding *Streptococcus suis* strains. *The Journal of Biological Chemistry* 1992;267(29):21105-11
47. Parkkinen J, Rogers GN, Korhonen T, Dahr W, Finne J. Identification of the O-linked sialyloligosaccharides of glycophorin A as the erythrocyte receptors for S-fimbriated *Escherichia coli*. *Infection and Immunity* 1986;54(1):37-42
48. Qadri SM, Donkor DA, Bhakta V, Eltringham-Smith LJ, Dwivedi DJ, Moore JC, et al. Phosphatidylserine externalization and procoagulant activation of erythrocytes induced by *Pseudomonas aeruginosa* virulence factor pyocyanin. *Journal of Cellular and Molecular Medicine* 2016;20(4):710-20
49. Piagnerelli M, Boudjeltia KZ, Rapotec A, Richard T, Brohé D, Babar S, et al. Neuraminidase alters red blood cells in sepsis. *Critical Care Medicine* 2009;37(4):1244-50
50. Gut H, King SJ, Walsh MA. Structural and functional studies of *Streptococcus pneumoniae* neuraminidase B: An intramolecular trans-sialidase. *FEBS Letters* 2008;582(23-24):3348-52
51. Durocher JR, Payne RC, Conrad ME. Role of sialic acid in erythrocyte survival. *Blood* 1975;45:11-20
52. Waugh RE. Effects of 2,3-diphosphoglycerate on the mechanical properties of erythrocyte membrane. *Blood* 1986;68(1):231-8
53. Suzuki Y, Nakajima T, Shiga T, Maeda N. Influence of 2,3-diphosphoglycerate on the deformability of human erythrocytes. *Biochimica et Biophysica Acta (BBA) – Biomembranes* 1990;1029(1):85-90
54. Lizcano A, Secundino I, Döhrmann S, Corriden R, Rohena C, Diaz S, et al. Erythrocyte sialoglycoproteins engage Siglec-9 on neutrophils to suppress activation. *Blood* 2017;129(23):3100-10
55. Dinkla S, Eijk LTV, Fuchs B, Schiller J, Joosten I, Brock R, et al. Inflammation-associated changes in lipid composition and the organization of the erythrocyte membrane. *BBA Clinical* 2016;5:186-92
56. Qadri SM, Bissinger R, Solh Z, Oldenburg P-A. Eryptosis in health and disease: A paradigm shift towards understanding the (patho)physiological implications of programmed cell death of erythrocytes. *Blood Reviews* 2017;31(6):349-61
57. Velásquez FC, Maté S, Bakás L, Herlax V. Induction of eryptosis by low concentrations of *E. coli* alpha-hemolysin. *Biochimica et Biophysica Acta (BBA) – Biomembranes* 2015;1848(11):2779-88
58. Föller M, Shumilina E, Lam R, Mohamed W, Kasinathan R, Huber S, et al. Induction of Suicidal Erythrocyte Death by Listeriolysin from *Listeria monocytogenes*. *Cellular Physiology and Biochemistry* 2007;20(6):1051-60
59. Föller M, Biswas R, Mahmud H, Akel A, Shumilina E, Wieder T, et al. Effect of peptidoglycans on erythrocyte survival. *International Journal of Medical Microbiology* 2009;299:75-85
60. Wang K, Mahmud H, Föller M, Biswas R, Lang KS, Bohn E, et al. Lipopeptides in the triggering of erythrocyte cell membrane scrambling. *Cellular Physiology and Biochemistry* 2008;22:381-6
61. Abed M, Towhid ST, Mia S, Pakladok T, Alesutan I, Borst O et al. Sphingomyelinase-induced adhesion of eryptotic erythrocytes to endothelial cells. *American Journal of Physiology-Cell Physiology* 2012;303:C991-C999
62. Abed M, Towhid ST, Pakladok T, Alesutan I, Gotz F, Gulbins E, et al. Effect of bacterial peptidoglycan on erythrocyte death and adhesion to endothelial cells. *International Journal of Medical Microbiology* 2013;303:182-9
63. Bogdanova A, Makhro A, Wang J, Lipp P, Kaestner L. Calcium in red blood cells-A perilous balance. *International Journal of Molecular Sciences* 2013;14:9848-9872
64. Todd JC, Mollitt DL. Effect of sepsis on

- erythrocyte intracellular calcium homeostasis. *Critical Care Medicine* 1995;23:459–465
65. Desai TK, Carlson RW, Geheb MA. Prevalence and clinical implications of hypocalcemia in acutely ill patients in a medical intensive care setting. *The American Journal of Medicine* 1988;84(2):209–14
 66. Snyder LM, Fortier NL, Trainor J, Jacobs J, Leb L, Lubin B, et al. Effect of hydrogen peroxide exposure on normal human erythrocyte deformability, morphology, surface characteristics, and spectrin-hemoglobin cross-linking. *Journal of Clinical Investigation* 1985;76(5):1971–7
 67. Jain SK, Mohandas N, Clark MR, Shohet SB. The effect of malonyldialdehyde, a product of lipid peroxidation, on the deformability, dehydration and ⁵¹Cr-survival of erythrocytes. *British Journal of Haematology* 1983;53(2):247–55
 68. Kempe DS, Akel A, Lang PA, Hermle T, Biswas R, Muresanu J, et al. Suicidal erythrocyte death in sepsis. *Journal of Molecular Medicine* 2007;85:273–81
 69. Saldanha C, Silva-Herdade A. Erythrocyte Nitric Oxide. *Novel Prospects in Oxidative and Nitrosative Stress* Pinar Atukeren, IntechOpen, DOI: 10.5772/intechopen.75931
 70. Bordin L, Brunati AM, Donella-Deana A, Baggio B, Toninello A, Clari G. Band 3 is an anchor protein and a target for SHP-2 tyrosine phosphatase in human erythrocytes. *Blood* 2002;100:276–278
 71. Bordin L, Fiore C, Bragadin M, Brunati AM, Clari G. Regulation of membrane band 3 Tyr-phosphorylation by proteolysis of p72Syk and possible involvement in senescence process. *Acta Biochimica et Biophysica Sinica* 2009;41(10):846–51
 72. Condon MR, Feketova E, Machiedo GW, Deitch EA, Spolarics Z. Augmented erythrocyte band-3 phosphorylation in septic mice. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* 2007;1772(5):580–6
 73. Lin X, Rogers S, Timm D, Angelo A, Jaya P, Melanie E et al. Sepsis Induced Red Cell Dysfunction (SiRD): Physiology and Mechanisms. *Blood* 2017;130(1):3469
 74. Spolarics Z, Condon MR, Siddiqi M, Machiedo GW, Deitch EA. Red blood cell dysfunction in septic glucose-6-phosphate dehydrogenase-deficient mice. *American Journal of Physiology-Heart and Circulatory Physiology* 2004;286(6)
 75. Zipser Y, Piade A, Barbul A, Korenstein R, Kosower NS. Ca²⁺ promotes erythrocyte band 3 tyrosine phosphorylation via dissociation of phosphotyrosine phosphatase from band 3. *Biochemical Journal* 2002;368(1):137–44
 76. Mallozzi C, Di Stasi AM, Minetti M. Peroxynitrite modulates tyrosine-dependent signal transduction pathway of human erythrocyte band 3. *FASEB J* 1997;11:1281–90
 77. Sega MF, Chu H, Christian J, Low PS. Interaction of deoxyhemoglobin with the cytoplasmic domain of murine erythrocyte band 3. *Biochemistry* 2012;51:3264–72
 78. Galli F, Rossi R, Simplicio PD, Floridi A, Canestrari F. Protein Thiols and Glutathione Influence the Nitric Oxide-Dependent Regulation of the Red Blood Cell Metabolism. *Nitric Oxide* 2002;6(2):186–99
 79. Clancy RM, Levartovsky D, Leszczynska-Piziak J, Yegudin J, Abramson SB. Nitric oxide reacts with intracellular glutathione and activates the hexose monophosphate shunt in human neutrophils: evidence for S-nitrosoglutathione as a bioactive intermediary. *Proceedings of the National Academy of Sciences* 1994;91(9):3680–4
 80. Jagger JE, Bateman RM, Ellsworth ML, Ellis CG. Role of erythrocyte in regulating local O₂ delivery mediated by hemoglobin oxygenation. *American Journal of Physiology-Heart and Circulatory Physiology* 2001;280(6)
 81. Bateman RM, Sharpe MD, Jagger JE, Ellis CG. Sepsis impairs microvascular autoregulation and delays capillary response within hypoxic capillaries. *Critical Care* 2015;19(1)
 82. Lopes de Almeida JP, Freitas-Santos T, Saldanha C. Evidence that the degree of band 3 phosphorylation modulates human erythrocytes nitric oxide efflux-in vitro model of fibrinogenemia. *Clinical Hemorheology and Microcirculation* 2011;49:407–416
 83. Silva-Herdade AS, Freitas T, Almeida JP, Saldanha C. Fibrinogen signaling in erythrocyte nitric oxide mobilization in presence of PI3-K and adenylyl cyclase inhibitors. *European Journal of Biomedical and Pharmaceutical Sciences* 2016;3:28–34
 84. Piagnerelli M., Cotton F, van Nuffelen M, Vincent JL, Gulbis B. Modifications in erythrocyte membrane protein content are not responsible for the alterations in rheology seen in sepsis. *Shock* 2012;37:17–21
 85. Chillar RK, Slawskyand P, Desforjes JF. Red Cell 2,3-Diphosphoglycerate and Adenosine Triphosphate in Patients with Shock. *British Journal of Haematology*. 1971;21(2):183–8.
 86. Watkins GM, Rabelo A, Plzak LF, Shel-

- don GF. The Left Shifted Oxyhemoglobin Curve in Sepsis. *Annals of Surgery* 1974;180(2):213–20
87. Myburgh JA, Webb RK, Worthley LG. The P50 is reduced in critically ill patients. *Intensive Care Medicine* 1991;17(6):355–8
88. Ibrahim EEDS, Mclellan SA, Walsh TS. Red blood cell 2,3-diphosphoglycerate concentration and in vivo P50 during early critical illness*. *Critical Care Medicine* 2005;33(10):2247–52
89. Schindler M, Koppel DE, Sheetz MP. Modulation of membrane protein lateral mobility by polyphosphates and polyamines. *Proceedings of the National Academy of Sciences* 1980;77(3):1457–61
90. Sheetz MP, Casaly J. Phosphate metabolite regulation of spectrin interactions. *Scandinavian Journal of Clinical and Laboratory Investigation* 1981;41(sup156):117–22
91. Muller, Becker, Kranzlin, Schachinger, Huber, Nylen, et al. Disordered calcium homeostasis of sepsis: association with calcitonin precursors. *European Journal of Clinical Investigation* 2000;30(9):823–31
92. Ruef P, Ehrhard M, Frommhold D, Koch L, Fritzsche B, Poeschl J. Lipid decreases human erythrocytes deformability by increasing intracellular Ca^{2+} : Effects of verapamil, staurosporine and the rho-kinase inhibitor Y-27632. *Clinical Hemorheology and Microcirculation* 2011;49:315–322
93. Deaciuc I, Spitzer J. Calcium content in liver and heart and its intracellular distribution in liver during endotoxemia and sepsis in rats. *Cell Calcium* 1987;8(5):365–76
94. Celes MRN, Malvestio LM, Suadicaní SO, Prado CM, Figueiredo MJ, et al. Disruption of Calcium Homeostasis in Cardiomyocytes Underlies Cardiac Structural and Functional Changes in Severe Sepsis. *PLOS ONE* 2013;8(7):e68809
95. Zaloga GP, Washburn D, Black KW, Prielipp R. Human sepsis increases lymphocyte intracellular calcium. *Critical Care Medicine* 1993;21(2):196–202
96. Smith BD, Celle PLL, Siefring GE, Lowe-Krentz L, Lorand L. Effects of the calcium-mediated enzymatic cross-linking of membrane proteins on cellular deformability. *The Journal of Membrane Biology* 1981;61(2):75–80
97. Lau YT, Hsieh CC, Liu MS, Hwang TL, Chen MF, Cheng HS. Erythrocyte Ca^{2+} pump is defective during sepsis. *Circulatory Shock* 1994;44:121–125
98. Todd JC, Mollitt DL. Leukocyte Modulation Inhibits Endotoxin-Induced Disruption Of Intracellular Calcium Homeostasis. *The Journal of Trauma: Injury, Infection, and Critical Care* 1994;37(6):1018–9
99. Helms CC, Gladwin MT, Kim-Shapiro DB. Erythrocytes and Vascular Function: Oxygen and Nitric Oxide. *Frontiers in Physiology* 2018;9
100. Hoffman JF. ATP compartmentation in human erythrocytes. *Current Opinion in Hematology* 1997;4(2):112–5
101. Bergfeld GR, Forrester T. Release of ATP from human erythrocytes in response to a brief period of hypoxia and hypercapnia. *Cardiovascular Research* 1992;26(1):40–7
102. Locovei S, Bao L, Dahl G. Pannexin 1 in erythrocytes: Function without a gap. *Proceedings of the National Academy of Sciences* 2006;103(20):7655–9
103. Ellsworth ML, Forrester T, Ellis CG, Dietrich HH. The erythrocyte as a regulator of vascular tone. *American Journal of Physiology-Heart and Circulatory Physiology* 1995;269(6)
104. Mathie R, Ralevic V, Alexander B, Burnstock G. Nitric oxide is the mediator of ATP-induced dilatation of the rabbit hepatic arterial vascular bed. *British Journal of Pharmacology* 1991;103(2):1602–6
105. Busse R, Ogilvie A, Pohl U. Vasomotor activity of diadenosine triphosphate and diadenosine tetraphosphate in isolated arteries. *American Journal of Physiology-Heart and Circulatory Physiology* 1988;254(5)
106. Sprague RS, Ellsworth ML, Stephenson AH, Lonigro AJ. ATP: the red blood cell link to NO and local control of the pulmonary circulation. *American Journal of Physiology-Heart and Circulatory Physiology* 1996;271(6)
107. Sprague RS, Stephenson AH, Dimmitt RA, Weintraub NL, Branch CA, Mcmurdo L, et al. Effect of L-NAME on pressure-flow relationships in isolated rabbit lungs: role of red blood cells. *American Journal of Physiology-Heart and Circulatory Physiology* 1995;269(6)
108. Serroukh Y, Djebara S, Lelubre C, Boudjeltia KZ, Biston P, Piagnerelli M. Alterations of the Erythrocyte Membrane during Sepsis. *Critical Care Research and Practice* 2012;2012:1–7
109. Forsyth AM, Wan J, Owrutsky PD, Abkarian M, Stone HA. Multiscale approach to link red blood cell dynamics, shear viscosity, and ATP release. *Proceedings of the National Academy of Sciences* 2011;108(27):10986–91
110. Rozier MD, Zata VJ, Ellsworth ML. Lactate

- interferes with ATP release from red blood cells. *American Journal of Physiology-Heart and Circulatory Physiology* 2007;292(6)
111. Greve A-S, Skals M, Fagerberg SK, Tonnus W, Ellermann-Eriksen S, Evans RJ, et al. P2X1, P2X4, and P2X7 Receptor Knock Out Mice Expose Differential Outcome of Sepsis Induced by α -Haemolysin Producing *Escherichia coli*. *Frontiers in Cellular and Infection Microbiology* 2017;7
 112. Skals M, Jorgensen NR, Leipziger J, Praetorius HA. Hemolysin from *Escherichia coli* uses endogenous amplification through P2X receptor activation to induce hemolysis. *Proceedings of the National Academy of Sciences* 2009;106(10):4030–5
 113. Skals M, Leipziger J, Praetorius HA. Hemolysis induced by α -toxin from *Staphylococcus aureus* requires P2X receptor activation. *Pflügers Archiv - European Journal of Physiology* 2011;462(5):669–79
 114. Nagahama M, Seike S, Shirai H, Takagishi T, Kobayashi K, Takehara M, et al. Role of P2X7 receptor in *Clostridium perfringens* beta-toxin-mediated cellular injury. *Biochimica et Biophysica Acta (BBA) - General Subjects* 2015;1850(11):2159–67
 115. Prauchner CA. Oxidative stress in sepsis: Pathophysiological implications justifying antioxidant co-therapy. *Burns* 2017;43(3):471–85
 116. Marik PE, Khangoora V, Rivera R, Hooper MH, Catravas J. Hydrocortisone, Vitamin C, and Thiamine for the Treatment of Severe Sepsis and Septic Shock. *Chest* 2017;151(6):1229–38
 117. Goode HF, Cowley HC, Walker BE, Howdle PD, Webster NR. Decreased antioxidant status and increased lipid peroxidation in patients with septic shock and secondary organ dysfunction. *Critical Care Medicine* 1995;23(4):646–51
 118. Metnitz PGH, Bartens C, Fischer M, Friedrich P, Steltzer H, Druml W. Antioxidant status in patients with acute respiratory distress syndrome. *Intensive Care Medicine* 1999;25(2):180–5
 119. Pascual C, Karzai W, Meier-Hellmann A, Oberhoffer M, Horn A, Bredle D, et al. Total plasma antioxidant capacity is not always decreased in sepsis. *Critical Care Medicine* 1998;26(4):705–9
 120. Richard C, Lemonnier F, Thibault M, Couturier M, Auzepy P. Vitamin E deficiency and lipoperoxidation during adult respiratory distress syndrome. *Critical Care Medicine* 1990;18(1):4–9
 121. Luca L, Rogobete AF, Bedreag OH. Oxidative Stress and Antioxidant Therapy in Critically Ill Polytrauma Patients with Severe Head Injury. *The Journal of Critical Care Medicine* 2015;1(3):83–91
 122. Dewas C, Dang PM-C, Gougerot-Pocidalo M-A, El-Benna J. TNF- Induces Phosphorylation of p47phox in Human Neutrophils: Partial Phosphorylation of p47phox Is a Common Event of Priming of Human Neutrophils by TNF- and Granulocyte-Macrophage Colony-Stimulating Factor. *The Journal of Immunology* 2003;171(8):4392–8
 123. Faivre B, Menu P, Labrude P, Vigneron C. Hemoglobin Autooxidation/Oxidation Mechanisms and Methemoglobin Prevention or Reduction Processes in the Bloodstream Literature review and outline of autooxidation reaction. *Artificial Cells, Blood Substitutes, and Biotechnology* 1998;26(1):17–26
 124. Baskurt OK, Temiz A, Meiselman HJ. Effect of Superoxide Anions on Red Blood Cell Rheologic Properties. *Free Radical Biology and Medicine* 1998;24(1):102–10
 125. Srouf MA, Bilto YY, Juma M, Irhimeh MR. Exposure of human erythrocytes to oxygen radicals causes loss of deformability, increased osmotic fragility, lipid peroxidation and protein degradation. *Clinical Hemorheology and Microcirculation* 2000;23:13–21
 126. Snyder LM, Fortier NL, Trainor J, Jacobs J, Leb L, Lubin B, et al. Effect of hydrogen peroxide exposure on normal human erythrocyte deformability, morphology, surface characteristics, and spectrin-hemoglobin cross-linking. *Journal of Clinical Investigation* 1985;76(5):1971–7
 127. Weiss SJ. The role of superoxide in the destruction of erythrocyte targets by human neutrophils. *Journal of Biological Chemistry* 1980;255:9912–9917
 128. Powell RJ, Machiedo GW, Rush BF Jr., Dikdan G. Effect of α -tocopherol on red cell deformability and survival in sepsis. *Current Surg* 1989;46:380–382
 129. Uyesaka N, Hasegawa S, Ishioka N, Ishioka R, Shio H, Schechter AN. Effects of superoxide anions on red cell deformability and membrane proteins. *Biorheology* 1992;29:217–229
 130. Jain SK, Ross JD, Levy GJ, Little RL, Duett J. The accumulation of malonyldialdehyde, an end product of membrane lipid peroxidation, can cause potassium leak in normal and sickle red blood cells. *Biochemical Medicine and Metabolic Biology* 1989;42(1):60–5
 131. Oliveira YPAD, Pontes-De-Carvalho LC,

- Couto RD, Noronha-Dutra AA. Oxidative stress in sepsis. Possible production of free radicals through an erythrocyte-mediated positive feedback mechanism. *The Brazilian Journal of Infectious Diseases* 2017;21(1):19–26
132. McKenney J, Valeri CR, Mohandas N, Fortier N, Giorgio A, Snyder LM. Decreased in vivo survival of hydrogen peroxide-damaged baboon red blood cells. *Blood* 1990;76:206–211
133. Butterfield D, Sun B, Bellary S, Arden WA, Anderson KW. Effect of endotoxin on lipid order and motion in erythrocyte membranes. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* 1994;1225(2):231–4
134. Yerer MB, Yapislar H, Aydogan S, Yalcin O, Baskurt O. Lipid peroxidation and deformability of red blood cells in experimental sepsis in rats: The protective effects of melatonin. *Clinical Hemorheology and Microcirculation* 2004;30(2):77–82
135. Snyder LM, Fortier NL, Trainor J, Jacobs J, Leb L, Lubin B, et al. Effect of hydrogen peroxide exposure on normal human erythrocyte deformability, morphology, surface characteristics, and spectrin-hemoglobin cross-linking. *Journal of Clinical Investigation* 1985;76(5):1975–7
136. Mal A, Chatterjee IB. Mechanism of autoxidation of oxyhaemoglobin. *Journal of Biochemistry* 1991;16(1-2):55–70
137. Bunn, HF, Forget BG. in: *Hemoglobin. Molecular and General Clinical Aspects*. Philadelphia:Saunders 1986:634–662
138. Kikugawa K, Sasahara T, Sasaki T, Kurechi T. Factors influencing the autoxidation of hemoglobin A. *Chemical & Pharmaceutical Bulletin* 1981;29(5):1382–9
139. Goldman D, Bateman RM, Ellis CG. Effect of sepsis on skeletal muscle oxygen consumption and tissue oxygenation: interpreting capillary oxygen transport data using a mathematical model. *American Journal of Physiology-Heart and Circulatory Physiology* 2004;287(6)
140. Spies CD, Reinhart K, Witt I, Meier-Hellmann A, Hannemann L, Bredle DL, et al. Influence of N-acetylcysteine on indirect indicators of tissue oxygenation in septic shock patients: Results from a prospective, randomized, double-blind study. *Critical Care Medicine* 1994;22:1738–46
141. Spapen H, Zhang H, Demanet C, Vleminckx W, Vincent JL, Huyghens L. Does N-acetyl-L-cysteine influence cytokine response during early human septic shock? *Chest* 1998;113:1616–24
142. Rank N, Michel C, Haertel C, Lenhart A, Welte M, Meier-Hellmann A, et al. N-acetylcysteine increases liver blood flow and improves liver function in septic shock patients: Results of a prospective, randomized, double-blind study. *Critical Care Medicine* 2000;28:3799–807
143. Ortolani O, Conte A, De Gaudio AR, Moraldi E, Cantini Q, Novelli G. The effect of glutathione and N-acetylcysteine on lipoperoxidative damage in patients with early septic shock. *American Journal of Respiratory and Critical Care Medicine* 2000;161:1907–11
144. Zimmermann T, Albrecht S, Kühne H, Vogelsang U, Grützmann R, Kopprasch S. Selenium administration in patients with sepsis syndrome. A prospective randomized study. *Medizinische Klinik* 1997;92(3):3–4
145. Angstwurm MW, Schottdorf J, Schopohl J, Gaertner R. Selenium replacement in patients with severe systemic inflammatory response syndrome improves clinical outcome. *Critical Care Medicine* 1999;27:1807–13
146. Mahmud H, Qadri SM, Föller M, Lang F. Inhibition of suicidal erythrocyte death by vitamin C. *Nutrition* 2010;26(6):671–6
147. Gadek JE, DeMichele SJ, Karlstad MD, Pacht ER, Donahoe M, Albertson TE, et al. Effect of enteral feeding with eicosapentaenoic acid, gammalinolenic acid, and antioxidants in patients with acute respiratory distress syndrome. *Critical Care Medicine* 1999;27:1409–20
148. Pontes-Arruda A, Albuquerque Aragão AM, Albuquerque JD. Effects of enteral feeding with eicosapentaenoic acid, gamma-linolenic acid, and antioxidants in mechanically ventilated patients with severe sepsis and septic shock. *Critical Care Medicine* 2006;34:2325–33
149. Hollenberg SM, Cunnion RE, Zimmerberg J. Nitric oxide synthase inhibition reverses arteriolar hyporesponsiveness to catecholamines in septic rats. *American Journal of Physiology* 1993;264:H660–3
150. Hollenberg SM, Piotrowski MJ, Parrillo JE. Nitric oxide synthase inhibition reverses arteriolar hyporesponsiveness to endothelin-1 in septic rats. *American Journal of Physiology* 1997;272:R969–74
151. Frost MT, Wang Q, Moncada S, Singer M. Hypoxia accelerates nitric oxide-dependent inhibition of mitochondrial complex I in activated macrophages. *American journal of physiology. Regulatory, integrative and*

- comparative physiology 2005;288:R394–400
152. Scott JA, Mehta S, Duggan M, Bihari A, McCormack DG. Functional inhibition of constitutive nitric oxide synthase in a rat model of sepsis. *American Journal of Respiratory and Critical Care Medicine* 2002;165:1426–32
 153. Starzyk D, Korbut R, Gryglewski RJ. The role of nitric oxide in regulation of deformability of red blood cells in acute phase of endotoxaemia in rats. *Journal of Physiology and Pharmacology* 1997;48:731–735
 154. Vincent J-L, Zhang H, Szabo C, Preiser J-C. Effects of Nitric Oxide in Septic Shock. *American Journal of Respiratory and Critical Care Medicine* 2000;161(6):1781–5
 155. Fernandes D, Assreuy J. Nitric oxide and vascular reactivity in sepsis. *Shock* 2008;30 Suppl 1:10-3
 156. Winkler MS, Kluge S, Holzmann M, Moritz E, Robbe L, Bauer A, et al. Markers of nitric oxide are associated with sepsis severity: an observational study. *Critical Care* 2017;21(1)
 157. Knowles RG, Moncada S. Nitric oxide synthases in mammals. *Biochemical Journal*. 1994;298(2):249–58
 158. Takatani Y, Ono K, Suzuki H, Inaba M, Sawada M, Matsuda N. Inducible nitric oxide synthase during the late phase of sepsis is associated with hypothermia and immune cell migration. *Laboratory Investigation* 2018;98(5):629–39
 159. Caironi P, Masson S, Mauri T, Bottazzi B, Leone R, Magnoli M, et al. Pentraxin 3 in patients with severe sepsis or shock: the ALBIOS trial. *European Journal of Clinical Investigation* 2016;47(1):73–83
 160. Carrizzo A, Lenzi P, Procaccini C, Damato A, Biagioni F, Ambrosio M et al. Pentraxin 3 Induces Vascular Endothelial Dysfunction Through a P-selectin/Matrix Metalloproteinase-1 Pathway. *Circulation* 2015;131:1495–505
 161. Lange M, Hamahata A, Traber DL, Nakano Y, Traber LD, Enkhbaatar P: Specific inhibition of nitric oxide synthases at different time points in a murine model of pulmonary sepsis. *Biochemical and Biophysical Research Communications* 2011;404(3):877–881
 162. Nardi GM, Scheschowitsch K, Ammar D, de Oliveira SK, Arruda TB, Assreuy J: Neuronal nitric oxide synthase and its interaction with soluble guanylate cyclase is a key factor for the vascular dysfunction of experimental sepsis. *Critical Care Medicine* 2014;42(6):e391–e400
 163. Gusarov I, Shatalin K, Starodubtseva M, Nudler E. Endogenous Nitric Oxide Protects Bacteria Against a Wide Spectrum of Antibiotics. *Science* 2009;325(5946):1380–4
 164. Gusarov I, Nudler E. NO-mediated cytoprotection: Instant adaptation to oxidative stress in bacteria. *Proceedings of the National Academy of Sciences* 2005;102(39):13855–60
 165. Nagababu E, Ramasamy S, Abernethy DR, Rifkind JM. Active nitric oxide produced in the red cell under hypoxic conditions by deoxyhemoglobin-mediated nitrite reduction. *The Journal of Biological Chemistry* 2003;278:46349–46356
 166. Jubelin B, Gierman J. Erythrocytes may synthesize their own nitric oxide. *American Journal of Hypertension* 1996;9(12):1214–9
 167. Carvalho A, Martins-Silva J, Saldanha C. Amperometric measurements of nitric oxide in erythrocytes. *Biosensors & Bioelectronics* 2004;20:505–508
 168. Kleinbongard P. Red blood cells express a functional endothelial nitric oxide synthase. *Blood* 2006 ;107(7):2943–51
 169. Silva-Herdade AS, Andolina G, Faggio C, Calado Â, Saldanha C. Erythrocyte deformability – A partner of the inflammatory response. *Microvascular Research* 2016;107:34–38
 170. Korbut R, Gryglewski RJ. The effect of prostacyclin and nitric oxide on deformability of red blood cells in septic shock in rats. *Journal of Physiology and Pharmacology* 1996;47:591–599
 171. Battinelli E, Willoughby SR, Foxall T, Valeri CR, Loscalzo J. Induction of platelet formation from megakaryocytoid cells by nitric oxide. *Proceedings of the National Academy of Sciences* 2001;98(25):14458–63
 172. Shiga T, Maeda N, Kon K. Erythrocyte rheology. *Critical Reviews in Oncology/Hematology* 1990;10(1):9–48
 173. Falkmarken NHD, Arihan O, Iskit AB. Comparison of endothelin and nitric oxide synthase blockerson hemorheological parameters in endotoxemic rats. *Turkish Journal Of Medical Sciences* 2017;47:1045–52
 174. Petrov V, Amery A, Lijnen P. Role of cyclic GMP in atrial-natriuretic-peptide stimulation of erythrocyte Na⁺/H exchange. *European Journal of Biochemistry* 1994;221(1):195–9
 175. Petrov V, Lijnen P. Regulation of human erythrocyte Na⁺/H exchange by soluble and particulate guanylate cyclase. *Ameri-*

- can Journal of Physiology-Cell Physiology 1996;271(5)
176. Caramelo C, Riesco A, Outeirino J, Millas I, Blum G, Monzu B, et al. Effects of Nitric Oxide on Red Blood Cells: Changes in Erythrocyte Resistance to Hypotonic Hemolysis and Potassium Efflux by Experimental Maneuvers That Decrease Nitric Oxide. *Biochemical and Biophysical Research Communications* 1994;199(2):447–54
177. Adragna N, Lauf P. Role of Nitrite, a Nitric Oxide Derivative, in K-Cl Cotransport Activation of Low-Potassium Sheep Red Blood Cells. *Journal of Membrane Biology* 1998;166(3):157–67
178. Grzelak A. Peroxynitrite Activates K⁺-Cl⁻ Cotransport In Human Erythrocytes. *Cell Biology International* 2001;25(11):1163–5
179. Elias MDO, Lima WTD, Vannuchi YB, Marcourakis T, Silva ZLD, Trezena AG, et al. Nitric oxide modulates Na⁺, K⁺-ATPase activity through cyclic GMP pathway in proximal rat trachea. *European Journal of Pharmacology* 1999;367(2-3):307–14
180. Pernollet M-G, Lantoine F, Devynck M-A. Nitric Oxide Inhibits ATP-Dependent CA2 Uptake into Platelet Membrane Vesicles. *Biochemical and Biophysical Research Communications* 1996;222(3):780–5
181. Korbut R, Gryglewski RJ. Nitric oxide from polymorphonuclear leukocytes modulates red blood cell deformability in vitro. *European Journal of Pharmacology* 1993;234(1):17–22
182. Avontuur JAM, Nolthenius RPT, Bodegom JWV, Bruining HA. Prolonged inhibition of nitric oxide synthesis in severe septic shock. *Critical Care Medicine* 1998;26(4):660–7
183. Andresen M, Dougnac A, Diaz O, Hernandez G, Castillo L, Bugedo G, et al. Use of methylene blue in patients with refractory septic shock: impact on hemodynamics and gas exchange. *Journal of Critical Care* 1998;13:164–8
184. Preiser JC, Lejeune P, Roman A, Carlier E, De Backer D, Leeman M, et al. Methylene blue administration in septic shock: a clinical trial. *Critical Care Medicine* 1995;23:259–64
185. Watson D, Grover R, Anzueto A, Lorente J, Smithies M, Bellomo R, et al. Cardiovascular effects of the nitric oxide synthase inhibitor NG-methyl-L-arginine hydrochloride (546C88) in patients with septic shock: results of a randomized, double-blind, placebo-controlled multicenter study (study no. 144-002). *Critical Care Medicine* 2004;32:13–20
186. Bakker J, Grover R, McLuckie A, Holzapfel L, Andersson J, Lodato R, et al. Administration of the nitric oxide synthase inhibitor NG-methyl-L-arginine hydrochloride (546C88) by intravenous infusion for up to 72 hours can promote the resolution of shock in patients with severe sepsis: results of a randomized, double-blind, placebo-controlled multicenter study (study no. 144-002). *Critical Care Medicine* 2004;32:1–12
187. Lorenz E. TLR2 and TLR4 expression during bacterial infections. *Current Pharmaceutical Design* 2006;12:4185–93
188. Loughran PA, Lei Z, Xu L, Deng M, Billiar TR. Nitric Oxide in Sepsis and Hemorrhagic Shock: Beneficial or Detrimental? *Nitric Oxide* 2017;289–300.
189. Boerma EC, Koopmans M, Konijn A, Kaiferova K, Bakker AJ, van Roon EN, et al. Effects of nitroglycerin on sublingual microcirculatory blood flow in patients with severe sepsis/septic shock after a strict resuscitation protocol: a double-blind randomized placebo controlled trial. *Critical Care Medicine* 2010;38:93–100
190. Spronk PE, Ince C, Gardien MJ, Mathura KR, Oudemans-van Straaten HM, Zandstra DF. Nitroglycerin in septic shock after intravascular volume resuscitation. *Lancet* 2002;360:1395–6