Letter to the Editor

Humectants in tobacco products on the US Market

EDITORS – The recent publication in this Journal by Rainey and co-workers [Quantitative Analysis of Humectants in Tobacco Products Using Gas Chromatography (GC) with Simultaneous Mass Spectrometry (MSD) and Flame Ionization Detection (FID), Beitr. Tabakforsch. Int. 25 (2013) 576–585] raises serious questions about the sampling, analyses, and reporting of results on commercial products currently on sale in a given country or region. Many articles that have appeared in this Journal over the years have dealt with experimental products and materials. While such products and materials may have been very similar to those in commerce, they were not identified as commercial samples. They may have been one-of-a-kind samples designed to show a specific phenomenon. Thus, it would be impractical to ask that the authors use representative samples and only use well-established test methods to characterize experimental materials. The situation is different with an article that presents analytical data on a product taken from the marketplace. Unless the authors are privy (through employment by the manufacturer, for example) to the formulas and/or specifications for such products, any analytical data reported are suspect unless the authors have taken special precautions. These special precautions include but are not limited to making sure that they have representative samples of the particular brand-styles under study, the analytical methods are really applicable to the brand-styles collected, and the data are presented accurately.

Based on the information presented in the article, it appears that Dr. Rainey and her team fell short of reasonable expectations for an article describing the analyses of commercial products, especially when one of the coauthors is associated with a regulatory and taxation authority (US Alcohol and Tobacco Tax and Trade Bureau, aka TTB) that is not a disinterested party to the publication of the research. The three deficiencies in the research presented were:

1. There appears to have been no effort to obtain a representative sample of each brand style tested. Tobacco

products are inherently variable and how the products are sampled at the points of manufacture, storage, and sale can affect the results of analytical and physical testing and the validity of those results. This variability has been recognized in international standards such as ISO 8243:2013 (and earlier versions) for cigarettes, ISO 15592-1:2001 for fine-cut tobacco (known to most US consumers as roll-your-own or RYO tobacco), CORESTA Recommended Method No. 47 Cigars – Sampling (2000), and CORESTA Recommended Method No. 71 Smokeless Tobacco Products – Sampling (2011). If the authors were not able to adhere to these standards in their entirety, then they should have made reasonable efforts to source products from several distinct geographical locations. This should not have been a problem since some of the authors were based in Indianapolis, IN, and the others were based in Beltsville, MD. The authors statement, “Three roll-your-own (RYO), thirteen cigar, eleven cigarette, ten moist snuff, and seven hookah tobacco products were purchased from tobacconists in Laurel, Maryland,” shows that they appear to have ignored good sampling practices.

2. The analyses of commercial products by a regulatory agency could lead to the analytical techniques used becoming *de facto* governmentally-mandated methods. This, in itself, would not necessarily be bad, if the techniques used by the regulatory were more effective and efficient than those used by others. However, the chromatography conditions used by Rainey and her team were worse than those used and published by others. One should not need gas chromatography – mass spectrometry (GC-MS) to determine tobacco humectants when GC with a Megabore column did the job way back in 1987 (see Figure 1, which was taken from FRANK and LIN, Tob. Chem. Res. Conf., 1987, 41, p. 36, abstr. 47, also see <http://legacy.library.ucsf.edu/tid/zwj19j00>). Figure 1 shows baseline separation between triethylene glycol and glycerin in the spiked extract of a tobacco matrix. The chromatogram generated in 1987 with 1987 instrumentation and columns showed faster and better chromatography than the authors obtained in 2013.

Clearly, the authors erred when they stated in the sentence of their summary, “Utilizing MSD greatly improves the reliability of quantitative results because

compensation for inadequate chromatographic resolution can be accomplished with mass selectivity in detection.” This error needs to be corrected since the use of the MSD needlessly drives up the cost of doing the determinations. All need to remember that the TTB has issued test methods for the analyses of tobacco products (http://www.ttb.gov/ssd/methods_tobacco.shtml). Is the methodology described by the authors now a *de facto* TTB method?

3. The analytical results shown in the authors’ Table 2, are of limited value because the authors chose not to provide at least generic descriptions of the products (RYO: type of blend, cut width; cigars: type of blend, dimensions, characterizing flavor; cigarettes: type of blend, dimensions, menthol/nonmenthol/additive-free; moist snuff: cut type, loose or pouched, straight, wintergreen, other characterizing flavor; hookah: characterizing flavor). Also, there was no determination of moisture (oven volatiles) or water in the products. For example, on an absolute basis is $1.59 \pm 0.01\%$ PG in Cigar Sample 4 the same as $1.56 \pm 0.03\%$ PG in Moist Snuff Sample 39? The authors should have determined water on

their samples using ISO 6488:2004 and then corrected their humectant values for water content of the products. Had they done this, not only would their data have been more meaningful, but they would have also found that Samples 38 to 44 were likely not moist snuff as indicated in Table 2, but were likely hookah tobacco. Perhaps the authors will issue a corrigendum that will clarify the identity of the samples, provide the water or oven-volatiles content of the samples, and provide the proper identification of Samples 38 to 44 in Table 2.

John H. Lauterbach, Ph.D., DABT
Sole Member & Principal, Chemistry & Toxicology
Lauterbach & Associates, LLC (Georgia Office)
211 Old Club Court
Macon, GA 31210-4708, USA
E-mail: john@lauterbachandassociates.net

FIGURE 1
TYPICAL CHROMATOGRAM

Figure 1. Chromatogram from FRANK and LIN; 41st Tobacco Chemists’ Research Conference, 1987, 41, p. 36, abstr. 47.