

# Partial Differentiation on Normed Linear Spaces $\mathcal{R}^n$

Noboru Endou  
Gifu National College of Technology  
Japan

Yasunari Shidama  
Shinshu University  
Nagano, Japan

Keiichi Miyajima  
Ibaraki University  
Hitachi, Japan

**Summary.** In this article, we define the partial differentiation of functions of real variable and prove the linearity of this operator [18].

MML identifier: PDIFF\_1, version: 7.8.05 4.84.971

The notation and terminology used here are introduced in the following papers: [21], [24], [25], [5], [26], [7], [6], [15], [13], [3], [1], [20], [11], [22], [23], [14], [8], [2], [4], [27], [28], [16], [9], [19], [17], [12], and [10].

## 1. PRELIMINARIES

Let  $i, n$  be elements of  $\mathbb{N}$ . The functor  $\text{proj}(i, n)$  yielding a function from  $\mathcal{R}^n$  into  $\mathbb{R}$  is defined by:

(Def. 1) For every element  $x$  of  $\mathcal{R}^n$  holds  $(\text{proj}(i, n))(x) = x(i)$ .

Next we state two propositions:

- (1)  $\text{dom proj}(1, 1) = \mathcal{R}^1$  and  $\text{rng proj}(1, 1) = \mathbb{R}$  and for every element  $x$  of  $\mathbb{R}$  holds  $(\text{proj}(1, 1))(\langle x \rangle) = x$  and  $(\text{proj}(1, 1))^{-1}(x) = \langle x \rangle$ .
- (2)(i)  $(\text{proj}(1, 1))^{-1}$  is a function from  $\mathbb{R}$  into  $\mathcal{R}^1$ ,
- (ii)  $(\text{proj}(1, 1))^{-1}$  is one-to-one,
- (iii)  $\text{dom}((\text{proj}(1, 1))^{-1}) = \mathbb{R}$ ,
- (iv)  $\text{rng}((\text{proj}(1, 1))^{-1}) = \mathcal{R}^1$ , and

- (v) there exists a function  $g$  from  $\mathbb{R}$  into  $\mathcal{R}^1$  such that  $g$  is bijective and  $(\text{proj}(1, 1))^{-1} = g$ .

One can check that  $\text{proj}(1, 1)$  is bijective.

Let  $g$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$ . The functor  $\langle g \rangle$  yields a partial function from  $\mathcal{R}^1$  to  $\mathcal{R}^1$  and is defined as follows:

(Def. 2)  $\langle g \rangle = (\text{proj}(1, 1))^{-1} \cdot g \cdot \text{proj}(1, 1)$ .

Let  $n$  be an element of  $\mathbb{N}$  and let  $g$  be a partial function from  $\mathcal{R}^n$  to  $\mathbb{R}$ . The functor  $\langle g \rangle$  yielding a partial function from  $\mathcal{R}^n$  to  $\mathcal{R}^1$  is defined as follows:

(Def. 3)  $\langle g \rangle = (\text{proj}(1, 1))^{-1} \cdot g$ .

Let  $i, n$  be elements of  $\mathbb{N}$ . The functor  $\text{Proj}(i, n)$  yielding a function from  $\langle \mathcal{E}^n, \|\cdot\| \rangle$  into  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  is defined as follows:

(Def. 4) For every point  $x$  of  $\langle \mathcal{E}^n, \|\cdot\| \rangle$  holds  $(\text{Proj}(i, n))(x) = \langle (\text{proj}(i, n))(x) \rangle$ .

Let  $i$  be an element of  $\mathbb{N}$  and let  $x$  be a finite sequence of elements of  $\mathbb{R}$ . The functor  $\text{reproj}(i, x)$  yielding a function is defined as follows:

(Def. 5)  $\text{dom reproj}(i, x) = \mathbb{R}$  and for every element  $r$  of  $\mathbb{R}$  holds  $(\text{reproj}(i, x))(r) = \text{Replace}(x, i, r)$ .

Let  $n, i$  be elements of  $\mathbb{N}$  and let  $x$  be an element of  $\mathcal{R}^n$ . Then  $\text{reproj}(i, x)$  is a function from  $\mathbb{R}$  into  $\mathcal{R}^n$ .

Let  $n, i$  be elements of  $\mathbb{N}$  and let  $x$  be a point of  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ . The functor  $\text{reproj}(i, x)$  yielding a function from  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  into  $\langle \mathcal{E}^n, \|\cdot\| \rangle$  is defined by the condition (Def. 6).

(Def. 6) Let  $r$  be an element of  $\langle \mathcal{E}^1, \|\cdot\| \rangle$ . Then there exists an element  $q$  of  $\mathbb{R}$  and there exists an element  $y$  of  $\mathcal{R}^n$  such that  $r = \langle q \rangle$  and  $y = x$  and  $(\text{reproj}(i, x))(r) = (\text{reproj}(i, y))(q)$ .

Let  $m, n$  be non empty elements of  $\mathbb{N}$ , let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ , and let  $x$  be an element of  $\mathcal{R}^m$ . We say that  $f$  is differentiable in  $x$  if and only if the condition (Def. 7) is satisfied.

(Def. 7) There exists a partial function  $g$  from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$  and there exists a point  $y$  of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  such that  $f = g$  and  $x = y$  and  $g$  is differentiable in  $y$ .

Let  $m, n$  be non empty elements of  $\mathbb{N}$ , let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ , and let  $x$  be an element of  $\mathcal{R}^m$ . Let us assume that  $f$  is differentiable in  $x$ . The functor  $f'(x)$  yields a function from  $\mathcal{R}^m$  into  $\mathcal{R}^n$  and is defined as follows:

(Def. 8) There exists a partial function  $g$  from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$  and there exists a point  $y$  of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  such that  $f = g$  and  $x = y$  and  $f'(x) = g'(y)$ .

We now state four propositions:

(3) Let  $I$  be a function from  $\mathbb{R}$  into  $\mathcal{R}^1$ . Suppose  $I = (\text{proj}(1, 1))^{-1}$ . Then

- (i) for every vector  $x$  of  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  and for every element  $y$  of  $\mathbb{R}$  such that  $x = I(y)$  holds  $\|x\| = |y|$ ,
  - (ii) for all vectors  $x, y$  of  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  and for all elements  $a, b$  of  $\mathbb{R}$  such that  $x = I(a)$  and  $y = I(b)$  holds  $x + y = I(a + b)$ ,
  - (iii) for every vector  $x$  of  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  and for every element  $y$  of  $\mathbb{R}$  and for every real number  $a$  such that  $x = I(y)$  holds  $a \cdot x = I(a \cdot y)$ ,
  - (iv) for every vector  $x$  of  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  and for every element  $a$  of  $\mathbb{R}$  such that  $x = I(a)$  holds  $-x = I(-a)$ , and
  - (v) for all vectors  $x, y$  of  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  and for all elements  $a, b$  of  $\mathbb{R}$  such that  $x = I(a)$  and  $y = I(b)$  holds  $x - y = I(a - b)$ .
- (4) Let  $J$  be a function from  $\mathcal{R}^1$  into  $\mathbb{R}$ . Suppose  $J = \text{proj}(1, 1)$ . Then
- (i) for every vector  $x$  of  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  and for every element  $y$  of  $\mathbb{R}$  such that  $J(x) = y$  holds  $\|x\| = |y|$ ,
  - (ii) for all vectors  $x, y$  of  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  and for all elements  $a, b$  of  $\mathbb{R}$  such that  $J(x) = a$  and  $J(y) = b$  holds  $J(x + y) = a + b$ ,
  - (iii) for every vector  $x$  of  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  and for every element  $y$  of  $\mathbb{R}$  and for every real number  $a$  such that  $J(x) = y$  holds  $J(a \cdot x) = a \cdot y$ ,
  - (iv) for every vector  $x$  of  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  and for every element  $a$  of  $\mathbb{R}$  such that  $J(x) = a$  holds  $J(-x) = -a$ , and
  - (v) for all vectors  $x, y$  of  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  and for all elements  $a, b$  of  $\mathbb{R}$  such that  $J(x) = a$  and  $J(y) = b$  holds  $J(x - y) = a - b$ .
- (5) Let  $I$  be a function from  $\mathbb{R}$  into  $\mathcal{R}^1$  and  $J$  be a function from  $\mathcal{R}^1$  into  $\mathbb{R}$ . Suppose  $I = (\text{proj}(1, 1))^{-1}$  and  $J = \text{proj}(1, 1)$ . Then
- (i) for every rest  $R$  of  $\langle \mathcal{E}^1, \|\cdot\| \rangle$ ,  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  holds  $J \cdot R \cdot I$  is a rest, and
  - (ii) for every linear operator  $L$  from  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  into  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  holds  $J \cdot L \cdot I$  is a linear function.
- (6) Let  $I$  be a function from  $\mathbb{R}$  into  $\mathcal{R}^1$  and  $J$  be a function from  $\mathcal{R}^1$  into  $\mathbb{R}$ . Suppose  $I = (\text{proj}(1, 1))^{-1}$  and  $J = \text{proj}(1, 1)$ . Then
- (i) for every rest  $R$  holds  $I \cdot R \cdot J$  is a rest of  $\langle \mathcal{E}^1, \|\cdot\| \rangle$ ,  $\langle \mathcal{E}^1, \|\cdot\| \rangle$ , and
  - (ii) for every linear function  $L$  holds  $I \cdot L \cdot J$  is a bounded linear operator from  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  into  $\langle \mathcal{E}^1, \|\cdot\| \rangle$ .

In the sequel  $f$  is a partial function from  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^1, \|\cdot\| \rangle$ ,  $g$  is a partial function from  $\mathbb{R}$  to  $\mathbb{R}$ ,  $x$  is a point of  $\langle \mathcal{E}^1, \|\cdot\| \rangle$ , and  $y$  is an element of  $\mathbb{R}$ .

We now state four propositions:

- (7) If  $f = \langle g \rangle$  and  $x = \langle y \rangle$  and  $f$  is differentiable in  $x$ , then  $g$  is differentiable in  $y$  and  $g'(y) = (\text{proj}(1, 1) \cdot f'(x) \cdot (\text{proj}(1, 1))^{-1})(1)$ .
- (8) If  $f = \langle g \rangle$  and  $x = \langle y \rangle$  and  $g$  is differentiable in  $y$ , then  $f$  is differentiable in  $x$  and  $f'(x)(\langle 1 \rangle) = \langle g'(y) \rangle$ .
- (9) If  $f = \langle g \rangle$  and  $x = \langle y \rangle$ , then  $f$  is differentiable in  $x$  iff  $g$  is differentiable in  $y$ .

- (10) If  $f = \langle g \rangle$  and  $x = \langle y \rangle$  and  $f$  is differentiable in  $x$ , then  $f'(x)(\langle 1 \rangle) = \langle g'(y) \rangle$ .

## 2. PARTIAL DIFFERENTIATION

For simplicity, we adopt the following rules:  $m, n$  are non empty elements of  $\mathbb{N}$ ,  $i, j$  are elements of  $\mathbb{N}$ ,  $f$  is a partial function from  $\langle \mathcal{E}^n, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^1, \|\cdot\| \rangle$ ,  $g$  is a partial function from  $\mathcal{R}^n$  to  $\mathbb{R}$ ,  $x$  is a point of  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ , and  $y$  is an element of  $\mathcal{R}^n$ .

Let  $n, m$  be non empty elements of  $\mathbb{N}$ , let  $i$  be an element of  $\mathbb{N}$ , let  $f$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ , and let  $x$  be a point of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$ . We say that  $f$  is partially differentiable in  $x$  w.r.t.  $i$  if and only if:

- (Def. 9)  $f \cdot \text{reproj}(i, x)$  is differentiable in  $(\text{Proj}(i, m))(x)$ .

Let  $m, n$  be non empty elements of  $\mathbb{N}$ , let  $i$  be an element of  $\mathbb{N}$ , let  $f$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ , and let  $x$  be a point of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$ . The functor  $\text{partdiff}(f, x, i)$  yielding a point of the real norm space of bounded linear operators from  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  into  $\langle \mathcal{E}^n, \|\cdot\| \rangle$  is defined as follows:

- (Def. 10)  $\text{partdiff}(f, x, i) = (f \cdot \text{reproj}(i, x))'((\text{Proj}(i, m))(x))$ .

Let  $n$  be a non empty element of  $\mathbb{N}$ , let  $i$  be an element of  $\mathbb{N}$ , let  $f$  be a partial function from  $\mathcal{R}^n$  to  $\mathbb{R}$ , and let  $x$  be an element of  $\mathcal{R}^n$ . We say that  $f$  is partially differentiable in  $x$  w.r.t.  $i$  if and only if:

- (Def. 11)  $f \cdot \text{reproj}(i, x)$  is differentiable in  $(\text{proj}(i, n))(x)$ .

Let  $n$  be a non empty element of  $\mathbb{N}$ , let  $i$  be an element of  $\mathbb{N}$ , let  $f$  be a partial function from  $\mathcal{R}^n$  to  $\mathbb{R}$ , and let  $x$  be an element of  $\mathcal{R}^n$ . The functor  $\text{partdiff}(f, x, i)$  yields a real number and is defined by:

- (Def. 12)  $\text{partdiff}(f, x, i) = (f \cdot \text{reproj}(i, x))'((\text{proj}(i, n))(x))$ .

We now state several propositions:

- (11)  $\text{Proj}(i, n) = (\text{proj}(1, 1))^{-1} \cdot \text{proj}(i, n)$ .  
(12) If  $x = y$ , then  $\text{reproj}(i, y) \cdot \text{proj}(1, 1) = \text{reproj}(i, x)$ .  
(13) If  $f = \langle g \rangle$  and  $x = y$ , then  $\langle g \cdot \text{reproj}(i, y) \rangle = f \cdot \text{reproj}(i, x)$ .  
(14) Suppose  $f = \langle g \rangle$  and  $x = y$ . Then  $f$  is partially differentiable in  $x$  w.r.t.  $i$  if and only if  $g$  is partially differentiable in  $y$  w.r.t.  $i$ .  
(15) If  $f = \langle g \rangle$  and  $x = y$  and  $f$  is partially differentiable in  $x$  w.r.t.  $i$ , then  $(\text{partdiff}(f, x, i))(\langle 1 \rangle) = \langle \text{partdiff}(g, y, i) \rangle$ .

Let  $m, n$  be non empty elements of  $\mathbb{N}$ , let  $i$  be an element of  $\mathbb{N}$ , let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ , and let  $x$  be an element of  $\mathcal{R}^m$ . We say that  $f$  is partially differentiable in  $x$  w.r.t.  $i$  if and only if the condition (Def. 13) is satisfied.

(Def. 13) There exists a partial function  $g$  from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$  and there exists a point  $y$  of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  such that  $f = g$  and  $x = y$  and  $g$  is partially differentiable in  $y$  w.r.t.  $i$ .

Let  $m, n$  be non empty elements of  $\mathbb{N}$ , let  $i$  be an element of  $\mathbb{N}$ , let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ , and let  $x$  be an element of  $\mathcal{R}^m$ . Let us assume that  $f$  is partially differentiable in  $x$  w.r.t.  $i$ . The functor  $\text{partdiff}(f, x, i)$  yielding an element of  $\mathcal{R}^n$  is defined as follows:

(Def. 14) There exists a partial function  $g$  from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$  and there exists a point  $y$  of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  such that  $f = g$  and  $x = y$  and  $\text{partdiff}(f, x, i) = (\text{partdiff}(g, y, i))(\langle 1 \rangle)$ .

One can prove the following four propositions:

- (16) Let  $m, n$  be non empty elements of  $\mathbb{N}$ ,  $F$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ ,  $G$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ ,  $x$  be a point of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$ , and  $y$  be an element of  $\mathcal{R}^m$ . Suppose  $F = G$  and  $x = y$ . Then  $F$  is partially differentiable in  $x$  w.r.t.  $i$  if and only if  $G$  is partially differentiable in  $y$  w.r.t.  $i$ .
- (17) Let  $m, n$  be non empty elements of  $\mathbb{N}$ ,  $F$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ ,  $G$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ ,  $x$  be a point of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$ , and  $y$  be an element of  $\mathcal{R}^m$ . Suppose  $F = G$  and  $x = y$  and  $F$  is partially differentiable in  $x$  w.r.t.  $i$ . Then  $(\text{partdiff}(F, x, i))(\langle 1 \rangle) = \text{partdiff}(G, y, i)$ .
- (18) Let  $g_1$  be a partial function from  $\mathcal{R}^n$  to  $\mathcal{R}^1$ . Suppose  $g_1 = \langle g \rangle$ . Then  $g_1$  is partially differentiable in  $y$  w.r.t.  $i$  if and only if  $g$  is partially differentiable in  $y$  w.r.t.  $i$ .
- (19) Let  $g_1$  be a partial function from  $\mathcal{R}^n$  to  $\mathcal{R}^1$ . Suppose  $g_1 = \langle g \rangle$  and  $g_1$  is partially differentiable in  $y$  w.r.t.  $i$ . Then  $\text{partdiff}(g_1, y, i) = \langle \text{partdiff}(g, y, i) \rangle$ .

### 3. LINEARITY OF PARTIAL DIFFERENTIAL OPERATOR

For simplicity, we use the following convention:  $X$  is a set,  $r$  is a real number,  $f, f_1, f_2$  are partial functions from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ ,  $g, g_1, g_2$  are partial functions from  $\mathcal{R}^n$  to  $\mathbb{R}$ ,  $h$  is a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ ,  $x$  is a point of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$ ,  $y$  is an element of  $\mathcal{R}^n$ , and  $z$  is an element of  $\mathcal{R}^m$ .

Let  $m, n$  be non empty elements of  $\mathbb{N}$ , let  $i, j$  be elements of  $\mathbb{N}$ , let  $f$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ , and let  $x$  be a point of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$ .

We say that  $f$  is partially differentiable in  $x$  w.r.t.  $i$  and  $j$  if and only if:

(Def. 15)  $\text{Proj}(j, n) \cdot f \cdot \text{reproj}(i, x)$  is differentiable in  $(\text{Proj}(i, m))(x)$ .

Let  $m, n$  be non empty elements of  $\mathbb{N}$ , let  $i, j$  be elements of  $\mathbb{N}$ , let  $f$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ , and let  $x$  be a point of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$ .

The functor  $\text{partdiff}(f, x, i, j)$  yields a point of the real norm space of bounded linear operators from  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  into  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  and is defined by:

$$\text{(Def. 16)} \quad \text{partdiff}(f, x, i, j) = (\text{Proj}(j, n) \cdot f \cdot \text{reproj}(i, x))'((\text{Proj}(i, m))(x)).$$

Let  $m, n$  be non empty elements of  $\mathbb{N}$ , let  $i, j$  be elements of  $\mathbb{N}$ , let  $h$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ , and let  $z$  be an element of  $\mathcal{R}^m$ . We say that  $h$  is partially differentiable in  $z$  w.r.t.  $i$  and  $j$  if and only if:

$$\text{(Def. 17)} \quad \text{proj}(j, n) \cdot h \cdot \text{reproj}(i, z) \text{ is differentiable in } (\text{proj}(i, m))(z).$$

Let  $m, n$  be non empty elements of  $\mathbb{N}$ , let  $i, j$  be elements of  $\mathbb{N}$ , let  $h$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ , and let  $z$  be an element of  $\mathcal{R}^m$ . The functor  $\text{partdiff}(h, z, i, j)$  yielding a real number is defined as follows:

$$\text{(Def. 18)} \quad \text{partdiff}(h, z, i, j) = (\text{proj}(j, n) \cdot h \cdot \text{reproj}(i, z))'((\text{proj}(i, m))(z)).$$

The following propositions are true:

(20) Let  $m, n$  be non empty elements of  $\mathbb{N}$ ,  $F$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ ,  $G$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ ,  $x$  be a point of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$ , and  $y$  be an element of  $\mathcal{R}^m$ . Suppose  $F = G$  and  $x = y$ . Then  $F$  is differentiable in  $x$  if and only if  $G$  is differentiable in  $y$ .

(21) Let  $m, n$  be non empty elements of  $\mathbb{N}$ ,  $F$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ ,  $G$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ ,  $x$  be a point of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$ , and  $y$  be an element of  $\mathcal{R}^m$ . If  $F = G$  and  $x = y$  and  $F$  is differentiable in  $x$ , then  $F'(x) = G'(y)$ .

(22) If  $f = h$  and  $x = z$ , then  $\text{Proj}(j, n) \cdot f \cdot \text{reproj}(i, x) = \langle \text{proj}(j, n) \cdot h \cdot \text{reproj}(i, z) \rangle$ .

(23) Suppose  $f = h$  and  $x = z$ . Then  $f$  is partially differentiable in  $x$  w.r.t.  $i$  and  $j$  if and only if  $h$  is partially differentiable in  $z$  w.r.t.  $i$  and  $j$ .

(24) If  $f = h$  and  $x = z$  and  $f$  is partially differentiable in  $x$  w.r.t.  $i$  and  $j$ , then  $(\text{partdiff}(f, x, i, j))(\langle 1 \rangle) = \langle \text{partdiff}(h, z, i, j) \rangle$ .

Let  $m, n$  be non empty elements of  $\mathbb{N}$ , let  $i$  be an element of  $\mathbb{N}$ , let  $f$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ , and let  $X$  be a set. We say that  $f$  is partially differentiable on  $X$  w.r.t.  $i$  if and only if:

$$\text{(Def. 19)} \quad X \subseteq \text{dom } f \text{ and for every point } x \text{ of } \langle \mathcal{E}^m, \|\cdot\| \rangle \text{ such that } x \in X \text{ holds } f \upharpoonright X \text{ is partially differentiable in } x \text{ w.r.t. } i.$$

We now state the proposition

(25) If  $f$  is partially differentiable on  $X$  w.r.t.  $i$ , then  $X$  is a subset of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$ .

Let  $m, n$  be non empty elements of  $\mathbb{N}$ , let  $i$  be an element of  $\mathbb{N}$ , let  $f$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ , and let us consider  $X$ . Let us assume that  $f$  is partially differentiable on  $X$  w.r.t.  $i$ . The functor  $f \upharpoonright^i X$  yielding a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to the real norm space of bounded linear operators from  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  into  $\langle \mathcal{E}^n, \|\cdot\| \rangle$  is defined by:

(Def. 20)  $\text{dom}(f \upharpoonright^i X) = X$  and for every point  $x$  of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  such that  $x \in X$  holds  $(f \upharpoonright^i X)_x = \text{partdiff}(f, x, i)$ .

The following propositions are true:

- (26)  $(f_1 + f_2) \cdot \text{reproj}(i, x) = f_1 \cdot \text{reproj}(i, x) + f_2 \cdot \text{reproj}(i, x)$  and  $(f_1 - f_2) \cdot \text{reproj}(i, x) = f_1 \cdot \text{reproj}(i, x) - f_2 \cdot \text{reproj}(i, x)$ .
- (27)  $r(f \cdot \text{reproj}(i, x)) = (r f) \cdot \text{reproj}(i, x)$ .
- (28) Suppose  $f_1$  is partially differentiable in  $x$  w.r.t.  $i$  and  $f_2$  is partially differentiable in  $x$  w.r.t.  $i$ . Then  $f_1 + f_2$  is partially differentiable in  $x$  w.r.t.  $i$  and  $\text{partdiff}(f_1 + f_2, x, i) = \text{partdiff}(f_1, x, i) + \text{partdiff}(f_2, x, i)$ .
- (29) Suppose  $g_1$  is partially differentiable in  $y$  w.r.t.  $i$  and  $g_2$  is partially differentiable in  $y$  w.r.t.  $i$ . Then  $g_1 + g_2$  is partially differentiable in  $y$  w.r.t.  $i$  and  $\text{partdiff}(g_1 + g_2, y, i) = \text{partdiff}(g_1, y, i) + \text{partdiff}(g_2, y, i)$ .
- (30) Suppose  $f_1$  is partially differentiable in  $x$  w.r.t.  $i$  and  $f_2$  is partially differentiable in  $x$  w.r.t.  $i$ . Then  $f_1 - f_2$  is partially differentiable in  $x$  w.r.t.  $i$  and  $\text{partdiff}(f_1 - f_2, x, i) = \text{partdiff}(f_1, x, i) - \text{partdiff}(f_2, x, i)$ .
- (31) Suppose  $g_1$  is partially differentiable in  $y$  w.r.t.  $i$  and  $g_2$  is partially differentiable in  $y$  w.r.t.  $i$ . Then  $g_1 - g_2$  is partially differentiable in  $y$  w.r.t.  $i$  and  $\text{partdiff}(g_1 - g_2, y, i) = \text{partdiff}(g_1, y, i) - \text{partdiff}(g_2, y, i)$ .
- (32) Suppose  $f$  is partially differentiable in  $x$  w.r.t.  $i$ . Then  $r f$  is partially differentiable in  $x$  w.r.t.  $i$  and  $\text{partdiff}(r f, x, i) = r \cdot \text{partdiff}(f, x, i)$ .
- (33) Suppose  $g$  is partially differentiable in  $y$  w.r.t.  $i$ . Then  $r g$  is partially differentiable in  $y$  w.r.t.  $i$  and  $\text{partdiff}(r g, y, i) = r \cdot \text{partdiff}(g, y, i)$ .

#### REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [2] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [3] Czesław Byliński. The complex numbers. *Formalized Mathematics*, 1(3):507–513, 1990.
- [4] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(3):529–536, 1990.
- [5] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [6] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [7] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [8] Czesław Byliński. The sum and product of finite sequences of real numbers. *Formalized Mathematics*, 1(4):661–668, 1990.
- [9] Agata Darmochwał. The Euclidean space. *Formalized Mathematics*, 2(4):599–603, 1991.
- [10] Noboru Endou and Yasunari Shidama. Completeness of the real Euclidean space. *Formalized Mathematics*, 13(4):577–580, 2005.
- [11] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [12] Hiroshi Imura, Morishige Kimura, and Yasunari Shidama. The differentiable functions on normed linear spaces. *Formalized Mathematics*, 12(3):321–327, 2004.
- [13] Jarosław Kotowicz. Partial functions from a domain to a domain. *Formalized Mathematics*, 1(4):697–702, 1990.

- [14] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(2):269–272, 1990.
- [15] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(1):223–230, 1990.
- [16] Jan Popiołek. Real normed space. *Formalized Mathematics*, 2(1):111–115, 1991.
- [17] Konrad Raczkowski and Paweł Sadowski. Real function differentiability. *Formalized Mathematics*, 1(4):797–801, 1990.
- [18] Laurent Schwartz. *Cours d'analyse*. Hermann, 1981.
- [19] Yasunari Shidama. Banach space of bounded linear operators. *Formalized Mathematics*, 12(1):39–48, 2004.
- [20] Andrzej Trybulec. Subsets of complex numbers. *To appear in Formalized Mathematics*.
- [21] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [22] Wojciech A. Trybulec. Pigeon hole principle. *Formalized Mathematics*, 1(3):575–579, 1990.
- [23] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(2):291–296, 1990.
- [24] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [25] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [26] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.
- [27] Hiroshi Yamazaki, Yoshinori Fujisawa, and Yatsuka Nakamura. On replace function and swap function for finite sequences. *Formalized Mathematics*, 9(3):471–474, 2001.
- [28] Hiroshi Yamazaki and Yasunari Shidama. Algebra of vector functions. *Formalized Mathematics*, 3(2):171–175, 1992.

*Received June 6, 2007*

---