

The Rank+Nullity Theorem

Jesse Alama
 Department of Philosophy
 Stanford University
 USA

Summary. The rank+nullity theorem states that, if T is a linear transformation from a finite-dimensional vector space V to a finite-dimensional vector space W , then $\dim(V) = \text{rank}(T) + \text{nullity}(T)$, where $\text{rank}(T) = \dim(\text{im}(T))$ and $\text{nullity}(T) = \dim(\ker(T))$. The proof treated here is standard; see, for example, [14]: take a basis A of $\ker(T)$ and extend it to a basis B of V , and then show that $\dim(\text{im}(T))$ is equal to $|B - A|$, and that T is one-to-one on $B - A$.

MML identifier: RANKNULL, version: 7.8.05 4.87.985

The articles [21], [11], [32], [22], [19], [33], [34], [7], [2], [17], [10], [18], [8], [9], [20], [1], [12], [3], [5], [6], [27], [29], [24], [31], [25], [13], [4], [30], [28], [26], [23], [15], [16], and [35] provide the notation and terminology for this paper.

1. PRELIMINARIES

One can prove the following three propositions:

- (1) For all functions f, g such that g is one-to-one and $f \upharpoonright \text{rng } g$ is one-to-one and $\text{rng } g \subseteq \text{dom } f$ holds $f \cdot g$ is one-to-one.
- (2) For every function f and for all sets X, Y such that $X \subseteq Y$ and $f \upharpoonright Y$ is one-to-one holds $f \upharpoonright X$ is one-to-one.
- (3) Let V be a 1-sorted structure and X, Y be subsets of V . Then X meets Y if and only if there exists an element v of V such that $v \in X$ and $v \in Y$.

In the sequel F is a field and V, W are vector spaces over F .

Let F be a field and let V be a finite dimensional vector space over F . One can verify that there exists a basis of V which is finite.

Let F be a field and let V, W be vector spaces over F . Note that there exists a function from V into W which is linear.

Next we state three propositions:

- (4) If Ω_V is finite, then V is finite dimensional.
- (5) For every finite dimensional vector space V over F such that $\overline{\overline{\Omega_V}} = 1$ holds $\dim(V) = 0$.
- (6) If $\overline{\overline{\Omega_V}} = 2$, then $\dim(V) = 1$.

2. BASIC FACTS OF LINEAR TRANSFORMATIONS

Let F be a field and let V, W be vector spaces over F . A linear transformation from V to W is a linear function from V into W .

In the sequel T is a linear transformation from V to W .

One can prove the following propositions:

- (7) For all non empty 1-sorted structures V, W and for every function T from V into W holds $\text{dom } T = \Omega_V$ and $\text{rng } T \subseteq \Omega_W$.
- (8) For all elements x, y of V holds $T(x) - T(y) = T(x - y)$.
- (9) $T(0_V) = 0_W$.

Let F be a field, let V, W be vector spaces over F , and let T be a linear transformation from V to W . The functor $\ker T$ yielding a strict subspace of V is defined as follows:

(Def. 1) $\Omega_{\ker T} = \{u; u \text{ ranges over elements of } V: T(u) = 0_W\}$.

We now state the proposition

- (10) For every element x of V holds $x \in \ker T$ iff $T(x) = 0_W$.

Let V, W be non empty 1-sorted structures, let T be a function from V into W , and let X be a subset of V . Then $T^\circ X$ is a subset of W .

Let F be a field, let V, W be vector spaces over F , and let T be a linear transformation from V to W . The functor $\text{im } T$ yielding a strict subspace of W is defined as follows:

(Def. 2) $\Omega_{\text{im } T} = T^\circ(\Omega_V)$.

The following propositions are true:

- (11) $0_V \in \ker T$.
- (12) For every subset X of V holds $T^\circ X$ is a subset of $\text{im } T$.
- (13) For every element y of W holds $y \in \text{im } T$ iff there exists an element x of V such that $y = T(x)$.
- (14) For every element x of $\ker T$ holds $T(x) = 0_W$.
- (15) If T is one-to-one, then $\ker T = \mathbf{0}_V$.
- (16) For every finite dimensional vector space V over F holds $\dim(\mathbf{0}_V) = 0$.

- (17) For all elements x, y of V such that $T(x) = T(y)$ holds $x - y \in \ker T$.
- (18) For every subset A of V and for all elements x, y of V such that $x - y \in \text{Lin}(A)$ holds $x \in \text{Lin}(A \cup \{y\})$.

3. SOME LEMMAS ON LINEARLY INDEPENDENT SUBSETS, LINEAR COMBINATIONS, AND LINEAR TRANSFORMATIONS

One can prove the following propositions:

- (19) For every subset X of V such that V is a subspace of W holds X is a subset of W .
- (20) For every subset A of V such that A is linearly independent holds A is a basis of $\text{Lin}(A)$.
- (21) For every subset A of V and for every element x of V such that $x \in \text{Lin}(A)$ and $x \notin A$ holds $A \cup \{x\}$ is linearly dependent.
- (22) For every subset A of V and for every basis B of V such that A is a basis of $\ker T$ and $A \subseteq B$ holds $T \upharpoonright (B \setminus A)$ is one-to-one.
- (23) Let A be a subset of V , l be a linear combination of A , x be an element of V , and a be an element of F . Then $l + \cdot (x, a)$ is a linear combination of $A \cup \{x\}$.

Let V be a 1-sorted structure and let X be a subset of V . The functor $V \setminus X$ yields a subset of V and is defined by:

(Def. 3) $V \setminus X = \Omega_V \setminus X$.

Let F be a field, let V be a vector space over F , let l be a linear combination of V , and let X be a subset of V . Then $l^\circ X$ is a subset of F .

In the sequel l is a linear combination of V .

Let F be a field and let V be a vector space over F . Note that there exists a subset of V which is linearly dependent.

Let F be a field, let V be a vector space over F , let l be a linear combination of V , and let A be a subset of V . The functor $l[A]$ yields a linear combination of A and is defined by:

(Def. 4) $l[A] = l \upharpoonright A + \cdot (V \setminus A \mapsto 0_F)$.

The following propositions are true:

- (24) $l = l[\text{the support of } l]$.
- (25) For every subset A of V and for every element v of V such that $v \in A$ holds $l[A](v) = l(v)$.
- (26) For every subset A of V and for every element v of V such that $v \notin A$ holds $l[A](v) = 0_F$.
- (27) For all subsets A, B of V and for every linear combination l of B such that $A \subseteq B$ holds $l = l[A] + l[B \setminus A]$.

Let F be a field, let V be a vector space over F , let l be a linear combination of V , and let X be a subset of V . Observe that $l^\circ X$ is finite.

Let V, W be non empty 1-sorted structures, let T be a function from V into W , and let X be a subset of W . Then $T^{-1}(X)$ is a subset of V .

We now state the proposition

- (28) For every subset X of V such that X misses the support of l holds $l^\circ X \subseteq \{0_F\}$.

Let F be a field, let V, W be vector spaces over F , let l be a linear combination of V , and let T be a linear transformation from V to W . The functor $T^\circ l$ yielding a linear combination of W is defined by:

(Def. 5) For every element w of W holds $(T^\circ l)(w) = \sum(l^\circ T^{-1}(\{w\}))$.

One can prove the following propositions:

- (29) $T^\circ l$ is a linear combination of $T^\circ(\text{the support of } l)$.
- (30) The support of $T^\circ l \subseteq T^\circ(\text{the support of } l)$.
- (31) Let l, m be linear combinations of V . Suppose the support of l misses the support of m . Then the support of $l + m = (\text{the support of } l) \cup (\text{the support of } m)$.
- (32) Let l, m be linear combinations of V . Suppose the support of l misses the support of m . Then the support of $l - m = (\text{the support of } l) \cup (\text{the support of } m)$.
- (33) For all subsets A, B of V such that $A \subseteq B$ and B is a basis of V holds V is the direct sum of $\text{Lin}(A)$ and $\text{Lin}(B \setminus A)$.
- (34) Let A be a subset of V , l be a linear combination of A , and v be an element of V . Suppose $T|_A$ is one-to-one and $v \in A$. Then there exists a subset X of V such that X misses A and $T^{-1}(\{T(v)\}) = \{v\} \cup X$.
- (35) For every subset X of V such that X misses the support of l and $X \neq \emptyset$ holds $l^\circ X = \{0_F\}$.
- (36) For every element w of W such that $w \in \text{the support of } T^\circ l$ holds $T^{-1}(\{w\})$ meets the support of l .
- (37) Let v be an element of V . Suppose $T|_{(\text{the support of } l)}$ is one-to-one and $v \in \text{the support of } l$. Then $(T^\circ l)(T(v)) = l(v)$.
- (38) Let G be a finite sequence of elements of V . Suppose $\text{rng } G = \text{the support of } l$ and $T|_{(\text{the support of } l)}$ is one-to-one. Then $T \cdot (l \cdot G) = (T^\circ l)(T \cdot G)$.
- (39) If $T|_{(\text{the support of } l)}$ is one-to-one, then $T^\circ(\text{the support of } l) = \text{the support of } T^\circ l$.
- (40) Let A be a subset of V , B be a basis of V , and l be a linear combination of $B \setminus A$. If A is a basis of $\ker T$ and $A \subseteq B$, then $T(\sum l) = \sum(T^\circ l)$.
- (41) Let X be a subset of V . Suppose X is linearly dependent. Then there exists a linear combination l of X such that the support of $l \neq \emptyset$ and

$$\sum l = 0_V.$$

Let F be a field, let V, W be vector spaces over F , let X be a subset of V , let T be a linear transformation from V to W , and let l be a linear combination of $T^\circ X$. Let us assume that $T \upharpoonright X$ is one-to-one. The functor $T \# l$ yields a linear combination of X and is defined as follows:

(Def. 6) $T \# l = l \cdot T + \cdot (V \setminus X \mapsto 0_F)$.

We now state two propositions:

- (42) Let X be a subset of V , l be a linear combination of $T^\circ X$, and v be an element of V . If $v \in X$ and $T \upharpoonright X$ is one-to-one, then $(T \# l)(v) = l(T(v))$.
- (43) For every subset X of V and for every linear combination l of $T^\circ X$ such that $T \upharpoonright X$ is one-to-one holds $T^\circ T \# l = l$.

4. THE RANK+NULLITY THEOREM

Let F be a field, let V, W be finite dimensional vector spaces over F , and let T be a linear transformation from V to W . The functor $\text{rank } T$ yielding a natural number is defined by:

(Def. 7) $\text{rank } T = \dim(\text{im } T)$.

The functor $\text{nullity } T$ yields a natural number and is defined by:

(Def. 8) $\text{nullity } T = \dim(\ker T)$.

Next we state two propositions:

- (44) Let V, W be finite dimensional vector spaces over F and T be a linear transformation from V to W . Then $\dim(V) = \text{rank } T + \text{nullity } T$.
- (45) Let V, W be finite dimensional vector spaces over F and T be a linear transformation from V to W . If T is one-to-one, then $\dim(V) = \text{rank } T$.

REFERENCES

- [1] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(2):377–382, 1990.
- [2] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [4] Grzegorz Bancerek and Andrzej Trybulec. Miscellaneous facts about functions. *Formalized Mathematics*, 5(4):485–492, 1996.
- [5] Czesław Byliński. Binary operations applied to finite sequences. *Formalized Mathematics*, 1(4):643–649, 1990.
- [6] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(3):529–536, 1990.
- [7] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [8] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [9] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Formalized Mathematics*, 1(3):521–527, 1990.
- [10] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.

- [11] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [12] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(1):165–167, 1990.
- [13] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Formalized Mathematics*, 1(2):335–342, 1990.
- [14] Serge Lang. *Algebra*. Springer, 3rd edition, 2005.
- [15] Robert Milewski. Associated matrix of linear map. *Formalized Mathematics*, 5(3):339–345, 1996.
- [16] Michał Muzalewski. Rings and modules – part II. *Formalized Mathematics*, 2(4):579–585, 1991.
- [17] Andrzej Trybulec. Subsets of complex numbers. *To appear in Formalized Mathematics*.
- [18] Andrzej Trybulec. Binary operations applied to functions. *Formalized Mathematics*, 1(2):329–334, 1990.
- [19] Andrzej Trybulec. Domains and their Cartesian products. *Formalized Mathematics*, 1(1):115–122, 1990.
- [20] Andrzej Trybulec. Function domains and Frænkel operator. *Formalized Mathematics*, 1(3):495–500, 1990.
- [21] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [22] Andrzej Trybulec. Tuples, projections and Cartesian products. *Formalized Mathematics*, 1(1):97–105, 1990.
- [23] Wojciech A. Trybulec. Basis of vector space. *Formalized Mathematics*, 1(5):883–885, 1990.
- [24] Wojciech A. Trybulec. Groups. *Formalized Mathematics*, 1(5):821–827, 1990.
- [25] Wojciech A. Trybulec. Linear combinations in real linear space. *Formalized Mathematics*, 1(3):581–588, 1990.
- [26] Wojciech A. Trybulec. Linear combinations in vector space. *Formalized Mathematics*, 1(5):877–882, 1990.
- [27] Wojciech A. Trybulec. Non-contiguous substrings and one-to-one finite sequences. *Formalized Mathematics*, 1(3):569–573, 1990.
- [28] Wojciech A. Trybulec. Operations on subspaces in vector space. *Formalized Mathematics*, 1(5):871–876, 1990.
- [29] Wojciech A. Trybulec. Pigeon hole principle. *Formalized Mathematics*, 1(3):575–579, 1990.
- [30] Wojciech A. Trybulec. Subspaces and cosets of subspaces in vector space. *Formalized Mathematics*, 1(5):865–870, 1990.
- [31] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(2):291–296, 1990.
- [32] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [33] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [34] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.
- [35] Mariusz Żynel. The Steinitz theorem and the dimension of a vector space. *Formalized Mathematics*, 5(3):423–428, 1996.

Received July 31, 2007
