Integrability Formulas. Part II

Bo Li
 Qingdao University of Science
 and Technology
 China

Na Ma
 Qingdao University of Science
 and Technology
 China

Xiquan Liang
 Qingdao University of Science
 and Technology
 China

Summary. In this article, we give several differentiation and integrability formulas of special and composite functions including trigonometric function, and polynomial function.

MML identifier: INTEGR13, version: 7.11.07 4.156.1112

The terminology and notation used here have been introduced in the following articles: [12], [13], [2], [3], [9], [1], [6], [11], [14], [4], [18], [7], [8], [5], [19], [10], [16], [17], and [15].

For simplicity, we use the following convention: a, x are real numbers, n is an element of $\mathbb{N}$, A is a closed-interval subset of $\mathbb{R}$, f, h, f_1, f_2 are partial functions from $\mathbb{R}$ to $\mathbb{R}$, and Z is an open subset of $\mathbb{R}$.

The following propositions are true:

- (1) Suppose that
  - (i) $A \subseteq Z$,
  - (ii) $f = \frac{1}{(\text{the function sin})(\text{the function cos})}$,
  - (iii) $Z \subseteq \text{dom}((\text{the function ln}) \cdot (\text{the function tan}))$,
  - (iv) $Z = \text{dom } f$, and
  - (v) f is continuous on A .

Then $\int_A f(x)dx = ((\text{the function ln}) \cdot (\text{the function tan}))(\sup A) - ((\text{the function ln}) \cdot (\text{the function tan}))(\inf A)$.

(2) Suppose that

- (i) $A \subseteq Z$,
- (ii) $f = -\frac{1}{(\text{the function sin})(\text{the function cos})}$,
- (iii) $Z \subseteq \text{dom}((\text{the function ln}) \cdot (\text{the function cot}))$,
- (iv) $Z = \text{dom } f$, and
- (v) f is continuous on A .

Then $\int_A f(x)dx = ((\text{the function ln}) \cdot (\text{the function cot}))(\sup A) - ((\text{the function ln}) \cdot (\text{the function cot}))(\inf A)$.

(3) Suppose that

- (i) $A \subseteq Z$,
- (ii) $f = 2((\text{the function exp})(\text{the function sin}))$,
- (iii) $Z \subseteq \text{dom}((\text{the function exp})((\text{the function sin}) - (\text{the function cos})))$,
- (iv) $Z = \text{dom } f$, and
- (v) f is continuous on A .

Then $\int_A f(x)dx = ((\text{the function exp})((\text{the function sin}) - (\text{the function cos}))) (\sup A) - ((\text{the function exp})((\text{the function sin}) - (\text{the function cos}))) (\inf A)$.

(4) Suppose that

- (i) $A \subseteq Z$,
- (ii) $f = 2((\text{the function exp})(\text{the function cos}))$,
- (iii) $Z \subseteq \text{dom}((\text{the function exp})((\text{the function sin}) + (\text{the function cos})))$,
- (iv) $Z = \text{dom } f$, and
- (v) f is continuous on A .

Then $\int_A f(x)dx = ((\text{the function exp})((\text{the function sin}) + (\text{the function cos}))) (\sup A) - ((\text{the function exp})((\text{the function sin}) + (\text{the function cos}))) (\inf A)$.

(5) Suppose $A \subseteq Z = \text{dom}((\text{the function cos}) - (\text{the function sin}))$ and $(\text{the function cos}) - (\text{the function sin})$ is continuous on A .

Then $\int_A ((\text{the function cos}) - (\text{the function sin}))(x)dx = ((\text{the function sin}) + (\text{the function cos}))(\sup A) - ((\text{the function sin}) + (\text{the function cos}))(\inf A)$.

(6) Suppose $A \subseteq Z = \text{dom}((\text{the function cos}) + (\text{the function sin}))$ and $(\text{the function cos}) + (\text{the function sin})$ is continuous on A .

Then $\int_A ((\text{the function cos}) + (\text{the function sin}))(x)dx = ((\text{the function sin}) - (\text{the function cos}))(\sup A) - ((\text{the function sin}) - (\text{the function cos}))(\inf A)$.

- (7) Suppose $Z \subseteq \text{dom}((-\frac{1}{2}) \frac{(\text{the function sin})+(\text{the function cos})}{\text{the function exp}})$. Then
- (i) $(-\frac{1}{2}) \frac{(\text{the function sin})+(\text{the function cos})}{\text{the function exp}}$ is differentiable on Z , and
  - (ii) for every x such that $x \in Z$ holds

$$((-\frac{1}{2}) \frac{(\text{the function sin})+(\text{the function cos})}{\text{the function exp}})'|_Z(x) = \frac{(\text{the function sin})(x)}{(\text{the function exp})(x)}.$$

(8) Suppose that

- (i) $A \subseteq Z$,
- (ii) $f = \frac{\text{the function sin}}{\text{the function exp}}$,
- (iii) $Z \subseteq \text{dom}((-\frac{1}{2}) \frac{(\text{the function sin})+(\text{the function cos})}{\text{the function exp}})$,
- (iv) $Z = \text{dom } f$, and
- (v) f is continuous on A .

$$\begin{aligned} \text{Then } \int_A f(x)dx &= ((-\frac{1}{2}) \frac{(\text{the function sin}) + (\text{the function cos})}{\text{the function exp}})(\sup A) - \\ &((-\frac{1}{2}) \frac{(\text{the function sin}) + (\text{the function cos})}{\text{the function exp}})(\inf A). \end{aligned}$$

- (9) Suppose $Z \subseteq \text{dom}(\frac{1}{2} \frac{(\text{the function sin})-(\text{the function cos})}{\text{the function exp}})$. Then
- (i) $\frac{1}{2} \frac{(\text{the function sin})-(\text{the function cos})}{\text{the function exp}}$ is differentiable on Z , and
  - (ii) for every x such that $x \in Z$ holds

$$(\frac{1}{2} \frac{(\text{the function sin})-(\text{the function cos})}{\text{the function exp}})'|_Z(x) = \frac{(\text{the function cos})(x)}{(\text{the function exp})(x)}.$$

(10) Suppose that

- (i) $A \subseteq Z$,
- (ii) $f = \frac{\text{the function cos}}{\text{the function exp}}$,
- (iii) $Z \subseteq \text{dom}(\frac{1}{2} \frac{(\text{the function sin})-(\text{the function cos})}{\text{the function exp}})$,
- (iv) $Z = \text{dom } f$, and
- (v) f is continuous on A .

$$\begin{aligned} \text{Then } \int_A f(x)dx &= (\frac{1}{2} \frac{(\text{the function sin}) - (\text{the function cos})}{\text{the function exp}})(\sup A) - \\ &(\frac{1}{2} \frac{(\text{the function sin}) - (\text{the function cos})}{\text{the function exp}})(\inf A). \end{aligned}$$

(11) Suppose that

- (i) $A \subseteq Z$,
- (ii) $f = (\text{the function exp}) ((\text{the function sin})+(\text{the function cos}))$,
- (iii) $Z \subseteq \text{dom}((\text{the function exp}) (\text{the function sin}))$,
- (iv) $Z = \text{dom } f$, and
- (v) f is continuous on A .

$$\begin{aligned} \text{Then } \int_A f(x)dx &= ((\text{the function exp}) (\text{the function sin}))(\sup A) - ((\text{the} \\ &\text{function exp}) (\text{the function sin}))(\inf A). \end{aligned}$$

(12) Suppose that

- (i) $A \subseteq Z$,

- (ii) $f = (\text{the function exp}) ((\text{the function cos}) - (\text{the function sin}))$,
- (iii) $Z \subseteq \text{dom}((\text{the function exp}) (\text{the function cos}))$,
- (iv) $Z = \text{dom } f$, and
- (v) f is continuous on A .

Then $\int_A f(x)dx = ((\text{the function exp}) (\text{the function cos}))(\sup A) - ((\text{the function exp}) (\text{the function cos}))(\inf A)$.

(13) Suppose that

- (i) $A \subseteq Z$,
- (ii) $f_1 = \square^2$,
- (iii) $f = -\frac{\frac{\text{the function sin}}{\text{the function cos}}}{f_1} + \frac{\frac{1}{\text{id}_Z}}{(\text{the function cos})^2}$,
- (iv) $Z \subseteq \text{dom}(\frac{1}{\text{id}_Z} (\text{the function tan}))$,
- (v) $Z = \text{dom } f$, and
- (vi) f is continuous on A .

Then $\int_A f(x)dx = (\frac{1}{\text{id}_Z} (\text{the function tan}))(\sup A) - (\frac{1}{\text{id}_Z} (\text{the function tan}))(\inf A)$.

(14) Suppose that

- (i) $A \subseteq Z$,
- (ii) $f = -\frac{\frac{\text{the function cos}}{\text{the function sin}}}{f_1} - \frac{\frac{1}{\text{id}_Z}}{(\text{the function sin})^2}$,
- (iii) $f_1 = \square^2$,
- (iv) $Z \subseteq \text{dom}(\frac{1}{\text{id}_Z} (\text{the function cot}))$,
- (v) $Z = \text{dom } f$, and
- (vi) f is continuous on A .

Then $\int_A f(x)dx = (\frac{1}{\text{id}_Z} (\text{the function cot}))(\sup A) - (\frac{1}{\text{id}_Z} (\text{the function cot}))(\inf A)$.

(15) Suppose that

- (i) $A \subseteq Z$,
- (ii) $f = \frac{\frac{\text{the function sin}}{\text{the function cos}}}{\text{id}_Z} + \frac{\text{the function ln}}{(\text{the function cos})^2}$,
- (iii) $Z \subseteq \text{dom}((\text{the function ln}) (\text{the function tan}))$,
- (iv) $Z = \text{dom } f$, and
- (v) f is continuous on A .

Then $\int_A f(x)dx = ((\text{the function ln}) (\text{the function tan}))(\sup A) - ((\text{the function ln}) (\text{the function tan}))(\inf A)$.

(16) Suppose that

- (i) $A \subseteq Z$,
- (ii) $f = \frac{\frac{\text{the function cos}}{\text{the function sin}}}{\text{id}_Z} - \frac{\text{the function ln}}{(\text{the function sin})^2}$,

- (iii) $Z \subseteq \text{dom}((\text{the function } \ln) (\text{the function } \cot)),$
- (iv) $Z = \text{dom } f,$ and
- (v) f is continuous on $A.$

Then $\int_A f(x)dx = ((\text{the function } \ln) (\text{the function } \cot))(\sup A) - ((\text{the function } \ln) (\text{the function } \cot))(\inf A).$

(17) Suppose that

- (i) $A \subseteq Z,$
- (ii) $f = \frac{\text{the function } \tan}{\text{id}_Z} + \frac{\text{the function } \ln}{(\text{the function } \cos)^2},$
- (iii) $Z \subseteq \text{dom}((\text{the function } \ln) (\text{the function } \tan)),$
- (iv) $Z \subseteq \text{dom}(\text{the function } \tan),$
- (v) $Z = \text{dom } f,$ and
- (vi) f is continuous on $A.$

Then $\int_A f(x)dx = ((\text{the function } \ln) (\text{the function } \tan))(\sup A) - ((\text{the function } \ln) (\text{the function } \tan))(\inf A).$

(18) Suppose that

- (i) $A \subseteq Z,$
- (ii) $f = \frac{\text{the function } \cot}{\text{id}_Z} - \frac{\text{the function } \ln}{(\text{the function } \sin)^2},$
- (iii) $Z \subseteq \text{dom}((\text{the function } \ln) (\text{the function } \cot)),$
- (iv) $Z \subseteq \text{dom}(\text{the function } \cot),$
- (v) $Z = \text{dom } f,$ and
- (vi) f is continuous on $A.$

Then $\int_A f(x)dx = ((\text{the function } \ln) (\text{the function } \cot))(\sup A) - ((\text{the function } \ln) (\text{the function } \cot))(\inf A).$

(19) Suppose that

- (i) $A \subseteq Z,$
- (ii) for every x such that $x \in Z$ holds $f_1(x) = 1,$
- (iii) $f = \frac{\text{the function } \arctan}{\text{id}_Z} + \frac{\text{the function } \ln}{f_1 + \square^2},$
- (iv) $Z \subseteq]-1, 1[,$
- (v) $Z = \text{dom } f,$ and
- (vi) f is continuous on $A.$

Then $\int_A f(x)dx = ((\text{the function } \ln) (\text{the function } \arctan))(\sup A) - ((\text{the function } \ln) (\text{the function } \arctan))(\inf A).$

(20) Suppose that

- (i) $A \subseteq Z,$
- (ii) for every x such that $x \in Z$ holds $f_1(x) = 1,$
- (iii) $f = \frac{\text{the function } \text{arccot}}{\text{id}_Z} - \frac{\text{the function } \ln}{f_1 + \square^2},$

- (iv) $Z \subseteq]-1, 1[$,
- (v) $Z = \text{dom } f$, and
- (vi) f is continuous on A .

Then $\int_A f(x)dx = ((\text{the function } \ln) \cdot (\text{the function } \operatorname{arccot}))(\sup A) - ((\text{the function } \ln) \cdot (\text{the function } \operatorname{arccot}))(\inf A)$.

- (21) Suppose $A \subseteq Z$ and $f = \frac{(\text{the function } \exp) \cdot (\text{the function } \tan)}{(\text{the function } \cos)^2}$ and $Z = \text{dom } f$ and f is continuous on A . Then $\int_A f(x)dx = ((\text{the function } \exp) \cdot (\text{the function } \tan))(\sup A) - ((\text{the function } \exp) \cdot (\text{the function } \tan))(\inf A)$.

- (22) Suppose $A \subseteq Z$ and $f = -\frac{(\text{the function } \exp) \cdot (\text{the function } \cot)}{(\text{the function } \sin)^2}$ and $Z = \text{dom } f$ and f is continuous on A . Then $\int_A f(x)dx = ((\text{the function } \exp) \cdot (\text{the function } \cot))(\sup A) - ((\text{the function } \exp) \cdot (\text{the function } \cot))(\inf A)$.

- (23) Suppose $Z \subseteq \text{dom}((\text{the function } \exp) \cdot (\text{the function } \cot))$. Then
- (i) $-(\text{the function } \exp) \cdot (\text{the function } \cot)$ is differentiable on Z , and
  - (ii) for every x such that $x \in Z$ holds $(-(\text{the function } \exp) \cdot (\text{the function } \cot))'|_Z(x) = \frac{(\text{the function } \exp)((\text{the function } \cot)(x))}{(\text{the function } \sin)(x)^2}$.

- (24) Suppose $A \subseteq Z$ and $f = \frac{(\text{the function } \exp) \cdot (\text{the function } \cot)}{(\text{the function } \sin)^2}$ and $Z = \text{dom } f$ and f is continuous on A . Then $\int_A f(x)dx = (-(\text{the function } \exp) \cdot (\text{the function } \cot))(\sup A) - (-(\text{the function } \exp) \cdot (\text{the function } \cot))(\inf A)$.

- (25) Suppose that

- (i) $A \subseteq Z$,
- (ii) $f = \frac{1}{\operatorname{id}_Z((\text{the function } \cos) \cdot (\text{the function } \ln))^2}$,
- (iii) $Z \subseteq \text{dom}((\text{the function } \tan) \cdot (\text{the function } \ln))$,
- (iv) $Z = \text{dom } f$, and
- (v) f is continuous on A .

Then $\int_A f(x)dx = ((\text{the function } \tan) \cdot (\text{the function } \ln))(\sup A) - ((\text{the function } \tan) \cdot (\text{the function } \ln))(\inf A)$.

- (26) Suppose that

- (i) $A \subseteq Z$,
- (ii) $f = -\frac{1}{\operatorname{id}_Z((\text{the function } \sin) \cdot (\text{the function } \ln))^2}$,
- (iii) $Z \subseteq \text{dom}((\text{the function } \cot) \cdot (\text{the function } \ln))$,
- (iv) $Z = \text{dom } f$, and
- (v) f is continuous on A .

Then $\int_A f(x)dx = ((\text{the function cot}) \cdot (\text{the function ln}))(\sup A) - ((\text{the function cot}) \cdot (\text{the function ln}))(\inf A)$.

(27) Suppose $Z \subseteq \text{dom}((\text{the function cot}) \cdot (\text{the function ln}))$. Then

- (i) $-(\text{the function cot}) \cdot (\text{the function ln})$ is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $(-(\text{the function cot}) \cdot (\text{the function ln}))'_Z(x) = \frac{1}{x \cdot (\text{the function sin})((\text{the function ln})(x))^2}$.

(28) Suppose that

- (i) $A \subseteq Z$,
- (ii) $f = \frac{1}{\text{id}_Z((\text{the function sin}) \cdot (\text{the function ln}))^2}$,
- (iii) $Z \subseteq \text{dom}((\text{the function cot}) \cdot (\text{the function ln}))$,
- (iv) $Z = \text{dom } f$, and
- (v) f is continuous on A .

Then $\int_A f(x)dx = (-(\text{the function cot}) \cdot (\text{the function ln}))(\sup A) - (-(\text{the function cot}) \cdot (\text{the function ln}))(\inf A)$.

(29) Suppose that

- (i) $A \subseteq Z$,
- (ii) $f = \frac{\text{the function exp}}{((\text{the function cos}) \cdot (\text{the function exp}))^2}$,
- (iii) $Z \subseteq \text{dom}((\text{the function tan}) \cdot (\text{the function exp}))$,
- (iv) $Z = \text{dom } f$, and
- (v) f is continuous on A .

Then $\int_A f(x)dx = ((\text{the function tan}) \cdot (\text{the function exp}))(\sup A) - ((\text{the function tan}) \cdot (\text{the function exp}))(\inf A)$.

(30) Suppose that

- (i) $A \subseteq Z$,
- (ii) $f = -\frac{\text{the function exp}}{((\text{the function sin}) \cdot (\text{the function exp}))^2}$,
- (iii) $Z \subseteq \text{dom}((\text{the function cot}) \cdot (\text{the function exp}))$,
- (iv) $Z = \text{dom } f$, and
- (v) f is continuous on A .

Then $\int_A f(x)dx = ((\text{the function cot}) \cdot (\text{the function exp}))(\sup A) - ((\text{the function cot}) \cdot (\text{the function exp}))(\inf A)$.

(31) Suppose $Z \subseteq \text{dom}((\text{the function cot}) \cdot (\text{the function exp}))$. Then

- (i) $-(\text{the function cot}) \cdot (\text{the function exp})$ is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $(-(\text{the function cot}) \cdot (\text{the function exp}))'_Z(x) = \frac{(\text{the function exp})(x)}{(\text{the function sin})((\text{the function exp})(x))^2}$.

(32) Suppose that

- (i) $A \subseteq Z$,

- (ii) $f = \frac{\text{the function exp}}{((\text{the function sin}) \cdot (\text{the function exp}))^2},$
- (iii) $Z \subseteq \text{dom}((\text{the function cot}) \cdot (\text{the function exp})),$
- (iv) $Z = \text{dom } f,$ and
- (v) f is continuous on $A.$

Then $\int_A f(x)dx = (-(\text{the function cot}) \cdot (\text{the function exp}))(\sup A) - (-(\text{the function cot}) \cdot (\text{the function exp}))(\inf A).$

(33) Suppose that

- (i) $A \subseteq Z,$
- (ii) for every x such that $x \in Z$ holds $f(x) = -\frac{1}{x^2 \cdot (\text{the function cos})(\frac{1}{x})^2},$
- (iii) $Z \subseteq \text{dom}((\text{the function tan}) \cdot \frac{1}{\text{id}_Z}),$
- (iv) $Z = \text{dom } f,$ and
- (v) f is continuous on $A.$

Then $\int_A f(x)dx = ((\text{the function tan}) \cdot \frac{1}{\text{id}_Z})(\sup A) - ((\text{the function tan}) \cdot \frac{1}{\text{id}_Z})(\inf A).$

(34) Suppose $Z \subseteq \text{dom}((\text{the function tan}) \cdot \frac{1}{\text{id}_Z}).$ Then

- (i) $-(\text{the function tan}) \cdot \frac{1}{\text{id}_Z}$ is differentiable on $Z,$ and
- (ii) for every x such that $x \in Z$ holds $(-(\text{the function tan}) \cdot \frac{1}{\text{id}_Z})'_Z(x) = \frac{1}{x^2 \cdot (\text{the function cos})(\frac{1}{x})^2}.$

(35) Suppose that

- (i) $A \subseteq Z,$
- (ii) for every x such that $x \in Z$ holds $f(x) = \frac{1}{x^2 \cdot (\text{the function cos})(\frac{1}{x})^2},$
- (iii) $Z \subseteq \text{dom}((\text{the function tan}) \cdot \frac{1}{\text{id}_Z}),$
- (iv) $Z = \text{dom } f,$ and
- (v) f is continuous on $A.$

Then $\int_A f(x)dx = (-(\text{the function tan}) \cdot \frac{1}{\text{id}_Z})(\sup A) - (-(\text{the function tan}) \cdot \frac{1}{\text{id}_Z})(\inf A).$

(36) Suppose that

- (i) $A \subseteq Z,$
- (ii) for every x such that $x \in Z$ holds $f(x) = \frac{1}{x^2 \cdot (\text{the function sin})(\frac{1}{x})^2},$
- (iii) $Z \subseteq \text{dom}((\text{the function cot}) \cdot \frac{1}{\text{id}_Z}),$
- (iv) $Z = \text{dom } f,$ and
- (v) f is continuous on $A.$

Then $\int_A f(x)dx = ((\text{the function cot}) \cdot \frac{1}{\text{id}_Z})(\sup A) - ((\text{the function cot}) \cdot \frac{1}{\text{id}_Z})(\inf A).$

- (37) Suppose that $A \subseteq Z$ and for every x such that $x \in Z$ holds $f_1(x) = 1$ and $(\text{the function } \arctan)(x) > 0$ and $f = \frac{1}{(f_1 + \square^2) \cdot \text{the function } \arctan}$ and $Z \subseteq]-1, 1[$ and $Z \subseteq \text{dom}((\text{the function } \ln) \cdot (\text{the function } \arctan))$ and $Z = \text{dom } f$ and f is continuous on A . Then $\int_A f(x)dx = ((\text{the function } \ln) \cdot (\text{the function } \arctan))(\sup A) - ((\text{the function } \ln) \cdot (\text{the function } \arctan))(\inf A)$.
- (38) Suppose that $A \subseteq Z$ and $f = n \frac{(\square^{n-1}) \cdot \text{the function } \arctan}{f_1 + \square^2}$ and for every x such that $x \in Z$ holds $f_1(x) = 1$ and $Z \subseteq]-1, 1[$ and $Z \subseteq \text{dom}((\square^n) \cdot \text{the function } \arctan)$ and $Z = \text{dom } f$ and f is continuous on A . Then $\int_A f(x)dx = ((\square^n) \cdot \text{the function } \arctan)(\sup A) - ((\square^n) \cdot \text{the function } \arctan)(\inf A)$.
- (39) Suppose that $A \subseteq Z$ and for every x such that $x \in Z$ holds $f_1(x) = 1$ and $f = -n \frac{(\square^{n-1}) \cdot \text{the function } \text{arccot}}{f_1 + \square^2}$ and $Z \subseteq]-1, 1[$ and $Z \subseteq \text{dom}((\square^n) \cdot \text{the function } \text{arccot})$ and $Z = \text{dom } f$ and f is continuous on A . Then $\int_A f(x)dx = ((\square^n) \cdot \text{the function } \text{arccot})(\sup A) - ((\square^n) \cdot \text{the function } \text{arccot})(\inf A)$.
- (40) Suppose $Z \subseteq \text{dom}((\square^n) \cdot \text{the function } \text{arccot})$ and $Z \subseteq]-1, 1[$. Then
- (i) $-(\square^n) \cdot \text{the function } \text{arccot}$ is differentiable on Z , and
  - (ii) for every x such that $x \in Z$ holds $(-(\square^n) \cdot \text{the function } \text{arccot})'_|Z(x) = \frac{n \cdot (\text{the function } \text{arccot})(x)^{n-1}}{1+x^2}$.
- (41) Suppose that $A \subseteq Z$ and for every x such that $x \in Z$ holds $f_1(x) = 1$ and $f = n \frac{(\square^{n-1}) \cdot \text{the function } \text{arccot}}{f_1 + \square^2}$ and $Z \subseteq]-1, 1[$ and $Z \subseteq \text{dom}((\square^n) \cdot \text{the function } \text{arccot})$ and $Z = \text{dom } f$ and f is continuous on A . Then $\int_A f(x)dx = (-(\square^n) \cdot \text{the function } \text{arccot})(\sup A) - (-(\square^n) \cdot \text{the function } \text{arccot})(\inf A)$.
- (42) Suppose that $A \subseteq Z$ and for every x such that $x \in Z$ holds $f_1(x) = 1$ and $f = \frac{\text{the function } \arctan}{f_1 + \square^2}$ and $Z \subseteq]-1, 1[$ and $Z \subseteq \text{dom}((\square^2) \cdot \text{the function } \arctan)$ and $Z = \text{dom } f$ and f is continuous on A . Then $\int_A f(x)dx = (\frac{1}{2} ((\square^2) \cdot \text{the function } \arctan))(\sup A) - (\frac{1}{2} ((\square^2) \cdot \text{the function } \arctan))(\inf A)$.
- (43) Suppose that $A \subseteq Z$ and for every x such that $x \in Z$ holds $f_1(x) = 1$ and $f = -\frac{\text{the function } \text{arccot}}{f_1 + \square^2}$ and $Z \subseteq]-1, 1[$ and $Z \subseteq \text{dom}((\square^2) \cdot \text{the function } \text{arccot})$ and $Z = \text{dom } f$ and f is continuous on A . Then $\int_A f(x)dx = (\frac{1}{2} ((\square^2) \cdot \text{the function } \text{arccot}))(\sup A) - (\frac{1}{2} ((\square^2) \cdot \text{the function } \text{arccot}))(\inf A)$.
- (44) Suppose $Z \subseteq \text{dom}((\square^2) \cdot \text{the function } \text{arccot})$ and $Z \subseteq]-1, 1[$. Then
- (i) $-\frac{1}{2} ((\square^2) \cdot \text{the function } \text{arccot})$ is differentiable on Z , and
  - (ii) for every x such that $x \in Z$ holds

$$(-\frac{1}{2}((\square^2) \cdot \text{the function arccot}))'_{|Z}(x) = \frac{(\text{the function arccot})(x)}{1+x^2}.$$

- (45) Suppose that $A \subseteq Z$ and for every x such that $x \in Z$ holds $f_1(x) = 1$ and $f = \frac{\text{the function arccot}}{f_1 + \square^2}$ and $Z \subseteq]-1, 1[$ and $Z \subseteq \text{dom}((\square^2) \cdot \text{the function arccot})$ and $Z = \text{dom } f$ and f is continuous on A . Then $\int_A f(x)dx = (-\frac{1}{2}((\square^2) \cdot \text{the function arccot}))(\sup A) - (-\frac{1}{2}((\square^2) \cdot \text{the function arccot}))(\inf A)$.

- (46) Suppose that

- (i) $A \subseteq Z$,
- (ii) for every x such that $x \in Z$ holds $f_1(x) = 1$,
- (iii) $f = (\text{the function arctan}) + \frac{\text{id}_Z}{f_1 + \square^2}$,
- (iv) $Z \subseteq]-1, 1[$,
- (v) $Z = \text{dom } f$, and
- (vi) f is continuous on A .

$$\text{Then } \int_A f(x)dx = (\text{id}_Z \text{ the function arctan})(\sup A) - (\text{id}_Z \text{ the function arctan})(\inf A).$$

- (47) Suppose that

- (i) $A \subseteq Z$,
- (ii) for every x such that $x \in Z$ holds $f_1(x) = 1$,
- (iii) $f = (\text{the function arccot}) - \frac{\text{id}_Z}{f_1 + \square^2}$,
- (iv) $Z \subseteq]-1, 1[$,
- (v) $Z = \text{dom } f$, and
- (vi) f is continuous on A .

$$\text{Then } \int_A f(x)dx = (\text{id}_Z \text{ the function arccot})(\sup A) - (\text{id}_Z \text{ the function arccot})(\inf A).$$

- (48) Suppose that

- (i) $A \subseteq Z$,
- (ii) $Z \subseteq]-1, 1[$,
- (iii) $f = \frac{(\text{the function exp}) \cdot (\text{the function arctan})}{f_1 + \square^2}$,
- (iv) for every x such that $x \in Z$ holds $f_1(x) = 1$,
- (v) $Z = \text{dom } f$, and
- (vi) f is continuous on A .

$$\text{Then } \int_A f(x)dx = ((\text{the function exp}) \cdot (\text{the function arctan}))(\sup A) - ((\text{the function exp}) \cdot (\text{the function arctan}))(\inf A).$$

- (49) Suppose that

- (i) $A \subseteq Z$,
- (ii) $Z \subseteq]-1, 1[$,

- (iii) $f = -\frac{(\text{the function exp}) \cdot (\text{the function arccot})}{f_1 + \square^2}$,
- (iv) for every x such that $x \in Z$ holds $f_1(x) = 1$,
- (v) $Z = \text{dom } f$, and
- (vi) f is continuous on A .

Then $\int_A f(x)dx = ((\text{the function exp}) \cdot (\text{the function arccot}))(\sup A) - ((\text{the function exp}) \cdot (\text{the function arccot}))(\inf A)$.

- (50) Suppose $Z \subseteq \text{dom}((\text{the function exp}) \cdot (\text{the function arccot}))$ and $Z \subseteq]-1, 1[$. Then

- (i) $-(\text{the function exp}) \cdot (\text{the function arccot})$ is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $(-(\text{the function exp}) \cdot (\text{the function arccot}))'_|Z(x) = \frac{(\text{the function exp})(\text{the function arccot})(x))}{1+x^2}$.

- (51) Suppose that

- (i) $A \subseteq Z$,
- (ii) $Z \subseteq]-1, 1[$,
- (iii) $f = \frac{(\text{the function exp}) \cdot (\text{the function arccot})}{f_1 + \square^2}$,
- (iv) for every x such that $x \in Z$ holds $f_1(x) = 1$,
- (v) $Z = \text{dom } f$, and
- (vi) f is continuous on A .

Then $\int_A f(x)dx = (-(\text{the function exp}) \cdot (\text{the function arccot}))(\sup A) - (-(\text{the function exp}) \cdot (\text{the function arccot}))(\inf A)$.

- (52) Suppose that $A \subseteq Z \subseteq \text{dom}((\text{the function ln}) \cdot (f_1 + f_2))$ and $f = \frac{\text{id}_Z}{f_1 + f_2}$ and $f_2 = \square^2$ and for every x such that $x \in Z$ holds $f_1(x) = 1$ and $Z = \text{dom } f$ and f is continuous on A . Then $\int_A f(x)dx = (\frac{1}{2}((\text{the function ln}) \cdot (f_1 + f_2)))(\sup A) - (\frac{1}{2}((\text{the function ln}) \cdot (f_1 + f_2)))(\inf A)$.

- (53) Suppose that $A \subseteq Z \subseteq \text{dom}((\text{the function ln}) \cdot (f_1 + f_2))$ and $f = \frac{\text{id}_Z}{a(f_1 + f_2)}$ and for every x such that $x \in Z$ holds $h(x) = \frac{x}{a}$ and $f_1(x) = 1$ and $a \neq 0$ and $f_2 = (\square^2) \cdot h$ and $Z = \text{dom } f$ and f is continuous on A . Then $\int_A f(x)dx = (\frac{a}{2}((\text{the function ln}) \cdot (f_1 + f_2)))(\sup A) - (\frac{a}{2}((\text{the function ln}) \cdot (f_1 + f_2)))(\inf A)$.

- (54) Suppose $Z \subseteq \text{dom}(\frac{1}{\text{id}_Z} \text{ the function arctan})$ and $Z \subseteq]-1, 1[$. Then

- (i) $-\frac{1}{\text{id}_Z} \text{ the function arctan}$ is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $(-\frac{1}{\text{id}_Z} \text{ the function arctan})'_|Z(x) = \frac{(\text{the function arctan})(x)}{x^2} - \frac{1}{x \cdot (1+x^2)}$.

- (55) Suppose $Z \subseteq \text{dom}(\frac{1}{\text{id}_Z} \text{ the function arccot})$ and $Z \subseteq]-1, 1[$. Then

- (i) $-\frac{1}{\text{id}_Z} \text{ the function arccot}$ is differentiable on Z , and

- (ii) for every x such that $x \in Z$ holds $(-\frac{1}{\text{id}_Z} \text{ the function arccot})'_|Z(x) = \frac{(\text{the function arccot})(x)}{x^2} + \frac{1}{x \cdot (1+x^2)}$.
- (56) Suppose that $A \subseteq Z$ and for every x such that $x \in Z$ holds $f_1(x) = 1$ and $f = \frac{\text{the function arctan}}{\square^2} - \frac{1}{\text{id}_Z (f_1 + \square^2)}$ and $Z \subseteq \text{dom}(\frac{1}{\text{id}_Z} \text{ the function arctan})$ and $Z \subseteq]-1, 1[$ and $Z = \text{dom } f$ and f is continuous on A . Then $\int_A f(x)dx = (-\frac{1}{\text{id}_Z} \text{ the function arctan})(\sup A) - (-\frac{1}{\text{id}_Z} \text{ the function arctan})(\inf A)$.
- (57) Suppose that $A \subseteq Z$ and for every x such that $x \in Z$ holds $f_1(x) = 1$ and $f = \frac{\text{the function arccot}}{\square^2} + \frac{1}{\text{id}_Z (f_1 + \square^2)}$ and $Z \subseteq \text{dom}(\frac{1}{\text{id}_Z} \text{ the function arccot})$ and $Z \subseteq]-1, 1[$ and $Z = \text{dom } f$ and f is continuous on A . Then $\int_A f(x)dx = (-\frac{1}{\text{id}_Z} \text{ the function arccot})(\sup A) - (-\frac{1}{\text{id}_Z} \text{ the function arccot})(\inf A)$.

REFERENCES

- [1] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [2] Noboru Endou and Artur Kornilowicz. The definition of the Riemann definite integral and some related lemmas. *Formalized Mathematics*, 8(1):93–102, 1999.
- [3] Noboru Endou, Katsumi Wasaki, and Yasunari Shidama. Definition of integrability for partial functions from $\mathbb{R}$ to $\mathbb{R}$ and integrability for continuous functions. *Formalized Mathematics*, 9(2):281–284, 2001.
- [4] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [5] Jarosław Kotowicz. Convergent real sequences. Upper and lower bound of sets of real numbers. *Formalized Mathematics*, 1(3):477–481, 1990.
- [6] Jarosław Kotowicz. Partial functions from a domain to a domain. *Formalized Mathematics*, 1(4):697–702, 1990.
- [7] Jarosław Kotowicz. Partial functions from a domain to the set of real numbers. *Formalized Mathematics*, 1(4):703–709, 1990.
- [8] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(2):269–272, 1990.
- [9] Jarosław Kotowicz. The limit of a real function at infinity. *Formalized Mathematics*, 2(1):17–28, 1991.
- [10] Xiquan Liang and Bing Xie. Inverse trigonometric functions arctan and arccot. *Formalized Mathematics*, 16(2):147–158, 2008, doi:10.2478/v10037-008-0021-3.
- [11] Konrad Raczkowski. Integer and rational exponents. *Formalized Mathematics*, 2(1):125–130, 1991.
- [12] Konrad Raczkowski and Paweł Sadowski. Real function continuity. *Formalized Mathematics*, 1(4):787–791, 1990.
- [13] Konrad Raczkowski and Paweł Sadowski. Real function differentiability. *Formalized Mathematics*, 1(4):797–801, 1990.
- [14] Konrad Raczkowski and Paweł Sadowski. Topological properties of subsets in real numbers. *Formalized Mathematics*, 1(4):777–780, 1990.
- [15] Yasunari Shidama. The Taylor expansions. *Formalized Mathematics*, 12(2):195–200, 2004.
- [16] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers. *Formalized Mathematics*, 1(3):445–449, 1990.
- [17] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [18] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.

- [19] Yuguang Yang and Yasunari Shidama. Trigonometric functions and existence of circle ratio. *Formalized Mathematics*, 7(**2**):255–263, 1998.

Received February 4, 2010