SYNTAXONOMIC AND NOMENCLATURAL NOTES ON THE SCREE VEGETATION OF CAUCASUS

Elena A. BELONOVSKAYA¹,
Ladislav MUCINA^{2*} & Jean-Paul THEURILLAT³

Abstract

We present description of a new class for the high-altitude screes of the Caucasus (the *Lamio tomentosi-Chaerophylletea humilis*) and two alliances such as the *Scrophulario minimae-Symphylomion graveolentis* (*Chaerophylletalia humilis*, *Lamio tomentosi-Chaerophylletea humilis*) and the *Scrophulario variegatae-Epilobion dodonaei* (*Epilobietalia fleischeri*, *Thlaspietalia rotundifolia*).

Key words: Georgia, alpine vegetation, Russian federation, riverine terraces, syntaxonomy.

Izvešček

V članku predstavljamo opis novega razreda na visoko ležečih meliščih na Kavkazu (*Lamio tomentosi-Lamietea tomentosi*) in dveh zvez – *Scrophulario minimae-Symphylomion graveolentis* (*Chaerophylletalia humilis*, *Lamio tomentosi-Chaerophylletea humilis*) in *Scrophulario variegatae-Epilobion dodonaei* (*Epilobietalia fleischeri*, *Thlaspietalia rotundifolia*).

Ključne besede: Gruzija, alpinska vegetacija, Ruska federacija, rečne terase, sintaksonomija.

1. INTRODUCTION

The communities of the *Thlaspietalia rotundifolia* (Braun-Blanquet 1948, Zollitsch 1966, Englisch et al. 1993, Valachovič et al. 1997 etc.) are a common appearance in scree habitats of temperate, boreal and subarctic zones of Europe, where they could be found in all vegetation altitudinal belts, from the planare and colline belts as high as the alpine and subnival belts. In the mediterranean-climate regions of the Iberian Peninsula they are replaced by the *Phagnalo-Rumicetea indurati* (Rivas Goday et Esteve 1972) Rivas-Martínez, Izco et Costa 1973 (e.g. Ortiz & Puglar 2000, Rivas-Martínez et al. 2011) and in the (sub)mediterranean regions of the Central and Eastern Mediterranean, they

give way to the *Drypidetalia spinosi* which, in our syntaxonomic concept, also includes the *Scrophulario-Helichrysetea italici* S. Brullo et al. 1998 (Brullo et al. 1998) and the *Onosmo polyphyllae-Ptilostemonetalia* Korzhenevsky 1990 (see Golub et al. 2011 for the latest syntaxonomic synthesis).

In Caucasus, Onipchenko (2002a: Table 3.1) described the *Chaerophylletalia humilis* and placed this order within the *Thlaspietalia rotundifolia*, although on page 18 he also noted that practically all diagnostic species of the class are missing in the Caucasian communities. Onipchenko (l.c.) also suggested that there were some diagnostic species of the European alliances and orders and therefore he decided to classify his new order (*Chaerophylletalia humilis*) within the *Thlaspietalia*

¹ Institute of Geography, Russian Academy of Sciences, Staromonetny 2, 109017 Moskva, Russian Federation; E-mail: belena53@mail.ru

² Iluka Chair in Vegetation Science & Biogeography, School of Plant Biology, The University of Western Australia, 35 Stirling Highway, Crawley WA 6009, Australia; E-mail: Laco.Mucina@uwa.edu.au

³ Centre alpin de Phytogéographie, Fondation J.-M. Aubert, Case postale 71, CH-1938 Champex-Lac, Switzerland; Laboratoire de Biogéographie, Section de Biologie, Université de Genève, 1, ch. de l'Impératrice, Case postale 60, CH-1292 Chambésy, Switzerland; E-mail: jean-paul.theurillat@unige.ch

* corresponding author

rotundifolii. In fact only *Oxyria digyna*, *Cryptogramma crispa* and *Epilobium dodonaei* would qualify at the latter group. Interestingly, these species occur in the Caucasian scree communities at lower altitudes (perhaps (sub)montane to subalpine zone). The high altitudes (alpine to subnivale zones) of the Caucasus range are home to endemic-rich scree communities that appear to be floristically very different from the *Thlaspietea rotundifolii* of Europe (see also Nakhutsrishvili & Gagnidze 1999; Nakhutsrishvili 2003).

Belonovskaya (2012; based on earlier paper by Belonovskaya & Korotkov 2002) developed the Onipchenko's (2002) remark on the poor floristic relationship of the Caucasus scree communities and those of Europe further and introduced a new syntaxonomic concept – the *Veronico telephiifoliae-Lamietea tomentosi*, including one new order (*Veronico telephiifoliae-Lamietalia tomentosi*), with two new alliances (*Drabo scabrae-Eunomion rotundifoliae*, *Scrophulario minimae-Symphylomion graveolentis*). Belonovskaya's (2012) paper however have not engaged fully with the syntaxonomy proposed by Onipchenko (2002) and all the suggested new syntaxonomic units (class, order and the alliances) are rendered either invalid according to the International Code Phytosociological Nomenclature; Weber et al. 2000) or (in one case) should be considered synonymous with an earlier syntaxonomic concept of Onipchenko (2002).

The prime aim of this paper is to validate the syntaxon names (or introduce a new one) and contribute to clarification of the syntaxonomic relationship of the Onipchenko's syntaxa and those suggested by Belonovskaya (l.c.) in the light of the new syntaxonomic-nomenclatural synthesis undertaken within so called 'EuroVeg-Checklist' (Mucina et al. in prep.).

2. NOMENCLATURE OF THE NAME 'VERONICO TELEPHIIFOLIAE-LAMIEA TOMENTOSI'

We agree that the endemic-rich, high-altitude scree communities of the Caucasus should be considered as a syntaxonomic concept different from the *Thlaspietea rotundifolii*. However, name '*Veronico telephiifoliae-Lamietea tomentosi*' (coined to accommodate these high-altitude Caucasian scree communities) has been invalidly (Art. 2b of the ICPN) published in Belonovskaya (2012:

967) because the name of the unique order – the '*Veronico telephiifoliae-Lamietalia tomentosi*' – include in the original diagnosis of the new class, was invalidly published at the time of the publication of the class name.

Indeed, the original diagnosis of the *Veronico telephiifoliae-Lamietalia tomentosi* contains two new alliances: the *Drabo scabrae-Eunomion rotundifoliae* and *Scrophulario minimae-Symphylomion graveolentis*. There are four new associations in the original diagnosis of the *Drabo scabrae-Eunomion rotundifoliae* and three new associations in the *Scrophulario minimae-Symphylomion graveolentis* (Belonovskaya 2012). All these seven associations are validly published because their original diagnosis contains only the type relevé (in Russian) presented in the Table 1 in Belonovskaya (2012) and each type relevé containing the name-giving taxa of respective association. On the other hand, both names of the new alliances are invalidly published because their type, indicated in the bottom of the Table 1 is given in Russian not matching the demand of the Art. 5 of the ICPN (Weber et al. 2000) that requires (*expressis verbis*) using the Latin terms *typus* or *holotypus*. Because of this technicality, the original diagnosis of the '*Veronico telephiifoliae-Lamietalia tomentosi*' containing no validly published alliances, the name remains invalidly published (Arts. 2b and 8 of the ICPN).

Besides, even if the new alliance *Drabo scabrae-Eunomion rotundifoliae* (Belonovskaya 2012) was validly published, the latter alliance name would have become automatically a superfluous name for the validly published *Chaerophyllion humilis* Onipchenko 2002 which was listed by Belonovskaya (2012) in the synonymy of the *Drabo scabrae-Eunomion rotundifoliae* on p. 968. Similarly, if the new order *Veronico telephiifoliae-Lamietalia tomentosi* were validly published, it would have been a homotypical synonym for the validly published *Chaerophylletalia humilis* Onipchenko 2002. It appears that syntaxonomically this order matches the Belonovskaya's '*Veronico telephiifoliae-Lamietalia tomentosi*'. Onipchenko (2002a: 168) listed both name-giving taxa (*Lamium tomentosum* and *Veronica telephiifolia*; syn. *V. minuta* C.A. Meyer; see Fig. 1) in the diagnosis of *Chaerophylletalia humilis*.

In order to avoid confusion in future, we introduce a new name for the invalid syntaxonomic concept of the '*Veronico telephiifoliae-Lamietea tomentosi*' and by recognising the role of *Chae-*

Figure 1: Selected character species of the *Lamio tomentosum*-*Chaerophylletea humilis*: A: *Lamium tomentosum*, B: *Veronica minuta*, C: *Chaerophyllum humile*. Broad valley of the Tergi (Terek) River near the town of Stepantsminda in Caucasus, Republic of Georgia with extensive terraces supporting communities of the *Murbeckiello huetii*-*Epilobion dodonaei* (D); *Epilobium dodonaei* photographed on the alluvial terraces of the Tergi (Terek) River, Georgia. Photo credits: A-C photos is Mikhail Dokukin (Nalchik, Kabardino-Balkaria, Russia), D-E: L. Mucina.

Slika 1: Izbrane značilnice razreda *Lamio tomentosum*-*Chaerophylletea humilis*: A: *Lamium tomentosum*, B: *Veronica minuta*, C: *Chaerophyllum humile*. Široka dolina reke Tergi (Terek) River v bližini mesta Stepantsminda na Kavkazu, Gruzija z ekstenzivnimi terasami, kjer se pojavljajo združbe zveze *Murbeckiello huetii*-*Epilobion dodonaei* (D); *Epilobium dodonaei* je fotografiran na aluvialni terasi reke Tergi (Terek), Gruzija. Foto: A-C Mikhail Dokukin (Nalchik, Kabardino-Balkaria, Russia), D-E: L. Mucina.

rophyllyum humilis as one of the local dominating species, we coin the name and concept of the *Lamio tomentosi-Chaerophylletea humilis* (for the full protologue see below).

3. THE CASE OF *MURBECKIELLION HUETII* ONIPCHENKO 2002

The concept and name *Murbeckiellion huetii* was introduced for the first time by Onipchenko (2002a) who classified two associations within this alliance – the *Hyalopoo ponticae-Oxyrietum digynae* Onipchenko 2002 from siliceous, snow-rich alpine moraine habitats and the *Scrophulario variegatae-Epilobietum dodonaei* which was designated as the *typus* of the alliance. The latter association, however, was not validly described because the *typus* of the association has not been assigned. The designation of the *typus* of the *Scrophulario variegatae-Epilobietum dodonaei* was attempted by Onipchenko (2002b) who selected (relevé) ‘26/95, Table 3.3’ in Onipchenko (2002a) as the *typus*. This attempt was, however, invalid since there are two relevés designated as 26/95: 0 26/95 and 1 26/95 in Table 3.3. Because the choice of the *typus* was not unequivocal, this validation cannot be recognised, hence both the *Scrophulario variegatae-Epilobietum dodonaei* and the *Murbeckiellion huetii* remain invalidly described.

The *Murbeckiellion huetii* is a syntaxonomically heterogeneous concept. The *Hyalopoo ponticae-Oxyrietum digynae* Onipchenko 2002 is floristically (and ecologically) closer to the *Hyalopoo ponticae* Rabotnova et Onipchenko in Onipchenko 2002 than any taxonomic unit of the *Lamio tomentosi-Chaerophylletea humilis*. The *Scrophulario variegatae-Epilobietum dodonaei*, on the other hand should be classified with the *Epilobietalia fleischeri*; it is characteristic of gravel alluvial banks of mountain streams and *Epilobium dodonaei* plays a major role in these community (see Fig. 1A-E). It is therefore not our intention to validate the *Murbeckiellion huetii*, but rather coin a new name that would better represent the vegetation of gravel terraces of Caucasus – the *Murbeckiello huetii-Epilobion dodonaei* (for the protologue see below).

The low-altitude *Murbeckiello huetii-Epilobion dodonaei* is classified here into the *Epilobietalia fleischeri* Moor 1958 nom. cons. propos. (Raus et al. in prep.) of the *Thlaspietea rotundifolia*.

4. DESCRIPTION OF THE NEW SYNTAXA AND THE NEW SYNTAXONOMIC SCHEME FOR THE CAUCASIAN SCREE VEGETATION

***Lamio tomentosi-Chaerophylletea humilis* Belonovskaya, Mucina et Theurillat classis nova hoc loco**
Holotypus: *Chaerophylletalia humilis* Onipchenko 2002 (Onipchenko 2002a: 18, 21)

Synonym: *Veronico telephiiifoliae-Lamietea tomentosi* Belonovskaya 2012 (Arts. 2b & 8 of the ICPN)
Diagnostic taxa: *Cerastium multiflorum*, *Corydalis alpestris*, *Jacobaea erucifolia* subsp. *arenaria*, *Jurinea humilis*, *Lamium tomentosum*, *Veronica telephiiifolia*

***Chaerophylletalia humilis* Onipchenko 2002**
Synonym: *Veronico telephiiifoliae-Lamietalia tomentosi* Belonovskaya 2012 (Arts. 2b & 8 of the ICPN)

***Chaerophyllion humilis* Onipchenko 2002**
Synonym: *Drabo scabrae-Eunomion rotundifoliae* Belonovskaya 2012 (Arts. 5 & 8 of the ICPN)

Diagnostic taxa: *Cerastium alpinum*, *Chaerophyllum humile*, *Delphinium caucasicum*, *Draba scabra*, *Eunomia rotundifolia*, *Potentilla gelida*, *Saxifraga moschata*, *Tephrosia integrifolia*, *Veronica minuta*

Veronico minutae-Chaerophylletum humilis Onipchenko 2002

Cruciato tauricae-Chaerophylletum humilis Belonovskaya 2012

Saxifraga sibirica-Alopecuretum sericeae Belonovskaya 2012

Myosotido alpestris-Potentilletum gelidae Belonovskaya 2012

***Scrophulario minimae-Symphylolomion graveolentis* Belonovskaya, Mucina et Theurillat all. nova hoc loco**

Holotypus: *Cruciato tauricae-Pseudovesicarietum digitatae* Belonovskaya 2012

Diagnostic taxa: *Cruciata taurica*, *Scrophularia minima*, *Symphyloloma graveolens*

Cruciato tauricae-Pseudovesicarietum digitatae Belonovskaya 2012

Ranunculetum arachnoidei Belonovskaya 2012

Anthemidetum creticae Belonovskaya 2012

Cerastietum alpini Belonovskaya 2012

***Thlaspietea rotundifolii* Br.-Bl. 1948**

***Epilobietalia fleischeri* Moor 1958 nom. cons. propos.**

***Murbeckiello huetii-Epilobion dodonaei* Belonovskaya, Mucina et Theurillat all. nova hoc loco**

Holotypus: *Scrophulario variegatae-Epilobietum dodonaei* Onipchenko ex Belonovskaya, Mucina et Theurillat ass. nova hoc loco; Holotypus: Onipchenko 2002a: Table 3.3, relevé 1 10 94

Synonym: *Murbeckiellion huetii* Onipchenko 2002 nom. inval. (Arts. 5 & 8 of the ICPN)

Diagnostic species: *Epilobium dodonaei*, *Murbeckiella huetii*

5. ACKNOWLEDGEMENTS

We thank Thomas Raus (Berlin, Germany) for checking the nomenclature and taxonomy of the diagnostic species mentioned in our paper. Marie Lochman (Perth, Australia) kindly scanned slides and Mikhail Dokukin (Nalchik, Russia) for the photos used in Figure 1. Zaza Kikvidze identified the locality in the Tergi River valley. LM thanks Gia Nakhutrishvili (Tbilisi, Georgia) and Georg Grabherr (Vienna, Austria) for making the visit to Caucasus in 1990 possible.

6. REFERENCES

- Belonovskaya, E. A. 2012: Klassifikatsiya rastitel'nykh gruppировок na osypakh v alpiiskom noyase Severnogo Kavkaza. (Classification of plant communities on screes in the alpine belt of the Northern Caucasus.) Izvestiya Samar-skogo Nauchnogo Tsentra Rossiiskoi Akademii Nauk 14: 967–970. (in Russian)
- Belonovskaya, E.A. & Korotkov, K.O. 2002: Raznoobrazi'e rastitel'nosti alpiiskogo poyasa Bol'shogo Kavkaza. (Alpine vegetation diversity on the Greater Caucasus.) Izvestiya Rossiiskoi Akademii Nauk, Seria geograficheskaya. N2: 89–96. (in Russian)
- Braun-Blanquet, J. 1948: La végétation alpine des Pyrénées orientales. Consejo Superior de Investigaciones Científicas, Barcelona, ES.
- Brullo, S., Scelsi, F. & Spampinato, G. 1998: Considerazioni sintassonomiche sulla vegetazione perenne pioniera dei substrati incoerenti dell'Italia meridionale e Sicilia. Itinera Geobotanica 11: 403–424.
- Englisch, T., Valachovič, M., Mucina, L., Grabherr, G. & Ellmauer, T. 1993: *Thlaspietea rotundifolii*. In: Grabherr, G. & Mucina, L. (eds.) Die Pflanzengesellschaften Österreichs. Teil I, pp. 276–342. G. Fischer Verlag, Jena,
- Golub, V.B., Greshushkina, N.A., Sorokin, A.N. & Nikolaichuk, L.F. 2011: Rastitel'nye soobshchestva klassa *Onosmo polyphyllae-Ptilosmetea* Korzhenevsky 1990 na territorii Chernomorskogo poberezhya Kavkaza i Krymskogo poluoostrova. (Vegetation of the class of *Onosmo polyphyllae-Ptilosmetea* Korzhenevsky 1990 on territory of Black Sea coast of Caucasus and of the Crimean Peninsula.) Rastitel'nost' Rossii 17–18: 3–16. (in Russian)
- Nakhutsrishvili, G. 2003: High mountain vegetation of the Caucasus region. In: Nagy, L., Grabherr, G., C. Körner & Thompson, D.B.A. (eds.) Alpine biodiversity in Europe. pp. 93–103. Springer, Heidelberg.
- Nakhutsrishvili, G. & Gagnidze, R. 1999: Die subnivale und nivale Hochgebirgsvegetation des Kaukasus. Phytocoenosis 11: 173–183.
- Onipchenko, V. G. 2002a: Alpine vegetation of the Teberda Reserve, the Northwestern Caucasus. Veröffentlichungen des Geobotanischen Institutes ETH Stiftung Rübel Zürich 130: 1–168.
- Onipchenko, V. G. 2002b: Tipifikatsiya nekotorikh assotsiatsii skalno-osypnoi rastitel'nosti Severo-Zapadnogo Kavkaza. (Typification of some associations of rock and scree vegetation of North-Western Caucasus.) Biulletin Moskovskogo Obshchestva Ispytatelei Prirody, Otdel Biologicheskii 107: 88–89. (in Russian)
- Ortiz, S. & Pulgar, I. 2000: As comunidades eurosiberianas da Clase *Phagnalio-Rumicetea indurati* en Galicia e Norte de Portugal (NO da Península Iberica). Nova Acta Cientifica Compostelana 10: 35–41.
- Rivas-Martínez, S. et coautores 2011: Mapa de series, geoseries y geopermaseries de vegetación de España. (Memoria del mapa de vegetación potencial de España). Parte II). Itinera Geobotanica N.S. 18: 5–424.
- Valachovič, M., Dierssen, K., Dimopoulos, P., Hadač, E., Loidi, J., Mucina, L., Rossi, G., Valle Tendero, F. & Tomaselli, M. 1997: The

vegetation on screes—a synopsis of higher syntaxa in Europe. *Folia Geobotanica et Phytotaxonomica* 32: 173–192.

Weber, H.E., Moravec, J. & Theurillat, J.-P. 2000: International Code of Phytosociological Nomenclature. 3rd edition. *Journal of Vegetation Science* 11: 739–768.

Zollitsch, B. 1966: Soziologische und Ökologische Untersuchungen auf Kalkschiefern in hochalpinen Gebieten. Teil 1: Die Steinschuttesellschaften der Alpen unter besonderer Berücksichtigung der Gesellschaften auf Kalkschiefern in den mittleren und östlichen Zentralalpen. *Berichte der Bayerischen Botanischen Gesellschaft* 40: 1–38.

Received 2.12.2013

Revision received: 6.2.2014

Accepted 6.2.2014