Helminth fauna of the Eurasian Marsh Frog, *Pelophylax ridibundus* (Pallas, 1771) (Anura: Ranidae), collected from Denizli Province, Inner-West Anatolia Region, Turkey

S. DÜŞEN^{1*}, M. ÖZ²

^{1*}Pamukkale University, Faculty of Arts and Sciences, Department of Biology, Kinikli Campus, 20017, Denizli, Turkey, E-mail: sdusen@pamukkale.edu.tr, serdar2290@yahoo.com; ²Akdeniz University Faculty of Science, Department of Biology, 07058, Antalya, Turkey

Summary

In this research, a total of 298 Eurasian marsh frogs, *Pelophylax ridibundus* (Pallas, 1771) were collected from different localities in Denizli province (Inner-west Anatolia Region, the eastern part of Aegean Region) Turkey between 2006 and 2009 and examined for helminths. Of 262 (87.91 %) *Pelophylax ridibundus* samples were infected with one or more helminths. *Pelophylax ridibundus* harbored eight species of digeneans (*Diplodiscus subclavatus*, *Gorgoderina vitelliloba*, *Gorgoderina cygnoides*, *Pleurogenoides medians*, *Prostotocus confusus*, *Skrjabinoeces breviansa*, *Encyclometra colubrimurorum* and *Ophistoglyphe ranae*), one species of cestode (*Nematotaenia dispar*), two species of acanthocephalans (*Acanthocephalus ranae* and *Pomphorhynchus laevis*), and six species of nematodes (*Rhabdias bufonis*, *Oswaldocruzia filiformis*, *Cosmocerca ornata*, *Oxysomatium brevicaudatum*, *Eustrongylides* sp. and *Abbreviata* sp.). *P. ridibundus* represents a host record for *Nematotaenia dispar* in Turkey.

Keywords: Amphibians, Denizli, Helminth, *Pelophylax ridibundus*, Turkey

Introduction

The Eurasian marsh frog, *Pelophylax ridibundus* (formerly known as *Rana ridibunda*), is a medium-sized semi-aquatic anuran species. *P. ridibundus* inhabits lakes, pools, or slowly flowing streams. It is a highly opportunistic diurnal amphibian, living in mixed and deciduous forests, forest steppe, and steppe and other grasslands, semi-desert and desert zones. Arid areas are largely colonized through river valleys and channels. *P. ridibundus* prefers open, well-warmed areas with abundant herbaceous vegetation (IUCN, 2012). In Turkey, this species is known in all suitable habitats except for a portion of the Lakes District (Baran & Atatür, 1997; Budak & Göçmen, 2008).

To our knowledge, the first helminthological study in Eurasian marsh frog, *P. ridibundus* (formerly known as *Rana ridibunda*) was reported by Schad *et al.* (1960) and they recorded only two nematodes in this species from Turkey. Other helminthological studies on *P. ridibundus* published by Saygı and Başibüyük (1990), Oğuz *et al.* (1994), Yıldırımhan *et al.* (1996, 1997, 2005a), Kır *et al.* (2001), Düşen and Öz (2006), Sağlam and Arıkan (2006), Unal *et al.* (2007), Düşen and Oğuz (2008, 2010), Düşen *et al.* (2010a), Heckmann *et al.* (2010, 2011) in Turkey.

So far, there has been no published detailed study on helminths of Eurasian marsh frog, *P. ridibundus* from Denizli province (Inner-west Anatolia Region - the eastern part of Aegean Region) in Turkey. This is the first detailed helminthological study, which has been done in this geographic area.

Materials and methods

Frog samples were collected by hand and dip net, between 2006 – 2009 from 17 different localities in Denizli province (38°29' – 38°52' N – 28°38' – 30°05' E). In total, 298 *P. ridibundus* (197♂♂, 101♀♀) were examined for helminth parasites. The mean ± SD snout-vent length (SVL) of specimens was 66.34 ± 14.91 mm, with a range from 29 to 101 mm; within 24 hr, toads were overdosed in ether-filled glass containers.

The samples body cavity was opened by a longitudinal ventral incision. The alimentary canal was excised and separated into stomach, small intestine, large intestine and rectum. The contents of each part and other organs (lungs, liver, gall bladder, kidneys and urinary bladder) were each mixed with 0.5 % saline solution and poured into petri dishes for examination under a stereomicroscope. The muscles, plus portions of peritoneum and spinal cord, were teased out with needles and examined under a stereomicro-

scope. Digeneans was immobilized by coverslip pressure in 70 % ethyl alcohol, fixed, and stored in 70 % ethanol. Nematodes was straightened by heat, fixed, and stored in 70 % ethyl alcohol with 5 % glycerol. Acanthocephalans were relaxed in saline and heat-fixed under slight coverslip pressure in warm ethyl alcohol-formalin-acetic acid. Digeneans, acanthocephalans and cestode samples were stained with acetocarmine, dehydrated, cleared in cedar oil or xylol, and mounted in Canada Balsam; nematodes were cleared in glycerol and examined. Intensities are presented as mean values (± 1 SE) followed by the range. Voucher host specimens and parasite specimens were deposited in Pamukkale University, Faculty of Sciences and Arts, Department of Biology, Denizli, Turkey (PAU-HELM-1-17/2009).

Results and discussion

Two-hundred and ninety eight *P. ridibundus* (197♂♂, 101♀♀) samples were collected between 2006 – 2009 years from Denizli province.

Trematoda

Family: Diplodiscidae

Diplodiscus subclavatus (Pallas, 1760) Diesing, 1836

Prevalence, intensity and range: Hosts infected, 31 of 298 (10.40 %, 4.03 ± 0.65 SE, 1 – 15).

Other reported hosts: *D. subclavatus* was observed in, *Triturus cristatus*, *T. vulgaris*, *Bufo viridis*, *Hyla arborea* (Shimalov *et al.*, 2001; Vojtková & Vojtek 1975, Buchvarov 1977, Kuc & Sulgostowska 1988b, Sey 1991); *H. savignyi* (Yıldırımhan *et al.*, 2012); *Rana temporaria*, *R. dalmatina*, *R. esculenta*, *R. graeca* (Vojtková & Vojtek 1975; Buchvarov, 1977; Kuc & Sulgostowska, 1988b; Sey, 1991; Düşen *et al.*, 2009), *Bombina bombina*, *Bo. variegata* (Vojtková & Vojtek, 1975; Grabda-Kazubska & Lewin, 1989), *Pelobates syriacus balcanicus* (Buchvarov, 1977), *T. alpestris*, *P. fuscus*, *R. lessonae* (Vojtková & Vojtek, 1975), *B. bufo*, *R. arvalis* (Vojtková & Vojtek, 1975; Sey 1991; Shimalov & Shimalov, 2001), *R. holtzi* (Topçu, 2002; Yıldırımhan *et al.*, 2006b), *Natrix natrix* (Sey, 1991; Buchvarov *et al.*, 2000; Shimalov & Shimalov, 2000), *N. tessellata* (Buchvarov *et al.*, 2000), *Vipera berus* (Shimalov & Shimalov, 2000) and *Esox lucius* (Öztürk *et al.*, 2000).

Geographic range: Europe (Yamaguti, 1958), Czech Republic (Vojtková & Vojtek, 1975), Bulgaria (Buchvarov *et al.*, 1975; Buchvarov, 1977), Poland (Kuc & Sulgostowska, 1988a, 1988b; Grabda-Kazubska & Lewin, 1989), Hungary (Sey, 1991), Turkey (Oğuz *et al.*, 1994), Iran (Masshaii *et al.*, 2000), Uzbekistan (Vashetko & Siddikov, 1999).

Specimens deposited: PAU-HELM-1/2009

Family: Gorgoderidae

Gorgoderina vitelliloba (Olsson, 1876) Loos, 1902

Prevalence, intensity and range: Hosts infected, 20 of 298 (6.71 %, 2.55 ± 0.81 SE, 1 – 13).

Other reported hosts: *G. vitelliloba* was reported in, *Bombinator igneus* (Yamaguti, 1958), *Bo. bombina* (Vojtková

& Vojtek, 1975), *Bo. variegata*, (Yamaguti, 1958; Vojtková & Vojtek, 1975; Buchvarov, 1977) *B. bufo* and *B. calamita* (Cedhagen, 1988), *B. viridis* (Buchvarov, 1977), *R. arvalis* (Yamaguti, 1958; Vojtková & Vojtek, 1975), *R. dalmatina* (Buchvarov, 1977), *R. esculenta*, (Vojtková & Vojtek, 1975), *R. temporaria* (Yamaguti, 1958; Vojtková & Vojtek, 1975; Buchvarov, 1977; Cedhagen, 1988), *R. macronemis* (Yıldırımhan *et al.*, 1997b; Yıldırımhan *et al.*, 2006a), *R. camerani* (Yıldırımhan *et al.*, 2006a) and *R. holtzi* (Yıldırımhan *et al.*, 2006b).

Geographic range: Europe and North America (Yamaguti, 1958), Czech Republic (Vojtková & Vojtek, 1975), Bulgaria (Buchvarov, 1977), Poland (Kuc & Sulgostowska, 1988a, 1988b), Turkey (Yıldırımhan *et al.*, 1996; Düşen & Öz, 2006).

Specimens deposited: PAU-HELM-2/2009

Family: Gorgoderidae

Gorgodera cygnoides (Zeder, 1800)

Prevalence, intensity and range: Hosts infected, 36 of 298 (12.08 %, 3.83 ± 0.70 SE, 1 – 17).

Other reported hosts: *G. cygnoides* is a common parasite of amphibians, *Bombinator igneus* (Yamaguti, 1958), *Bo. Bombina* and *Bo. variegata* (Vojtková & Vojtek, 1975), *B. viridis* (Buchvarov, 1977; Shimalov & Shimalov, 2001), *H. arborea* (Yamaguti, 1958), *R. arvalis* (Vojtková & Vojtek, 1975), *R. dalmatina* (Buchvarov, 1983), *R. esculenta* (Yamaguti, 1958; Vojtková & Vojtek, 1975; Buchvarov, 1977; Kuc & Sulgostowska, 1988b), *R. lessonae* (Vojtková & Vojtek, 1975), *R. clamitans* (Yamaguti, 1958), *R. temporaria* (Yamaguti, 1958; Vojtková & Vojtek, 1975), *R. camerani* (Yıldırımhan *et al.*, 2006a), *R. holtzi* (Yıldırımhan *et al.*, 2006b), *R. kurtmuelleri* (Hristovski *et al.*, 2006) and *Pelodytes caucasicus* (Yıldırımhan *et al.*, 2009).

Geographic range: Western Europe (Yamaguti, 1958) Turkey (Yıldırımhan *et al.*, 1996; Düşen & Öz, 2006).

Specimens deposited: PAU-HELM-3/2009

Family: Lecithodendriidae

Pleurogenoides medians (Olsson, 1876) Travassos, 1921

Prevalence, intensity and range: Hosts infected, 47 of 298 (15.77 %, 27.70 ± 4.57 SE, 1 – 126).

Other reported hosts: *P. medians* has been reported in various amphibians and reptiles species, *T. cristatus* (Shimalov *et al.*, 2001), *T. vulgaris* (Vojtková & Vojtek, 1975; Shimalov *et al.*, 2001); *Bo. bombina*, (Vojtková & Vojtek, 1975), *Bo. variegata*, (Vojtková & Vojtek, 1975), *B. bufo* (Shimalov & Shimalov, 2001), *B. calamita*, (Vojtková & Vojtek, 1975), *B. vulgaris* (Yamaguti, 1958), *B. viridis* (Düşen *et al.*, 2010; Düşen & Oğuz, 2010), *H. arborea* (Vojtková & Vojtek, 1975; Düşen & Öz, 2004), *H. savignyi* (Yıldırımhan *et al.*, 2012), *R. arvalis* (Vojtková & Vojtek, 1975), *R. camerani* (Yıldırımhan *et al.*, 2006a; Düşen, 2007), *R. dalmatina* (Buchvarov, 1977; Düşen *et al.*, 2009), *R. esculenta* (Vojtková & Vojtek, 1975; Buchvarov, 1977; Kuc & Sulgostowska, 1988b), *R. macronemis* (Yıldırımhan *et al.*, 2006b; Düşen, 2007), *R. arvalis* (Vojtková & Vojtek, 1975), *R. temporaria* (Vojtková &

Vojtek, 1975; Cedhagen, 1977), *R. holtzi* (Topçu, 2002), *Lacerta trilineata* (Yamaguti, 1963; Yıldırımhan *et al.*, 2011), *N. natrix* (Kirin, 2000b) and *Atheris hispida* (Hassl, 2010; Hassl *et al.*, 2010).

Geographic range: Europe and Asia (Yamaguti, 1958); Austrolasian Regions (Prudhoe & Bray, 1982).

Specimens deposited: PAU-HELM-4/2009

Family: Lecithodendriidae

Prosotocus confusus (Loos, 1894) Loos, 1899

Prevalence, intensity and range: Hosts infected, 23 of 298 (7.72 %, 5.82 ± 1.33 SE, 1 – 33).

Other reported hosts: *R. esculenta* (Yamaguti, 1958; Vojtková & Vojtek, 1975; Buchvarov, 1977; Kuc & Sulgostowska, 1988b), *B. viridis* (Yamaguti, 1958; Vojtková & Vojtek, 1975), *B. vulgaris* (Yamaguti, 1958), and *Bo. variegata*, *B. bufo*, *B. calamita*, *H. arborea*, *R. temporaria*, *R. arvalis* (Vojtková & Vojtek, 1975; Buchvarov, 1977). This species is only observed in *R. ridibunda* (Yıldırımhan *et al.*, 1996, 2005a; Düşen & Öz 2006) from Turkey. **Geographic range:** Europe and Asia (Yamaguti, 1958), Turkey (Yıldırımhan *et al.*, 1996, 2005a; Düşen & Öz 2006).

Specimens deposited: PAU-HELM-5/2009

Family: Plagiorchiidae

Skrjabinoeces breviansa (Sudarikov, 1950)

Prevalence, intensity and range: Hosts infected, 29 of 298 (9.73 %, 5.34 ± 1.10 SE, 1 – 35).

Other reported hosts: Sey and Eory (1992) were reported *Skrjabinoeces breviansa* in *R. ridibunda* from Hungary, Mashai *et al.* (2000) were reported *S. breviansa* in *R. ridibunda* from Northern parts of Iran. Kovalenko (2007) was recorded this digenean in *R. ridibunda* and *R. esculenta* from Ukrain. Yıldırımhan *et al.* (2005a), Düşen and Öz (2006), Düşen and Oğuz (2010) were recorded this helminth species in *R. ridibunda* from Turkey.

Geographic range: Europe, Russia (Yamaguti, 1958), Turkey (Yıldırımhan *et al.*, 2005a; Düşen and Öz (2006), and Iran (Mashai *et al.*, 2000).

Specimens deposited: PAU-HELM-6/2009

Family: Plagiorchiidae

Encyclometra colubrimurorum (Rudolphi, 1819) Dollfus, 1922 (Metacercariae)

Prevalence, intensity and range: Hosts infected, 7 of 298 (2.35 %, 2.43 ± 1.11 SE, 1 – 8).

Other reported hosts: *R. tigrina*, *Xenochhnopis piscator*, *Coluber gemonensis*, *Ptyas mucosus*, (Yamaguti, 1958); *N. natrix* (Yamaguti, 1958; Shimalov & Shimalov, 2000; Mihalca *et al.*, 2007); *Bo. bombina*, *P. fuscus*, *B. viridis*, *R. dalmatina*, *R. temporaria* (Vojtková & Vojtek, 1975); *R. esculenta* (Buchvarov, 1977); *H. arborea* (Vojtková & Vojtek, 1975; Düşen & Öz, 2004). The definitive host of *E. colubrimurorum* is a colubrid snake, *N. natrix* (grass snake) (Liang-Sheng, 1958), *E. colubrimurorum* parasitised *P. ridibundus* as an intermediate host. Düşen and Öz (2006) observed this digenean from *R. ridibunda* in An-

talya region in Turkey.

Geographic range: Europe, Asia (Yamaguti, 1958); Czech Republic, France, Italy, Russia, The Netherlands (Vojtková & Vojtek, 1975); Iran (Masshai *et al.*, 2000); Turkey (Düşen & Öz, 2004; Düşen & Öz, 2006).

Specimens deposited: PAU-HELM-7/2009

Family: Telorchidae

Opisthioglyphe ranae (Froelich, 1791)

Prevalence, intensity and range: Hosts infected, 4 of 298 (1.34 %, 4.75 ± 0.85 SE, 3 – 7).

Other reported hosts: The reported hosts of *O. ranae* were, *Salamandra maculosa* (Vojtková & Vojtek, 1975; Buchvarov, 1977); *T. cristatus* (Yamaguti, 1958; Shimalov *et al.*, 2001), *T. vulgaris* (Shimalov *et al.*, 2001), *Bo. bombina* (Vojtková & Vojtek, 1975; Yıldırımhan *et al.*, 2001a), *Bo. variegata* (Vojtková & Vojtek, 1975), *B. bufo* (Shimalov & Shimalov, 2001), *B. calamita* (Yamaguti, 1958; Shimalov & Shimalov, 2001), *B. vulgaris*, *B. variabilis*, (Yamaguti, 1958), *B. viridis*, (Vojtková & Vojtek, 1975; Buchvarov, 1977), *H. arborea* (Vojtková & Vojtek, 1975; Buchvarov, 1977), *P. fuscus* (Vojtková & Vojtek, 1975); *R. dalmatina* (Vojtková & Vojtek, 1975; Buchvarov, 1977), *R. esculenta* (Yamaguti, 1958; Vojtková & Vojtek, 1975; Buchvarov, 1977), *R. lessonae* (Vojtková & Vojtek, 1975), *R. temporaria* (Yamaguti, 1958; Vojtková & Vojtek, 1975; Buchvarov, 1977), *R. arvalis* (Vojtková & Vojtek, 1975), *R. camerani* (Yıldırımhan *et al.*, 2006a), *R. holtzi* (Topçu, 2002), *N. Natrix* and *V. berus* (Shimalov & Shimalov, 2000).

Geographic range: Europe (Yamaguti, 1958), Turkey (Yıldırımhan *et al.*, 1996; Düşen & Öz, 2006), Saudi Arabia (Fernando, 1989), Iran (Masshai *et al.*, 2000).

Specimens deposited: PAU-HELM-8/2009

Cestoda

Family: Nematotaeniidae

Nematotaenia dispar (Goeze, 1782) Lühe, 1899

Prevalence, intensity and range: 2 of 298 hosts infected (0.67 %, 2, 2).

Other reported hosts: *N. dispar* is commonly parasitic in the intestines of amphibians, but rarely reptiles of the Oriental, Nearctic, and Palearctic regions (Prudhoe & Bray, 1982), including *Bo. bombina* (Buchvarov, 1977), *Bo. variegata* (Buchvarov, 1977; Prudhoe & Bray, 1982), *B. bufo* (Hristovski & Riggio, 1971; Prudhoe & Bray, 1982), *B. alvarius* (Goldberg & Bursey, 1991), *H. arborea* (Hristovski & Riggio, 1974; Buchvarov, 1977; Prudhoe & Bray, 1982; Vashetko & Siddikkov, 1999; Saeed *et al.*, 2007), *H. savignyi* (Al-Sorkhy & Amr, 2003; Mashai, 2005; Yıldırımhan *et al.* 2012), *R. temporaria* (Hristovski & Lees, 1973; Buchvarov, 1997), *R. dalmatina* (Buchvarov, 1977), *Mertensiella caucasica* (Yıldırımhan *et al.*, 2001b; 2005b), *Amietophrynus (Bufo) regularis* (Ibrahim, 2008), *B. viridis* (Hristovski & Riggio, 1971; Buchvarov, 1977; Yıldırımhan, 1999; Düşen, 2003, 2011; Saeed *et al.*, 2007; Masshai *et al.*, 2008; Düşen *et al.*, 2010a; Düşen & Oğuz,

2010), *R. camerani* (Yıldırımhan *et al.*, 2006a). Buchvarov (1977) recorded this cestode in *R. ridibunda* from, Bulgaria.

Geographic range: Palaearctic, Oriental and Australian regions (Prudhoe & Bray, 1982; Al-Sorkhy & Amr, 2003). *Nematotaenia dispar* is first time observed in *P. ridibundus* from Turkey.

Specimens deposited: PAU-HELM-9/2009

Nematoda

Family: Rhabdiasidae

Rhabdias bufonis (Schrunk, 1788) Stiles and Hassal, 1905
Prevalence, intensity and range: 19 of 298 hosts infected (6.37 %, 2.05 ± 0.23 SE, 1 – 10).

Other reported hosts: *R. bufonis* is known from various amphibian species including, *Bufo* sp., *Rana* sp., *Pelobates* sp., *Bombinator* sp., *Anguis fragilis* (Yamaguti, 1961; Düşen *et al.*, 2010b), *Bo. bombina* (Grabda-Kazubska & Lewin, 1989; Yıldırımhan *et al.*, 2001a), *Pe. caucasicus* (Yıldırımhan *et al.*, 2009), *R. esculenta* (Buchvarov, 1977; Kuc & Sulgostowska, 1988b), *Bo. variegata*, *P. syriacus* (Buchvarov, 1977), *B. viridis* (Buchvarov *et al.*, 1975; Buchvarov, 1977; Yıldırımhan, 1999a; Shimalov & Shimalov, 2001; Düşen, 2011), *B. bufo* (Buchvarov, 1977; Shimalov & Shimalov, 2001; Yıldırımhan & Karadeniz, 2007; Düşen, 2011), *B. calamita* (Shimalov & Shimalov, 2001), *Amietophrynus (Bufo) regularis* (Ibrahim, 2008), *R. temporaria*, *R. arvalis* (Kuc & Sulgostowska, 1988b; Cedhagen, 1988), *R. dalmatina* (Buchvarov *et al.*, 1975; Buchvarov, 1977; Düşen *et al.*, 2009), *R. camerani* (Yıldırımhan *et al.*, 2006a), *R. macrocnemis* (Yıldırımhan *et al.*, 2006b).

Geographic range: Europe, Siberia, China, Canada, U.S.A. (Yamaguti, 1961).

Specimens deposited: PAU HEL-10/2009

Family: Molineidae

Oswaldocruzia filiformis (Goeze, 1782) Travassos, 1917
Prevalence, intensity and range: 17 of 298 hosts infected (5.70 %, 2.47 ± 0.70 SE, 1 – 10).

Other reported hosts: *O. filiformis* is recorded from various amphibian and reptile species, including *Salamandra salamandra* (Buchvarov, 1977), *T. alpestris* and *T. karelinii* (Buchvarov, 1977; Cedhagen 1988; Kirin & Buchvarov, 2002), *T. vulgaris* (Buchvarov, 1977; Shimalov *et al.*, 2001), *T. vittatus* (Yıldırımhan, 2008), *Bo. bombina* and *Bo. variegata* (Buchvarov, 1977; Kirin & Buchvarov, 2002), *B. regularis* (probably *Bufo viridis*) (Schad *et al.*, 1960), *B. viridis* (Buchvarov, 1977; Yıldırımhan, 1999; Shimalov & Shimalov, 2001; Topçu & Bayrak, 2000; Düşen *et al.*, 2010a; Düşen & Oğuz, 2010), *B. bufo* (Buchvarov, 1977; Yıldırımhan & Karadeniz, 2007; Düşen & Oğuz, 2010; Düşen, 2011); *Pe. caucasicus* (Yıldırımhan *et al.*, 2009), *H. arborea* (Buchvarov, 1977; Yıldırımhan *et al.*, 2006c), *R. camerani* (Yıldırımhan *et al.*, 2006a); *H. savignyi* (Yıldırımhan *et al.*, 2012); *R. dalmatina*, (Buchvarov *et al.*, 1975; Buchvarov, 1977; Kirin & Buchvarov, 2002; Düşen *et al.*, 2009), *R. kurtmuelleri* (Hristovski *et*

al., 2006), *R. macrocnemis* (Schad *et al.*, 1960; Yıldırımhan *et al.*, 1997b; Yıldırımhan *et al.*, 2006b), *R. temporaria*, (Buchvarov, 1977; Cedhagen, 1988; Kirin & Buchvarov, 2002), *R. graeca* (Božkov & Stojkova, 1970; Buchvarov, 1977), *L. agilis*, (Sharpilo *et al.*, 2001; Shimalov *et al.*, 2000; Mihalca *et al.*, 2007), *L. trilineata* (Yıldırımhan *et al.*, 2011), *L. viridis* (Biserkov & Kostadinova, 1998; Yıldırımhan, 1999b; Kirin, 2002a; Borkovcová & Kopřiva, 2005), *L. vivipara* (Shimalov *et al.*, 2000), *A. fragilis* (Schad *et al.*, 1960; Bertman & Okulewicz, 1987; Shimalov *et al.*, 2000; Borkovcová & Kopřiva 2005; Düşen *et al.*, 2010b), *Zootoca vivipara* (Sanchis *et al.*, 2000), *N. natrrix* (Bertman & Okulewicz, 1987; Shimalov & Shimalov, 2000; Kirin 2002b) and *V. berus* (Shimalov & Shimalov, 2000).

Geographic range: Europe and Asia (Yamaguti, 1961).

Specimens deposited: PAU-HELM-11/2009

Family: Cosmocercidae

Cosmocerca ornata (Dujardin, 1845)

Prevalence, intensity and range: 114 of 298 hosts were infected (38.25 %, 5.28 ± 0.75 SE, 1 – 47).

Other reported hosts: There are several papers reporting *C. ornata* from many species of amphibians and reptiles, including *Bufo*, *Hyla*, *Rana*, *Triturus* (Yamaguti, 1961), *T. alpestris* (Walton, 1933; Buchvarov, 1977; Shimalov *et al.*, 2000), *T. cristatus* (Walton, 1933; Shimalov *et al.*, 2001), *T. vulgaris* (Shimalov *et al.*, 2001), *Bo. bombina*, (Buchvarov, 1977; Grabda-Kazubska & Lewin, 1989); *Bo. variegata* (Buchvarov, 1977; Grabda-Kazubska & Lewin, 1989; Kirin & Buchvarov, 2002), *B. viridis* (Schad *et al.*, 1960; Buchvarov *et al.*, 1975; Buchvarov, 1977; Vashetko & Siddikov, 1999; Masshah, 2005; Düşen & Oğuz, 2010; Düşen *et al.*, 2010; Düşen 2011), *B. bufo* (Buchvarov, 1977; Shimalov & Shimalov, 2001; Yıldırımhan & Karadeniz, 2007; Düşen & Oğuz, 2010; Düşen *et al.*, 2010; Düşen 2011); *Pe. caucasicus* (Yıldırımhan *et al.*, 2009), *H. arborea* (Buchvarov, 1977; Yıldırımhan *et al.*, 2006c), *P. syriacus* (Buchvarov, 1977; Shimalov *et al.*, 2000), *R. esculenta* (Walton, 1933; Buchvarov, 1977), *R. arvalis* (Cedhagen, 1988; Kuc & Sulgostowska, 1988b), *R. temporaria*, (Walton, 1933; Buchvarov, 1977; Kuc & Sulgostowska, 1988b), *R. graeca* (Božkov & Stojkova, 1970; Buchvarov, 1977), *R. holtzi* (Yıldırımhan *et al.*, 2006b), *R. macrocnemis* (Schad *et al.*, 1960; Yıldırımhan *et al.*, 2006b; Düşen, 2007), *R. camerani* (Yıldırımhan *et al.*, 2006a; Düşen, 2007), *R. dalmatina* (Buchvarov, 1977; Düşen *et al.*, 2009), *R. tavasensis* (Düşen, 2012); *Chiasmocleis capixaba* (Van Sluys *et al.*, 2006) and *A. fragilis* (Shimalov *et al.*, 2000; Düşen *et al.*, 2010b).

Geographic range: New and Old Worlds (Baker, 1987).

Specimens deposited: PAU-HELM-12/2009

Oxysomatium brevicaudatum (Zeder, 1800) Railliet and Henry, 1916

Prevalence, intensity and range: 22 of 298 hosts were infected (7.38 %, 8.86 ± 2.19 SE, 1 – 32).

Other reported hosts: *O. brevicaudatum* was reported from

different hosts of amphibians and reptiles in Europe and Asia, including *Bombina*, *Bufo*, *Hyla*, *Pelobates*, *Rana*, *Salamandra* (Yamaguti, 1961), *S. salamandra*, *S. atra*, *Pseudotriton ruber* and *Bo. bombina* (Walton, 1933), *T. vittatus*, *T. karelinii* (Yıldırımhan, 2008), *P. syriacus* (Yıldırımhan *et al.*, 1997a; Yıldırımhan & Bursey, 2010), *B. viridis* (Walton, 1933; Schad *et al.*, 1960; Buchvarov, 1977; Yıldırımhan, 1999a; Topçu & Bayrak, 2000; Düşen

& Oğuz, 2010; Düşen *et al.*, 2010; Düşen, 2011), *B. bufo* (Düşen & Oğuz, 2010; Düşen *et al.*, 2010; Düşen, 2011), *B. regularis* (probably *B. viridis*) (Schad *et al.*, 1960), *H. arborea* (Walton, 1933), *R. dalmatina* (Buchvarov, 1977; Düşen *et al.*, 2009), *R. graeca* (Božkov & Stojkova, 1970; Buchvarov, 1977), *R. esculenta* (Walton, 1933), *R. kurtmuelleri* (Hristovski *et al.*, 2006), *R. macrocnemis* (Schad *et al.*, 1960), *R. temporaria* (Walton, 1933; Buch-

Table 1. Prevalence, intensity, infection sites and range of helminths in *P. ridibundus*

Identified helminth group	Developmental stage	Site of infection	No. of infected (%)	Mean intensity ($\pm$ SE)	Range
<i>Diplodiscus subclavatus</i> (Pallas, 1760) Diesing, 1836	Adult	LI	31 (10.40)	4.03 $\pm$ 0.65	1 – 15
<i>Gorgoderina vitelliloba</i> (Olsson, 1876) Loos, 1902	Adult	UB	20 (6.71)	2.55 $\pm$ 0.81	1 – 13
<i>Gorgodera cygnoides</i> (Zeder, 1800)	Adult	UB	36(12.08)	3.83 $\pm$ 0.70	1 – 17
<i>Pleurogenoides medians</i> (Olsson, 1876) Travassos, 1921	Adult	SI	47 (15.77)	27.70 $\pm$ 4.57	1 – 126
<i>Prosotocus confusus</i> (Loos, 1894) Loos, 1899	Adult	SI	23 (7.72)	5.82 $\pm$ 1.33	1 – 33
<i>Skrjabinoces breviansa</i> (Sudarikov, 1950)	Adult	L	29 (9.73)	5.34 $\pm$ 1.10	1 – 35
<i>Encyclometra colubrimurorum</i> (Rudolphi, 1819) Dollfus, 1922	Metacercariae	M, OS, BC	7 (2.35)	2.43 $\pm$ 1.11	1 – 8
<i>Opisthioglyphe ranae</i> (Froelich, 1791)	Adult	SI	4 (1.34)	4.75 $\pm$ 0.85	3 – 7
<i>Nematotaenia dispar</i> (Goeze, 1782) Lühe, 1899	Adult	SI	2 (0.67)	2	2
<i>Rhabdias bufonis</i> (Schränk, 1788) Stiles and Hassal, 1905	Adult	L	19 (6.37)	2.05 $\pm$ 0.23	1 – 10
<i>Oswaldocruzia filiformis</i> (Goeze, 1782) Travassos, 1917	Adult	SI	17 (5.70)	2.47 $\pm$ 0.70	1 – 10
<i>Cosmocerca ornata</i> (Dujardin, 1845)	Adult	LI, SI	114 (38.25)	5.28 $\pm$ 0.75	1 – 47
<i>Oxysomatium brevicaudatum</i> (Zeder, 1800) Railliet and Henry, 1916	Adult	LI, SI	22 (7.38)	8.86 $\pm$ 2.19	1 – 32
<i>Eustrongylides</i> sp.	Larvae	LI, SI	3 (1.01)	1	1
<i>Abbreviata</i> sp. (Encapsulated larvae in submucosa of stomach and small intestine)	Larvae	S, LI, OS	35 (11.74)	1	Uncount.
<i>Acanthocephalus ranae</i> (Schränk, 1788) Lühe, 1911	Adult	SI, LI	63 (21.14)	4.92 $\pm$ 0.67	1 – 26
<i>Pomphorhynchus laevis</i> (Zoega in Müller 1776)	Larvae and cysts	M, B, LI, SI	55 (18.45)	21.36 $\pm$ 1.33	1 – 79

BC: Body cavities, L: Lung, LI: Large intestine, M: Mesentery, OS: Organ surfaces, R: Rectum, S: Stomach, SI: Small intestine; UB: Urinary bladder

varov, 1977; Kirin & Buchvarov, 2002), *P. fuscus* (Walton, 1933), *R. kurtmuelleri* (Hristovski *et al.*, 2006), *A. fragilis* (Schad *et al.*, 1960; Sharpilo, 2003; Düşen *et al.*, 2010b), *N. natrix* (Schad *et al.*, 1960).

Geographic range: Europe and Asia (Yamaguti, 1961). Schad *et al.* (1960) were first time reported *O. brevicaudatum* in *Bufo regularis* (probably *B. viridis*), *B. viridis*, *R. macrocnemis*, *R. ridibunda* and *A. fragilis* from Turkey. Sharpilo (2003) pointed out that *O. brevicaudatum* has a great disperse in *A. fragilis* in the Caucasian Region.

Specimens deposited: PAU-HELM-13/2009

Family: Dioctophymatidae

***Eustrongylides* sp. (larvae)**

Prevalence, intensity and range: Hosts infected, 3 of 298 (1.01 %, 1, 1).

Other reported hosts: *Eustrongylides* sp. is parasitic in glands of proventriculus of aquatic birds. Larvae normally inhabited connective tissue and body cavities of fishes (Yamaguti, 1961). Yıldırımhan *et al.* (1996b) and Aydoğdu *et al.* (1996a, 1996b, 1997 and 2000) recorded this nematode larvae in *N. natrix*, roach (*Rutilus frisii*), sheatfish (*Silurus glanis*), and tench (*Tinca tinca*) from Lake Izmir (Bursa). Mihalca *et al.* (2007) were recorded this nematode in *N. natrix* and *N. tessellata* in Romania. Anderson (2000) stated that the larvae of this nematode is observed in *R. ridibunda*. Düşen and Öz (2006) reported this nematode larvae in *R. ridibunda* from Antalya Region in Turkey.

Geographic range: Europe south-east Asia, the Middle East and Australia (Anderson, 2000).

Specimens deposited: PAU-HELM-14/2009

Family: Physalopteridae

***Abbreviata* sp. (larvae - Encapsulated larvae in submucosa of stomach and small intestine).**

Prevalence, intensity and range: Hosts infected, 35 of 298 (11.74 %, Uncountable).

Other reported hosts: Fernando (1989) observed this nematode larvae in *R. ridibunda* from Saudi Arabia; Fernando (1989) reported that parasite larvae in brown cysts where located subcutaneously on body wall, heavy infestations also occurred in the submucosae of stomach, small intestine and mesenteries and embedded deeply in them muscle. Similarly, Düşen and Öz (2006) observed this encapsulated larvae in submucosa of stomach and small intestines of *R. ridibunda* from Antalya region in Turkey. Borkovcová and Kopřiva (2005) recorded this nematode species in some lizards (*A. fragilis*, *L. viridis* and *L. agilis*) and one snake (*Coronella austriaca*) species in South Moravia (Czech Republic). In this study, the larvae of *Abbreviata* sp. were observed numerous (embedded deeply in submucosa of the stomach and small intestine) therefore counting was difficult.

Geographic range: Europe Asia, and Western Australia (Anderson, 2000).

Specimens deposited: PAU-HELM-15/2009

Acanthocephala

Family: Echinorhynchidae

***Acanthocephalus ranae* (Schränk, 1788) Lühe, 1911**

Prevalence, intensity and range: 63 of 298 hosts were infected (21.14 %, 4.92 ± 0.67 SE, 1 – 26).

Other reported hosts: *Rana* sp., *Bombinator* sp., *Hyla* sp., *Triturus* sp., *Salamandra* sp., *Diemictylus viridescens* (Yamaguti, 1963), *Bo. bombina* (Buchvarov, 1977; Grabda-Kazubska & Lewin, 1989; Yıldırımhan *et al.*, 2001a), *Bo. variegata* (Buchvarov, 1977; Grabda-Kazubska & Lewin, 1989), *B. viridis* (Buchvarov, 1977; Yıldırımhan, 1999a; Vashetko & Siddikov, 1999; Shimalov & Shimalov, 2001; Düşen & Oğuz, 2010), *B. bufo* (Yıldırımhan & Karadeniz, 2007; Düşen, 2011), *B. calamita* (Shimalov & Shimalov, 2001); *H. arborea* (Düşen & Öz, 2004), *R. arvalis*, *R. dalmatina* (Buchvarov, 1977; Düşen *et al.*, 2009); *R. temporaria* (Buchvarov, 1977; Cedhagen, 1988; Kuc & Sulgostowska, 1988b), *R. esculenta* (Buchvarov, 1977; Kuc & Sulgostowska, 1988b), *R. macrocnemis* (Yıldırımhan *et al.*, 1997b; Yıldırımhan *et al.*, 2006b; Düşen, 2007), *R. camerani* (Yıldırımhan *et al.*, 2006a), *M. caucasica* (Yıldırımhan *et al.*, 2001b; 2005b), *R. kurtmuelleri* (Hristovski *et al.*, 2006), *R. tavasensis* (Düşen, 2012), *A. fragilis* (Shimalov *et al.*, 2000), *N. natrix* (Yamaguti, 1963; Shimalov & Shimalov, 2000).

Geographic range: Europe, U.S.A., Russia (Yamaguti, 1963); Turkey (Oğuz *et al.*, 1994).

Specimens deposited: PAU-HELM-16/2009

Family: Pomphorhynchidae

***Pomphorhynchus laevis* (Zoega in Müller 1776) (larvae and cysts)**

Prevalence, intensity and range: Hosts infected, 55 of 298 (18.45 %, 21.36 ± 1.33 SE, 1 – 79).

Other reported hosts: In general, *Pomphorhynchus laevis* occurs predominantly in cyprinid and also in salmonid fishes with the Palaearctic distribution (Dudiňák & Šnábel, 2001). *P. laevis*, is observed in several fish species in Turkey: barbel (*Barbus plebejus escherichii*), bleak (*Alburnus alburnus*), chub (*Leuciscus cephalus*), common carp (*Cyprinus carpio*), crucian carp (*Carassius carassius*), loach (*Nemachilus* sp.), nase (*Chondrostoma nasus*), pike (*Esox lucius*), sheatfish (*Silurus glanis*), (Öktener, 2003), and tench (*Tinca tinca*) (Yıldız, 2003). A number of marine and freshwater fishes have been reported as definitive hosts of acanthocephalan *P. laevis* (Ziolkowska & Rokicki, 2003). Buchvarov (1977) and Düşen and Oğuz (2008) observed in *Rana ridibunda* and Veith and Erpelding (1995) in fire salamander (*S. salamandra*) are reported *P. laevis*.

Geographic distribution: Europe Buchvarov (1977)

Specimens deposited: PAU-HELM-17/2009

Seventeen helminth species were found infecting *P. ridibundus*. The site of infection in the Eurasian marsh frog and the data on infection parameters for each host and helminth species are shown in Table 1. According the data

obtained 262 (87.91 %) *P. ridibundus* harbored the one or more species of helminths and the remaining 36 (%)12.08 were uninfected. In summary, 4311 individuals of 17 helminth species were collected from the 298 *P. ridibundus* were examined. Digeneans and nematodes were observed in both large-small intestines, lungs, rectum, mesenteries and organ surfaces; cestode and acanthocephalans were also observed in small intestine of *P. ridibundus*. There were 2.57 ± 1.56 (1 – 7 with a range from) helminth species per infected host and there were 14.46 helminth individuals per infected host (except *Abbreviata* sp., uncountable). The helminth species that were observed in *P. ridibundus*, are common parasites of European anurans and also, some reptiles (e.g. *E. colubrimurorum*), aquatic birds (e.g. *Eustrongylides* sp.) fishes (e.g. *P. laevis*) (Yamaguthi, 1958, 1961, 1963; Liang-Sheng, 1958; Buchvarov, 1977; Yıldırımhan, 1999; Anderson, 2000; Dudiňák & Šnábel, 2001; Düşen & Öz, 2006; Yıldırımhan *et al.*, 2007; Düşen *et al.*, 2010a; Düşen, 2012). *P. ridibundus* represents a new host record for *Nematotaenia dispar* in Turkey.

The present study on the helminthfauna on *P. ridibundus*, Eurasian marsh frog in Denizli province (inner-west Anatolia region Turkey), offers a contribution to the present knowledge in this field.

Acknowledgements

This study was supported by TÜBİTAK (The Scientific and Technical Research Council of Turkey) Project no: 107T917, and partly supported by the Pamukkale University Scientific Research Projects Unit Project no: 2008BSP005. We also thank, for permission and helps, the Department of National Parks and Wildlife of the Ministry of Environment and Forestry, of the Republic of Turkey.

References

AL-SORKHY, M. K., AMR, Z. (2003): Helminth parasites of some amphibians in Jordan. *Tr. J. of Zool.*, 27: 89 – 93

ANDERSON, R. C. (2000): *Nematode Parasites of Vertebrates: Their Development and Transmission* (2nd Edition). CABI Publishing, Wallingford, Oxon, U.K., 650 pp.

AYDOĞDU, A., YILDIRIMHAN, H. S., ALTUNEL, F. N. (1996a): An investigation on catfish (*Silurus glanis* L.) in Lake Iznik and its around, XIIIth National Biology Congress, 17 – 20 Eylül 1996. Cilt I, 63 – 70 (In Turkish)

AYDOĞDU, A., YILDIRIMHAN, H. S., ALTUNEL, F. N. (1996b): A Study on the parasites of Tench fishes of Lake Iznik (*Tinca tinca* L., 1758). *Türkiye Parazitoloji Dergisi (Acta Parasitol. Tur.)* 20: 261 – 270 (In Turkish, with English abstract)

AYDOĞDU, A., YILDIRIMHAN, H. S., ALTUNEL, F. N. (1997): An investigation on the ecto and endoparasites of the Akbalık (*Rutilus frisii*), collected from Lake Iznik, IXth Aquaculture Congress, 17 – 19 Eylül 1997 Eğirdir/Isparta. 431 – 443 (In Turkish)

AYDOĞDU, A., YILDIRIMHAN, H. S., ALTUNEL, F. N. (2000): The Helminth Fauna of the Adriatic Roach (*Rutilus rubilio*)

in İznik Lake. *Bull. Eur. Ass. Fish Pathol.*, 20 (3), 170

BAKER, M. R. (1987): *Synopsis of the Nematoda parasitic in amphibians and reptiles*. Memorial University of Newfoundland, *Occas. Pap. in Biol.*, 1: 1 – 325

BARAN, I., ATATÜRK, M. K. (1997): *Turkish Herpetofauna*. The Republic of Turkey, Ministry of Environment Publications, Ankara. 214 pp.

BERTMAN, M., OKULEWICZ, A. (1987): Lizards (*Anguis fragilis* L.) and snakes (*Natrix natrix* (L.)) as new hosts of *Oswaldocruzia filiformis* (Goeze, 1782) Travassos 1917 (Nematoda). *Wiadomosci Parazytol.*, 33 (2): 209 – 212 (In Polish)

BISERKOV, V., KOSTADINOVA, A. (1998): Intestinal helminth communities in the green lizard, *Lacerta viridis*, from Bulgaria. *J. Helminthol.*, 72 (3): 267 – 271. DOI: 10.1017/S0022149X00016540

BORKOVCOVÁ, M., KOPŘIVA, J. (2005): Parasitic helminths of reptiles (Reptilia) South Moravia (Czech Republic). *Parasitol. Res.*, 96: 77 – 78. DOI: 10.1007/s00436-004-1258-6

BOŽKOV, D., STOJKOVA, R. (1970): Beitrag zur untersuchung der Helminthenfauna der *Rana graeca* in Bulgarien. *Bulletin de L'institute de Zoologie et Musée, Academie Bulgare des Sciences*, 69 – 75 (In Bulgarian)

BUCHVAROV, G. K. (1977): *Catalogue des helminthes des Amphibies en Bulgarie* - Universite de Plovdiv "P Hilendarski", Plovdiv, Bulgaria, 53 pp. (In Bulgarian).

BUCHVAROV, G. K., PETROV, P., CHOCHEV, B. (1975): To the question about helminthofauna of Amphibious Eucaudate (Amphibia-Eucaudata) of Velingrad's District. Universite de Plovdiv "P Hilendarski", Travaux Scientifiques, *Biologie*, 13 (4): 53 – 64

BUDAK, A., GÖÇMEN, B. (2008): *Herpetology (Herpetoloji)* Ege University Press, Science Faculty Publ. no: 194 (2nd Edition) 226 pp. (In Turkish)

CEDHAGEN, T. (1988): Endoparasites in some Swedish amphibians. *Acta Parasitol.*, 33: 107 – 113

DUDIŇÁK, V., ŠNÁBEL, V. (2001): Comparative Analysis of Slovak and Czech Populations of *Pomphorhynchus laevis* (Acanthocephala) Using Morphological and Isoenzyme Analyses. *Acta Zool. Univ. Comenianae*, 44: 41 – 50

DÜŞEN, S. (2003): *The helminth fauna of tailless frog species (Ordo: Anura) distributed in Antalya*. PhD thesis, Akdeniz University, Antalya, Turkey. 116 pp. (in Turkish, with English abstract)

DÜŞEN S., ÖZ, M. (2004): Helminth parasites of the tree frog, *Hyla arborea* (Linnaeus, 1758) (Anura: Hylidae) from Southwest Turkey. *Comp. Parasitol.*, 71 (2): 258 – 261. DOI: 10.1654/4123

DÜŞEN S., ÖZ, M. (2006): Parasitic helminths of the marsh frog, *Rana ridibunda* Pallas, 1771 (Anura: Ranidae), from Antalya Province, south-west Turkey. *Comp. Parasitol.*, 73 (1): 121 – 129. DOI: 10.1654/4162.1

DÜŞEN, S. (2007): Helminths of the two mountain frogs, Banded frog, *Rana camerani* Boulenger, 1886 and Uludağ Frog *Rana macrocnemis* Boulenger, 1885 (Anura: Ranidae), collected from Antalya Province. *Acta Parasitol. Tur.*, 31 (1): 84 – 88

- DÜŞEN, S., OĞUZ, M. C. (2008): Occurrence of *Pomphorhynchus laevis* (Acanthocephala) in the Marsh Frog (*Rana ridibunda* Pallas, 1771), from Turkey. *Helminthologia*, 45 (3): 154 – 156
- DÜŞEN, S., UĞURTAŞ, İ. H., AYDOĞDU, A., OĞUZ, M. C. (2009): The helminth community of the Agile frog, *Rana dalmatina* Bonaparte, 1839 (Anura: Ranidae) Collected from Northwest of Turkey. *Helminthologia*, 46 (3): 177 – 182. DOI: 10.2478/s11687-009-0033-8
- DÜŞEN, S., OĞUZ, M. C., BARTON, D., P. ARAL, A., ŞULEKOĞLU, S., TEPE, Y. (2010a): A metazoan parasitological research of three species of Anurans collected from the Çanakkale Province, Northwestern Turkey. *North-West J. Zool.*, 6 (1): 25 – 35
- DÜŞEN, S., UĞURTAŞ, İ. H., AYDOĞDU, A. (2010b): Nematode parasites of the two limbless lizards: Turkish worm lizard, *Blanus strauchi* (Bedriaga, 1884) (Squamata: Amphisbaenidae), and slow worm, *Anguis fragilis* Linnaeus 1758 (Squamata: Anguidae), from Turkey. *Helminthologia*, 47 (3): 158 – 163. DOI: 10.2478/s11687-010-0024-9
- DÜŞEN, S., OĞUZ, M. C. (2010): Metazoan endoparasites of three species of anurans collected from the Middle Black Sea region of Turkey. *Helminthologia*, 47 (4): 226 – 232. DOI: 10.2478/s11687-010-0035-6
- DÜŞEN, S. (2011): The helminth parasites of the bufonid toads, European Common Toad, *Bufo bufo* (Linnaeus, 1758) and European Green toad, *Bufo (Pseudepidalea) viridis* Laurenti, 1768 (Anura: Bufonidae), collected from Denizli Province, Inner-West Anatolia Region, Turkey *Helminthologia*. 48 (2): 101 – 107
- DÜŞEN, S. (2012): First data on helminthfauna of the locally distributed a mountain frog, “Tavas Frog”, *Rana tavasensis* Baran and Atatür, 1986 (Anura: Ranidae), from Inner-West Anatolia Region, Turkey. *Tr J. of Zoology*. 36(4): 496 – 502. DOI:10.3906/zoo-0909-15
- FERNANDO, M. M. (1989): The parasitic burden of the frog *Rana ridibunda* Pallas, 1771 from Saudia Arabia. A preliminary list of parasitic helminths. *Herpetol. J.*, 1: 415 – 417
- GRABDA-KAZUBSKA, B., LEWIN, J. (1989): The helminth fauna of *Bombina bombina* (L.) and *Bombina variegata* (L.) in Poland. *Acta Parasitol.*, 34: 273 – 279
- HASSL, A. R (2010): A case of a facultative life-cycle diversification in the fluke *Pleurogenoides* sp. (Lecithodendriidae, Plagiorchiida). *Wien Klin. Wochenschr. Suppl.* 3: 91 – 93
- HASSL, A. R., RICHTER, B., KÜBBERHEISS, A. (2010): Captive snake's care as parasite's life cycle conservation program. *Herpetozoa.*, 23 (1/2): 83 – 85
- HECKMANN, R. A., OGUZ, M. C., AMIN, O. M., DÜŞEN, S., TEPE, Y., ASLAN, B. (2010): Host and geographical distribution of *Pomphorhynchus spindletuncatus* (Acanthocephala: Pomphorhynchidae) in Turkey, with enhanced description from new fish and amphibian hosts using SEM, and histopathological notes. *Sci. Parasitol.*, 11: 129 – 139
- HECKMANN, R. A., AMIN, O. M., TEPE, Y., DÜŞEN, S., OGUZ, M. C. (2011): *Acanthocephalus ranae* (Acanthocephala: Echinorhynchidae) from amphibians in Turkey, with special reference to new morphological features revealed by SEM, and histopathology. *Sci. Parasitol.*, 12: 23 – 32
- HRISTOVSKI, N. D., RIGGIO, S. (1971): On some helminths from two toad species (*Bufo bufo* and *B. viridis*) collected in localities of Macedonia, Serbia (Yugoslavia), and Sicily (Italy). *Fragmenta Balcanica Mus. Macedonica Sci. Nat.*, 8: 17 – 25
- HRISTOVSKI, N. D., LEES, E. (1973): The helminth fauna of *Rana temporaria* in relation to that of Europe generally. *Acta Parasitol. Iugosl.* 4: 93 – 98
- HRISTOVSKI, N. D., RIGGIO, S. (1974): Some data on the parasitic helminthofaunas in populations of *Hyla arborea* Laur. From Yugoslavia, Greece and the island of Corsica. *Acta Parasitol. Iugosl.* 5: 21 – 23
- HRISTOVSKI, N., SMILKOV, S., TOMOVSKA, D., POPOVICH E., KOSTICH, D. (2006): *Helminthofauna of Rana kurtmueleri* Gayda, 1940 syn. *Rana balcanica* Shneider et Sinish, 1992 (Anura:Ranidae) from Dojran Lake, Makedonia and Greece. 10th International Congress on the Zoogeography and Ecology of Greece and Adjacent Regions Patras, Greece. June 26 – 30.
- IUCN (2012): *Pelophylax ridibundus* IUCN Red List of Threatened Species. Downloaded on 21 October 2012 from www.iucnredlist.org.
- IBRAHIM, M. M. I. (2008): Helminth infracommunities of the maculated toad *Amietophrynus regularis* (Anura: Bufonidae) from Ismailia, Egypt. *Dis. Aquat. Org.*, 82: 19 – 26
- KIRIN, D. (2002a): New data on the helminth fauna of *Lacerta viridis* Laurenti, 1768, and *Podarcis muralis* (Laurenti, 1768) (Reptilia: Lacertidae) in Bulgaria. *Acta Zool. Bulg.*, 54 (1): 43 – 48
- KIR, İ., YILDIRIM, M. Z., BECER, A. AND İKİZ, R. (2001): The feeding habits and parasites of the lake frogs (*Rana ridibunda* Pallas, 1771; Anura: Ranidae) of Lake Eğirdir. *Acta Parasitol. Tur.*, 25: 83 – 87
- KIRIN, D. (2002b): New records of the helminth fauna from grass snake, *Natrix natrix* L., 1758 and dice snake, *Natrix tessellata* Laurenti, 1768 (Colubridae: Reptilia) in South Bulgaria. *Acta Zool. Bul.*, 54: 49 – 53
- KIRIN, D., BUCHVAROV, G. (2002): Biodiversity of of the helminth communities of acaudated Amphibians (Amphibia: Ecaudata) from Bistritsa Riverside (Gotse Delchev Region). *Exp. Pathol. Parasitol.*, 5(8): 13 – 16
- KOVALENKO, M. V. (2007): First Ukrainian record of the frog lung fluke *Skrjabinoeces breviansa* Sudarikov, 1950 (Trematoda: Plagiorchiidae). *Seriesbiology. Univ. Kharkov.* no: 595.122(477). 93-96
- KUC, I., SULGOSTOWSKA, T. (1988a): Helminth fauna of *Rana ridibunda* Pallas, 1771 from Goclawski Canal in Warsaw (Poland). *Acta Parasitol.*, 33: 101 – 105
- KUC, I., SULGOSTOWSKA, T. (1988b): Helminth fauna of frogs in the forest of Kampinos near Warszawa. *Acta Parasitol.*, 33: 267 – 272
- LIANG-SHENG, Y. (1958): A review of the trematode genus *Encylometra* Baylis and Cannon, 1924. *J. Helminthol.*, 32: 99 – 144
- MASSHAI, N., BALOUCH, M., MOBEDI, I. (2000): New records about helminth parasites of the marsh frog, *Rana ridibunda* (Anura: Ranidae), from the North of Iran. *Iran. J. Fish. Sci.*, 2: 77 – 88

- MASHAI, N. (2005): Helminth Parasites of green toad, *Bufo viridis* (Anura: Bufonidae), Tree frog, *Hyla arborea savignyi* (Anura: Hylidae) and Marsh Frog, *Rana ridibunda ridibunda* (Anura: Ranidae) from Southwest of Iran. *Iran. J. Vet. Res.*, 6 (3): 67 – 73
- MASSHAI, N., BALOUCH M., MOBEDI, I. (2008): Report about helminth parasites of some Amphibians (Anura: Ranidae, Bufonidae) from the North and Northeast of Iran. *J. Sci. Univ. Tehran.*, 33 (4): 9 – 13
- MIHALCA, A. D., GHERMAN, C., GHIRA, I., COZMA, V. (2007): Helminth parasites of reptiles (Reptilia) in Romania. *Parasitol. Res.*, 95: 77 – 78. DOI: 10.1007/s00436-007-0486-y
- OĞUZ, M. C., ALTUNEL, F. N., UĞURTAŞ, İ. H. (1994): An investigation of the species of Plathelminthes and *Acanthocephalus ranae* (Schrank, 1788, Acanthocephala) of marsh frogs (*Rana ridibunda* Pallas, 1771) which were collected from the Bursa and Edirne Regions. *Turk. J. Zool.*, 18: 47 – 51 (In Turkish, with English abstract)
- ÖKTENER, A. (2003): A checklist of Metazoan Parasites Recorded in Freshwater Fish from Turkey. *Zootaxa*, 394: 1 – 28
- PRUDHOE, S., A. BRAY, S. (1982): *Platyhelminth Parasites of Amphibia*. British Museum of Natural History. Oxford University Press, London. 217 pp
- SAEED, I. E., AL-BARWARI S., AL-HARMNI, K. I. (2007): A metazoan parasitological research of some Iraqi Amphibians. *Acta Parasitol. Tur.*, 3 (4): 337 – 345
- SAĞLAM, N., ARIKAN, H. (2006): Endohelminth parasites of the Marsh Frog *Rana ridibunda* from Hazar Lake, Turkey. *Dis. Aquat. Org.*, 72: 253 – 260
- SANCHIS, V., ROIG, J. M., CARRETERO, M. A., ROCA, V., LLORENTE, G. A. (2000): Host – parasite relationships of *Zootoca vivipara* (Sauria: Lacertidae), in Pyrenees, North Spain. *Folia Parasitol.*, 47: 118 – 122
- SAYGI, G., BAŞIBÜYÜK, H. H. (1990): Protozoon and metazoan parasites of *Rana ridibunda*. *Acta Parasitol. Tur.*, 15: 105 – 118
- SCHAD, G. A., KUNTZ, R. E., WELLS, W. H. (1960): Nematode parasites from Turkish vertebrates: an annotated list. *Can. J. Zoolog.*, 38: 949 – 963
- SHARPILO V. P., BISERKOV, V., KOSTADINOVA, A., BEHNKE, J. M., KUZMIN, Y. I. (2001): Helminths of the sand lizard, *Lacerta agilis* (Reptilia, Lacertidae), in the Palaearctic: faunal diversity and spatial patterns of variation in the composition and structure of component communities. *Parasitology*, 123: 389 – 400. DOI: 10.1017/S0031182001008587
- SHARPILO, V. P. (2003): Rare and locally distributed pale-arctic species of the reptile parasitic worms: *Oxysomatium caucasicum* (Nematoda, Cosmocertidae). *Vesnik Zool.*, 37 (5): 69 – 72
- SHIMALOV, V. V., SHIMALOV, V. T. (2000): Helminth fauna of snakes (Reptilia, Serpentes) in Belorussian Polesye. *Parasitol. Res.*, 86: 340 – 341. DOI: 10.1007/s004360050055
- SHIMALOV, V. T., SHIMALOV, V. V., SHIMALOV, A. V. (2000): Helminth fauna of lizards in the southern part of Belarus. *Parasitol. Res.*, 86: 343. DOI: 10.1007/s004360050057
- SHIMALOV, V. V., SHIMALOV V. T. (2001): Helminth fauna of toads in Belorussian Polesie. *Parasitol. Res.*, 87: 84. DOI: 10.1007/s004360000266
- SHIMALOV, V. V., SHIMALOV, V. T., SHIMALOV, A. V. (2001): Helminth fauna of newts in Belarussian Polesie. *Parasitol. Res.*, 87: 356. DOI: 10.1007/PL00008592
- TOPÇU, A., BAYRAK, N. (2000): The digestive system helminths of the *Rana ridibunda* 1771, Pallas and *Bufo* (Pseudepidalea) *viridis* Laurenti, 1768 (Anura) distributed in Nigde Province and its vicinity, XVth National Biology Congress, 5 – 9 Eylül 2000. Ankara. (In Turkish)
- TOPÇU, A. (2002): The digestive system helminths found in the Taurus frog (*Rana holtzi* Werner, 1898), XVIth National Biology Congress, 4 – 7 Eylül 2002. Malatya. (In Turkish)
- VAN SLUYS, M., SCHITTINI, G., MARRA, R. V., AZEVEDO, A., VICENTE J. J., VRCIBRADIC, D., (2006): Body size, diet and endoparasites of the microhylid frog *Chiasmocleis capixaba* in an Atlantic Forest Area of Southern Bahia-state, Brazil. *Brazil. J. Biol.*, 66 (1A): 167 – 173
- VASHETKO, E. V., SIDDIKOV, B. H. (1999): The effect of the ecology of toads on the distribution of helminths. *Turk. J. Zool.*, 23: 107 – 110
- VEITH, M., ERPELDING, G. (1995): Presence of *Pomphorhynchus laevis* in *Salamandra salamandra*. *J. Helminthol.*, 69: 267 – 268. DOI: 10.1017/S0022149X00014267
- WALTON, A. C. (1933): The nematoda as parasites of Amphibia. *J. Parasitol.* 20: (1) 1 – 33
- YAMAGUTI, S. (1958): *Systema Helminthum*. Vol. I: *The Digenetic Trematodes of Vertebrates*. Interscience Publishers, New York & London 1575 pp
- YAMAGUTI, S. (1961): *Systema Helminthum: The nematodes of vertebrates*. Vol III., Part II. *Nematodes of Amphibians*. Intersciences Publishers Ltd. London, England 679 pp
- YAMAGUTI, S. (1963): *Systema Helminthum*. Acanthocephala. Vol. V. Intersciences Publishers, London, England. 423 pp
- YILDIRIMHAN, H. S., UĞURTAŞ, İ. H., ALTUNEL, F. N. (1996): An investigation on helminths of *Rana ridibunda* Pallas, 1771 (marsh frog). *Acta Parasitol. Tur.*, 20: 113 – 130 (in Turkish, with English abstract)
- YILDIRIMHAN, H. S., OĞUZ, M. C., UĞURTAŞ, İ. H. (1997a): An investigation on the nematodes of some tailless frogs (*Rana ridibunda*, *Bufo bufo* and *Pelobates syriacus*) collected from the Bursa regions. *Hacettepe Fen ve Müh. Bil. Derg.*, 18: 45 – 58 (in Turkish with English abstract)
- YILDIRIMHAN, H. S., UĞURTAŞ İ. H., ALTUNEL, F. N. (1997b): An investigation on parasitic helminths of *Rana macrocnemis* Boulenger, 1885 (Uludag frog). *Turk. J. Zool.* 21: 467 – 473
- YILDIRIMHAN, H. S. (1999a): Researches on parasitic helminths of *Bufo viridis* Laurenti, 1768 (Anura: Amphibia). *Turk. J. Zool.* 23: 177 – 195
- YILDIRIMHAN, H. S. (1999b): *Helminth fauna of lizards species belonging to Lacertidae (Reptilia) family distributed in Bursa and its around* (PhD thesis). Uludağ Univer-

- sity, Institute of Science. 120 pp, Bursa (In Turkish, with English abstract)
- YILDIRIMHAN, H. S., AYDOĞDU, A., UĞURTAŞ, I. H., ALTUNEL, F. N. (2001a): Helminth fauna of *Bombina bombina* (Linnaeus, 1767) (Fire-Bellied Toad) collected from Sakarya and Edirne (Turkey). *Acta Parasitol. Tur.*, 25 (3): 308 – 311 (in Turkish, with English abstract)
- YILDIRIMHAN, H. S., AYDOĞDU, A., UĞURTAŞ, I. H., ALTUNEL, F. N. (2001b): An investigation on Plathelminth and Acanthocephala of *Mertensiella caucasica* (Caucasian Salamander). *Acta Parasitol. Tur.*, 25: 393 – 397 (In Turkish, with English abstract)
- YILDIRIMHAN, H. S., KARADENİZ, E., GÜRKAN, E., KOYUN, M. (2005a): Metazoon parasites of the Marsh frog (*Rana ridibunda* Pallas, 1771; Anura) collected from the different regions in Turkey. *Acta Parasitol. Tur.*, 29 (2): 135 – 139 (In Turkish with English abstract)
- YILDIRIMHAN, H. S., BURSEY, C. R., GOLDBERG, S. R. (2005b): Helminth parasites of the Caucasian salamander, *Mertensiella caucasica*, from Turkey. *Comp. Parasitol.*, 72(1): 75 – 87. DOI: 10.1654/4152
- YILDIRIMHAN, H. S., GOLDBERG, S. R., BURSEY, C. R. (2006a): Helminth parasites of the Banded Frog, *Rana camerani* (Ranidae) from Turkey. *Comp. Parasitol.*, 73(2): 222 – 236. DOI: 10.1654/4229.1
- YILDIRIMHAN, H. S., BURSEY, C. R., GOLDBERG, S. R. (2006b): Helminth parasites of the Taurus frog, *Rana holtzi*, and the Uludag frog, *Rana macrocnemis*, with remarks on the helminth community of Turkish anurans. *Comp. Parasitol.*, 73 (2): 237 – 248. DOI: 10.1654/4191.1
- YILDIRIMHAN, H. S., ALTUNEL, F. N., UĞURTAŞ, I. H. (2006c): Helminth parasites of *Hyla arborea* (Linnaeus, 1758) (Tree Frog) collected from Bursa, Edirne and Sakarya. *Acta Parasitol. Tur.*, 30 (1): 56 – 59 (In Turkish with English abstract)
- YILDIRIMHAN, H. S., KARADENİZ, E. (2007): Helminth Parasites of the Common Toad, *Bufo bufo* (Linnaeus, 1758) (Anura: Bufonidae) from Northeast Turkey. *Comp. Parasitol.*, 74 (1): 176 – 178. DOI: 10.1654/4246.1
- YILDIRIMHAN, H. S. (2008): A Preliminary investigation on helminth fauna of *Triturus vittatus* (Jenyns, 1835) and *T. karelinii* (Strauch, 1870). *Acta Parasitol. Tur.*, 32 (2): 158 – 160 (In Turkish with English abstract)
- YILDIRIMHAN, H. S., BURSEY, C. R., GOLDBERG, S. R. (2009): Helminth Parasites of the Caucasian Parsley Frog, *Pelodytes caucasicus*, from Turkey. *Comp. Parasitol.*, 76(2): 247 – 257. DOI: 10.1654/4376.1
- YILDIRIMHAN, H. S., BURSEY, C. R. (2010): Helminth parasites of the eastern spadefoot toad, *Pelobates syriacus* (Pelobatidae), from Turkey. *Turk. J. Zool.*, 34: 11 – 319
- YILDIRIMHAN, H. S., BURSEY, C. R., ALTUNEL, F. N. (2011): Helminth parasites of the Balkan green lizard, *Lacerta trilineata* Bedriaga 1886, from Bursa, Turkey. *Turk. J. Zool.*, 35(4): 519 – 535
- YILDIRIMHAN, H. S., NURHAN, S., İNCEDOĞAN, S., BURSEY, C. R. (2012): Helminth parasites of the lemon-yellow tree Anura and Urodela, *Hyla savignyi* (Hylidae), from Turkey. *Turk. J. Zool.*, 36: 171 – 184
- YILDIZ, K. (2003): Helminth Infections in Tench (*Tinca tinca*) from Kapulukaya Dam Lake. *Turk. J. Vet. Anim. Sci.*, 27: 671 – 675 (In Turkish)
- ZIOLKOWSKA, M., ROKICKI, J. (2003): An Attempt to Determine the Intermediate Host for *Pomphorhynchus laevis* (Acanthocephala) in the Baltic Sea. *Acta Ichthyol. Pisc.*, 37 – 43

RECEIVED NOVEMBER 2, 2012

ACCEPTED DECEMBER 4, 2012