

Reconstruction of the pre-compactional thickness of the Zechstein Main Dolomite in northwest Poland

Grażyna Semyrka^{1*}, Marzena Gancarz², Zbigniew Mikołajewski³

¹AGH University of Science and Technology, Faculty of Geology, Geophysics and Environment Protection, Department of Fossil Fuels, Mickiewicza 30, 30-059 Kraków, Poland

²AGH University of Science and Technology, Faculty of Drilling, Oil and Gas, Department of Gas Engineering, Mickiewicza 30, 30-059 Kraków, Poland

³Polish Oil and Gas Co. SA, Branch of Geology and Exploitation, Department of Projects in Piła, Staszica 9, 64-920 Piła, Poland

* corresponding author, e-mail: gsemyrka@agh.edu.pl

Abstract

Our reconstruction of the pre-compactional thickness of the Main Dolomite strata from the so-called Grotów Peninsula (northwest Poland) was based on macroscopic observations of drill cores from three wells: Mokrzec-1, Sieraków-4 and Międzychód-5. These wells are located in various palaeogeographical zones of the Main Dolomite and cored rocks represent a range of microfacies. The amount of compactional reduction in thickness of the Main Dolomite was estimated by summing the total heights (W_{st}) of all stylolites encountered in logs of these wells. For calculations, a generalised model of a drill core was developed, which embraced all types of stylolite seams present in the Main Dolomite succession studied. Also the method of stylolite dimensioning was demonstrated. The number of stylolites in the drill cores studied varied from 511 in the Sieraków-4 well to 1,534 in the Międzychód-5 well. In all cores studied low-amplitude macrostylolites predominated, but the reduction of thickness was controlled mostly by the low- and medium-amplitude macrostylolites. The largest number of stylolites was found in the grainstone/packstone microfacies. The turnout of stylolites depends of microfacies. The highest density of stylolites was documented in mudstones/wackestones (24 stylolites per metre of rock thickness) and the lowest in boundstones (14 stylolites per metre of rock thickness). The low-amplitude stylolites appear most frequently in the mudstone/wackestone microfacies (15 stylolites per metre of rock thickness); in grainstones/packstones, rudstones/floatstones and boundstones middle-amplitude stylolites are rare (3 stylolites per metre of rock thickness). The degree of compaction of the Main Dolomite succession studied varied from 6 to 10%; hence, its calculated initial thickness also varied in the wells studied: from 41.3 m in the Sieraków-4 well to 56.9 m in the Mokrzec-1 well and to 97.1 m in the Międzychód-5 well. The volumes of reservoir fluids expelled during compaction of 1 m³ of Main Dolomite carbonates were estimated as 56 l in the Sieraków-4 well, 90 l in the Mokrzec-1 well and 97 l in the Międzychód-5 well.

Keywords: compaction, solution seam, stylolitic seam, thickness reduction

1. Introduction

The initial thickness of sediments can be reconstructed with various methods and from such reconstructions the amount of compaction can be estimated (see e.g., Shinn & Robin, 1983; Aplin et al., 1995; Goldhammer, 1997; Westphal, 1998;

Westphal & Munnecke, 1997; Broichhausen et al., 2005; Katsman & Aharonov, 2006; Kochman, 2006). However, the fastest and simplest method is macroscopic analysis of drill cores (Stockdale, 1926; Mossop, 1972, Waschs & Hein, 1974; Kaplan, 1976; Peacock & Azzam, 2006; Vandeginste & John, 2013).




Fig. 1. The study area plotted on a palaeogeographical map of the Main Dolomite in the Polish Zechstein Basin (modified after Kotarba & Wagner, 2007)

During compaction, the load from the overburden results in pressure solution of carbonate rocks and formation of stylolite seams. Hence, the thickness reduction of rocks (which is a measure of the amount of compaction) can be estimated by summing the heights of all stylolites existing in the interval studied.

The stylolitisation process plays a double role. First, it records the intensity of compaction and reflects the expulsion of reservoir fluids during primary migration and, second, it controls the transformation of reservoir from porous to dual, porous-fractured. The systems of tectonic fractures together with the systems of stylolite seams both influence the migration of reservoir fluids (Stockdale, 1926; Ramsden, 1952; Radlicz, 1966; Dunnington, 1967; Semyrka, 1985; Strzetelski, 1977; Koepnick, 1988; Aydin, 2000; Agosta et al., 2009; Agosta et al., 2010; Heap et al., 2014).

In the Polish Zechstein Basin stylolites are common in the Main Dolomite succession. The so-called "Grotów Peninsula" was selected as a study area because of the large number of core samples and analytical data and assessment reports available (Fig. 1).

In order to reconstruct the pre-compactional thickness of the Main Dolomite succession, a generalised model of a drill core was developed from the macroscopic observations of lithostratigraphic columns from 15 wells. The wells represent various palaeogeographical zones of the Main Dolomite ba-

sin, from toe-of-slope of a carbonate platform with a bay shoal to a barrier and a lagoon (Fig. 1).

There are five methods for estimating the thickness of material dissolved along a stylolite: the maximum stylolite height (Stockdale, 1926; Glover, 1968; Mossop, 1972; Kaplan, 1976; Bathurst, 1984), the thickness of a stylolite seam (Stockdale, 1926; Heald, 1955; Barrett, 1964), the condensation of heavy minerals along a stylolite seam (Young, 1945; Heald, 1955), the displacement of pre-existing veins (Conybeare, 1949; Bushinskyi, 1961; Wasch & Hein, 1974) and the reconstruction of truncated fossils and oolites (Bushinskyi, 1961; Wasch & Hein, 1974). One of the commonest methods is to measure maximum stylolite height. In the present paper we propose to measure the average height and thickness of stylolites as a means of determining thickness reduction and calculating compaction.

2. Compaction

Compaction is a diagenetic densification of sediment under the pressure exerted by the load of overburden. The diagenesis advances due to a combined effect of geometric (mechanical compaction) and mineralogical (chemical compaction) changes within the sediment controlled by its primary lithological composition and sedimentary conditions (Dunnington, 1967; Coogan, 1970; Bathurst, 1975, 1987, 1995; Moore, 2001; Flügel, 2004; Tucker &

Wright, 1990; Katsman et al., 2005; Ehrenberg, 2006; Twardowski & Traple, 2008).

The mechanical compaction proceeds with various intensity during early diagenesis, down to burial depth of about 200 m and results in dense packing of grains, expulsion of water and reduction of porosity/permeability (Dunington, 1967; Bathurst, 1975, 1987, 1995; Shinn & Robbin, 1983; Ricken, 1987; Choquette & James, 1990; Clari & Martire, 1996; Goldhammer, 1997; Moore, 2001; Katsman et al., 2005).

The chemical compaction starts at a burial depth of about 200-300 m and includes a variety of processes: cementation, dissolution and recrystallisation, which change the structure of sediment. Controlling factors are: temperature, type of porosity, degree of infilling of pore space, susceptibility of particular minerals to dissolution and the presence of clay minerals (Wanless, 1979; Buxton & Sibley, 1981; Scholle & Halley, 1985; Leythaeuser et al., 1995).

Dissolution of minerals under the load of overburden is named "pressure solution" and results in the formation of stylolitic seams, and solution seams. Usually, pressure solution develops along various discontinuities (e.g., bedding planes, fractures, clay laminae) and is controlled by the presence of pore fluids. Pressure solution is effective when maximum stress appears at grain-to-grain contacts and results in stylolitisation, removal of water from clay minerals and expulsion of reservoir fluids. Solutes are then transported to the zones of lower stress, in consistence with the Riecke principle (Füchtbauer, 1974). As a result, the stylolite seams can be filled with residue composed of e.g., clay minerals and/or organic matter (Park & Schot, 1968; Neugenbauer, 1973; Bathurst, 1975; Larsen & Chilingar, 1979; Choquette & James, 1990; Matyszkiewicz, 1996; Środoń, 1996; Agosta & Kirschner, 2003; Ehrenberg, 2006; Ben-Itzhak et al., 2012; Rustichelli et al., 2012).

Compaction affects all geological formations but its results are different in particular rocks depending of lithology and susceptibility of rocks to this process. It is generally believed that susceptibility of pure carbonates to mechanical compaction is insignificant. Calcium carbonate can be dissolved at the contacts of lithologically different rocks buried at depths of some hundreds of metres and then the secondary calcium carbonate can be precipitated as a pore cement. Hence, such rocks become lithified and thus resistant to mechanical compaction. Consequently, their initial thickness is preserved (Bathurst, 1975, 1987; Ricken, 1987; Matyszkiewicz, 1996; Moore, 2001; Kieft, 2002; Flügel, 2004; Tucker & Wright, 1990). Lithification is accelerated by un-

stable carbonate minerals and increasing temperature.

Considering details of the lithification process, the amount of compaction in the Main Dolomite was calculated only from the effects of stylolitisation.

3. Geological setting

The so-called "Grotów Peninsula" is located in the western part of the Polish Zechstein Basin (Fig. 1). The basin reveals highly diverse sea floor morphology which controlled facies development of the Main Dolomite rocks. In particular, we interpret deep- and shallow-marine, high- and low-energy environments varying in bathymetry and in microfacies. The Main Dolomite was deposited on carbonate platforms and microplatforms (barriers and lagoons), platform slope and toe-of-slope, and a carbonate ramp (for details see e.g., Peryt & Dyjaczynski, 1991; Wagner, 1994; Protas & Wojtkowiak, 2000; Jaworowski & Mikołajewski, 2007; Kotarba & Wagner, 2007; Słowakiewicz & Mikołajewski, 2009; Czeakański et al., 2010).

In the carbonate platform environment, the high-energy zones produced mostly the grainstones with boundstone horizons (mainly sublittoral carbonate muddy sands; also carbonate sands and carbonate sandy muds), whereas the low-energy zones were dominated by mudstones, wackestones and packstones with abundant bioclasts (mainly dark grey sublittoral carbonate sandy muds and carbonate muds; carbonate muddy sands and microbial sediments being frequent). In the carbonate barriers, the high-energy environments gave rise to grainstone and boundstone formation, rarely to packstones, wackestones, floatstones and rudstones. Peri- and sublittoral carbonate sands and microbial sediments are predominant. Carbonate muddy sands occur fairly frequently. Carbonate sandy muds and carbonate conglomerates are rare. The platform slopes, occupying the border zone between the shallow-marine, high-energy platform and the deep-marine, low-energy basinal plain produced a variety of microfacies: mudstones, wackestones, packstones, grainstones, floatstones, rudstones and boundstones. There is a co-occurrence of sublittoral carbonate sands and muddy sands, carbonate sandy muds and muds. Carbonate conglomerates, sedimentary breccia and microbial sediments have also been observed. Typical of the toe-of-slope environment were mudstones and packstones intercalated by floatstones. Mostly the same sediments as those known from both the plat-




Fig. 2. Evolution sequence of stylolites

form slope and basin floor. Especially characteristic is the presence of carbonate sands with a carbonate mud admixture, carbonate muds and interbeds of carbonate conglomerates. Anhydrite conglomerates are observed in the lower portion of the section. Finally, the low-energy basinal plain environment led to the formation of mostly mudstones and laminated strata with bio-sedimentary structures (sublittoral dark grey carbonate muds and carbonate sandy muds; occasional carbonate muddy sands, thin microbial sediments) (Wagner, 1994; Kotarba & Wagner, 2007; Jaworowski & Mikołajewski, 2007).

4. Methods

Macroscopic observations of drill cores from the Grotów Peninsula revealed the common presence of stylolite seams in the Main Dolomite succession. The stylolites differ in origin and morphology: from almost simple through wave-like to columnar and seismogram-like, according to the sequence of stylolite evolution (Fairbridge, 1968; Park & Schot, 1968; Strzetelski, 1977; Andrews & Railsbak, 1997; Sheppard, 2002; Sinha-Roy, 2002; Renard et al., 2004; Schmittbuhl et al., 2004; Brouste et al., 2007; Ben-Itzhak et al., 2012) (Fig. 2).

Stylolites observed in the Main Dolomite successions fall into three morphological types: wave-like, columnar and serrate with the two last-named most often encountered in samples studied. Apart from horizontal stylolites, concordant with the bedding planes and produced by increasing load from the overburden, we observed also slicolites, i.e., stylolites inclined to the bedding planes, which were produced by tectonic stress (Bushinskiy, 1961; Radlicz, 1966; Kijewski & Kaszper, 1973; Strzetelski, 1977; Peryt, 1978; Dadlez & Jaroszewski, 1994; Ebner et al., 2009; Bonnetier et al., 2009; Krzesińska et al., 2010).

Based on macroscopic observations, we developed a generalised model of a drill core, which presents various types of stylolite seams and methods of their dimensioning (Fig. 3).

The reduction of the length of a drill core (R_{st}) resulting from evolution of a single stylolite is a sum of all its elements (h), i.e., amplitude of leading wave and heights of columns, teeth and secondary peaks, and average aperture of stylolite (g_{st}), which gives the total height of a stylolite (W_{st}). Taking into account the limited resolution of the human eye for stylolite apertures smaller than 1 mm, we applied in calculations the double value of the average aperture of a stylolite ($2g_{st}$). We also assumed that stylolites of < 2 mm amplitudes are microstylolites and those of > 2 mm macrostylolites. Among the latter we distinguished low-amplitude (2–10 mm), medium-amplitude (10–50 mm) and high-amplitude (> 50 mm) stylolites.

In the drill-core model (Fig. 3) of recent length $H_{ob} = 15$ cm and diameter 3.5 cm, we included eight types of stylolite:

- wave-like, smooth macrostylolite - total height (W_{st1}) is the sum of average aperture ($g_{st} = 1$ mm) and amplitude of leading wave ($h_f = 3.5$ mm):

$$W_{st1} = g_{st} + h_f = 1 + 3.5 = 4.5 \text{ mm};$$

- simple, serrate microstylolite - total height (W_{st2}) is the sum of double aperture ($2g_{st} = 2 \cdot 0.5$ mm) and average height of teeth ($h_z = 1$ mm):

$$W_{st2} = 2g_{st} + h_z = 1 + 1 = 2 \text{ mm};$$

- wave-like, serrate macrostylolite - total height (W_{st3}) is the sum of double aperture ($2g_{st} = 2 \cdot 0.2$ mm), average amplitude of leading wave ($h_f = 5$ mm) and average height of teeth ($h_z = 1.5$ mm):

$$W_{st3} = 2g_{st} + h_f + h_z = 0.4 + 5 + 1.5 = 6.9 \text{ mm};$$

- wave-like, smooth, irregular macrostylolite - total height (W_{st4}) is the sum of double aperture ($2g_{st} = 2 \cdot 0.2$ mm), average height of secondary peaks ($h_o = 1.6$ mm) and amplitude of leading wave ($h_f = 4.5$ mm):

$$W_{st4} = 2g_{st} + h_o + h_f = 0.4 + 1.6 + 4.5 = 6.5 \text{ mm};$$




Fig. 3. Model of a drill core showing eight types of stylolite seams used for demonstration of a calculation method of compactional reduction of thickness.

1 - wave-like, smooth; 2 - simple, serrate; 4 - wave-like, smooth, irregular; 5 - seismogram-like; 6 - columnar, serrate; 7 - irregularly serrate; 8 - kink-like; h_f - amplitude of leading wave, g_{st} - aperture of stylolite, h_z - height of teeth, h_o - height of secondary peaks, h_{zm} - height of small teeth, h_{zd} - height of large teeth, h_k - height of column, h_w - height of kinks

- seismogram-like macrostylolite - total height (W_{st5}) is the sum of double aperture ($2g_{st} = 2 \cdot 0.6$ mm), average height of large teeth ($h_{zd} = 8.7$ mm) and average height of small teeth ($h_{zm} = 4.5$ mm) measured vertically:

$$W_{st5} = 2g_{st} + h_{zd} + h_{zm} = 1.2 + 8.7 + 4.5 = 14.4 \text{ mm};$$

- columnar, serrate macrostylolite - total height (W_{st6}) is the sum of double aperture ($2g_{st} = 2 \cdot 0.6$ mm), average height of columns ($h_k = 10.1$ mm) and average height of teeth ($h_z = 1.5$ mm):

$$W_{st6} = 2g_{st} + h_k + h_z = 1.2 + 10.1 + 1.5 = 12.8 \text{ mm};$$

- irregularly serrate microstylolite - total height (W_{st7}) is the sum of double aperture ($2g_{st} = 2 \cdot 0.2$ mm) and average height of teeth ($h_z = 1.3$ mm) measured vertically:

$$W_{st7} = 2g_{st} + h_z = 0.4 + 1.3 = 1.7 \text{ mm};$$

- kink-like macrostylolite - total height (W_{st8}) is the sum of average aperture ($g_{st} = 1.2$ mm) and average height of kinks ($h_w = 2$ mm) measured vertically:

$$W_{st8} = g_{st} + h_w = 1.2 + 2 = 3.2 \text{ mm}.$$

As concluded from the descriptions, proper identification and measurements of stylolite components are crucial for their dimensioning because average values of component heights are added in order to obtain the total height of a given stylolite (W_{st}) (Fig. 3).

The amount of stylolite-induced reduction of drill core length (R_{st}) is calculated as the sum of the heights of all stylolites in the core interval studied, according to the formula:

$$R_{st} = \sum W_{st1-8} = 0.45 + 0.2 + 0.69 + 0.65 + 1.44 + 1.28 + 0.17 + 0.32 = 5.2 \text{ cm}$$

Taking into account the present length of the core (H_{ob}) and the amount of reduction (R_{st}), we can calculate the approximate initial core length (H_p) as well as the degree of compaction K_{st} and the compaction coefficient k_{st} :

$$H_p = H_{ob} + R_{st} = 15 + 5.2 = 20.2 \text{ cm}$$

$$K_{st} = R_{st}/H_p \cdot 100\% = 5.2/20.2 \cdot 100\% = 26\%$$

$$k_{st} = R_{st}/H_p = 5.2/20.2 = 0.257$$

When the value of compaction coefficient (k_{calc}) is known we can approximate the volume of reservoir fluids (W_{pzi}) expelled from 1 m³ of rock during compaction. In the model presented this is $W_{\text{pzi}} = 2571/\text{m}^3$.

5. Results

The methodology of reconstruction of pre-compactional thickness of sediments and determination of the volume of reservoir fluids expelled during primary migration presented here was applied to the samples derived from three wells: Mokrzec-1, Sieraków-4 and Międzychód-5. These wells are located in specific palaeogeographical zones of the Grotów Peninsula. The Mokrzec-1 well represents a toe-of-slope of carbonate platform, the Międzychód-5 well is located in the barrier zone and the Sieraków-4 well represents a shoal within the carbonate ramp (Fig. 1). These wells were selected due to high diversity and greater number of stylolite seams contained in drill cores.

In the lithostratigraphic columns of the wells studied, the following microfacies groups were distinguished in the Main Dolomite succession: mud-dominated (mudstones/wackestones), grain-dominated (grainstones/packstones and rudstones/floatstones) and biogenic (boundstones) (Dunham, 1962; Gradziński et al., 1986; Jaworowski & Mikołajewski, 2007; Mikołajewski & Słowakiewicz, 2008).

5.1. The Mokrzec-1 well

The Mokrzec-1 well is located at the margin zone of the carbonate platform slope and the basinal plain (Fig.1). The Main Dolomite was found at a depth of 3,313.5–3,261.8 m (thickness: 51.7 m). Three microfacies groups were identified: mud-dominated, grain-dominated with prevailing




Fig. 4. Lithological and microfacies column of the Mokrzec-1 well (after Mikołajewski, 2007)

Table 1. Numbers and total stylolite heights in particular microfacies of the Main Dolomite in the Mokrzec-1 well

Microfacies	Total number of stylolites [n]	Total height of stylolites [mm]	Mokrzec-1 well			
			Micro-	Macrostylolites		
			< 2 mm	2–10 mm	10–50 mm	> 50 mm
Grainstone/packstone	428	2371	124	245	59	–
Mudstone/wackestone	555	2389	62	1315	994	–
Rudstone/floatstone	55	361	171	342	42	–
Total	1038	5121	85	1596	708	–
			137	31	11	–
			158	196	–	–
			308	618	112	–
			154	3069	1898	–




Fig. 5. Lithological and microfacies column of the Sieraków-4 well (after Mikołajewski, 2007)

Table 2. Numbers and total stylolite heights in particular microfacies of the Main Dolomite in the Sieraków-4 well

Microfacies	Total number of stylolites [n]	Total height of stylolites [mm]	Sieraków-4 well			
			Micro-		Macrostylolites	
			< 2 mm	2–10 mm	10–50 mm	> 50 mm
			Number of stylolites [n]		Total height of stylolites [mm]	
Boundstone	263	1082	87	156	20	-
Grainstone/ Packstone	209	859	44	720	318	-
Rudstone/ floatstone	39	390	68	122	19	-
			34	537	288	-
			10	14	15	-
			5	99	286	-
Total	511	2331	165	292	54	-
			83	1356	892	-

grainstones/packstones, and boundstones containing biosedimentary laminae (Fig. 4).

In the column analysed 1,038 stylolites were measured. Their total height is 5,121 mm, which corresponds to their compactional reduction of thickness due to stylolitisation. In detail, 308 microstylolites of a total height of 154 mm and 730 macrostylolites of a total height of 4,967 mm were analysed. Among macrostylolites, 618 belonged to the low-amplitude group (total height 3,069 mm) and 112 represented the medium-amplitude group (total height 1,898 mm) (Table 1). The largest number of stylolites was observed in the mudstone/wackestone microfacies. These are mostly low-amplitude stylolites and they contribute decisively to thickness reduction of this facies (and also to that of the grainstone/packstone microfacies), whereas medium-amplitude stylolites predominate in thickness reduction of the rudstone/floatstone microfacies (Table 1).

The Main Dolomite succession observed in the Mokrzec-1 well is dominated by grainstone/packstone and mudstone/wackestone microfacies (Fig. 4). Low-amplitude stylolites prevail and they control the overall thickness reduction of succession analysed (Table 1). Our observations revealed also that the highest amplitudes of stylolites occurred in mudstone/wackestone (33.1–42.3 mm) and grainstone/packstone (35.5–41.8 mm) microfacies.

The stylolite-generated thickness reduction of the Main Dolomite succession in the Mokrzec-1 well is $R_{st} = 5.1$ m and its reconstructed initial thickness is $H_p = 56.85$ m. The calculated coefficient of compaction is $k_{calc} = 0.090$, which enables us to estimate the volume of reservoir fluids expelled from the Main Dolomite due to compaction to be $W_{pzi} = 90$ l/m³ of rock.

5.2. The Sieraków-4 well

The Sieraków-4 belongs to the lagoonal zone of carbonate platform and is located within a local,

high-energy shoal (Fig.1). The Main Dolomite succession was encountered at a depth of 3,284–3,245 m (thickness: 39 m). A characteristic feature is the predominance of boundstones over grain-dominated microfacies (grainstones/packstones and rudstone/floatstone) (Fig. 5).

In the succession analysed 521 stylolites were measured, of total height 2,331 mm. Among them were 165 microstylolites of a total height of 83 mm and 346 macrostylolites of a total height of 2,248 mm. The macrostylolites included 292 low-amplitude (total height 1,356 mm) and 54 medium-amplitude (total height 892 mm) stylolites (Table 2). The largest number of stylolites was encountered in boundstones. Most of them were low-amplitude microstylolites, which controlled the reduction of thickness within this microfacies. Similar relationships were found in grainstone/packstone microfacies, but in rudstones/floatstones medium-amplitude macrostylolites were decisive for thickness reduction (Table 2). In the Sieraków-4 well, the Main Dolomite succession is dominated by the boundstone microfacies (Fig. 5) and low-amplitude macrostylolites, which controlled thickness reduction of the core interval studied (Table 2). Macroscopic observations showed that stylolites of highest amplitudes occur in boundstones (31.8–32.2 mm).

The amount of stylolite-induced thickness reduction of Main Dolomite strata in the Sieraków-4 well is $R_{st} = 2.3$ m and the reconstructed initial thickness is $H_p = 41.3$ m. The calculated coefficient of compaction is $k_{calc} = 0.056$, which allowed us to estimate the volume of reservoir fluids expelled due to compaction from the Main Dolomite at this locality as $W_{pzi} = 56$ l/m³ of rock.




Fig. 6. Lithological and microfacies column of the Międzychód-5 well (after Mikołajewski, 2007)

Table 3. Numbers and total stylolite heights in microfacies of the Main Dolomite in the Międzychód-5 well

Microfacies	Total number of stylolites [n]	Total height of stylolites [mm]	Międzychód-5 well			
			Micro- < 2 mm	Macrostylolites		
			2–10 mm	10–50 mm	> 50 mm	
			Number of stylolites [n]			
			Total height of stylolites [mm]			
Grainstone/ Packstone	1147	6281	323	680	140	4
			162	3505	2401	213
Rudstone/ floatstone	284	2339	43	168	71	2
			22	948	1267	102
Boundstone	80	718	16	37	27	-
			8	255	455	-
Mudstone/ wackestone	23	144	7	12	4	-
			4	81	59	-
Total	1534	9482	389	897	242	6
			196	4789	4182	315

5.3. The Międzychód-5 well

The Międzychód-5 well is located in the barrier zone of the Main Dolomite (Fig. 1). Its succession was found at a depth of 3,217.5–3,129.8 m (thickness: 87.7 m). A diverse assemblage of microfacies was identified, inclusive of grain-dominated (grainstones/packstones, rudstones/floatstones) and biogenic (boundstones) ones. Rarely, a mud-dominated microfacies (mudstones/wackestones) was observed as well (Fig. 6).

In the succession analysed, we measured 1,534 stylolites of a total height of 9,482 mm. Among them were 389 microstylolites of a total height of 196 mm and 1,145 macrostylolites of a total height of 9,286 mm. Macrostylolites included 897 low-amplitude examples (total height 4,789 mm), 242 medium-amplitude ones (total height 4,182 mm) and six high-amplitude stylolites (total height 315 mm) (Table 3). The largest number of stylolites (mostly low-amplitude macrostylolites) was observed in grainstone/packstone microfacies and these forms contributed decisively to the reduction of its thickness. In both boundstones and rudstones/floatstones, medium-amplitude macrostylolites prevailed and controlled thickness reduction (Table 3). In the Międzychód-5 well, the Main Dolomite succession is dominated by grainstone/packstone microfacies (Fig. 6). Commonest are low-amplitude macrostylolites, but low- and medium-amplitude macrostylolites similarly contributed to thickness reduction (Table 3). Macroscopic observations revealed that stylolites of highest amplitudes occur in grainstones/packstones (52–55.5 mm) and rudstones/floatstones (50.4–51.8 mm) microfacies.

The stylolite-induced reduction of Main Dolomite thickness in the Międzychód-5 well is $R_{st} = 9.4$ m and the reconstructed initial thickness is $H_p = 97.1$ m. The calculated coefficient of compaction $k_{calc} = 0.097$ enabled us to estimate the volume of reservoir fluids expelled from the sediments studied during compaction to be $W_{pzi} = 97$ l/m³ of rock.

6. Discussion and conclusions

In the study area stylolitisation of Main Dolomite sediments is a common feature. The amount of thickness reduction corresponding to the amount of compaction was determined in the three wells; Mokrzec-1, Sieraków-4 and Międzychód-5, which represent various microfacies types reflecting a range of depositional environments.

The number of stylolites observed in the Main Dolomite succession studied varied from 511 in

the Sieraków-4 well to 1,534 in the Międzychód-5 well. The highest density of stylolites was found in the Mokrzec-1 well: 20 stylolites per metre of rock thickness and the lowest density of stylolites in the Sieraków-4 well: 15 stylolites.

The turnout of stylolites depends of microfacies. The highest density of stylolites was found in mudstones/wackestones (24 stylolites per metre of rock thickness) and the lowest density in boundstones (14 stylolites). The low-amplitude stylolites appear most frequently in mudstone/wackestone microfacies (15 stylolites per metre of rock thickness); rare are middle-amplitude stylolites in grainstones/packstones, rudstones/floatstones and boundstones (3 stylolites per metre of rock thickness).

The analysis of drill cores revealed that the occurrence of stylolites is not restricted to any particular palaeogeographical zone of the Main Dolomite carbonate platform (toe-of-slope, barrier, carbonate ramp) and does not depend of depth. All types of stylolites distinguished are more or less regular and show variable amplitudes and apertures. In all three wells, low-amplitude macrostylolites predominated, but thickness reduction was controlled mostly by low- and medium-amplitude macrostylolites (e.g., in the Międzychód well the stylolite-induced reduction thickness is $R_{st} = 9.4$ m and total height of low-amplitude stylolite is 4.789 m and total height of medium-amplitude stylolite is 4.182 m). Hence, we conclude that the amount of thickness reduction depends of the amplitudes of stylolites and on their number in the successions analysed. This is confirmed by the relatively great thickness reduction (315 mm) estimated in the Międzychód-5 well which resulted from the action of merely six high-amplitude macrostylolites. In the wells studied, the amount of thickness reduction changed from 2.3 m in the Sieraków-4 well, through 5.1 m in the Mokrzec-1 well to 9.4 m in the Międzychód-5 well. The calculated degrees of compaction were: 6%, 9% and 10%, respectively. The amounts of thickness reduction calculated in our study, 6 to 10%, correspond very well to values published by Stockdale (1926) for the Columbus Limestone (approx. 5%), by Mossop (1972) for the Reef Limestone (5.3 to 7.8%), by Waschs & Hein (1974) for the Franciscan Limestone (10%), by Kaplan (1976) for limestones (approx. 7–8%), by Peacock & Azzam (2006) for limestones and dolomites (3.8 to 7%) and by Vandeginste & John (2013) for limestone (7–12%).

The initial thickness of Main Dolomite sediments calculated from the amount of thickness reduction was 41.3 m in the Sieraków-4 well, 56.9 m in the Mokrzec-1 well and 97.1 m in the Międzychód-5 well.

The volume of reservoir fluids expelled by compaction during primary migration of Main Dolomite strata depended mostly of stylolite density and on their heights (amplitudes). Our calculations revealed that 1 m³ of Main Dolomite carbonate supplied variable volumes of reservoir fluids; from 56 l in the Sieraków-4 well, through 90 l in the Mokrzec-1 well to 97 l in the Międzychód-5 well.

Acknowledgements

The results presented here were obtained in a research project entitled "The importance of the stylolitisation process for petroleum potential of carbonates from the Main Dolomite", which was financed by the National Centre for Science, project No. ODW – 0601/B/P01/2011/40 (AGH UST No. 18.18.140.083) managed by one of us (GS).

References

- Agosta, F. & Kirschner, D.L., 2003. Fluid conduits in carbonate-hosted seismogenic normal faults of Central Italy. *Journal of Geophysical Research* 108, B4, 1–13.
- Agosta, F., Alessandrini, M., Tondi, E. & Aydin, A., 2009. Oblique normal faulting along the northern edge of the Majella anticline, central Italy: inferences on hydrocarbon migration and accumulation. *Journal of Structural Geology* 31, 674–690.
- Agosta, F., Alessandrini, M., Antonellini, M., Tondi, E. & Giorgioni, M., 2010. From fractures to flow: a field-based quantitative analysis of an outcropping carbonate reservoir. *Tectonophysics* 490, 197–213.
- Andrews, L.M. & Railsbak, L.B., 1997. Controls on stylolite development: morphologic, lithologic, and temporal evidence from bedding-parallel and transverse stylolites from the US Appalachians. *Journal of Geology* 105, 59–73.
- Aplin, A.C., Yang, Y. & Hansen, S., 1995. Assessment of the compression coefficient of mudstones and its relationship with detailed lithology. *Marine and Petroleum Geology* 12, 995–963.
- Aydin, A., 2000. Fractures, faults, and hydrocarbon entrapment, migration and flow. *Marine and Petroleum Geology* 17, 797–814.
- Barrett, P. J., 1964. Residual Seams and Cementation in Oligocene Shell Calcarenites, Te Kuiti Group. *Journal of Sedimentary Petrology* 34, 524–531.
- Bathurst, R.G.C., 1975. *Carbonate Sediments and their Diagenesis*. Elsevier, Amsterdam, 658.
- Bathurst, R.G.C., 1984. The integration of pressure solution with mechanical compaction and cementation. In: Yahya, F.A. (Ed.), *Stylolites and associated phenomena. Relevance to Hydrocarbon Reservoirs*. Abu Dhabi National Reserves. Found., 41–55.
- Bathurst, R.G.C., 1987. Diagenetically enhanced bedding in argillaceous platform limestone: stratified cementation and selective compaction. *Sedimentology* 34, 749–779.
- Bathurst, R.G.C., 1995. Burial diagenesis of limestones under simple overburden. Stylolites, cementation, and feedback: *Bulletin de La Societe Geologique de France* 166, 181–192.
- Ben-Itzhak, L.L., Aharonov, E., Toussaint, R. & Sagy, A., 2012. Upper bound on stylolite roughness as indicator for amount of dissolution. *Earth and Planetary Science Letters* 337–338, 186–196.
- Bonnetier, E., Misbah, C., Renard, F., Toussaint, R. & Gratier, J. P., 2009. Does roughening of rock-fluid-rock interfaces emerge from a stress-induced instability? *European Physical Journal B* 67, 121–131.
- Broichhausen, H., Littke, R. & Hantschel, T., 2005. Mudstone compaction and its influence on overpressure generation, elucidated by 3D case study in the North Sea. *International Journal of Earth Sciences* 94, 956–978.
- Brouste, A., Renard, F., Gratier, J.P. & Schmittbuhl, J., 2007. Variety of stylolites morphologies and statistical characterization of the amount of heterogeneities in the rock. *Journal of Structural Geology* 29, 422–434.
- Bushinskiy, G.I., 1961. Stylolites. *Izvestiya Akademia Nauk S.S.S.R., Serie Correlación Geológica* 8, 31–46.
- Buxton, T.M. & Sibley, D.F., 1981. Pressure solution features in a shallow buried limestone. *Journal of Sedimentary Petrology* 51, 19–26.
- Choquette, P.W. & James, N.P., 1990. Limestones – The Burial Diagenetic Environment. *Geoscience, Canada*, 75–112.
- Clari, P. & Martire, L., 1996. Interplay of cementation, mechanical compaction, and chemical compaction in nodular limestones of the Rosso Ammonitico Veronese (middle-upper Jurassic, northeastern Italy). *Journal of Sedimentary Research* 66, 447–458.
- Conybeare, C.E.B., 1949. Stylolites in Pre-Cambrian quartzite. *Journal of Geology* 57, 83–85.
- Coogan, A.H., 1970. Measurement of compaction in oolitic grainstone. *Journal of Sedimentary Petrology* 40, 921–929.
- Czekański, E., Kwolek, K. & Mikołajewski, Z., 2010. Złoża węglowodorów w utworach cechsztyńskiego dolomitu głównego (Ca₂) na bloku Gorzowa [Hydrocarbon fields in the Zechstein Main Dolomite (Ca₂) of the Gorzów Block (NW Poland)]. *Przegląd Geologiczny* 58, 695–703.
- Dadlez, R. & Jaroszewski, W., 1994. *Tektonika [Tectonics]* Wydawnictwo Naukowe PWN, Warszawa, 743.
- Dunham, R.J., 1962. Classification of carbonate rocks according to depositional texture. In: Ham, W.E., (Ed.): *Classification of carbonate rocks*. A Symposium of American Association of Petroleum Geology, 1, 108–121.
- Dunnington, H.V., 1967. Aspects of diagenesis and shape change in stylolitic limestone reservoirs. Proceedings of the 7th World Petroleum Congress. *Journal of the Middle East Petroleum Geosciences* 339–352.
- Ebner, M., Koehn, D., Toussaint, R., Renard, F. & Schmittbuhl, J., 2009. Stress sensitivity of stylolite

- morphology. *Earth and Planetary Science Letters* 277, 394–398.
- Ehrenberg, S.M., 2006. Porosity destruction in carbonate platforms. *Journal of Petroleum Geology* 29, 41–55.
- Fairbridge, R.W., 1968. *Encyclopedia of Geomorphology*. Dowden, Hutchinson and Ross, Pennsylvania, 1295 pp.
- Flügel, E., 2004. *Microfacies of carbonate rocks. Analysis, Interpretation and Application*. Springer, New York, 983.
- Füchtbauer, H., 1974. *Sediments and Sedimentary Rocks*, 1. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, 1–464.
- Glover, J. E., 1968. Significance of stylolites in dolomitic limestones. *Nature* 217, 835–836.
- Goldhammer, R.K., 1997. Compaction and decompaction algorithms for sedimentary carbonates. *Journal of Sedimentary Research* 67, 26–35.
- Gradziński, R., Kostecka, A., Radomski, A. & Unrug, R., 1986. *Zarys sedimentologii [Outline of Sedimentology]*. Wydawnictwa Geologiczne, Warszawa, 628 pp.
- Heald, M.T., 1955. Stylolites in sandstone. *Journal of Geology* 63, 101–114.
- Heap, M.J., Baud, P., Reuschlé, T. & Meredith, P.G., 2014. Stylolites in limestones: Barriers to fluid flow? *Geology* 42, 51–54.
- Jaworowski, K. & Mikołajewski, Z., 2007. Oil- and gas-bearing sediments of the Main Dolomite (Ca₂) in the Międzychód region: a depositional model and the problem of the boundary between the second and third depositional sequences in the Polish Zechstein Basin. *Przeгляд Geologiczny* 55, 1017–1024.
- Kaplan, M.Ye., 1976. Origin of stylolites. *Earth Science Section* 211, 205–207.
- Katsman, R. & Aharonov, E., 2006. A study of compaction bands originating from crack, notches, and compacted defects. *Journal of Structural Geology* 28, 508–518.
- Katsman, R., Aharonov, E. & Scher, H., 2005. Numerical simulation of compaction bands in high-porosity sedimentary rock. *Mechanics of Materials* 37, 143–162.
- Kiełt, M., 2002. *Geofizyka wiertnicza w poszukiwaniach węglowodorów. Strukturalne i sedimentologiczne zastosowanie otworowych profilowań geofizycznych [Well-log geophysics in hydrocarbon exploration. Structural and sedimentological application of geophysical logs]*. Adam Marszałek Publishing House, Toruń, 543.
- Kijewski, P. & Kaszper, J., 1973. Tekstury stylolitowe w cechsztyńskich skałach węglanowych poziomu W1 monokliny przedsudeckiej [Stylolitic textures in the Zechstein carbonate rocks of the horizon W1 of the Fore-Sudetic Monocline]. *Geological Quarterly* 17, 497–506.
- Kochman, A., 2006. Wybrane metody szacowania kompaktacji w osadach węglanowych [Different methods for reconstruction of compaction applied in limestones]. *Technika Poszukiwań Geologicznych: Geotermia, Zrównoważony Rozwój* 45, 35–43.
- Koepnick, R.B., 1988. Significance of Stylolite Development in Hydrocarbon Reservoirs with an Emphasis on the Lower Cretaceous of the Middle East. *Geological Society of Malaysia, Bulletin* 22, 23–43.
- Kotarba, M. & Wagner, R., 2007. Generation potential of the Zechstein Main Dolomite (Ca₂) carbonates in the Gorzów Wielkopolski-Międzychód-Lubiatów area: geological and geochemical approach to microbial-algal source rock. *Przeгляд Geologiczny* 55, 1025–1036.
- Krzesińska, A., Redlińska-Marczyńska, A., Wilkosz, P. & Żelaźniewicz, A., 2010. Struktury hydratacyjne i deformacyjne w skałach czapy gipsowej wysadu solnego Dębiny w rowie Kleszczowa [Deformation and hydrational structures in cap rocks of the *Dębina Salt Dome*, the Kleszczów Graben, central Poland]. *Przeгляд Geologiczny* 58, 522–530.
- Larsen, G. & Chilingar, G.V., 1979. *Diagenesis in Sediments and Sedimentary Rocks*. Elsevier, Amsterdam, 579 pp.
- Leythaeuser, D., Borromeo, O., Mosca, F., Primio, R., Radke, M. & Schaefer, R.G., 1995. Pressure solution in carbonate source rocks and its control on petroleum generation and migration. *Marine and Petroleum Geology* 12, 711–733.
- Matyszkiewicz, J., 1996. Wybrane problemy diagenety osadów węglanowych [Selected problems of diagenesis of carbonate rocks]. *Przeгляд Geologiczny* 44, 596–603.
- Mikołajewski, Z. & Słowakiewicz, M., 2008. Microfacies and diagenesis of the Main Dolomite (Ca₂) strata in the Międzychód barrier area (Grotów Peninsula, Western Poland). *Biuletyn Państwowego Instytutu Geologicznego* 429, 191–198.
- Moore, C.H., 2001. *Carbonate Reservoirs: Porosity Evolution and Diagenesis in a Sequence Stratigraphic Framework*. Elsevier, Amsterdam, 444 pp.
- Mossop, G.D., 1972. Origin of the peripheral rim, Redwater Reef, Alberta. *Bulletin of Canadian Petroleum Geology* 20, 238–280.
- Neugenbauer, J., 1973. The diagenetic problem of chalk the role of pressure solution and pore fluid. *Neues Jahrbuch für Geologie und Paläontologie* 143, 223–245.
- Park, W.C. & Schot, E.K., 1968. Stylolites: Their nature and origin. *Journal of Sedimentary Petrology* 38, 175–191.
- Peacock, D.C.P. & Azzam, I.N., 2006. Development and scaling relationships of a stylolite population. *Journal of Structural Geology* 28, 1883–1889.
- Peryt, T. M., 1978. Charakterystyka mikrofacjalna cechsztyńskich osadów węglanowych cyklotemu pierwszego i drugiego na obszarze Monokliny Przedsudeckiej [Microfacies of the carbonate sediments of the Zechstein Werra and Stassfurt cyclothems in the Fore-Sudetic Monocline]. *Studia Geologica Polonica* 54, 1–88.
- Peryt, T.M. & Dyjaczynski, K., 1991. An isolated carbonate bank in the Zechstein Main Dolomite basin, Western Poland. *Journal of Petroleum Geology* 14, 445–458.
- Protas, A., Wojtkowiak, Z., 2000. Blok Gorzowa. Geologia dolnego cechsztynu [The Gorzów Block. Geology of the Lower Zechstein]. *Guide to 71st Congress of the Polish Geological Society*, 163–171.
- Radlicz, K., 1966. Tekstury stylolitowe [The structures of stylolites]. *Geological Quarterly* 10, 367–382.
- Ramsden, R.M., 1952. Stylolites and oil migration. *American Association of Petroleum Geologists Bulletin* 36, 2185–2192.

- Renard, F., Schmittbuhl, J., Gratier, J.P., Meakin, P. & Merino, E.M., 2004. Three-dimensional roughness of stylolites in limestones. *Journal of Geophysical Research* 109, B3, 1–12.
- Ricken, W., 1987. The carbonate compaction law: a new tool. *Sedimentology* 34, 571–584.
- Rustichelli, A., Tondi, E., Agosta, F., Cilona, A. & Giorgioni, M., 2012. Development and distribution of bed-parallel compaction bands and pressure solution seams in carbonates (Bolognano Formation, Majella Mountain, Italy). *Journal of Structural Geology* 37, 181–199.
- Schmittbuhl, J., Renard, F., Gratier, J.P. & Toussaint, R., 2004. Roughness of Stylolites: Implications of 3D High Resolution Topography Measurements. *The American Physical Society* 93, 1–4.
- Scholle, P.A. & Halley, R.B., 1985. Burial diagenesis: out of sight, out of mind. In: Carbonate Cements. *Society of Economic Paleontologists and Mineralogists Special Publication* 36, 135–160.
- Semyrka, R., 1985. Uwarunkowania roponośności dolomitu głównego na obszarze Pomorza Zachodniego [Dependences of oil-bearing capacity of Main Dolomite in the region of Pomorze Zachodnie]. *Prace Geologiczne Polskiej Akademii Nauk* 129, 1–113.
- Sheppard, T.H., 2002. Stylolite development at sites of primary and diagenetic fabric contrast within the Sutton Stone (Lower Lias), Ogmores-by-Sea, Glamorgan, UK. *Proceedings of the Geologists Association* 113, 97–109.
- Shinn, E.A. & Robbin, D.M., 1983. Mechanical and chemical compaction in fine-grained shallow-water limestones. *Journal of Petroleum Geology* 53, 595–618.
- Sinha-Roy, S., 2002. Kinetics of differentiated stylolite formation. *Current Science* 82, 1038–1046.
- Słowakiewicz, M. & Mikołajewski, Z., 2009. Sequence stratigraphy of the Upper Permian Zechstein Main Dolomite carbonates in Western Poland: a new approach. *Journal of Petroleum Geology* 32, 215–234.
- Stockdale, P.B., 1926. The stratigraphic significance of solution in rocks. *Journal of Geology* 34, 399–414.
- Strzetelski, W., 1977. Rozwój procesów stylolityzacji i deformacji epigenetycznych w aspekcie roponośności piaskowców kwarcytowych kambru środkowego w rejonie Żarnowca [The evolution of stylolitization and epigenetic deformations in the Middle Cambrian oil-bearing quartzose sandstones in the area of Żarnowiec (Northern Poland)]. *Rocznik Polskiego Towarzystwa Geologicznego* 47, 559–584.
- Środoń, J., 1996. Minerale ilaste w procesach diagenetyzacji [Clay minerals in diagenetic processes]. *Przegląd Geologiczny* 44, 604–607.
- Tucker, M.E. & Wright, V.P., 1990. *Carbonate Sedimentology*. Blackwell, Oxford, 482 pp.
- Twardowski, K. & Traple, J., 2008. O kompaktacji utworów geologicznych. [Compaction of geologic formations]. *Wiertnictwo, Nafta, Gaz* 25, 53–62.
- Vandeginste, V. & John, C.M., 2013. Diagenetic implications of stylolitization in pelagic carbonates, Canterbury Basin, Offshore New Zealand. *Journal of Sedimentary Research* 83, 226–240.
- Wagner, R., 1994. Stratigraphy and evolution of the Zechstein basin in the Polish Lowland. *Prace Państwowego Instytutu Geologicznego* 166, 1–71.
- Wanless, H.R., 1979. Limestone response to stress: pressure solution and dolomitization. *Journal of Sedimentary Petrology* 49, 437–462.
- Waschs, D. & Hein, J.R., 1974. Petrography and diagenesis of Franciscan limestone. *Journal of Sedimentary Petrology* 44, 1217–1231.
- Westphal, H., 1998. *Carbonate platform slopes – a record of changing conditions. The Pliocene of the Bahamas*. Lecture Notes in Earth Sciences 75, Springer, Heidelberg, 197.
- Westphal, H. & Munnecke, A., 1997. Mechanical compaction versus early cementation in fine-grained limestones: differentiation by the presentation of organic microfossils. *Sedimentary Geology* 112, 33–42.
- Young, R.B., 1945. Stylolitic solution in Witwatersrand quartzites. *Transactions of Geological Society of South Africa* 47, 137–142.

Manuscript received: 20 March 2015
Revision accepted: 15 September 2015