

## On a $(p, k)$ -analogue of the Gamma function and some associated Inequalities

K. NANTOMAH<sup>a</sup>, E. PREMPEH<sup>b</sup> AND S. B. TWUM<sup>a</sup>

---

**ABSTRACT.** In this paper, we introduce a new two-parameter deformation of the classical Gamma function, which we call a  $(p, k)$ -analogue of the Gamma function. We also provide some identities generalizing those satisfied by the classical Gamma function. Furthermore, we establish some inequalities involving this new function.

**2010 Mathematics Subject Classification.** 33B15, 26D07, 26D15.

**Key words and phrases.** Gamma function,  $(p, k)$ -analogue, inequality.

---

### 1. Introduction

The classical Euler's Gamma function,  $\Gamma(x)$  is usually defined for  $x > 0$  by

$$\Gamma(x) = \int_0^{\infty} t^{x-1} e^{-t} dt = \lim_{n \rightarrow \infty} \frac{n! n^x}{x(x+1)(x+2) \dots (x+n)}$$

It is well-known that  $\Gamma(x)$  satisfies the following basic relations.

$$\begin{aligned} \Gamma(n+1) &= n!, \quad n \in \mathbb{Z}^+ \cup \{0\}, \\ \Gamma(x+1) &= x\Gamma(x), \quad x \in \mathbb{R}^+. \end{aligned}$$

---

Received February 20, 2016 - Accepted June 08, 2016.

©The Author(s) 2016. This article is published with open access by Sidi Mohamed Ben Abdallah University

<sup>a</sup>Department of Mathematics, University for Development Studies, Navrongo Campus, P. O. Box 24, Navrongo, UE/R, Ghana.

e-mail: mykwarasoft@yahoo.com, knantomah@uds.edu.gh  
 stwum@uds.edu.gh

<sup>b</sup>Department of Mathematics, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana.

e-mail: eprempeh.cos@knust.edu.gh

Closely associated with the Gamma function is the Digamma or Psi function  $\psi(x)$ , which is defined for  $x > 0$  as the logarithmic derivative of the Gamma function. That is,

$$\begin{aligned}\psi(x) &= \frac{d}{dx} \ln \Gamma(x) = \frac{\Gamma'(x)}{\Gamma(x)}, \\ &= -\gamma + (x-1) \sum_{n=0}^{\infty} \frac{1}{(n+1)(n+x)}, \\ &= -\gamma - \frac{1}{x} + \sum_{n=1}^{\infty} \frac{x}{n(n+x)}\end{aligned}$$

where  $\gamma = \lim_{n \rightarrow \infty} (\sum_{k=1}^n \frac{1}{k} - \ln n) = 0.577215664\dots$  is the Euler-Mascheroni's constant. The Polygamma functions,  $\psi^{(m)}(x)$  are defined for  $x > 0$  and  $m \in \mathbb{N}$  as

$$\begin{aligned}\psi^{(m)}(x) &= \frac{d^m}{dx^m} \psi(x) = \frac{d^{m+1}}{dx^{m+1}} \ln \Gamma(x) \\ &= (-1)^{m+1} m! \sum_{n=0}^{\infty} \frac{1}{(n+x)^{m+1}}\end{aligned}$$

where  $\psi^{(0)}(x) \equiv \psi(x)$ .

The  $p$ -analogue (also known as  $p$ -extension or  $p$ -deformation) of the Gamma function is defined for  $p \in \mathbb{N}$  and  $x > 0$  as

$$\Gamma_p(x) = \frac{p! p^x}{x(x+1)\dots(x+p)}$$

where  $\lim_{p \rightarrow \infty} \Gamma_p(x) = \Gamma(x)$ . See [1, p. 270]. It satisfies the identities:

$$\begin{aligned}\Gamma_p(x+1) &= \frac{px}{x+p+1} \Gamma_p(x), \\ \Gamma_p(1) &= \frac{p}{p+1}.\end{aligned}$$

The  $p$ -analogues of the Digamma and Polygamma functions are defined for  $x > 0$  as

$$\begin{aligned}\psi_p(x) &= \frac{d}{dx} \ln \Gamma_p(x) = \ln p - \sum_{n=0}^p \frac{1}{n+x}, \\ \psi_p^{(m)}(x) &= \frac{d^m}{dx^m} \psi_p(x) = (-1)^{m-1} m! \sum_{n=0}^p \frac{1}{(n+x)^{m+1}}\end{aligned}$$

where  $\psi_p^{(0)}(x) \equiv \psi_p(x)$ .

In 2007, Díaz and Pariguan [2] also defined the  $k$ -analogue of the Gamma function for  $k > 0$  and  $x \in \mathbb{C} \setminus k\mathbb{Z}^-$  as

$$\Gamma_k(x) = \int_0^\infty t^{x-1} e^{-\frac{t^k}{k}} dt = \lim_{n \rightarrow \infty} \frac{n! k^n (nk)^{\frac{x}{k}-1}}{(x)_{n,k}}$$

where  $\lim_{k \rightarrow 1} \Gamma_k(x) = \Gamma(x)$  and  $(x)_{n,k} = x(x+k)(x+2k)\dots(x+(n-1)k)$  is the Pochhammer  $k$ -symbol. The  $k$ -analogue also satisfies the identities:

$$\begin{aligned} \Gamma_k(x+k) &= x\Gamma_k(x), \quad x \in \mathbb{R}^+ \\ \Gamma_k(k) &= 1 \end{aligned}$$

Similarly, the  $k$ -analogues of the Digamma and Polygamma functions are defined for  $x > 0$  as

$$\begin{aligned} \psi_k(x) &= \frac{d}{dx} \ln \Gamma_k(x) = \frac{\ln k - \gamma}{k} - \frac{1}{x} + \sum_{n=1}^{\infty} \frac{x}{nk(nk+x)}, \\ \psi_k^{(m)}(x) &= \frac{d^m}{dx^m} \psi_k(x) = (-1)^{m+1} m! \sum_{n=0}^{\infty} \frac{1}{(nk+x)^{m+1}} \end{aligned}$$

where  $\psi_k^{(0)}(x) \equiv \psi_k(x)$ .

The purpose of this paper is to introduce a new two-parameter deformation of the classical Gamma function, called a  $(p, k)$ -analogue of the Gamma function. In addition, we provide some identities and inequalities involving this function. We present our results in the following section.

## 2. Results and Discussion

**Definition 2.1.** Let  $p \in \mathbb{N}$  and  $k > 0$ . Then the  $(p, k)$ -analogue (also called the  $(p, k)$ -deformation or  $(p, k)$ -generalization) of the Gamma function is defined as

$$\Gamma_{p,k}(x) = \int_0^p t^{x-1} \left(1 - \frac{t^k}{pk}\right)^p dt \quad (1)$$

$$= \frac{(p+1)! k^{p+1} (pk)^{\frac{x}{k}-1}}{x(x+k)(x+2k)\dots(x+pk)} \quad (2)$$

for  $x \in \mathbb{R}^+$ . It satisfies the identities:

$$\Gamma_{p,k}(x+k) = \frac{pkx}{x+pk+k} \Gamma_{p,k}(x), \quad (3)$$

$$\Gamma_{p,k}(ak) = \frac{p+1}{p} k^{a-1} \Gamma_p(a), \quad a \in \mathbb{R}^+ \quad (4)$$

$$\Gamma_{p,k}(k) = 1. \quad (5)$$

Also, observe that  $\Gamma_{p,k}(x)$  satisfies the following commutative diagram.

$$\begin{array}{ccc} \Gamma_{p,k}(x) & \xrightarrow{p \rightarrow \infty} & \Gamma_k(x) \\ k \rightarrow 1 \downarrow & & \downarrow k \rightarrow 1 \\ \Gamma_p(x) & \xrightarrow{p \rightarrow \infty} & \Gamma(x) \end{array}$$

The  $(p, k)$ -analogue of the Digamma function is defined as the logarithmic derivative of  $\Gamma_{p,k}(x)$ . That is

$$\psi_{p,k}(x) = \frac{d}{dx} \ln \Gamma_{p,k}(x) = \frac{\Gamma'_{p,k}(x)}{\Gamma_{p,k}(x)}.$$

The function  $\psi_{p,k}(x)$  satisfies the following series and integral representations.

$$\psi_{p,k}(x) = \frac{1}{k} \ln(pk) - \sum_{n=0}^p \frac{1}{(nk+x)} \quad (6)$$

$$= \frac{1}{k} \ln(pk) - \int_0^\infty \frac{1 - e^{-k(p+1)t}}{1 - e^{-kt}} e^{-xt} dt \quad (7)$$

The  $(p, k)$ -analogue of the Polygamma functions are defined as

$$\psi_{p,k}^{(m)}(x) = \frac{d^m}{dx^m} \psi_{p,k}(x) = (-1)^{m+1} m! \sum_{n=0}^p \frac{1}{(nk+x)^{m+1}} \quad (8)$$

$$= (-1)^{m+1} \int_0^\infty \left( \frac{1 - e^{-k(p+1)t}}{1 - e^{-kt}} \right) t^m e^{-xt} dt \quad (9)$$

for  $m \in \mathbb{N}$ , where  $\psi_{p,k}^{(0)}(x) \equiv \psi_{p,k}(x)$ . It follows easily from (8) that,

$$\psi_{p,k}^{(m)}(x) = \begin{cases} > 0 & \text{if } m \text{ is odd} \\ < 0 & \text{if } m \text{ is even,} \end{cases} \quad (10)$$

which means that the function  $\psi'_{p,k}(x)$  is a completely monotonic function of  $x$ , for  $x \in \mathbb{R}^+$ .

**Remark 2.1.** From the identity (3), we obtain the relations

$$\psi_{p,k}(x+k) - \psi_{p,k}(x) = \frac{1}{x} - \frac{1}{x+pk+k}, \quad (11)$$

$$\psi_{p,k}^{(m)}(x+k) - \psi_{p,k}^{(m)}(x) = \frac{(-1)^m m!}{x^{m+1}} - \frac{(-1)^m m!}{(x+pk+k)^{m+1}}, \quad m \in \mathbb{N}. \quad (12)$$

Also from (6), we obtain the relation

$$\psi_{p,k}(k) = \frac{1}{k} [\ln(pk) - H(p+1)]$$

where  $H(n)$  is the  $n$ th harmonic number.

The  $(p, k)$ -analogue of the classical Beta function is defined as

$$B_{p,k}(x, y) = \frac{\Gamma_{p,k}(x)\Gamma_{p,k}(y)}{\Gamma_{p,k}(x+y)}, \quad x > 0, y > 0.$$

**Lemma 2.1.** *The function  $\psi_{p,k}(x)$  satisfies the following limit properties.*

- (i)  $\psi_{p,k}(x) \rightarrow \psi_k(x)$  as  $p \rightarrow \infty$ ,
- (ii)  $\psi_{p,k}(x) \rightarrow \psi_p(x)$  as  $k \rightarrow 1$ ,
- (iii)  $\psi_{p,k}(x) \rightarrow \psi(x)$  as  $p \rightarrow \infty$  and  $k \rightarrow 1$ .

*Proof.* (i) By (6), we have

$$\begin{aligned} \lim_{p \rightarrow \infty} \psi_{p,k}(x) &= \lim_{p \rightarrow \infty} \left[ \frac{1}{k} \ln(pk) - \frac{1}{x} - \sum_{n=1}^p \frac{1}{(nk+x)} - \sum_{n=1}^p \frac{1}{nk} + \sum_{n=1}^p \frac{1}{nk} \right] \\ &= \lim_{p \rightarrow \infty} \left[ \frac{1}{k} \ln(pk) - \sum_{n=1}^p \frac{1}{nk} - \frac{1}{x} + \sum_{n=1}^p \frac{1}{nk} - \sum_{n=1}^p \frac{1}{(nk+x)} \right] \\ &= \lim_{p \rightarrow \infty} \left[ \frac{1}{k} \ln k + \frac{1}{k} \ln p - \frac{1}{k} \sum_{n=1}^p \frac{1}{n} - \frac{1}{x} + \sum_{n=1}^p \frac{x}{nk(nk+x)} \right] \\ &= \frac{1}{k} \ln k + \frac{1}{k} \lim_{p \rightarrow \infty} \left[ \ln p - \sum_{n=1}^p \frac{1}{n} \right] - \frac{1}{x} + \sum_{n=1}^{\infty} \frac{x}{nk(nk+x)} \\ &= \frac{1}{k} \ln k - \frac{\gamma}{k} - \frac{1}{x} + \sum_{n=1}^{\infty} \frac{x}{nk(nk+x)} \\ &= \psi_k(x). \end{aligned}$$

(ii) Also by (6), we have

$$\lim_{k \rightarrow 1} \psi_{p,k}(x) = \lim_{k \rightarrow 1} \left[ \frac{1}{k} \ln(pk) - \sum_{n=0}^p \frac{1}{(nk+x)} \right] = \ln p - \sum_{n=0}^p \frac{1}{(n+x)} = \psi_p(x).$$

(iii) This follows easily from (i) or (ii). That is,

$$\begin{aligned} \lim_{k \rightarrow 1} \left( \lim_{p \rightarrow \infty} \psi_{p,k}(x) \right) &= \lim_{k \rightarrow 1} \psi_k(x) = \psi(x), \quad \text{or} \\ \lim_{p \rightarrow \infty} \left( \lim_{k \rightarrow 1} \psi_{p,k}(x) \right) &= \lim_{p \rightarrow \infty} \psi_p(x) = \psi(x). \end{aligned}$$

□

**Lemma 2.2.** *Let  $\gamma_{p,k} = -\psi_{p,k}(1)$  be the  $(p, k)$ -analogue of the Euler-Mascheroni's constant. Then  $\gamma_{p,k} \rightarrow \gamma$  as  $p \rightarrow \infty$  and  $k \rightarrow 1$ .*

*Proof.* Proceed as follows

$$\lim_{k \rightarrow 1} \psi_{p,k}(1) = \lim_{k \rightarrow 1} \left[ \ln(pk) - \sum_{n=0}^p \frac{1}{(nk+1)} \right]$$

$$= \ln p - \sum_{n=0}^p \frac{1}{n+1}.$$

Then,

$$\begin{aligned} \lim_{p \rightarrow \infty} \left( \lim_{k \rightarrow 1} \gamma_{p,k} \right) &= \lim_{p \rightarrow \infty} \left( - \lim_{k \rightarrow 1} \psi_{p,k}(1) \right) \\ &= - \lim_{p \rightarrow \infty} \left[ \ln p - \sum_{n=0}^p \frac{1}{n+1} \right] \\ &= - \lim_{p \rightarrow \infty} \left[ \ln p - \sum_{n=1}^p \frac{1}{n} + \sum_{n=1}^p \frac{1}{n} - 1 - \sum_{n=1}^p \frac{1}{n+1} \right] \\ &= - \lim_{p \rightarrow \infty} \left[ \ln p - \sum_{n=1}^p \frac{1}{n} \right] + 1 - \sum_{n=1}^{\infty} \left[ \frac{1}{n} - \frac{1}{n+1} \right] \\ &= \gamma. \end{aligned}$$

□

**Definition 2.2.** A function  $f$  is said to be logarithmically convex if the following inequality holds for all  $x, y > 0$ .

$$\log f(\alpha x + \beta y) \leq \alpha \log f(x) + \beta \log f(y)$$

or equivalently

$$f(\alpha x + \beta y) \leq (f(x))^\alpha (f(y))^\beta$$

where  $\alpha, \beta > 0$  such that  $\alpha + \beta = 1$ .

**Theorem 2.1.** The function,  $\Gamma_{p,k}(x)$  is logarithmically convex.

*Proof.* We want to show that for  $x, y > 0$  and  $\alpha, \beta > 0$  such that  $\alpha + \beta = 1$ ,

$$\Gamma_{p,k}(\alpha x + \beta y) \leq (\Gamma_{p,k}(x))^\alpha (\Gamma_{p,k}(y))^\beta. \quad (13)$$

Recall that the Young's inequality is given by

$$x^\alpha y^\beta \leq \alpha x + \beta y \quad (14)$$

where  $x, y > 0$  and  $\alpha, \beta > 0$  such that  $\alpha + \beta = 1$ . By this, we obtain

$$\left(k + \frac{x}{t}\right)^\alpha \left(k + \frac{y}{t}\right)^\beta \leq k + \frac{\alpha x + \beta y}{t}. \quad (15)$$

Next, taking  $\prod_{t=1}^p$  on (15), we have

$$\prod_{t=1}^p \left(k + \frac{x}{t}\right)^\alpha \left(k + \frac{y}{t}\right)^\beta \leq \prod_{t=1}^p \left(k + \frac{\alpha x + \beta y}{t}\right)$$

which implies

$$\left( \frac{(x+k)(x+2k)\dots(x+pk)}{1 \times 2 \times \dots \times p} \right)^\alpha \left( \frac{(y+k)(y+2k)\dots(y+pk)}{1 \times 2 \times \dots \times p} \right)^\beta$$

$$\leq \frac{(\alpha x + \beta y + k)(\alpha x + \beta y + 2k) \dots (\alpha x + \beta y + pk)}{1 \times 2 \times \dots \times p}$$

which further implies

$$\begin{aligned} & \frac{p!}{(\alpha x + \beta y + k)(\alpha x + \beta y + 2k) \dots (\alpha x + \beta y + pk)} \\ & \leq \left( \frac{p!}{(x+k)(x+2k) \dots (x+pk)} \right)^\alpha \left( \frac{p!}{(y+k)(y+2k) \dots (y+pk)} \right)^\beta. \end{aligned} \quad (16)$$

Then, by multiplying (16) by the identities:

$$\begin{aligned} \frac{1}{\alpha x + \beta y} & \leq \frac{1}{x^\alpha y^\beta}, \\ (p+1) & = (p+1)^{\alpha+\beta}, \\ k^{p+1} & = (k^{p+1})^{\alpha+\beta}, \\ (pk)^{\frac{\alpha x + \beta y}{k} - 1} & = (pk)^{\frac{\alpha x}{k} - \alpha} (pk)^{\frac{\beta y}{k} - \beta} \end{aligned}$$

we obtain

$$\begin{aligned} & \frac{(p+1)! k^{p+1} (pk)^{\frac{\alpha x + \beta y}{k} - 1}}{(\alpha x + \beta y)(\alpha x + \beta y + k)(\alpha x + \beta y + 2k) \dots (\alpha x + \beta y + pk)} \\ & \leq \left( \frac{(p+1)! k^{p+1} (pk)^{\frac{x}{k} - 1}}{x(x+k)(x+2k) \dots (x+pk)} \right)^\alpha \left( \frac{(p+1)! k^{p+1} (pk)^{\frac{y}{k} - 1}}{y(y+k)(y+2k) \dots (y+pk)} \right)^\beta \end{aligned}$$

which is (13). That completes the proof.  $\square$

**Remark 2.2.** *Alternatively, a compact proof of Theorem 2.1 could have been as follows. By using the definition of  $\psi_{p,k}(x)$  and the fact that  $\psi'_{p,k}(x) > 0$ , it follows immediately that  $\Gamma_{p,k}(x)$  is logarithmically convex. Then, from the definition 2.2 for  $x, y > 0$ ,  $\alpha, \beta > 0$  such that  $\alpha + \beta = 1$ , we obtain the inequality (13).*

**Corollary 2.1.** *Let  $p \in \mathbb{N}$  and  $k > 0$ . Then the inequality*

$$\Gamma_{p,k}\left(\frac{x+y}{2}\right) \leq \sqrt{\Gamma_{p,k}(x)\Gamma_{p,k}(y)} \quad (17)$$

*holds for  $x, y > 0$ .*

*Proof.* This follows directly from Theorem 2.1 by letting  $\alpha = \beta = \frac{1}{2}$ .  $\square$

**Theorem 2.2.** *Let  $p \in \mathbb{N}$  and  $k > 0$ . Then the inequality*

$$\Gamma_{p,k}(nx) \leq (pk)^{\frac{x}{k}(n-1)} \Gamma_{p,k}(x) \quad (18)$$

*holds for  $n \in \mathbb{N}$  and  $x > 0$ .*

*Proof.* It follows easily from (2) that

$$\frac{\Gamma_{p,k}(nx)}{\Gamma_{p,k}(x)} = (pk)^{\frac{x}{k}(n-1)} \frac{x(x+k)(x+2k) \dots (x+pk)}{nx(nx+k)(nx+2k) \dots (nx+pk)}$$

which completes the proof.  $\square$

**Corollary 2.2.** *Let  $p \in \mathbb{N}$  and  $k > 0$ . Then the inequality*

$$\Gamma_{p,k}(x+y) \leq (pk)^{\frac{x+y}{2k}} \sqrt{\Gamma_{p,k}(x)\Gamma_{p,k}(y)} \quad (19)$$

holds for  $x, y > 0$ .

*Proof.* From (17), and by using (18) for  $n = 2$ , we obtain

$$\begin{aligned} \Gamma_{p,k}(x+y) &\leq \sqrt{\Gamma_{p,k}(2x)\Gamma_{p,k}(2y)} \\ &\leq (pk)^{\frac{x+y}{2k}} \sqrt{\Gamma_{p,k}(x)\Gamma_{p,k}(y)}. \end{aligned}$$

□

**Remark 2.3.** *Results similar to (13), (17), (18) and (19) for the  $(q, k)$ -analogue of the Gamma function can also be found in [3].*

**Lemma 2.3** ([9]). *Let  $f : (0, \infty) \rightarrow (0, \infty)$  be a differentiable, logarithmically convex function. Then the function*

$$g(x) = \frac{(f(x))^\alpha}{f(\alpha x)}, \quad \alpha \geq 1$$

is decreasing on its domain.

**Theorem 2.3.** *Let  $p \in \mathbb{N}$ ,  $k > 0$  and  $\alpha \geq 1$ . Then the inequality*

$$\frac{[\Gamma_{p,k}(y)]^\alpha}{\Gamma_{p,k}(\alpha y)} \leq \frac{[\Gamma_{p,k}(x)]^\alpha}{\Gamma_{p,k}(\alpha x)} \leq \frac{p}{p+1} k^{1-\alpha} \frac{1}{\Gamma_p(\alpha)} \quad (20)$$

is valid for  $k \leq x \leq y$ .

*Proof.* Recall from Theorem 2.1 that  $\Gamma_{p,k}(x)$  is logarithmically convex. Then by Lemma 2.3, the function  $H(x) = \frac{[\Gamma_{p,k}(x)]^\alpha}{\Gamma_{p,k}(\alpha x)}$  is decreasing. Hence for  $k \leq x \leq y$ , we have  $H(y) \leq H(x) \leq H(k)$  yielding the result (20). □

**Theorem 2.4.** *Let  $p \in \mathbb{N}$ ,  $k > 0$  and  $\alpha \geq 1$ . Then the inequality*

$$\frac{[\Gamma_{p,k}(1+k)]^\alpha}{\Gamma_{p,k}(\alpha+k)} \leq \frac{[\Gamma_{p,k}(x+k)]^\alpha}{\Gamma_{p,k}(\alpha x+k)} \leq 1 \quad (21)$$

is valid for  $x \in [0, 1]$ .

*Proof.* Define  $Q$  by  $Q(x) = \frac{[\Gamma_{p,k}(x+k)]^\alpha}{\Gamma_{p,k}(\alpha x+k)}$  for  $p \in \mathbb{N}$ ,  $k > 0$  and  $\alpha \geq 1$ . Let  $\lambda(x) = \ln Q(x)$ . Then

$$\begin{aligned} \lambda'(x) &= \alpha \frac{\Gamma'_{p,k}(x+k)}{\Gamma_{p,k}(x+k)} - \alpha \frac{\Gamma'_{p,k}(\alpha x+k)}{\Gamma_{p,k}(\alpha x+k)} \\ &= \alpha [\psi_{p,k}(x+k) - \psi_{p,k}(\alpha x+k)] \\ &\leq 0 \end{aligned}$$

since  $\psi_{p,k}(x)$  is increasing for  $x > 0$ . Hence  $Q(x)$  is decreasing on  $[0, \infty)$ . Then for  $x \in [0, 1]$ , we obtain  $Q(1) \leq Q(x) \leq Q(0)$  yielding the result (21). □



**Remark 2.4.** By letting  $p \rightarrow \infty$  as  $k \rightarrow 1$  in (21), we recover the results of [10] as a special case.

**Theorem 2.5.** Let  $p \in \mathbb{N}$ ,  $k > 0$ ,  $a > 1$ ,  $\frac{1}{a} + \frac{1}{b} = 1$  and  $m, n \in \mathbb{N}$  such that  $\frac{m}{a} + \frac{n}{b} \in \mathbb{N}$ . Then, the inequality

$$\left| \psi_{p,k}^{\left(\frac{m}{a} + \frac{n}{b}\right)} \left( \frac{x}{a} + \frac{y}{b} \right) \right| \leq \left| \psi_{p,k}^{(m)}(x) \right|^{\frac{1}{a}} \left| \psi_{p,k}^{(n)}(y) \right|^{\frac{1}{b}} \quad (22)$$

holds for  $x > 0$  and  $y > 0$ .

*Proof.* From the integral representation (9), we obtain

$$\begin{aligned} \left| \psi_{p,k}^{\left(\frac{m}{a} + \frac{n}{b}\right)} \left( \frac{x}{a} + \frac{y}{b} \right) \right| &= \int_0^\infty \left( \frac{1 - e^{-k(p+1)t}}{1 - e^{-kt}} \right) t^{\frac{m}{a} + \frac{n}{b}} e^{-\left(\frac{x}{a} + \frac{y}{b}\right)t} dt \\ &= \int_0^\infty \left( \frac{1 - e^{-k(p+1)t}}{1 - e^{-kt}} \right)^{\frac{1}{a} + \frac{1}{b}} t^{\frac{m}{a} + \frac{n}{b}} e^{-\left(\frac{x}{a} + \frac{y}{b}\right)t} dt \\ &= \int_0^\infty \left( \frac{1 - e^{-k(p+1)t}}{1 - e^{-kt}} \right)^{\frac{1}{a}} t^{\frac{m}{a}} e^{-\frac{xt}{a}} \cdot \left( \frac{1 - e^{-k(p+1)t}}{1 - e^{-kt}} \right)^{\frac{1}{b}} t^{\frac{n}{b}} e^{-\frac{yt}{b}} dt \\ &\leq \left[ \int_0^\infty \left( \frac{1 - e^{-k(p+1)t}}{1 - e^{-kt}} \right) t^m e^{-xt} dt \right]^{\frac{1}{a}} \\ &\quad \times \left[ \int_0^\infty \left( \frac{1 - e^{-k(p+1)t}}{1 - e^{-kt}} \right) t^n e^{-yt} dt \right]^{\frac{1}{b}} \\ &= \left| \psi_{p,k}^{(m)}(x) \right|^{\frac{1}{a}} \left| \psi_{p,k}^{(n)}(y) \right|^{\frac{1}{b}} \end{aligned}$$

which concludes the proof.  $\square$

Note: The absolute signs in (22) are not required if  $m$  and  $n$  are positive odd integers such that  $\frac{m+1}{a}, \frac{n+1}{b} \in \mathbb{N}$ .

**Corollary 2.3.** Let  $p \in \mathbb{N}$ ,  $k > 0$  and  $m \in \mathbb{N}$ . Then the inequality

$$\left| \psi_{p,k}^{(m)}(x) \right| \left| \psi_{p,k}^{(m+2)}(x) \right| - \left| \psi_{p,k}^{(m+1)}(x) \right|^2 \geq 0$$

holds for  $x > 0$ .

*Proof.* This follows from Theorem 2.5 by letting  $x = y$ ,  $a = b = 2$  and  $n = m + 2$ .  $\square$

**Remark 2.5.** By letting  $p \rightarrow \infty$  in Theorem 2.5, we obtain the  $k$ -analogue of (22). Also, by letting  $k \rightarrow 1$  in Theorem 2.5, we obtain the  $p$ -analogue of (22) as presented in Theorem 2.1 of [5].

**Remark 2.6.** By letting  $p \rightarrow \infty$  as  $k \rightarrow 1$  in Theorem 2.5, we obtain Theorem 2.5 of [11] as a special case.

**Remark 2.7.** Let  $x = y$  and  $a = b = 2$  in Theorem 2.5. Then, by letting  $p \rightarrow \infty$  as  $k \rightarrow 1$ , we obtain Theorem 2.1 of [4].

**Theorem 2.6.** *Let  $m, n, p \in \mathbb{N}$  and  $k > 0$ . Then, the inequality*

$$\left[ \left| \psi_{p,k}^{(m)}(x) \right| + \left| \psi_{p,k}^{(n)}(y) \right| \right]^{\frac{1}{u}} \leq \left| \psi_{p,k}^{(m)}(x) \right|^{\frac{1}{u}} + \left| \psi_{p,k}^{(n)}(y) \right|^{\frac{1}{u}} \quad (23)$$

holds for  $x > 0$  and  $y > 0$ , where  $u$  is a positive integer.

*Proof.* We employ the Minkowski's inequality for finite sums, and the fact that  $a^u + b^u \leq (a + b)^u$ , for  $a, b \geq 0$  and  $u$  a positive integer. From (8), we obtain

$$\begin{aligned} \left[ \left| \psi_{p,k}^{(m)}(x) \right| + \left| \psi_{p,k}^{(n)}(y) \right| \right]^{\frac{1}{u}} &= \left[ \sum_{i=0}^p \frac{m!}{(ik+x)^{m+1}} + \sum_{i=0}^p \frac{n!}{(ik+y)^{n+1}} \right]^{\frac{1}{u}} \\ &= \left[ \sum_{i=0}^p \left( \left( \frac{(m!)^{\frac{1}{u}}}{(ik+x)^{\frac{m+1}{u}}} \right)^u + \left( \frac{(n!)^{\frac{1}{u}}}{(ik+y)^{\frac{n+1}{u}}} \right)^u \right) \right]^{\frac{1}{u}} \\ &\leq \left[ \sum_{i=0}^p \left( \left( \frac{(m!)^{\frac{1}{u}}}{(ik+x)^{\frac{m+1}{u}}} \right) + \left( \frac{(n!)^{\frac{1}{u}}}{(ik+y)^{\frac{n+1}{u}}} \right) \right)^u \right]^{\frac{1}{u}} \\ &\leq \left[ \sum_{i=0}^p \left( \frac{(m!)^{\frac{1}{u}}}{(ik+x)^{\frac{m+1}{u}}} \right)^u \right]^{\frac{1}{u}} + \left[ \sum_{i=0}^p \left( \frac{(n!)^{\frac{1}{u}}}{(ik+y)^{\frac{n+1}{u}}} \right)^u \right]^{\frac{1}{u}} \\ &= \left| \psi_{p,k}^{(m)}(x) \right|^{\frac{1}{u}} + \left| \psi_{p,k}^{(n)}(y) \right|^{\frac{1}{u}} \end{aligned}$$

which concludes the proof.  $\square$

Note: The absolute signs in (23) are not required if  $m$  and  $n$  are positive odd integers.

**Remark 2.8.** *By letting  $p \rightarrow \infty$  in Theorems 2.5 and 2.6, we obtain the  $k$ -analogues of (22) and (23).*

**Remark 2.9.** *By letting  $k \rightarrow 1$  in Theorems 2.5 and 2.6, we obtain the  $p$ -analogues of (22) and (23) as presented in [5].*

**Remark 2.10.** *The  $q$ -analogues,  $(p, q)$ -analogues and  $(q, k)$ -analogues of the inequalities (22) and (23) can be found in [12], [6] and [7] respectively.*

**Theorem 2.7.** *Let  $p \in \mathbb{N}$ ,  $k > 0$  and  $m \in \mathbb{N}$ . Then, the inequalities*

$$\left( \exp \psi_{p,k}^{(m)}(x) \right)^2 \geq \exp \psi_{p,k}^{(m+1)}(x) \cdot \exp \psi_{p,k}^{(m-1)}(x), \quad \text{if } m \text{ is odd} \quad (24)$$

$$\left( \exp \psi_{p,k}^{(m)}(x) \right)^2 \leq \exp \psi_{p,k}^{(m+1)}(x) \cdot \exp \psi_{p,k}^{(m-1)}(x), \quad \text{if } m \text{ is even} \quad (25)$$

are satisfied for  $x > 0$ .

*Proof.* By relation (8), we obtain

$$\psi_{p,k}^{(m)}(x) - \frac{1}{2} \left[ \psi_{p,k}^{(m+1)}(x) + \psi_{p,k}^{(m-1)}(x) \right]$$

$$\begin{aligned}
&= \sum_{n=0}^p \frac{(-1)^{m+1} m!}{(nk+x)^{m+1}} - \frac{1}{2} \sum_{n=0}^p \frac{(-1)^{m+2} (m+1)!}{(nk+x)^{m+2}} - \frac{1}{2} \sum_{n=0}^p \frac{(-1)^m (m-1)!}{(nk+x)^m} \\
&= \frac{(-1)^m}{2} \left[ 2 \sum_{n=0}^p \frac{-m!}{(nk+x)^{m+1}} - \sum_{n=0}^p \frac{(m+1)!}{(nk+x)^{m+2}} - \sum_{n=0}^p \frac{(m-1)!}{(nk+x)^m} \right] \\
&= \frac{(-1)^{m+1}}{2} \left[ \sum_{n=0}^p \frac{2m!}{(nk+x)^{m+1}} + \sum_{n=0}^p \frac{(m+1)!}{(nk+x)^{m+2}} + \sum_{n=0}^p \frac{(m-1)!}{(nk+x)^m} \right] \\
&= \frac{(-1)^{m+1}}{2} \sum_{n=0}^p \frac{(m-1)!}{(nk+x)^m} \left[ \frac{2m}{nk+x} + \frac{(m+1)m}{(nk+x)^2} + 1 \right] \\
&= \frac{(-1)^{m+1}}{2} \sum_{n=0}^p \frac{(m-1)!}{(nk+x)^{m+2}} [(m+nk+x)^2 + m] \\
&= \begin{cases} \geq 0, & m \text{ odd} \\ \leq 0, & m \text{ even.} \end{cases}
\end{aligned}$$

That implies,

$$2\psi_{p,k}^{(m)}(x) \geq \psi_{p,k}^{(m+1)}(x) + \psi_{p,k}^{(m-1)}(x) \quad (26)$$

and

$$2\psi_{p,k}^{(m)}(x) \leq \psi_{p,k}^{(m+1)}(x) + \psi_{p,k}^{(m-1)}(x) \quad (27)$$

respectively for odd  $m$  and even  $m$ . Then, by exponentiating the inequalities (26) and (27), we obtain the desired results.  $\square$

**Remark 2.11.** By letting  $p \rightarrow \infty$  in Theorem 2.7, we obtain the  $k$ -analogues of (24) and (25).

**Remark 2.12.** By letting  $k \rightarrow 1$  in Theorem 2.7, we obtain the  $p$ -analogues of (24) and (25) as presented in Theorem 2.5 of [5] as a special case.

**Remark 2.13.** By letting  $p \rightarrow \infty$  as  $k \rightarrow 1$  in Theorem 2.7, we obtain Theorem 3.2 of [8] as a special case.

### 3. Conclusion

We have introduced a new two-parameter deformation of the classical Gamma function, called the  $(p, k)$ -analogue. In addition, we have established some identities and inequalities involving this new function. The established results provide generalizations of some known results in the literature.

### Acknowledgements

The authors are very grateful to the anonymous referees for their useful comments and suggestions, which helped in improving the quality of this paper.

## References

- [1] T. M. Apostol, *Introduction to Analytic Number Theory*, Springer-Verlag, 1976.
- [2] R. Díaz and E. Pariguan, *On hypergeometric functions and Pachhammer  $k$ -symbol*, *Divulgaciones Matemáticas*, 15(2)(2007), 179-192.
- [3] C. G. Kokologiannaki, *Some Properties of  $\Gamma_{q,k}(t)$  and Related Functions*, *International Journal of Contemporary Mathematical Sciences*, 11(1)(2016), 1-8.
- [4] A. Laforgia and P. Natalini, *Turan type inequalities for some special functions*, *J. Ineq. Pure Appl. Math.*, 7(1)(2006), Art. 22, 1-5.
- [5] F. Merovci, *Turan type inequalities for  $p$ -polygamma functions*, *Le Matematiche*, Fasc II (2013), 99-106.
- [6] F. Merovci, *Turan type inequalities for  $(p, q)$ -Gamma function*, *Scientia Magna*, 9(1) (2013), 25-30.
- [7] F. Merovci, *Turan type inequalities for some  $(q, k)$ -Special functions*, *Acta Universitatis Apulensis*, 34 (2013), 69-76.
- [8] C. Mortici, *Turan type inequalities for the Gamma and Polygamma functions*, *Acta Universitatis Apulensis*, 23 (2010), 117-121.
- [9] E. Neuman, *Inequalities involving a logarithmically convex function and their applications to special functions*, *J. Inequal. Pure Appl. Math.*, 7(1)(2006) Art. 16.
- [10] J. Sándor, *A note on certain inequalities for the gamma function*, *J. Ineq. Pure Appl. Math.*, 6(3)(2005), Art. 61.
- [11] W. T. Sulaiman, *Turan inequalities for the digamma and polygamma functions*, *South Asian Journal of Mathematics*, 1(2)(2011), 49-55.
- [12] W. T. Sulaiman, *Turan type inequalities for some special functions*, *The Australian Journal of Mathematical Analysis and Applications*, 9(1)(2012), 1-7.