BONE METASTASIS SECONDARY TO VULVAR CANCER

doi: 10.2478/rojost-2018-0049

C. Berceanu¹, E. Brătilă², M.M. Cîrstoiu³, S. Berceanu¹, A.V. Tetileanu⁴, N. L. Voicu¹, C. Mehedințu⁵

¹Department of Obstetrics and Gynecology, Clinical University Emergency Hospital, Craiova, Romania; University of Medicine and Pharmacy, Craiova, Romania

²Department of Obstetrics, Gynecology and Neonatology, "Panait Sârbu" Clinical Obstetrics and Gynecology Hospital, Bucharest, Romania; "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

³Department of Obstetrics, Gynecology and Neonatology, Clinical University Emergency Hospital, Bucharest, Romania; "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

⁴PhD Studies School, County Emergency Hospital, Târgu Jiu, Romania; Craiova University of Medicine and Pharmacy, Craiova, Romania

⁵Department of Obstetrics, Gynecology and Neonatology, "Nicolae Malaxa" Clinical Hospital, Bucharest, Romania; "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

Bone metastases from a vulvar cancer are extremely rare with only few reported cases in literature. Some cases were reported, in which patients developed pathologic fractures within eight months following radical surgery for advanced vulvar carcinoma. It is unanimously recognized that metastatic vulvar cancer is a rare condition. Therefore, information on metastatic determinations and corresponding prognosis or therapeutic approaches is very low. This paper is a review on some extremely rare conditions, such as bone metastases in vulvar cancer.

Keywords: rare condition, metastases, vulvar carcinoma