

MANAGEMENT OF FRAILTY AT INDIVIDUAL LEVEL: NARRATIVE REVIEW OF PHYSICAL ACTIVITY FROM THE EUROPEAN PERSPECTIVE OF JOINT ACTION ON FRAILTY - JA ADVANTAGE

MANAGEMENT KRHKOSTI NA RAVNI POSAMEZNIKA: PREGLED LITERATURE IZ EVROPSKE PERSPEKTIVE PROJEKTA SKUPNEGA UKREPANJA ADVANTAGE

Vojko STROJNIK¹, Branko GABROVEC^{2*}

¹University of Ljubljana, Faculty of Sport, Gortanova 22, 1000 Ljubljana, Slovenia

²National Institute of Public Health, Trubarjeva 2, 1000 Ljubljana, Slovenia

Received: Mr 12, 2018
Accepted: Feb 6, 2019

Review

ABSTRACT

Introduction: This paper aimed to review the effect of physical activity and exercise in frail older persons. As the process which leads to frailty and disability can be slowed down or even completely reversed, it can be appropriate for early interventions.

Keywords:

physical activity, exercise, frailty

Methodology: A literature search was conducted in the following databases: PubMed, Cochrane, Embase, Cinahl and UpToDate. The criterion in selecting the literature was that articles were published from 2002 to 2017. From 620,043 initial hits, 25 publications were selected.

Results: Physical activity and exercise in frail elderly are effective and relatively safe and may reverse frailty.

Conclusion: Different exercise interventions in frail elderly persons can increase strength and power, improve balance and reduce fall incidence resulting in greater quality of life. From this perspective, physical exercise interventions should become daily routine in frail elderly persons.

IZVLEČEK

Uvod: Namen tega prispevka je pregledati učinek fizične aktivnosti in telovadbe pri krhkih starostnikih. Ker se lahko proces, ki vodi h krhkosti in odvisnosti od drugih upočasni ali popolnoma zavre, je primeren za zgodnje intervencije.

Ključne besede:

telesna aktivnost, vadba, krhkost

Metode: Za to raziskavo je bil uporabljen pregled literature v naslednjih podatkovnih bazah: PubMed, Cochrane, Embase, Cinahl in UpToDate. Vključitveni kriterij je bil objava člankov v letih 2002 do 2017. Od 620.043 zadetkov je bilo izbranih 25 publikacij.

Rezultati: Fizična aktivnost in vadba starejših krhkih oseb je učinkovita in relativno varna ter lahko odpravlja krhkost.

Zaključki: Različne vadbene intervencije pri krhkih starejših osebah lahko povečajo telesno moč, izboljšajo ravnotežje in zmanjšujejo nevarnost padcev, kar izboljšuje kakovost življenja. Iz te perspektive naj intervencije telesne aktivnosti postanejo del dnevne rutine starejših oseb.

*Corresponding author: Tel. + 386 40 381 470; E-mail: branko.gabrovec@nijz.si

1 INTRODUCTION

Sedentary lifestyle is the sole most important factor in the development of chronic diseases (1). Sedentary behaviour has a strong negative impact on health outcomes among older adults, including higher risk of all-cause mortality, metabolic syndrome, high triglycerides, high blood pressure, overweight, abdominal obesity, mental disorders, etc. (2). Maintaining this lifestyle for years may accelerate some aspects of secondary ageing, such as speeding the reduction in bone mineral density, maximal oxygen consumption, and skeletal muscle strength and power (3), which are all used as determinants of frailty. In the elderly population, sedentary lifestyle is even more pronounced (4) which additionally increases the risk of becoming frail in older age.

The number of older persons is increasing, however, the number of the “oldest” old is rising even more rapidly. Older age is related to physical and cognitive decline, which contributes to an increased number of frail persons with increasing age (5). Frailty affects many domains including muscle strength, mobility, balance, endurance, co-ordination and, in general, there is a decrease in the level of physical and functional activity (6).

Reduced physical functioning is the most dominant sign of frailty (7, 8). The ageing associated loss of muscle mass seems to be one of the major causes for reduced physical abilities in older age and, consequently, disability and frailty (9). There are many influences contributing to this process, for example, motor neuron death, and hormonal and immunological changes as a normal part of ageing (10). On the other hand, there are additional behavioural influences such as poor nutrition and reduced physical activity that affect muscle mass reduction and are more pronounced with ageing. This is a very important observation since nutrition and physical activity can be changed by adopting an active and healthy lifestyle.

Frailty among older persons is a dynamic process (11). During the observed period (51 months), transitions to states of greater frailty were more common (rates up to 43.3%) than transitions to states of lesser frailty (rates up to 23.0%), and the probability of transitioning from being frail to non-frail was very low (rates between 0%-0.9%). With shorter observation intervals, the transition rates are even higher (12). Disability lasting only 1 or 2 months is strongly associated with the development of future disability and death (13). This strongly suggests that preventing frailty is a key factor for maintaining an active and healthy lifestyle in older age. However, even if frailty occurs, it can still be reversed. In the above-mentioned studies no specific treatments were performed to reverse frailty.

There is abundant evidence from prospective and clinical studies that physical activity not only delays but

also prevents or reverses frailty. For instance, a recent observational study (14) showed that physical activity might attenuate frailty. Mild physical activity was insufficient to significantly slow the progression of frailty, moderate physical activity reduced the progression of frailty in some age groups (particularly ages 65 and above) and vigorous activity significantly reduced the trajectory of frailty progression in all older adults.

A dose-response relationship has been shown in several studies. Higher cardiorespiratory fitness showed improved survival with higher function of metabolic equivalents (METs) across all age groups (15).

2 METHODS

Descriptive research methodology was used to review peer-reviewed medical literature. A narrative literature review was conducted as it enables the obtainment of data from various sources and ensures a holistic understanding of the research subject. The literature search was conducted using the following databases: PubMed, The Cochrane Library, Embase, UpToDate, Cumulative Index of Nursing and Allied Health Literature (CINAHL), by means of several combinations of selected search words in the English language and their synonyms were prepared and used with Boolean operators AND or: Frail Muscle strength *() OR Frailty Activity *() OR Elderly Exercise *() OR Older adult Functional ability *() OR Aged functional decline *() OR Older person Mobility *() OR Geriatric Disability *() OR Inactivity Vulnerable Elderly *() OR Physical activity Aged Function *() OR Training Aged *() OR Functional outcomes Geriatric *() OR Physical interventions Vulnerable *() OR Sports Older person *() OR Patterns of activity Older adult *() OR Leisure activity Elderly *(); searching in title, key words and in abstract.

The selection criterion for articles to be included in the review was that they were published during the last 15 years, i.e. between 2002 and 2017. Key words were selected from a proposal of key words that was prepared by the task leader and the working group focusing on Physical activity as part of the European Commission project “Joint Action on Frailty prevention - JA ADVANTAGE”, Work Package 6 - Management of Frailty at Individual Level. Final paper selection was also performed by the working group focusing on Physical activity.

The inclusion criteria were based on scientific facts, contextual relevance and full-text availability. Articles regarding current policies and guidelines on frailty prevention in older people that were published in peer-reviewed scientific journals. Information from editorials, letters, interviews, posters and articles with no access to full text were not included in the study. The total number of all search results was 620,043. After excluding

duplicates and taking inclusion criteria into account, a total of 25 articles/sources remained for analysis (Table 1). Initial selection using database search engines was performed by the Joint Action Advantage Work package 6 working group (selection of 119 sources). Further selection and analysis was performed by authors.

The process of the Literature Review is displayed in the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analysis) diagram, as shown in Figure 1.

With our approach, we found papers of different evidence levels of research. Between selected papers of different evidence levels of research, we included 4 qualitative and 21 quantitative researches. Data synthesis was conducted using the descriptive method.




Figure 1. Flowchart of search strategy and literature selection process - PRISMA diagram.

3 RESULTS

Table 1. Search table.

	Key word	No. of hits	Chosen hits	Repeated chosen hits	Final selection
PubMed	Frail Muscle strength	627	16	6	1
	Frailty Activity	1261	12	1	1
	Elderly Exercise	119309	11	11	3
	Older adult Functional ability	57636	11	11	2
	Aged functional decline	6779	3	2	1
	Older person Mobility	4694	2	2	1
	Geriatric Disability	4277	4	4	1
	Inactivity Vulnerable Elderly	52	1	1	1
	Physical activity Aged Function	92075	14	11	2
	Training Aged	245729	16	10	3
	Functional outcomes Geriatric	1724	2	1	1
	Physical interventions Vulnerable	741	4	1	1
	Sports Older person	8246	4	4	1
	Patterns of activity Older adult	11744	6	6	1
	Leisure activity Elderly	54789	6	6	1
Cochrane	Frail Muscle strength	171	5	5	1
	Frailty Activity	186	0	0	0
	Elderly Exercise	2321			1
	Older adult Functional ability	229	0	0	0
	Aged functional decline	1279	0	0	0
Embase	Older person Mobility	328	0	0	0
	Geriatric Disability	511	0	0	0
	Inactivity vulnerable Elderly	9	0	0	0
	Physical activity Aged Function	791	0	0	0
UpToDate	Training Aged	2680	1	1	1
	Functional outcomes Geriatric	1180	0	0	0
	Physical interventions Vulnerable	101	0	0	0
	Sports Older person	483	0	0	0
Cinahl	Patterns of activity Older adult	47	1	1	1
	Leisure activity Elderly	30	0	0	0
	Frail Muscle strength	14	0	0	0
Other sources		620043	119		25

4 DISCUSSION

Physical activity and exercise in frail elderly are effective and relatively safe and may reverse frailty. Most studies researched the effects of interventions on fall prevention and functional outcomes. Different exercise interventions in frail elderly persons can increase strength and power, have the potential to maintain or even slightly increase fat-free mass, and are effective in improving aerobic capacity and balance.

4.1 Strength and Power

The main reason behind strength and power decline is sarcopenia, loss of muscle mass with age due to motor neuron death, immunological factors, hormonal change, increased sedentary lifestyle and malnutrition (10). Supervised centre-based interventions seem to be more effective than home ones in improving strength in frail older persons (16-19) but not in all examples (20-23).

An important parameter of strength gain is exercise load, i.e. intensity, usually expressed in % of 1RM (one repetition maximum). Siegrist et al. (23) reported no strength gains after 16 weeks of a supervised exercise training program (1 hour/week) with strength and power training, challenging balance and gait training with increasing, but in general low, levels of difficulty. Tai-chi based and low-level strength exercise programs (24) produced a small but significant improvement in strength. With fitness machines and loads of 60% of 1RM substantial strength improvements were obtained (about 20% in isometric exercises and about 100% in lifting weights). Similar effects were seen in a study by Binder et al. (17) with exercise loads of 70-80% of 1RM. In the oldest old persons, 70% of 1RM load managed to improve leg press strength by 20% after 8 weeks of hypertrophy type strength training. Weight training with loads between 60-80% of 1RM increases muscle mass even in very old persons (25). The gains in strength and muscle mass may be similar in young and older women (26). These results are consistent with findings in healthy older persons where greater loads are related to greater increases in strength and power parameters (27) supporting a dose-response relationship. Exercise interventions were of different durations, ranging from 8 weeks up to 2 years. Already the shortest trial duration was enough to increase strength (28).

One interesting exercise principle is integrating exercise into everyday routines; however, high levels of improvisation and motivation are required. Results from the LiFE project (20) showed a limited effect of this approach on strength improvement. Its major limitation seemed to be low exercise loads, as high loads were hard to achieve in everyday routines.

Amino acid supplementation (AAS) may promote muscle growth but does not necessarily improve strength and power in healthy older adults (29). In older sarcopenic women, knee strength was improved in the exercise + AAS group but not in the exercise only group (30). This implies that AAS may augment muscle strength in this population. In another study (31) supplementation with milk fat globule membrane had no effect on muscle mass and strength gains. Similarly, supplementing vitamin D did not show any effect on strength (32). Supplementation with iron, folate, vitamin B6 and B12, calcium and Vitamin D (33) showed no additional improvement in strength, however, it increased overall physical activity and energy.

4.2 Endurance

Loss of aerobic capacity may be due to decreased muscle mass or lower cardiac output (35). Ehsani et al. (36) studied cardiovascular adaptation in older mild-to-moderate frail subjects after endurance exercise at 78% of peak heart rate. They found 14% increase in peak VO₂ after 9 months of intervention and that the main adaptation was increase

in heart rate and probably stroke volume. It is not possible to draw conclusions on the optimal regime to improve endurance and VO₂ max.

4.3 Balance and Risk of Falling

Exercise programs are effective in reducing falls and fall-related injuries in healthy older persons (37, 38). El-Khoury et al. (39) showed that exercise can reduce fall risk (including for serious falls) by 19% in older women already at risk of falls. Similar results (22% reduction) were seen in a study by Lord et al. (40) where, in the group with a previous fall, the incidence was reduced by 31%. Tai-chi and low-level exercise (24) reduced fall incident rate ratio by 58% overall. In a study by Siegrist et al. (23), the fall incidence in the exercise group was roughly half compared to the control group which received no treatment. This shows that fall prevention interventions in frail older persons are effective, however, they generally have a smaller effect on fall prevention compared to healthy older persons (37). Additionally, exercise reduced fear of falling (23, 41). Faber et al. (22) showed that frail persons, compared to pre-frail, benefit more from exercise intervention in terms of fall reduction. The opposite was found for improving balance and mobility.

There is abundant evidence that exercise intervention improves balance in frail elderly persons (19, 20, 22-24, 36, 39, 41-43), even in very old persons. A combination of strength and balance training further improves balance outcomes (17, 42, 43, 45). When strength and balance were complemented with gait and functional exercises (23, 39, 42) no additional effect on balance outcomes was observed.

Lifestyle integrated exercise was similarly effective as structured strength and balance training in promoting balance (20). This makes it possible to integrate balance exercises into daily activities.

4.4 Adverse Effects and Risks

Although exercise is generally safe for older people (46), this may not be valid for frail older persons. Some studies reported adverse effects, including falls during exercise sessions (36); groin strain and pelvic stress fracture (20); a wrist fracture, a twisted ankle, and two bruises (39); two reports of back pains (45); 9 (not identified) reported events related to intervention or testing (19); 23 reports of knee and back pain (32); 107 (5.0%) adverse events including IADL-ADL dependency, hospitalization or any fall (33); 11 reports of non-specified aches and pains (24). There was no pattern observed in adverse events related to interventions. However, there are some concerns that high-intensity exercise might pose a greater risk of injury than a program of lower intensity (32).

5 CONCLUSION

Physical activity and exercise in frail elderly are effective and relatively safe and may reverse frailty. Different exercise interventions in frail elderly persons can increase strength and power, have the potential to maintain or even slightly increase fat-free mass, are effective in improving aerobic capacity and balance, reduce fall incidence and improve quality of life. From this perspective, physical exercise interventions should become daily routine in frail elderly persons and supported by long-term care legislation.

The aim of this research was to present the results of a narrative literature review and data analysis focusing on physical activity in the context of managing frailty at an individual level. For the purposes of this research, a literature review method was used. The method proved to be appropriate and the aim was achieved.

CONFLICTS OF INTEREST

None.

FUNDING

This publication arises from the Joint Action ‘724099 / ADVANTAGE’, which has received funding from the European Union’s Health Programme (2014-2020). The content of this report represents the views of the author only and is his/her sole responsibility; it can’t be considered to reflect the views of the European Commission and/or the Consumers, Health, Agriculture and Food Executive Agency or any other body of the European Union. The European Commission and the Agency do not accept any responsibility for use that may be made of the information it contains.

ETHICAL APPROVAL

Not applicable.

REFERENCES

- Booth FW, Roberts CK, Laye MJ. Lack of exercise is a major cause of chronic diseases. *Compr Physiol*. 2012;2:1143-211. doi: 10.1002/cphy.c110025.
- Rezende LFM de, Rey-López JP, Matsudo VKR, Luiz O do C. Sedentary behavior and health outcomes among older adults: a systematic review. *BMC Public Health*. 2014;14:333. doi: 10.1186/1471-2458-14-333.
- Booth FW, Laye MJ, Roberts MD. Lifetime sedentary living accelerates some aspects of secondary aging. *J Appl Physiol*. 2011;111:1497-504. doi: 10.1152/jappphysiol.00420.2011.
- Davis MG, Fox KR, Hillsdon M, Sharp DJ, Coulson JC, Thompson JL. Objectively measured physical activity in a diverse sample of older urban UK adults. *Med Sci Sport Exerc*. 2011;43:647-54. doi: 10.1249/MSS.0b013e3181f36196.
- Kehler DS, Ferguson T, Stammers AN, Bohm C, Arora RC, Duhamel TA, et al. Prevalence of frailty in Canadians 18-79 years old in the Canadian Health Measures Survey. *BMC Geriatr*. 2017;17:28. doi: 10.1186/s12877-017-0423-6.
- Gobbens RJ, Luijckx KG, Wijnen-Sponselee MT, Schols JM. Toward a conceptual definition of frail community dwelling older people. *Nurs Outlook*. 2010;58:76-86. doi: 10.1016/j.outlook.2009.09.005.
- Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, et al. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci*. 2001;56:M146-56.
- Veninšek G, Gabrovec B. Management of frailty at individual level - clinical management: systematic literature review. *Zdr Varst*. 2018;57:106-15. doi: 10.2478/sjph-2018-0014.
- Roubenoff R. Sarcopenia: a major modifiable cause of frailty in the elderly. *J Nutr Health Aging*. 2000;4:140-2.
- Narici M V, Maganaris CN. Adaptability of elderly human muscles and tendons to increased loading. *J Anat*. 2006;208:433-43. doi: 10.1111/j.1469-7580.2006.00548.x.
- Gill TM, Gahbauer EA, Allore HG, Han L. transitions between frailty states among community-living older persons. *Arch Intern Med*. 2006;166:418. doi: 10.1001/archinte.166.4.418.
- Hardy SE, Gill TM. Recovery from disability among community-dwelling older persons. *JAMA*. 2004;291:1596. doi: 10.1001/jama.291.13.1596.
- Gill TM, Kurland BF. Prognostic effect of prior disability episodes among nondisabled community-living older persons. *Am J Epidemiol*. 2003;158:1090-6. doi: 10.1093/aje/kwg237.
- Rogers NT, Marshall A, Roberts CH, Demakakos P, Steptoe A, Scholes S. Physical activity and trajectories of frailty among older adults: evidence from the English longitudinal study of ageing. *PLoS One*. 2017;12:e0170878. doi: 10.1371/journal.pone.0170878.
- Feldman DI, Al-Mallah MH, Keteyian SJ, Brawner CA, Feldman T, Blumenthal RS, et al. No evidence of an upper threshold for mortality benefit at high levels of cardiorespiratory fitness. *J Am Coll Cardiol*. 2015;65(6):629-30. doi: 10.1016/j.jacc.2014.11.030.
- Binder EF, Schechtman KB, Ehsani AA, Steger-May K, Brown M, Sinacore DR, et al. Effects of exercise training on frailty in community-dwelling older adults: results of a randomized, controlled trial. *J Am Geriatr Soc*. 2002;50:1921-8. doi: 10.1046/j.1532-5415.2002.50601.x.
- Binder EF, Yarasheski KE, Steger-May K, Sinacore DR, Brown M, Schechtman KB, et al. Effects of progressive resistance training on body composition in frail older adults: results of a randomized, controlled trial. *J Gerontol A Biol Sci Med Sci*. 2005;60:1425-31.
- LIFE Study Investigators, Pahor M, Blair SN, Espeland M, Fielding R, Gill TM, et al. Effects of a physical activity intervention on measures of physical performance: results of the lifestyle interventions and independence for Elders Pilot (LIFE-P) study. *J Gerontol A Biol Sci Med Sci*. 2006;61:1157-65.
- King MB, Whipple RH, Gruman CA, Judge JO, Schmidt JA, Wolfson LI. The Performance Enhancement Project: improving physical performance in older persons. *Arch Phys Med Rehabil*. 2002;83:1060-9.
- Clemson L, Fiatarone Singh MA, Bundy A, Cumming RG, Manolagas K, O’Loughlin P, et al. Integration of balance and strength training into daily life activity to reduce rate of falls in older people (the LiFE study): randomised parallel trial. *BMJ*. 2012;345:e4547. doi: 10.1136/bmj.e4547.
- Lustosa LP, Silva JP, Coelho FM, Pereira DS, Parentoni AN, Pereira LSM. Impact of resistance exercise program on functional capacity and muscular strength of knee extensor in pre-frail community-dwelling older women: a randomized crossover trial. *Rev Bras Fisioter*. 2011;15:318-24.

22. Faber MJ, Bosscher RJ, Chin A Paw MJ, van Wieringen PC. Effects of exercise programs on falls and mobility in frail and pre-frail older adults: a multicenter randomized controlled trial. *Arch Phys Med Rehabil.* 2006;87:885-96. doi: 10.1016/j.apmr.2006.04.005.
23. Siegrist M, Freiburger E, Geilhof B, Salb J, Hentschke C, Landendoerfer P, et al. Fall prevention in a primary care setting. *Dtsch Arztebl Int.* 2016;113:365-72. doi: 10.3238/arztebl.2016.0365.
24. Taylor D, Hale L, Schluter P, Waters DL, Binns EE, McCracken H, et al. Effectiveness of tai chi as a community-based falls prevention intervention: a randomized controlled trial. *J Am Geriatr Soc.* 2012;60:841-8. doi: 10.1111/j.1532-5415.2012.03928.x.
25. Harridge SD, Kryger A, Stensgaard A. Knee extensor strength, activation, and size in very elderly people following strength training. *Muscle Nerve.* 1999;22:831-9.
26. Cannon J, Kay D, Tarpennin KM, Marino FE. Comparative effects of resistance training on peak isometric torque, muscle hypertrophy, voluntary activation and surface EMG between young and elderly women. *Clin Physiol Funct Imaging.* 2007;27:91-100. doi: 10.1111/j.1475-097X.2007.00719.x.
27. Steib S, Schoene D, Pfeifer K. Dose-response relationship of resistance training in older adults. *Med Sci Sport Exerc.* 2010;42:902-14. doi: 10.1249/MSS.0b013e3181c34465.
28. Serra-Rexach JA, Bustamante-Ara N, Hierro Villarán M, González Gil P, Sanz Ibáñez MJ, Blanco Sanz N, et al. Short-term, light- to moderate-intensity exercise training improves leg muscle strength in the oldest old: a randomized controlled trial. *J Am Geriatr Soc.* 2011;59:594-602. doi: 10.1111/j.1532-5415.2011.03356.x.
29. Finger D, Goltz FR, Umpierre D, Meyer E, Rosa LHT, Schneider CD. Effects of protein supplementation in older adults undergoing resistance training: a systematic review and meta-analysis. *Sport Med.* 2015;45:245-55. doi: 10.1007/s40279-014-0269-4.
30. Kim HK, Suzuki T, Saito K, Yoshida H, Kobayashi H, Kato H, et al. Effects of exercise and amino acid supplementation on body composition and physical function in community-dwelling elderly Japanese sarcopenic women: a randomized controlled trial. *J Am Geriatr Soc.* 2012;60:16-23. doi: 10.1111/j.1532-5415.2011.03776.x.
31. Kim I-Y, Schutzler S, Schrader A, Spencer H, Kortebein P, Deutz NEP, et al. Quantity of dietary protein intake, but not pattern of intake, affects net protein balance primarily through differences in protein synthesis in older adults. *Am J Physiol Endocrinol Metab.* 2015;308:E21-8. doi: 10.1152/ajpendo.00382.2014.
32. Latham NK, Anderson CS, Lee A, Bennett DA, Moseley A, Cameron ID, et al. A randomized, controlled trial of quadriceps resistance exercise and vitamin D in frail older people: the Frailty Interventions Trial in Elderly Subjects (FITNESS). *J Am Geriatr Soc.* 2003;51:291-9.
33. Ng TP, Feng L, Nyunt MSZ, Feng L, Niti M, Tan BY, et al. Nutritional, physical, cognitive, and combination interventions and frailty reversal among older adults: a randomized controlled trial. *Am J Med.* 2015;128:1225-36.e1. doi: 10.1016/j.amjmed.2015.06.017.
34. Fleg JL, Lakatta EG. Role of muscle loss in the age-associated reduction in VO2 max. *J Appl Physiol.* 1988;65:1147-51. doi: 10.1152/jappl.1988.65.3.1147.
35. Ogawa T, Spina RJ, Martin WH, Kohrt WM, Schechtman KB, Holloszy JO, et al. Effects of aging, sex, and physical training on cardiovascular responses to exercise. *Circulation.* 1992;86:494-503.
36. Ehsani AA, Spina RJ, Peterson LR, Rinder MR, Glover KL, Villareal DT, et al. Attenuation of cardiovascular adaptations to exercise in frail octogenarians. *J Appl Physiol.* 2003;95:1781-8. doi: 10.1152/jappphysiol.00194.2003.
37. El-Khoury F, Cassou B, Charles M-A, Dargent-Molina P. The effect of fall prevention exercise programmes on fall induced injuries in community dwelling older adults: systematic review and meta-analysis of randomised controlled trials. *BMJ.* 2013;347:f6234. doi: 10.1136/bmj.f6234.
38. Gillespie LD, Robertson MC, Gillespie WJ, Sherrington C, Gates S, Clemson LM, et al. Interventions for preventing falls in older people living in the community. *Cochrane Database Syst Rev.* 2012;(9):CD007146. doi: 10.1002/14651858.CD007146.pub3.
39. El-Khoury F, Cassou B, Latouche A, Aegerter P, Charles M-A, Dargent-Molina P. Effectiveness of two year balance training programme on prevention of fall induced injuries in at risk women aged 75-85 living in community: Ossébo randomised controlled trial. *BMJ.* 2015;351:h3830. doi: 10.1136/bmj.h3830.
40. Lord SR, Castell S, Corcoran J, Dayhew J, Matters B, Shan A, et al. The effect of group exercise on physical functioning and falls in frail older people living in retirement villages: a randomized, controlled trial. *J Am Geriatr Soc.* 2003;51:1685-92.
41. Giné-Garriga M, Guerra M, Unnithan VB. The effect of functional circuit training on self-reported fear of falling and health status in a group of physically frail older individuals: a randomized controlled trial. *Aging Clin Exp Res.* 2013;25:329-36. doi: 10.1007/s40520-013-0048-3.
42. Freiburger E, Häberle L, Spirduso WW, Rixt Zijlstra GA. Long-term effects of three multicomponent exercise interventions on physical performance and fall-related psychological outcomes in community-dwelling older adults: a randomized controlled trial. *J Am Geriatr Soc.* 2012;60:437-46. doi: 10.1111/j.1532-5415.2011.03859.x.
43. Giné-Garriga M, Guerra M, Pagès E, Manini TM, Jiménez R, Unnithan VB. The effect of functional circuit training on physical frailty in frail older adults: a randomized controlled trial. *J Aging Phys Act.* 2010;18:401-24.
44. Cadore EL, Casas-Herrero A, Zambom-Ferraresi F, Idoate F, Millor N, Gómez M, et al. Multicomponent exercises including muscle power training enhance muscle mass, power output, and functional outcomes in institutionalized frail nonagenarians. *Age (Omaha).* 2014;36:773-85. doi: 10.1007/s11357-013-9586-z.
45. Fairhall N, Sherrington C, Lord SR, Kurrle SE, Langron C, Lockwood K, et al. Effect of a multifactorial, interdisciplinary intervention on risk factors for falls and fall rate in frail older people: a randomised controlled trial. *Age Ageing.* 2014;43:616-22. doi:10.1093/ageing/afz204.
46. Global recommendations on physical activity for health. Copenhagen: World Health Organisation, 2015.